

# A raising operator formula for Macdonald polynomials via LLT polynomials in the elliptic Hall algebra

George H. Seelinger

joint work with J. Blasiak, M. Haiman, J. Morse, and A. Pun

[ghseeli@umich.edu](mailto:ghseeli@umich.edu)

Loyola University Chicago TACO Seminar

Based on [arXiv:2112.07063](https://arxiv.org/abs/2112.07063) and [arXiv:2307.06517](https://arxiv.org/abs/2307.06517)

October 4, 2023

Glad to be back



Graduation May 2015


- ① **Background on symmetric functions and Macdonald polynomials**
- ② A new formula for Macdonald polynomials
- ③ LLT polynomials in the elliptic Hall algebra

# Symmetric Group

- Permutations  $\sigma: \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$ :

# Symmetric Group


- Permutations  $\sigma: \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$ :

$$\begin{pmatrix} 1 & 2 & 3 & 4 \\ 2 & 3 & 1 & 4 \end{pmatrix} =$$


The diagram illustrates the permutation  $\sigma$  as a mapping between two rows of four nodes. The top row represents the domain  $\{1, 2, 3, 4\}$  and the bottom row represents the codomain  $\{1, 2, 3, 4\}$ . Lines connect the nodes as follows: a line from the first node to the second, from the second to the third, from the third to the first, and a vertical line from the fourth to the fourth. This represents the permutation  $(1\ 2\ 3)$ .

# Symmetric Group

- Permutations  $\sigma: \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$ :

$$\begin{pmatrix} 1 & 2 & 3 & 4 \\ 2 & 3 & 1 & 4 \end{pmatrix} =$$


- For  $f \in \mathbb{Q}[x_1, \dots, x_n]$  multivariate polynomial,  $\sigma \in S_n$  acts as  $\sigma.f(x_1, x_2, \dots, x_n) = f(x_{\sigma(1)}, x_{\sigma(2)}, \dots, x_{\sigma(n)})$

$$\begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix} (5x_1^2 + 5x_2^2 + 8x_3^2) = 8x_1^2 + 5x_2^2 + 5x_3^2$$

# Symmetric Polynomials

- Polynomials  $f \in \mathbb{Q}[x_1, \dots, x_n]$  satisfying  $\sigma.f = f$  for all  $\sigma \in S_n$ .

# Symmetric Polynomials

- Polynomials  $f \in \mathbb{Q}[x_1, \dots, x_n]$  satisfying  $\sigma.f = f$  for all  $\sigma \in S_n$ .

## Generators

$$e_r = \sum_{i_1 < i_2 < \dots < i_r} x_{i_1} x_{i_2} \cdots x_{i_r} \text{ or } h_r = \sum_{i_1 \leq i_2 \leq \dots \leq i_r} x_{i_1} x_{i_2} \cdots x_{i_r}$$



# Symmetric Polynomials

- Polynomials  $f \in \mathbb{Q}[x_1, \dots, x_n]$  satisfying  $\sigma.f = f$  for all  $\sigma \in S_n$ .

## Generators

$$e_r = \sum_{i_1 < i_2 < \dots < i_r} x_{i_1} x_{i_2} \cdots x_{i_r} \text{ or } h_r = \sum_{i_1 \leq i_2 \leq \dots \leq i_r} x_{i_1} x_{i_2} \cdots x_{i_r}$$

- E.g. for  $n = 3$ ,

$$e_1 = x_1 + x_2 + x_3 = h_1$$

$$e_2 = x_1 x_2 + x_1 x_3 + x_2 x_3 \quad h_2 = x_1^2 + x_1 x_2 + x_1 x_3 + x_2^2 + x_2 x_3 + x_3^2$$

$$e_3 = x_1 x_2 x_3 \quad h_3 = x_1^3 + x_1^2 x_2 + x_1^2 x_3 + x_1 x_2^2 + \dots$$

# Symmetric Polynomials

- Polynomials  $f \in \mathbb{Q}[x_1, \dots, x_n]$  satisfying  $\sigma.f = f$  for all  $\sigma \in S_n$ .

## Generators

$$e_r = \sum_{i_1 < i_2 < \dots < i_r} x_{i_1} x_{i_2} \cdots x_{i_r} \text{ or } h_r = \sum_{i_1 \leq i_2 \leq \dots \leq i_r} x_{i_1} x_{i_2} \cdots x_{i_r}$$

- E.g. for  $n = 3$ ,

$$e_1 = x_1 + x_2 + x_3 = h_1$$

$$e_2 = x_1 x_2 + x_1 x_3 + x_2 x_3 \quad h_2 = x_1^2 + x_1 x_2 + x_1 x_3 + x_2^2 + x_2 x_3 + x_3^2$$

$$e_3 = x_1 x_2 x_3 \quad h_3 = x_1^3 + x_1^2 x_2 + x_1^2 x_3 + x_1 x_2^2 + \dots$$

- Let  $\Lambda = \mathbb{Q}[e_1, e_2, \dots] = \mathbb{Q}[h_1, h_2, \dots]$ . Call these “symmetric functions.”

# Symmetric Polynomials

- Polynomials  $f \in \mathbb{Q}[x_1, \dots, x_n]$  satisfying  $\sigma.f = f$  for all  $\sigma \in S_n$ .

## Generators

$$e_r = \sum_{i_1 < i_2 < \dots < i_r} x_{i_1} x_{i_2} \cdots x_{i_r} \text{ or } h_r = \sum_{i_1 \leq i_2 \leq \dots \leq i_r} x_{i_1} x_{i_2} \cdots x_{i_r}$$

- E.g. for  $n = 3$ ,

$$e_1 = x_1 + x_2 + x_3 = h_1$$

$$e_2 = x_1 x_2 + x_1 x_3 + x_2 x_3 \quad h_2 = x_1^2 + x_1 x_2 + x_1 x_3 + x_2^2 + x_2 x_3 + x_3^2$$

$$e_3 = x_1 x_2 x_3 \quad h_3 = x_1^3 + x_1^2 x_2 + x_1^2 x_3 + x_1 x_2^2 + \dots$$

- Let  $\Lambda = \mathbb{Q}[e_1, e_2, \dots] = \mathbb{Q}[h_1, h_2, \dots]$ . Call these “symmetric functions.”
- $\Lambda$  is a  $\mathbb{Q}$ -algebra.

# Bases for symmetric functions

Dimension of degree  $d$  symmetric functions?

# Bases for symmetric functions

Dimension of degree  $d$  symmetric functions? Number of partitions of  $d$ .

# Bases for symmetric functions

Dimension of degree  $d$  symmetric functions? Number of partitions of  $d$ .

## Definition

$n \in \mathbb{Z}_{>0}$ , a *partition of  $n$*  is  $\lambda = (\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_\ell > 0)$  such that  $\lambda_1 + \lambda_2 + \cdots + \lambda_\ell = n$ .

# Bases for symmetric functions

Dimension of degree  $d$  symmetric functions? Number of partitions of  $d$ .

## Definition

$n \in \mathbb{Z}_{>0}$ , a *partition of  $n$*  is  $\lambda = (\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_\ell > 0)$  such that  $\lambda_1 + \lambda_2 + \cdots + \lambda_\ell = n$ .

$$5 \rightarrow \begin{array}{|c|c|c|c|c|}\hline\hline\hline\hline\hline\hline\end{array}$$

$$4 + 1 \rightarrow \begin{array}{|c|c|c|c|}\hline\hline\hline\hline\hline\end{array}$$

$$3 + 2 \rightarrow \begin{array}{|c|c|c|}\hline\hline\hline\hline\hline\end{array}$$

$$3 + 1 + 1 \rightarrow \begin{array}{|c|c|c|}\hline\hline\hline\hline\hline\end{array}$$

$$2 + 2 + 1 \rightarrow \begin{array}{|c|c|c|}\hline\hline\hline\hline\hline\end{array}$$

$$2 + 1 + 1 + 1 \rightarrow \begin{array}{|c|c|c|}\hline\hline\hline\hline\hline\end{array}$$

$$1 + 1 + 1 + 1 + 1 \rightarrow \begin{array}{|c|}\hline\hline\hline\hline\hline\end{array}$$

# Bases for symmetric functions


Dimension of degree  $d$  symmetric functions? Number of partitions of  $d$ .


## Definition

$n \in \mathbb{Z}_{>0}$ , a *partition of  $n$*  is  $\lambda = (\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_\ell > 0)$  such that  $\lambda_1 + \lambda_2 + \cdots + \lambda_\ell = n$ .


5 → 

--	--	--	--	--


$2 + 2 + 1 \rightarrow$  

$4 + 1 \rightarrow$  

$$2 + 1 + 1 + 1 \rightarrow \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}$$

$3 + 2 \rightarrow$  

$$1 + 1 + 1 + 1 + 1 \rightarrow$$

$3 + 1 + 1 \rightarrow$  

$\implies$  any basis of degree  $d$  symmetric functions can be indexed by partitions of  $d$ .



# Semistandard Young Tableaux

## Definition

Filling of partition diagram of  $\lambda$  with numbers such that

# Semistandard Young Tableaux

## Definition

Filling of partition diagram of  $\lambda$  with numbers such that

- 1 strictly increasing up columns

# Semistandard Young Tableaux

## Definition

Filling of partition diagram of  $\lambda$  with numbers such that

- 1 strictly increasing up columns
- 2 weakly increasing along rows

# Semistandard Young Tableaux

## Definition

Filling of partition diagram of  $\lambda$  with numbers such that

- ① strictly increasing up columns
- ② weakly increasing along rows

Collection is called  $\text{SSYT}(\lambda)$ .

# Semistandard Young Tableaux

## Definition

Filling of partition diagram of  $\lambda$  with numbers such that

- ① strictly increasing up columns
- ② weakly increasing along rows

Collection is called  $\text{SSYT}(\lambda)$ .

For  $\lambda = (2, 1)$ ,

2		3		3		2		3		3		2		3	
1	1	1	1	2	2	1	2	1	3	2	3	1	3	1	2

# Polynomials from tableaux

Associate a polynomial to  $\text{SSYT}(\lambda)$ .

# Polynomials from tableaux

Associate a polynomial to  $\text{SSYT}(\lambda)$ .

2		3		3		2		3		3		2		3	
1	1	1	1	2	2	1	2	1	3	2	3	1	3	1	2

# Polynomials from tableaux

Associate a polynomial to  $\text{SSYT}(\lambda)$ .

2		3		3		2		3		3		2		3	
1	1	1	1	2	2	1	2	1	3	2	3	1	3	1	2

$$s_{(2,1)}(x_1, x_2, x_3) = x_1^2 x_2 + x_1^2 x_3 + x_2^2 x_3 + x_1 x_2^2 + x_1 x_3^2 + x_2 x_3^2 + 2x_1 x_2 x_3$$



# Polynomials from tableaux

Associate a polynomial to  $\text{SSYT}(\lambda)$ .

2		3		3		2		3		3		2		3	
1	1	1	1	2	2	1	2	1	3	2	3	1	3	1	2

$$s_{(2,1)}(x_1, x_2, x_3) = x_1^2 x_2 + x_1^2 x_3 + x_2^2 x_3 + x_1 x_2^2 + x_1 x_3^2 + x_2 x_3^2 + 2x_1 x_2 x_3$$

## Definition

For  $\lambda$  a partition, set

$$s_\lambda = \sum_{T \in \text{SSYT}(\lambda)} \mathbf{x}^T \text{ for } \mathbf{x}^T = \prod_{i \in T} x_i$$

# Polynomials from tableaux

Associate a polynomial to  $\text{SSYT}(\lambda)$ .

2		3		3		2		3		3		2		3	
1	1	1	1	2	2	1	2	1	3	2	3	1	3	1	2

$$s_{(2,1)}(x_1, x_2, x_3) = x_1^2 x_2 + x_1^2 x_3 + x_2^2 x_3 + x_1 x_2^2 + x_1 x_3^2 + x_2 x_3^2 + 2x_1 x_2 x_3$$

## Definition

For  $\lambda$  a partition, set

$$s_\lambda = \sum_{T \in \text{SSYT}(\lambda)} \mathbf{x}^T \text{ for } \mathbf{x}^T = \prod_{i \in T} x_i$$

- $s_\lambda$  is a symmetric function.

# Polynomials from tableaux

Associate a polynomial to  $\text{SSYT}(\lambda)$ .

2		3		3		2		3		3		2		3	
1	1	1	1	2	2	1	2	1	3	2	3	1	3	1	2

$$s_{(2,1)}(x_1, x_2, x_3) = x_1^2 x_2 + x_1^2 x_3 + x_2^2 x_3 + x_1 x_2^2 + x_1 x_3^2 + x_2 x_3^2 + 2x_1 x_2 x_3$$

## Definition

For  $\lambda$  a partition, set

$$s_\lambda = \sum_{T \in \text{SSYT}(\lambda)} \mathbf{x}^T \text{ for } \mathbf{x}^T = \prod_{i \in T} x_i$$

- $s_\lambda$  is a symmetric function.
- $\{s_\lambda\}_\lambda$  forms a basis for  $\Lambda$ .

# Symmetric functions and Schur functions

- Convention:  $h_0 = 1$  and  $h_d = 0$  for  $d < 0$ .
- For any  $\gamma = (\gamma_1, \dots, \gamma_n) \in \mathbb{Z}^n$ , set

$$s_\gamma = \det(h_{\gamma_i + j - i})_{1 \leq i, j \leq n}$$

# Symmetric functions and Schur functions

- Convention:  $h_0 = 1$  and  $h_d = 0$  for  $d < 0$ .
- For any  $\gamma = (\gamma_1, \dots, \gamma_n) \in \mathbb{Z}^n$ , set

$$s_\gamma = \det(h_{\gamma_i + j - i})_{1 \leq i, j \leq n}$$

Then,  $s_\gamma = \pm s_\lambda$  or 0 for some partition  $\lambda$ .

# Symmetric functions and Schur functions

- Convention:  $h_0 = 1$  and  $h_d = 0$  for  $d < 0$ .
- For any  $\gamma = (\gamma_1, \dots, \gamma_n) \in \mathbb{Z}^n$ , set

$$s_\gamma = \det(h_{\gamma_i + j - i})_{1 \leq i, j \leq n}$$

Then,  $s_\gamma = \pm s_\lambda$  or 0 for some partition  $\lambda$ .

Precisely, for  $\rho = (n-1, n-2, \dots, 1, 0)$ ,

$$s_\gamma = \begin{cases} \operatorname{sgn}(\gamma + \rho) s_{\operatorname{sort}(\gamma + \rho) - \rho} & \text{if } \gamma + \rho \text{ has distinct nonnegative parts,} \\ 0 & \text{otherwise,} \end{cases}$$

- $\operatorname{sort}(\beta) =$  weakly decreasing sequence obtained by sorting  $\beta$ ,
- $\operatorname{sgn}(\beta) =$  sign of the shortest permutation taking  $\beta$  to  $\operatorname{sort}(\beta)$ .

Example:  $s_{201} = 0$ ,  $s_{2-11} = -s_{200}$ .

# Representation theory and Schur functions

Frobenius characteristc,  $\text{Frob}: \text{Rep}(S_n) \rightarrow \Lambda$ .

# Representation theory and Schur functions

Frobenius characteristc,  $\text{Frob}: \text{Rep}(S_n) \rightarrow \Lambda$ .

- Irreducible representations of  $S_n$  are labeled by partitions of  $n$ .



# Representation theory and Schur functions

Frobenius characteristc,  $\text{Frob}: \text{Rep}(S_n) \rightarrow \Lambda$ .

- Irreducible representations of  $S_n$  are labeled by partitions of  $n$ .
- Irreducible  $S_n$ -representation  $V_\lambda$  has  $\text{Frob}(V_\lambda) = s_\lambda$

# Representation theory and Schur functions

Frobenius characteristc,  $\text{Frob}: \text{Rep}(S_n) \rightarrow \Lambda$ .

- Irreducible representations of  $S_n$  are labeled by partitions of  $n$ .
- Irreducible  $S_n$ -representation  $V_\lambda$  has  $\text{Frob}(V_\lambda) = s_\lambda$
- $U \cong V \oplus W \implies \text{Frob}(U) = \text{Frob}(V) + \text{Frob}(W)$

# Representation theory and Schur functions

Frobenius characterstic,  $\text{Frob}: \text{Rep}(S_n) \rightarrow \Lambda$ .

- Irreducible representations of  $S_n$  are labeled by partitions of  $n$ .
- Irreducible  $S_n$ -representation  $V_\lambda$  has  $\text{Frob}(V_\lambda) = s_\lambda$
- $U \cong V \oplus W \implies \text{Frob}(U) = \text{Frob}(V) + \text{Frob}(W)$
- $\text{Ind}_{S_m \times S_n}^{S_{m+n}}(V \times W) \mapsto \text{Frob}(V) \cdot \text{Frob}(W)$

# Representation theory and Schur functions

Frobenius characterstic,  $\text{Frob}: \text{Rep}(S_n) \rightarrow \Lambda$ .

- Irreducible representations of  $S_n$  are labeled by partitions of  $n$ .
- Irreducible  $S_n$ -representation  $V_\lambda$  has  $\text{Frob}(V_\lambda) = s_\lambda$
- $U \cong V \oplus W \implies \text{Frob}(U) = \text{Frob}(V) + \text{Frob}(W)$
- $\text{Ind}_{S_m \times S_n}^{S_{m+n}}(V \times W) \mapsto \text{Frob}(V) \cdot \text{Frob}(W)$
- Upshot:  $S_n$ -representations go to symmetric functions in structure preserving way.

# Representation theory and Schur functions

Frobenius character map,  $\text{Frob}: \text{Rep}(S_n) \rightarrow \Lambda$ .

- Irreducible representations of  $S_n$  are labeled by partitions of  $n$ .
- Irreducible  $S_n$ -representation  $V_\lambda$  has  $\text{Frob}(V_\lambda) = s_\lambda$
- $U \cong V \oplus W \implies \text{Frob}(U) = \text{Frob}(V) + \text{Frob}(W)$
- $\text{Ind}_{S_m \times S_n}^{S_{m+n}}(V \times W) \mapsto \text{Frob}(V) \cdot \text{Frob}(W)$
- Upshot:  $S_n$ -representations go to symmetric functions in structure preserving way.

## Hidden Guide: Schur Positivity

“Naturally occurring” symmetric functions which are non-negative (coefficients in  $\mathbb{N}$ ) linear combinations in Schur polynomial basis are interesting since they could have representation-theoretic models.

# An Example: Harmonic polynomials

Harmonic polynomials

$M =$  polynomials killed by all symmetric differential operators.

# An Example: Harmonic polynomials

## Harmonic polynomials

$M$  = polynomials killed by all symmetric differential operators.

Explicitly, for

$$\Delta = \det \begin{vmatrix} x_1^2 & x_1 & 1 \\ x_2^2 & x_2 & 1 \\ x_3^2 & x_3 & 1 \end{vmatrix} = x_1^2(x_2 - x_3) - x_2^2(x_1 - x_3) + x_3^2(x_1 - x_2)$$

# An Example: Harmonic polynomials

## Harmonic polynomials

$M$  = polynomials killed by all symmetric differential operators.

Explicitly, for

$$\Delta = \det \begin{vmatrix} x_1^2 & x_1 & 1 \\ x_2^2 & x_2 & 1 \\ x_3^2 & x_3 & 1 \end{vmatrix} = x_1^2(x_2 - x_3) - x_2^2(x_1 - x_3) + x_3^2(x_1 - x_2)$$

$M$  is the vector space given by



# An Example: Harmonic polynomials

## Harmonic polynomials

$M$  = polynomials killed by all symmetric differential operators.

Explicitly, for

$$\Delta = \det \begin{vmatrix} x_1^2 & x_1 & 1 \\ x_2^2 & x_2 & 1 \\ x_3^2 & x_3 & 1 \end{vmatrix} = x_1^2(x_2 - x_3) - x_2^2(x_1 - x_3) + x_3^2(x_1 - x_2)$$

$M$  is the vector space given by

$$\begin{aligned} M &= \text{sp} \left\{ \left( \partial_{x_1}^a \partial_{x_2}^b \partial_{x_3}^c \right) \Delta \mid a, b, c \geq 0 \right\} \\ &= \text{sp} \{ \Delta, 2x_1(x_2 - x_3) - x_2^2 + x_3^2, 2x_2(x_3 - x_1) - x_3^2 + x_1^2, \\ &\quad x_3 - x_1, x_2 - x_3, 1 \} \end{aligned}$$

# Harmonic polynomials

$$\text{sp}\{\Delta, 2x_1(x_2 - x_3) - x_2^2 + x_3^2, 2x_2(x_3 - x_1) - x_3^2 + x_1^2, x_3 - x_1, x_2 - x_3, 1\}$$

# Harmonic polynomials

$$\text{sp}\{\Delta, 2x_1(x_2 - x_3) - x_2^2 + x_3^2, 2x_2(x_3 - x_1) - x_3^2 + x_1^2, x_3 - x_1, x_2 - x_3, 1\}$$

- 1 Break  $M$  up into irreducible  $S_n$ -representations (smallest  $S_n$  fixed subspaces).

# Harmonic polynomials

$$\text{sp}\{\Delta, 2x_1(x_2 - x_3) - x_2^2 + x_3^2, 2x_2(x_3 - x_1) - x_3^2 + x_1^2, x_3 - x_1, x_2 - x_3, 1\}$$

- ① Break  $M$  up into irreducible  $S_n$ -representations (smallest  $S_n$  fixed subspaces).

$$\underbrace{\text{sp}\{\Delta\}}_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}} \oplus \underbrace{\text{sp}\{2x_1(x_2 - x_3) - x_2^2 + x_3^2, 2x_2(x_3 - x_1) - x_3^2 + x_1^2\}}_{\begin{array}{|c|} \hline \square \\ \hline \square \square \\ \hline \end{array}} \oplus \underbrace{\text{sp}\{x_3 - x_1, x_2 - x_3\}}_{\begin{array}{|c|} \hline \square \\ \hline \square \square \\ \hline \end{array}} \oplus \underbrace{\text{sp}\{1\}}_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}}$$

# Harmonic polynomials

$$\text{sp}\{\Delta, 2x_1(x_2 - x_3) - x_2^2 + x_3^2, 2x_2(x_3 - x_1) - x_3^2 + x_1^2, x_3 - x_1, x_2 - x_3, 1\}$$

- ① Break  $M$  up into irreducible  $S_n$ -representations (smallest  $S_n$  fixed subspaces).

$$\underbrace{\text{sp}\{\Delta\}}_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}} \oplus \underbrace{\text{sp}\{2x_1(x_2 - x_3) - x_2^2 + x_3^2, 2x_2(x_3 - x_1) - x_3^2 + x_1^2\}}_{\begin{array}{|c|} \hline \square \\ \hline \square \square \\ \hline \end{array}} \oplus \underbrace{\text{sp}\{x_3 - x_1, x_2 - x_3\}}_{\begin{array}{|c|} \hline \square \\ \hline \square \square \\ \hline \end{array}} \oplus \underbrace{\text{sp}\{1\}}_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}}$$

- ② How many times does an irreducible  $S_n$ -representation occur?

# Harmonic polynomials

$$\text{sp}\{\Delta, 2x_1(x_2 - x_3) - x_2^2 + x_3^2, 2x_2(x_3 - x_1) - x_3^2 + x_1^2, x_3 - x_1, x_2 - x_3, 1\}$$

- ① Break  $M$  up into irreducible  $S_n$ -representations (smallest  $S_n$  fixed subspaces).

$$\underbrace{\text{sp}\{\Delta\}}_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}} \oplus \underbrace{\text{sp}\{2x_1(x_2 - x_3) - x_2^2 + x_3^2, 2x_2(x_3 - x_1) - x_3^2 + x_1^2\}}_{\begin{array}{|c|} \hline \square \\ \hline \square \square \\ \hline \end{array}} \oplus \underbrace{\text{sp}\{x_3 - x_1, x_2 - x_3\}}_{\begin{array}{|c|} \hline \square \\ \hline \square \square \\ \hline \end{array}} \oplus \underbrace{\text{sp}\{1\}}_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}}$$

- ② How many times does an irreducible  $S_n$ -representation occur?  
Frobenius:

# Harmonic polynomials

$$\text{sp}\{\Delta, 2x_1(x_2 - x_3) - x_2^2 + x_3^2, 2x_2(x_3 - x_1) - x_3^2 + x_1^2, x_3 - x_1, x_2 - x_3, 1\}$$

- ① Break  $M$  up into irreducible  $S_n$ -representations (smallest  $S_n$  fixed subspaces).

$$\underbrace{\text{sp}\{\Delta\}}_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}} \oplus \underbrace{\text{sp}\{2x_1(x_2 - x_3) - x_2^2 + x_3^2, 2x_2(x_3 - x_1) - x_3^2 + x_1^2\}}_{\begin{array}{|c|} \hline \square \\ \hline \square \square \\ \hline \end{array}} \oplus \underbrace{\text{sp}\{x_3 - x_1, x_2 - x_3\}}_{\begin{array}{|c|} \hline \square \\ \hline \square \square \\ \hline \end{array}} \oplus \underbrace{\text{sp}\{1\}}_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}}$$

- ② How many times does an irreducible  $S_n$ -representation occur?  
Frobenius:

$$e_1^3 = (x_1 + x_2 + x_3)^3 = s_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}} + s_{\begin{array}{|c|} \hline \square \\ \hline \square \square \\ \hline \end{array}} + s_{\begin{array}{|c|} \hline \square \\ \hline \square \square \\ \hline \end{array}} + s_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}}$$

# Harmonic polynomials

$$\text{sp}\{\Delta, 2x_1(x_2 - x_3) - x_2^2 + x_3^2, 2x_2(x_3 - x_1) - x_3^2 + x_1^2, x_3 - x_1, x_2 - x_3, 1\}$$

- ① Break  $M$  up into irreducible  $S_n$ -representations (smallest  $S_n$  fixed subspaces).

$$\underbrace{\text{sp}\{\Delta\}}_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}} \oplus \underbrace{\text{sp}\{2x_1(x_2 - x_3) - x_2^2 + x_3^2, 2x_2(x_3 - x_1) - x_3^2 + x_1^2\}}_{\begin{array}{|c|} \hline \square \\ \hline \square \square \\ \hline \end{array}} \oplus \underbrace{\text{sp}\{x_3 - x_1, x_2 - x_3\}}_{\begin{array}{|c|} \hline \square \\ \hline \square \square \\ \hline \end{array}} \oplus \underbrace{\text{sp}\{1\}}_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}}$$

- ② How many times does an irreducible  $S_n$ -representation occur?  
Frobenius:

$$e_1^3 = (x_1 + x_2 + x_3)^3 = s_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}} + s_{\begin{array}{|c|} \hline \square \square \\ \hline \end{array}} + s_{\begin{array}{|c|} \hline \square \\ \hline \square \square \\ \hline \end{array}} + s_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}}$$

Remark:  $M \cong \mathbb{C}[x_1, x_2, x_3]/(\mathbb{C}[x_1, x_2, x_3]_{+}^{S_3})$ .



## Getting more information

# Getting more information

Break  $M$  up into irreducible representations.

$$\underbrace{\text{sp}\{\Delta\}}_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}} \oplus \underbrace{\text{sp}\{2x_1(x_2-x_3)-x_2^2+x_3^2, 2x_2(x_3-x_1)-x_3^2+x_1^2\}}_{\begin{array}{|c|c|} \hline \square & \\ \hline \square & \square \\ \hline \end{array} \text{deg}=2} \oplus \underbrace{\text{sp}\{x_3-x_1, x_2-x_3\}}_{\begin{array}{|c|c|} \hline \square & \\ \hline \square & \square \\ \hline \end{array} \text{deg}=1} \oplus \underbrace{\text{sp}\{1\}}_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}}$$

# Getting more information

Break  $M$  up into irreducible representations.

$$\underbrace{\text{sp}\{\Delta\}}_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}} \oplus \underbrace{\text{sp}\{2x_1(x_2-x_3)-x_2^2+x_3^2, 2x_2(x_3-x_1)-x_3^2+x_1^2\}}_{\begin{array}{|c|c|} \hline \square & \\ \hline \square & \square \\ \hline \end{array} \text{deg}=2} \oplus \underbrace{\text{sp}\{x_3-x_1, x_2-x_3\}}_{\begin{array}{|c|c|} \hline \square & \\ \hline \square & \square \\ \hline \end{array} \text{deg}=1} \oplus \underbrace{\text{sp}\{1\}}_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}}$$

Solution: irreducible  $S_n$ -representation of polynomials of degree  $d \mapsto q^d s_\lambda$   
(graded Frobenius)

$$?? = q^3 s_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}} + q^2 s_{\begin{array}{|c|c|} \hline \square & \\ \hline \square & \square \\ \hline \end{array}} + q s_{\begin{array}{|c|c|} \hline \square & \\ \hline \square & \square \\ \hline \end{array}} + s_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}}$$

# Getting more information

Break  $M$  up into irreducible representations.

$$\underbrace{\text{sp}\{\Delta\}}_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}} \oplus \underbrace{\text{sp}\{2x_1(x_2-x_3)-x_2^2+x_3^2, 2x_2(x_3-x_1)-x_3^2+x_1^2\}}_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \text{deg}=2} \oplus \underbrace{\text{sp}\{x_3-x_1, x_2-x_3\}}_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \text{deg}=1} \oplus \underbrace{\text{sp}\{1\}}_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}}$$

Solution: irreducible  $S_n$ -representation of polynomials of degree  $d \mapsto q^d s_\lambda$   
(graded Frobenius)

$$?? = q^3 s_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}} + q^2 s_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} + q s_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} + s_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}}$$

Answer: Hall-Littlewood polynomial  $H_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}}(X; q)$ .

# A Problem

- In 1988, Macdonald introduces a family of symmetric polynomials with coefficients in  $\mathbb{Q}(q, t)$  generalizing Hall-Littlewood polynomials (and many other famous bases).

# A Problem

- In 1988, Macdonald introduces a family of symmetric polynomials with coefficients in  $\mathbb{Q}(q, t)$  generalizing Hall-Littlewood polynomials (and many other famous bases).
- Defined by orthogonality and triangularity under a certain inner-product. (Indirect)

# A Problem

- In 1988, Macdonald introduces a family of symmetric polynomials with coefficients in  $\mathbb{Q}(q, t)$  generalizing Hall-Littlewood polynomials (and many other famous bases).
- Defined by orthogonality and triangularity under a certain inner-product. (Indirect)
- Garsia modifies these polynomials so

$$\tilde{H}_\lambda(X; q, t) = \sum_{\mu} \tilde{K}_{\lambda\mu}(q, t) s_{\mu} \text{ conjecturally satisfies } \tilde{K}_{\lambda\mu}(q, t) \in \mathbb{N}[q, t]$$

# A Problem

- In 1988, Macdonald introduces a family of symmetric polynomials with coefficients in  $\mathbb{Q}(q, t)$  generalizing Hall-Littlewood polynomials (and many other famous bases).
- Defined by orthogonality and triangularity under a certain inner-product. (Indirect)
- Garsia modifies these polynomials so

$$\tilde{H}_\lambda(X; q, t) = \sum_{\mu} \tilde{K}_{\lambda\mu}(q, t) s_{\mu} \text{ conjecturally satisfies } \tilde{K}_{\lambda\mu}(q, t) \in \mathbb{N}[q, t]$$

- $\tilde{H}_\lambda(X; 1, 1) = e_1^{|\lambda|}$ .



# A Problem

- In 1988, Macdonald introduces a family of symmetric polynomials with coefficients in  $\mathbb{Q}(q, t)$  generalizing Hall-Littlewood polynomials (and many other famous bases).
- Defined by orthogonality and triangularity under a certain inner-product. (Indirect)
- Garsia modifies these polynomials so

$$\tilde{H}_\lambda(X; q, t) = \sum_{\mu} \tilde{K}_{\lambda\mu}(q, t) s_{\mu} \text{ conjecturally satisfies } \tilde{K}_{\lambda\mu}(q, t) \in \mathbb{N}[q, t]$$

- $\tilde{H}_\lambda(X; 1, 1) = e_1^{|\lambda|}$ .
- Does there exist a family of  $S_n$ -regular representations whose bigraded Frobenius characteristics equal  $\tilde{H}_\lambda(X; q, t)$ ?

# Garsia-Haiman modules

- $\mathbb{Q}[x_1, \dots, x_n, y_1, \dots, y_n]$  with  $\sigma(x_i) = x_{\sigma(i)}$ ,  $\sigma(y_j) = y_{\sigma(j)}$ .

# Garsia-Haiman modules

- $\mathbb{Q}[x_1, \dots, x_n, y_1, \dots, y_n]$  with  $\sigma(x_i) = x_{\sigma(i)}$ ,  $\sigma(y_j) = y_{\sigma(j)}$ .
- Garsia-Haiman (1993):  $M_\mu = \text{span of partial derivatives of}$   
 $\Delta_\mu = \det_{(i,j) \in \mu, k \in [n]} (x_k^{i-1} y_k^{j-1})$

# Garsia-Haiman modules

- $\mathbb{Q}[x_1, \dots, x_n, y_1, \dots, y_n]$  with  $\sigma(x_i) = x_{\sigma(i)}$ ,  $\sigma(y_j) = y_{\sigma(j)}$ .
- Garsia-Haiman (1993):  $M_\mu = \text{span of partial derivatives of}$   
 $\Delta_\mu = \det_{(i,j) \in \mu, k \in [n]} (x_k^{i-1} y_k^{j-1})$

$$\Delta_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} = \det \begin{vmatrix} 1 & y_1 & x_1 \\ 1 & y_2 & x_2 \\ 1 & y_3 & x_3 \end{vmatrix} = x_3 y_2 - y_3 x_2 - y_1 x_3 + y_1 x_2 + y_3 x_1 - y_2 x_1$$

# Garsia-Haiman modules

- $\mathbb{Q}[x_1, \dots, x_n, y_1, \dots, y_n]$  with  $\sigma(x_i) = x_{\sigma(i)}$ ,  $\sigma(y_j) = y_{\sigma(j)}$ .
- Garsia-Haiman (1993):  $M_\mu = \text{span of partial derivatives of } \Delta_\mu = \det_{(i,j) \in \mu, k \in [n]} (x_k^{i-1} y_k^{j-1})$

$$\Delta_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} = \det \begin{vmatrix} 1 & y_1 & x_1 \\ 1 & y_2 & x_2 \\ 1 & y_3 & x_3 \end{vmatrix} = x_3 y_2 - y_3 x_2 - y_1 x_3 + y_1 x_2 + y_3 x_1 - y_2 x_1$$

$$M_{2,1} = \underbrace{\text{sp}\{\Delta_{2,1}\}}_{\deg=(1,1)} \oplus \underbrace{\text{sp}\{y_3 - y_1, y_1 - y_2\}}_{\deg=(0,1)} \oplus \underbrace{\text{sp}\{x_3 - x_1, x_1 - x_2\}}_{\deg=(1,0)} \oplus \underbrace{\text{sp}\{1\}}_{\deg=(0,0)}$$

# Garsia-Haiman modules

- $\mathbb{Q}[x_1, \dots, x_n, y_1, \dots, y_n]$  with  $\sigma(x_i) = x_{\sigma(i)}$ ,  $\sigma(y_j) = y_{\sigma(j)}$ .
- Garsia-Haiman (1993):  $M_\mu = \text{span of partial derivatives of } \Delta_\mu = \det_{(i,j) \in \mu, k \in [n]} (x_k^{i-1} y_k^{j-1})$

$$\Delta_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} = \det \begin{vmatrix} 1 & y_1 & x_1 \\ 1 & y_2 & x_2 \\ 1 & y_3 & x_3 \end{vmatrix} = x_3 y_2 - y_3 x_2 - y_1 x_3 + y_1 x_2 + y_3 x_1 - y_2 x_1$$

$$M_{2,1} = \underbrace{\text{sp}\{\Delta_{2,1}\}}_{\deg=(1,1)} \oplus \underbrace{\text{sp}\{y_3 - y_1, y_1 - y_2\}}_{\deg=(0,1)} \oplus \underbrace{\text{sp}\{x_3 - x_1, x_1 - x_2\}}_{\deg=(1,0)} \oplus \underbrace{\text{sp}\{1\}}_{\deg=(0,0)}$$

Irreducible  $S_n$ -representation with bidegree  $(a, b) \mapsto q^a t^b s_\lambda$

# Garsia-Haiman modules

- $\mathbb{Q}[x_1, \dots, x_n, y_1, \dots, y_n]$  with  $\sigma(x_i) = x_{\sigma(i)}$ ,  $\sigma(y_j) = y_{\sigma(j)}$ .
- Garsia-Haiman (1993):  $M_\mu = \text{span of partial derivatives of } \Delta_\mu = \det_{(i,j) \in \mu, k \in [n]} (x_k^{i-1} y_k^{j-1})$

$$\Delta_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} = \det \begin{vmatrix} 1 & y_1 & x_1 \\ 1 & y_2 & x_2 \\ 1 & y_3 & x_3 \end{vmatrix} = x_3 y_2 - y_3 x_2 - y_1 x_3 + y_1 x_2 + y_3 x_1 - y_2 x_1$$

$$M_{2,1} = \underbrace{\text{sp}\{\Delta_{2,1}\}}_{\deg=(1,1)} \oplus \underbrace{\text{sp}\{y_3 - y_1, y_1 - y_2\}}_{\deg=(0,1)} \oplus \underbrace{\text{sp}\{x_3 - x_1, x_1 - x_2\}}_{\deg=(1,0)} \oplus \underbrace{\text{sp}\{1\}}_{\deg=(0,0)}$$

Irreducible  $S_n$ -representation with bidegree  $(a, b) \mapsto q^a t^b s_\lambda$

$$\tilde{H}_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} = qts \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array} + ts \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} + qs \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} + s \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}$$

# Garsia-Haiman modules

## Theorem (Haiman, 2001)

*The Garsia-Haiman module  $M_\lambda$  has bigraded Frobenius characteristic given by  $\tilde{H}_\lambda(X; q, t)$ .*



## Theorem (Haiman, 2001)

*The Garsia-Haiman module  $M_\lambda$  has bigraded Frobenius characteristic given by  $\tilde{H}_\lambda(X; q, t)$ .*

- Proved via connection to the Hilbert Scheme  $\text{Hilb}^n(\mathbb{C}^2)$ .

# Garsia-Haiman modules

## Theorem (Haiman, 2001)

*The Garsia-Haiman module  $M_\lambda$  has bigraded Frobenius characteristic given by  $\tilde{H}_\lambda(X; q, t)$ .*

- Proved via connection to the Hilbert Scheme  $\text{Hilb}^n(\mathbb{C}^2)$ .

## Corollary

$\tilde{H}_\lambda(X; q, t) = \sum_\mu \tilde{K}_{\lambda\mu}(q, t) s_\mu$  satisfies  $\tilde{K}_{\lambda\mu}(q, t) \in \mathbb{N}[q, t]$ .

# Garsia-Haiman modules

## Theorem (Haiman, 2001)

*The Garsia-Haiman module  $M_\lambda$  has bigraded Frobenius characteristic given by  $\tilde{H}_\lambda(X; q, t)$ .*

- Proved via connection to the Hilbert Scheme  $\text{Hilb}^n(\mathbb{C}^2)$ .

## Corollary

$\tilde{H}_\lambda(X; q, t) = \sum_\mu \tilde{K}_{\lambda\mu}(q, t) s_\mu$  satisfies  $\tilde{K}_{\lambda\mu}(q, t) \in \mathbb{N}[q, t]$ .

- No combinatorial description of  $\tilde{K}_{\lambda\mu}(q, t)$ .

- ① Background on symmetric functions and Macdonald polynomials
- ② **A new formula for Macdonald polynomials**
- ③ LLT polynomials in the elliptic Hall algebra

# Root ideals

$R_+ = \{\alpha_{ij} \mid 1 \leq i < j \leq n\}$  denotes the set of positive roots for  $GL_n$ , where  $\alpha_{ij} = \epsilon_i - \epsilon_j$ .

	(12)	(13)	(14)	(15)
		(23)	(24)	(25)
			(34)	(35)
				(45)

# Root ideals

$R_+ = \{\alpha_{ij} \mid 1 \leq i < j \leq n\}$  denotes the set of positive roots for  $GL_n$ , where  $\alpha_{ij} = \epsilon_i - \epsilon_j$ .

	(12)	(13)	(14)	(15)
		(23)	(24)	(25)
			(34)	(35)
				(45)

A root ideal  $\Psi \subseteq R_+$  is an upper order ideal of positive roots.

	(12)	(13)	(14)	(15)
		(23)	(24)	(25)
			(34)	(35)
				(45)

$\Psi = \text{Roots above Dyck path}$

# Weyl symmetrization

Define the *Weyl symmetrization operator*  $\sigma: \mathbb{Q}[z_1^{\pm 1}, \dots, z_n^{\pm 1}] \rightarrow \Lambda(X)$  by linearly extending

$$\mathbf{z}^\gamma \mapsto s_\gamma(X)$$

where  $\mathbf{z}^\gamma = z_1^{\gamma_1} \cdots z_n^{\gamma_n}$ .

# Weyl symmetrization

Define the *Weyl symmetrization operator*  $\sigma: \mathbb{Q}[z_1^{\pm 1}, \dots, z_n^{\pm 1}] \rightarrow \Lambda(X)$  by linearly extending

$$\mathbf{z}^\gamma \mapsto s_\gamma(X)$$

where  $\mathbf{z}^\gamma = z_1^{\gamma_1} \cdots z_n^{\gamma_n}$ .

## Definition

A *Catalan function* is a symmetric function indexed by a root ideal  $\Psi \subseteq R_+$  and  $\gamma \in \mathbb{Z}^n$



# Weyl symmetrization

Define the *Weyl symmetrization operator*  $\sigma: \mathbb{Q}[z_1^{\pm 1}, \dots, z_n^{\pm 1}] \rightarrow \Lambda(X)$  by linearly extending

$$\mathbf{z}^\gamma \mapsto s_\gamma(X)$$

where  $\mathbf{z}^\gamma = z_1^{\gamma_1} \cdots z_n^{\gamma_n}$ .

## Definition

A *Catalan function* is a symmetric function indexed by a root ideal  $\Psi \subseteq R_+$  and  $\gamma \in \mathbb{Z}^n$  given by

$$H(\Phi; \gamma) = \sigma \left( \frac{\mathbf{z}^\gamma}{\prod_{(i,j) \in \Psi} (1 - tz_i/z_j)} \right)$$

Denominator factors are understood as geometric series  
 $(1 - tz_i/z_j)^{-1} = 1 + tz_i/z_j + t^2(z_i/z_j)^2 + \cdots$


# Catalan functions

## Definition

A *Catalan function* is a symmetric function indexed by a root ideal  $\Psi \subseteq R_+$  and  $\gamma \in \mathbb{Z}^n$  given by

$$H(\Phi; \gamma) = \sigma \left( \frac{z^\gamma}{\prod_{(i,j) \in \Psi} (1 - tz_i/z_j)} \right)$$

Example:

$\Psi =$    $\gamma = (1, 1, 1)$


# Catalan functions

## Definition

A *Catalan function* is a symmetric function indexed by a root ideal  $\Psi \subseteq R_+$  and  $\gamma \in \mathbb{Z}^n$  given by

$$H(\Psi; \gamma) = \sigma \left( \frac{\mathbf{z}^\gamma}{\prod_{(i,j) \in \Psi} (1 - tz_i/z_j)} \right)$$

Example:

$\Psi =$    $\gamma = (1, 1, 1)$

$$\begin{aligned} H(\Psi; \gamma) &= \sigma \left( \left(1 + t \frac{z_1}{z_2} + t^2 \frac{z_1^2}{z_2^2} + \dots\right) \left(1 + t \frac{z_1}{z_3} + t^2 \frac{z_1^2}{z_3^2} + \dots\right) z_1 z_2 z_3 \right) \\ &= s_{111} + t(s_{201} + s_{210}) + t^2(s_{3-10} + s_{300} + s_{31-1}) + \dots \\ &= s_{111} + ts_{210} \end{aligned}$$


# A Catalan function for modified Hall-Littlewoods

$B_\mu$  = set of roots above block diagonal matrix with block sizes  $\mu_{\ell(\mu)}, \dots, \mu_1$



# A Catalan function for modified Hall-Littlewoods

$B_\mu$  = set of roots above block diagonal matrix with block sizes  $\mu_{\ell(\mu)}, \dots, \mu_1$



Theorem (Weyman, Shimozono-Weyman)

$$\tilde{H}_\mu(X; 0, t) = \omega \sigma \left( \frac{z_1 \cdots z_n}{\prod_{\alpha \in B_\mu} (1 - t z^\alpha)} \right),$$

where  $z^\alpha = z_i / z_j$ .

$\omega(s_\lambda) = s_{\lambda'}$  for  $\lambda'$  the transpose partition of  $\lambda$ .


# Catalan functions for modified Hall-Littlewoods

$b_1$		
$b_2$	$b_3$	
$b_4$	$b_5$	$b_6$
$b_7$	$b_8$	$b_9$

row reading order

$$b_1 \prec b_2 \prec \cdots \prec b_n$$

$$R_\mu := \{ \alpha_{ij} \in R_+ \mid \text{south}(b_i) \preceq b_j \}.$$

$$R_{3321} =$$



# Catalan functions for modified Hall-Littlewoods

$b_1$		
$b_2$	$b_3$	
$b_4$	$b_5$	$b_6$
$b_7$	$b_8$	$b_9$

row reading order

$$b_1 \prec b_2 \prec \cdots \prec b_n$$

$$R_\mu := \{ \alpha_{ij} \in R_+ \mid \text{south}(b_i) \preceq b_j \}.$$



$$\begin{aligned} \tilde{H}_\mu(X; 0, t) &= \omega \sigma \left( \frac{z_1 \cdots z_n}{\prod_{\alpha \in B_\mu} (1 - t z^\alpha)} \right), \\ &= \omega \sigma \left( \frac{z_1 \cdots z_n}{\prod_{\alpha \in R_\mu} (1 - t z^\alpha)} \right) \end{aligned}$$

# A formula for $\tilde{H}_\mu(X; q, t)$

$b_1$	
$b_2$	
$b_3$	$b_4$
$b_5$	$b_6$
$b_7$	$b_8$

row reading order

$$b_1 \prec b_2 \prec \cdots \prec b_n$$

$$R_\mu := \{ \alpha_{ij} \in R_+ \mid \text{south}(b_i) \preceq b_j \},$$

$$\hat{R}_\mu := \{ \alpha_{ij} \in R_+ \mid \text{south}(b_i) \prec b_j \}.$$



# A formula for $\tilde{H}_\mu(X; q, t)$

$b_1$	
$b_2$	
$b_3$	$b_4$
$b_5$	$b_6$
$b_7$	$b_8$

row reading order

$$b_1 \prec b_2 \prec \cdots \prec b_n$$

$$R_\mu := \{\alpha_{ij} \in R_+ \mid \text{south}(b_i) \preceq b_j\},$$

$$\hat{R}_\mu := \{\alpha_{ij} \in R_+ \mid \text{south}(b_i) \prec b_j\}.$$

## Theorem (Blasiak-Haiman-Morse-Pun-S.)


The modified Macdonald polynomial  $\tilde{H}_\mu = \tilde{H}_\mu(X; q, t)$  is given by

$$\tilde{H}_\mu = \omega \sigma \left( z_1 \cdots z_n \frac{\prod_{\alpha_{ij} \in R_\mu \setminus \hat{R}_\mu} (1 - q^{\text{arm}(b_i)+1} t^{-\text{leg}(b_i)} z_i / z_j) \prod_{\alpha \in \hat{R}_\mu} (1 - q t z^\alpha)}{\prod_{\alpha \in R_+} (1 - q z^\alpha) \prod_{\alpha \in R_\mu} (1 - t z^\alpha)} \right).$$

# Example



partition  $\mu = 22211$



# Example

$1 - q^{\frac{z_1}{z_2}}$	
$1 - qt^{-1} \frac{z_2}{z_3}$	
$1 - q^2 t^{-2} \frac{z_3}{z_5}$	$1 - q^{\frac{z_4}{z_6}}$
$1 - q^2 t^{-3} \frac{z_5}{z_7}$	$1 - qt^{-1} \frac{z_6}{z_8}$

numerator factors  $1 - q^{\text{arm}+1} t^{-\text{leg}} z_i/z_j$



●  $R_\mu \setminus \hat{R}_\mu$  ( $t$  factors)

■  $\hat{R}_\mu$  ( $t$  and  $qt$  factors)

$q = t = 1$  specialization

$$\begin{aligned}
 & \omega \sigma \left( z_1 \cdots z_n \frac{\prod_{\alpha_{ij} \in R_\mu \setminus \hat{R}_\mu} (1 - q^{\text{arm}(b_i)+1} t^{-\text{leg}(b_i)} z_i / z_j) \prod_{\alpha \in \hat{R}_\mu} (1 - q t z^\alpha)}{\prod_{\alpha \in R_+} (1 - q z^\alpha) \prod_{\alpha \in R_\mu} (1 - t z^\alpha)} \right) \\
 & \xrightarrow{q=t=1} \omega \sigma \left( z_1 \cdots z_n \frac{\prod_{\alpha \in R_\mu \setminus \hat{R}_\mu} (1 - z^\alpha) \prod_{\alpha \in \hat{R}_\mu} (1 - z^\alpha)}{\prod_{\alpha \in R_+} (1 - z^\alpha) \prod_{\alpha \in R_\mu} (1 - z^\alpha)} \right) \\
 & = \omega \sigma \left( \frac{z_1 \cdots z_n}{\prod_{\alpha \in R_+} (1 - z^\alpha)} \right) \\
 & = \omega h_1^n \\
 & = e_1^n
 \end{aligned}$$

$q = 0$  specialization

$$\begin{aligned}
 & \omega\sigma \left( z_1 \cdots z_n \frac{\prod_{\alpha_{ij} \in R_\mu \setminus \widehat{R}_\mu} (1 - q^{\text{arm}(b_i)+1} t^{-\text{leg}(b_i)} z_i/z_j) \prod_{\alpha \in \widehat{R}_\mu} (1 - q t z^\alpha)}{\prod_{\alpha \in R_+} (1 - q z^\alpha) \prod_{\alpha \in R_\mu} (1 - t z^\alpha)} \right) \\
 & \xrightarrow{q=0} \omega\sigma \left( \frac{z_1 \cdots z_n}{\prod_{\alpha \in R_\mu} (1 - t z^\alpha)} \right) \\
 & = \tilde{H}_\mu(X; 0, t)
 \end{aligned}$$

# Proof of formula for $\tilde{H}_\mu$

## Definition

$\nabla$  is the linear operator on symmetric functions satisfying  $\nabla \tilde{H}_\mu = t^{n(\mu)} q^{n(\mu^*)} \tilde{H}_\mu$ , where  $n(\mu) = \sum_i (i-1)\mu_i$ .

# Proof of formula for $\tilde{H}_\mu$

## Definition

$\nabla$  is the linear operator on symmetric functions satisfying  $\nabla \tilde{H}_\mu = t^{n(\mu)} q^{n(\mu^*)} \tilde{H}_\mu$ , where  $n(\mu) = \sum_i (i-1)\mu_i$ .

- Start with the Haglund-Haiman-Loehr formula for  $\tilde{H}_\mu$  as a sum of LLT polynomials  $\mathcal{G}_\nu(X; q)$ .

# Proof of formula for $\tilde{H}_\mu$

## Definition

$\nabla$  is the linear operator on symmetric functions satisfying  $\nabla \tilde{H}_\mu = t^{n(\mu)} q^{n(\mu^*)} \tilde{H}_\mu$ , where  $n(\mu) = \sum_i (i-1)\mu_i$ .

- Start with the Haglund-Haiman-Loehr formula for  $\tilde{H}_\mu$  as a sum of LLT polynomials  $\mathcal{G}_\nu(X; q)$ .
- Apply  $\omega \nabla$  to both sides.



# Proof of formula for $\tilde{H}_\mu$

## Definition

$\nabla$  is the linear operator on symmetric functions satisfying  $\nabla \tilde{H}_\mu = t^{n(\mu)} q^{n(\mu^*)} \tilde{H}_\mu$ , where  $n(\mu) = \sum_i (i-1)\mu_i$ .

- Start with the Haglund-Haiman-Loehr formula for  $\tilde{H}_\mu$  as a sum of LLT polynomials  $\mathcal{G}_\nu(X; q)$ .
- Apply  $\omega \nabla$  to both sides.
- Use Catalan-like (“Catalanimal”) formula for  $\omega \nabla \mathcal{G}_\nu(X; q)$  and collect terms.

# LLT Polynomials

Let  $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$  be a tuple of skew shapes.

$$\nu = \left( \begin{array}{|c|c|c|} \hline \square & \square & \\ \hline \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array} \right)$$



# LLT Polynomials

Let  $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$  be a tuple of skew shapes.

- The *content* of a box in row  $y$ , column  $x$  is  $x - y$ .

$$\nu = \left( \begin{array}{|c|c|c|} \hline \square & \square & \\ \hline \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array} \right)$$

-4	-3	-2	-1	0	1
-3	-2	-1	0	1	2
-2	-1	0	1	2	3
-1	0	1	2	3	4
0	1	2	3	4	5

# LLT Polynomials

Let  $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$  be a tuple of skew shapes.

- The *content* of a box in row  $y$ , column  $x$  is  $x - y$ .
- *Reading order*: label boxes  $b_1, \dots, b_n$  by scanning each diagonal from southwest to northeast, in order of increasing content.

$$\nu = \left( \begin{array}{|c|c|c|} \hline \square & \square & \\ \hline \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array} \right)$$

				$b_3$	$b_6$
				$b_5$	$b_8$
$b_1$	$b_2$				
	$b_4$	$b_7$			

# LLT Polynomials

Let  $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$  be a tuple of skew shapes.

- The *content* of a box in row  $y$ , column  $x$  is  $x - y$ .
- *Reading order*: label boxes  $b_1, \dots, b_n$  by scanning each diagonal from southwest to northeast, in order of increasing content.
- A pair  $(a, b) \in \nu$  is *attacking* if  $a$  precedes  $b$  in reading order and
  - $\text{content}(b) = \text{content}(a)$ , or
  - $\text{content}(b) = \text{content}(a) + 1$  and  $a \in \nu_{(i)}, b \in \nu_{(j)}$  with  $i > j$ .

$$\nu = \left( \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right)$$

				$b_3$	$b_6$
				$b_5$	$b_8$
$b_1$	$b_2$				
	$b_4$	$b_7$			

Attacking pairs:  $(b_2, b_3), (b_3, b_4), (b_4, b_5), (b_4, b_6), (b_5, b_7), (b_6, b_7), (b_7, b_8)$

# LLT Polynomials

Let  $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$  be a tuple of skew shapes.

- The *content* of a box in row  $y$ , column  $x$  is  $x - y$ .
- *Reading order*: label boxes  $b_1, \dots, b_n$  by scanning each diagonal from southwest to northeast, in order of increasing content.
- A pair  $(a, b) \in \nu$  is *attacking* if  $a$  precedes  $b$  in reading order and
  - $\text{content}(b) = \text{content}(a)$ , or
  - $\text{content}(b) = \text{content}(a) + 1$  and  $a \in \nu_{(i)}, b \in \nu_{(j)}$  with  $i > j$ .

$$\nu = \left( \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right)$$

				$b_3$	$b_6$
				$b_5$	$b_8$
$b_1$	$b_2$				
	$b_4$	$b_7$			

Attacking pairs:  $(b_2, b_3), (b_3, b_4), (b_4, b_5), (b_4, b_6), (b_5, b_7), (b_6, b_7), (b_7, b_8)$

# LLT Polynomials

Let  $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$  be a tuple of skew shapes.

- The *content* of a box in row  $y$ , column  $x$  is  $x - y$ .
- *Reading order*: label boxes  $b_1, \dots, b_n$  by scanning each diagonal from southwest to northeast, in order of increasing content.
- A pair  $(a, b) \in \nu$  is *attacking* if  $a$  precedes  $b$  in reading order and
  - $\text{content}(b) = \text{content}(a)$ , or
  - $\text{content}(b) = \text{content}(a) + 1$  and  $a \in \nu_{(i)}, b \in \nu_{(j)}$  with  $i > j$ .

$$\nu = \left( \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right)$$

				$b_3$	$b_6$
				$b_5$	$b_8$
$b_1$	$b_2$				
	$b_4$	$b_7$			

Attacking pairs:  $(b_2, b_3), (b_3, b_4), (b_4, b_5), (b_4, b_6), (b_5, b_7), (b_6, b_7), (b_7, b_8)$

# LLT Polynomials

Let  $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$  be a tuple of skew shapes.

- The *content* of a box in row  $y$ , column  $x$  is  $x - y$ .
- *Reading order*: label boxes  $b_1, \dots, b_n$  by scanning each diagonal from southwest to northeast, in order of increasing content.
- A pair  $(a, b) \in \nu$  is *attacking* if  $a$  precedes  $b$  in reading order and
  - $\text{content}(b) = \text{content}(a)$ , or
  - $\text{content}(b) = \text{content}(a) + 1$  and  $a \in \nu_{(i)}, b \in \nu_{(j)}$  with  $i > j$ .

$$\nu = \left( \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right)$$

				$b_3$	$b_6$
				$b_5$	$b_8$
$b_1$	$b_2$				
	$b_4$	$b_7$			

Attacking pairs:  $(b_2, b_3), (b_3, b_4), (b_4, b_5), (b_4, b_6), (b_5, b_7), (b_6, b_7), (b_7, b_8)$



# LLT Polynomials

Let  $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$  be a tuple of skew shapes.

- The *content* of a box in row  $y$ , column  $x$  is  $x - y$ .
- *Reading order*: label boxes  $b_1, \dots, b_n$  by scanning each diagonal from southwest to northeast, in order of increasing content.
- A pair  $(a, b) \in \nu$  is *attacking* if  $a$  precedes  $b$  in reading order and
  - $\text{content}(b) = \text{content}(a)$ , or
  - $\text{content}(b) = \text{content}(a) + 1$  and  $a \in \nu_{(i)}, b \in \nu_{(j)}$  with  $i > j$ .

$$\nu = \left( \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right)$$

				$b_3$	$b_6$
				$b_5$	$b_8$
$b_1$	$b_2$				
	$b_4$	$b_7$			

Attacking pairs:  $(b_2, b_3), (b_3, b_4), (b_4, b_5), (b_4, b_6), (b_5, b_7), (b_6, b_7), (b_7, b_8)$

# LLT Polynomials

Let  $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$  be a tuple of skew shapes.

- The *content* of a box in row  $y$ , column  $x$  is  $x - y$ .
- *Reading order*: label boxes  $b_1, \dots, b_n$  by scanning each diagonal from southwest to northeast, in order of increasing content.
- A pair  $(a, b) \in \nu$  is *attacking* if  $a$  precedes  $b$  in reading order and
  - $\text{content}(b) = \text{content}(a)$ , or
  - $\text{content}(b) = \text{content}(a) + 1$  and  $a \in \nu_{(i)}, b \in \nu_{(j)}$  with  $i > j$ .

$$\nu = \left( \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right)$$

					$b_3$	$b_6$
					$b_5$	$b_8$
$b_1$	$b_2$					
	$b_4$	$b_7$				

Attacking pairs:  $(b_2, b_3), (b_3, b_4), (b_4, b_5), (b_4, b_6), (b_5, b_7), (b_6, b_7), (b_7, b_8)$

# LLT Polynomials

Let  $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$  be a tuple of skew shapes.

- The *content* of a box in row  $y$ , column  $x$  is  $x - y$ .
- *Reading order*: label boxes  $b_1, \dots, b_n$  by scanning each diagonal from southwest to northeast, in order of increasing content.
- A pair  $(a, b) \in \nu$  is *attacking* if  $a$  precedes  $b$  in reading order and
  - $\text{content}(b) = \text{content}(a)$ , or
  - $\text{content}(b) = \text{content}(a) + 1$  and  $a \in \nu_{(i)}, b \in \nu_{(j)}$  with  $i > j$ .

$$\nu = \left( \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right)$$

				$b_3$	$b_6$
				$b_5$	$b_8$
$b_1$	$b_2$				
	$b_4$	$b_7$			

Attacking pairs:  $(b_2, b_3), (b_3, b_4), (b_4, b_5), (b_4, b_6), (b_5, b_7), (b_6, b_7), (b_7, b_8)$

# LLT Polynomials

Let  $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$  be a tuple of skew shapes.

- The *content* of a box in row  $y$ , column  $x$  is  $x - y$ .
- *Reading order*: label boxes  $b_1, \dots, b_n$  by scanning each diagonal from southwest to northeast, in order of increasing content.
- A pair  $(a, b) \in \nu$  is *attacking* if  $a$  precedes  $b$  in reading order and
  - $\text{content}(b) = \text{content}(a)$ , or
  - $\text{content}(b) = \text{content}(a) + 1$  and  $a \in \nu_{(i)}, b \in \nu_{(j)}$  with  $i > j$ .

$$\nu = \left( \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right)$$

				$b_3$	$b_6$
				$b_5$	$b_8$
$b_1$	$b_2$				
	$b_4$	$b_7$			

Attacking pairs:  $(b_2, b_3), (b_3, b_4), (b_4, b_5), (b_4, b_6), (b_5, b_7), (b_6, b_7), (b_7, b_8)$

# LLT Polynomials

- A *semistandard tableau* on  $\nu$  is a map  $T: \nu \rightarrow \mathbb{Z}_+$  which restricts to a semistandard tableau on each  $\nu_{(i)}$ .
- An *attacking inversion* in  $T$  is an attacking pair  $(a, b)$  such that  $T(a) > T(b)$ .

The *LLT polynomial* indexed by a tuple of skew shapes  $\nu$  is

$$\mathcal{G}_\nu(\mathbf{x}; q) = \sum_{T \in \text{SSYT}(\nu)} q^{\text{inv}(T)} \mathbf{x}^T,$$

where  $\text{inv}(T)$  is the number of attacking inversions in  $T$  and  $\mathbf{x}^T = \prod_{a \in \nu} x_{T(a)}$ .

$T =$

				5	6
				1	1
2	4				
	3	5			

# LLT Polynomials

- A *semistandard tableau* on  $\nu$  is a map  $T: \nu \rightarrow \mathbb{Z}_+$  which restricts to a semistandard tableau on each  $\nu_{(i)}$ .
- An *attacking inversion* in  $T$  is an attacking pair  $(a, b)$  such that  $T(a) > T(b)$ .

The *LLT polynomial* indexed by a tuple of skew shapes  $\nu$  is

$$\mathcal{G}_\nu(\mathbf{x}; q) = \sum_{T \in \text{SSYT}(\nu)} q^{\text{inv}(T)} \mathbf{x}^T,$$

where  $\text{inv}(T)$  is the number of attacking inversions in  $T$  and  $\mathbf{x}^T = \prod_{a \in \nu} x_{T(a)}$ .

$T =$

				5	6
				1	1
2	4				
	3	5			

non-inversion

# LLT Polynomials

- A *semistandard tableau* on  $\nu$  is a map  $T: \nu \rightarrow \mathbb{Z}_+$  which restricts to a semistandard tableau on each  $\nu_{(i)}$ .
- An *attacking inversion* in  $T$  is an attacking pair  $(a, b)$  such that  $T(a) > T(b)$ .

The *LLT polynomial* indexed by a tuple of skew shapes  $\nu$  is

$$\mathcal{G}_\nu(\mathbf{x}; q) = \sum_{T \in \text{SSYT}(\nu)} q^{\text{inv}(T)} \mathbf{x}^T,$$

where  $\text{inv}(T)$  is the number of attacking inversions in  $T$  and  $\mathbf{x}^T = \prod_{a \in \nu} x_{T(a)}$ .

$T =$

				5	6	
				1	1	
2	4					
	3	5				

inversion

# LLT Polynomials

- A *semistandard tableau* on  $\nu$  is a map  $T: \nu \rightarrow \mathbb{Z}_+$  which restricts to a semistandard tableau on each  $\nu_{(i)}$ .
- An *attacking inversion* in  $T$  is an attacking pair  $(a, b)$  such that  $T(a) > T(b)$ .

The *LLT polynomial* indexed by a tuple of skew shapes  $\nu$  is

$$\mathcal{G}_\nu(\mathbf{x}; q) = \sum_{T \in \text{SSYT}(\nu)} q^{\text{inv}(T)} \mathbf{x}^T,$$

where  $\text{inv}(T)$  is the number of attacking inversions in  $T$  and  $\mathbf{x}^T = \prod_{a \in \nu} x_{T(a)}$ .

$T =$

				5	6
				1	1
2	4				
	3	5			

inversion



# LLT Polynomials

- A *semistandard tableau* on  $\nu$  is a map  $T: \nu \rightarrow \mathbb{Z}_+$  which restricts to a semistandard tableau on each  $\nu_{(i)}$ .
- An *attacking inversion* in  $T$  is an attacking pair  $(a, b)$  such that  $T(a) > T(b)$ .

The *LLT polynomial* indexed by a tuple of skew shapes  $\nu$  is

$$\mathcal{G}_\nu(\mathbf{x}; q) = \sum_{T \in \text{SSYT}(\nu)} q^{\text{inv}(T)} \mathbf{x}^T,$$

where  $\text{inv}(T)$  is the number of attacking inversions in  $T$  and  $\mathbf{x}^T = \prod_{a \in \nu} x_{T(a)}$ .

$T =$

				5	6	
				1	1	
2	4					
	3	5				

non-inversion

# LLT Polynomials

- A *semistandard tableau* on  $\nu$  is a map  $T: \nu \rightarrow \mathbb{Z}_+$  which restricts to a semistandard tableau on each  $\nu_{(i)}$ .
- An *attacking inversion* in  $T$  is an attacking pair  $(a, b)$  such that  $T(a) > T(b)$ .

The *LLT polynomial* indexed by a tuple of skew shapes  $\nu$  is

$$\mathcal{G}_\nu(\mathbf{x}; q) = \sum_{T \in \text{SSYT}(\nu)} q^{\text{inv}(T)} \mathbf{x}^T,$$

where  $\text{inv}(T)$  is the number of attacking inversions in  $T$  and  $\mathbf{x}^T = \prod_{a \in \nu} x_{T(a)}$ .

$T =$

				5	6	
				1	1	
2	4					
	3	5				

non-inversion

# LLT Polynomials

- A *semistandard tableau* on  $\nu$  is a map  $T: \nu \rightarrow \mathbb{Z}_+$  which restricts to a semistandard tableau on each  $\nu_{(i)}$ .
- An *attacking inversion* in  $T$  is an attacking pair  $(a, b)$  such that  $T(a) > T(b)$ .

The *LLT polynomial* indexed by a tuple of skew shapes  $\nu$  is

$$\mathcal{G}_\nu(\mathbf{x}; q) = \sum_{T \in \text{SSYT}(\nu)} q^{\text{inv}(T)} \mathbf{x}^T,$$

where  $\text{inv}(T)$  is the number of attacking inversions in  $T$  and  $\mathbf{x}^T = \prod_{a \in \nu} x_{T(a)}$ .

$T =$

				5	6	
				1	1	
2	4					
	3	5				

inversion

# LLT Polynomials

- A *semistandard tableau* on  $\nu$  is a map  $T: \nu \rightarrow \mathbb{Z}_+$  which restricts to a semistandard tableau on each  $\nu_{(i)}$ .
- An *attacking inversion* in  $T$  is an attacking pair  $(a, b)$  such that  $T(a) > T(b)$ .

The *LLT polynomial* indexed by a tuple of skew shapes  $\nu$  is

$$\mathcal{G}_\nu(\mathbf{x}; q) = \sum_{T \in \text{SSYT}(\nu)} q^{\text{inv}(T)} \mathbf{x}^T,$$

where  $\text{inv}(T)$  is the number of attacking inversions in  $T$  and  $\mathbf{x}^T = \prod_{a \in \nu} x_{T(a)}$ .

$T =$

				5	6	
				1	1	
2	4					
	3	5				

inversion

$$\text{inv}(T) = 4, \quad \mathbf{x}^T = x_1^2 x_2 x_3 x_4 x_5^2 x_6$$

## Definition

The *Catalanimal* indexed by  $R_q, R_t, R_{qt} \subseteq R_+$  and  $\lambda \in \mathbb{Z}^n$  is

## Definition

The *Catalanimal* indexed by  $R_q, R_t, R_{qt} \subseteq R_+$  and  $\lambda \in \mathbb{Z}^n$  is

$$H(R_q, R_t, R_{qt}, \lambda) = \sigma \left( \frac{z^\lambda \prod_{\alpha \in R_{qt}} (1 - qtz^\alpha)}{\prod_{\alpha \in R_q} (1 - qz^\alpha) \prod_{\alpha \in R_t} (1 - tz^\alpha)} \right).$$

## Definition

The *Catalanimal* indexed by  $R_q, R_t, R_{qt} \subseteq R_+$  and  $\lambda \in \mathbb{Z}^n$  is

$$H(R_q, R_t, R_{qt}, \lambda) = \sigma \left( \frac{z^\lambda \prod_{\alpha \in R_{qt}} (1 - qt z^\alpha)}{\prod_{\alpha \in R_q} (1 - q z^\alpha) \prod_{\alpha \in R_t} (1 - t z^\alpha)} \right).$$

With  $n = 3$ ,

$$\begin{aligned} H(R_+, R_+, \{\alpha_{13}\}, (111)) &= \sigma \left( \frac{z^{111} (1 - qt z_1/z_3)}{\prod_{1 \leq i < j \leq 3} (1 - q z_i/z_j) (1 - t z_i/z_j)} \right) \\ &= s_{111} + (q + t + q^2 + qt + t^2) s_{21} + (qt + q^3 + q^2 t + qt^2 + t^3) s_3 \\ &= \omega \nabla e_3. \end{aligned}$$

# LLT Catalanimals

For a tuple of skew shapes  $\nu$ , the *LLT Catalanimal*  $H_\nu = H(R_q, R_t, R_{qt}, \lambda)$  is determined by

- $R_+ \supseteq R_q \supseteq R_t \supseteq R_{qt}$ ,



# LLT Catalananimals

For a tuple of skew shapes  $\nu$ , the *LLT Catalananimal*  $H_\nu = H(R_q, R_t, R_{qt}, \lambda)$  is determined by

- $R_+ \supseteq R_q \supseteq R_t \supseteq R_{qt}$ ,
- $R_+ \setminus R_q$  = pairs of boxes in the same diagonal,
- $R_q \setminus R_t$  = the attacking pairs,
- $R_t \setminus R_{qt}$  = pairs going between adjacent diagonals,

# LLT Catalananimals

For a tuple of skew shapes  $\nu$ , the *LLT Catalananimal*  $H_\nu = H(R_q, R_t, R_{qt}, \lambda)$  is determined by


- $R_+ \supseteq R_q \supseteq R_t \supseteq R_{qt}$ ,
- $R_+ \setminus R_q$  = pairs of boxes in the same diagonal,
- $R_q \setminus R_t$  = the attacking pairs,
- $R_t \setminus R_{qt}$  = pairs going between adjacent diagonals,
- $\lambda$ : fill each diagonal  $D$  of  $\nu$  with  $1 + \chi(D \text{ contains a row start}) - \chi(D \text{ contains a row end})$ .  
Listing this filling in reading order gives  $\lambda$ .

# LLT Catalanimals


- $R_+ \setminus R_q =$  pairs of boxes in the same diagonal,
- $R_q \setminus R_t =$  the attacking pairs,
- $R_t \setminus R_{qt} =$  pairs going between adjacent diagonals,
- $R_{qt} =$  all other pairs,

$\lambda$ : fill each diagonal  $D$  of  $\nu$  with

$1 + \chi(D \text{ contains a row start}) - \chi(D \text{ contains a row end})$ .



$\nu$




# LLT Catalanimals


- $R_+ \setminus R_q =$  pairs of boxes in the same diagonal,
- $R_q \setminus R_t =$  the attacking pairs,
- $R_t \setminus R_{qt} =$  pairs going between adjacent diagonals,
- $R_{qt} =$  all other pairs,

$\lambda$ : fill each diagonal  $D$  of  $\nu$  with

$1 + \chi(D \text{ contains a row start}) - \chi(D \text{ contains a row end})$ .



$\lambda$ , as a filling of  $\nu$



## Theorem (Blasiak-Haiman-Morse-Pun-S.)

Let  $\nu$  be a tuple of skew shapes and let  $H_\nu = H(R_q, R_t, R_{qt}, \lambda)$  be the associated LLT Catalanimal. Then

$$\begin{aligned}\nabla \mathcal{G}_\nu(X; q) &= c_\nu \omega H_\nu \\ &= c_\nu \omega \sigma \left( \frac{z^\lambda \prod_{\alpha \in R_{qt}} (1 - qt z^\alpha)}{\prod_{\alpha \in R_q} (1 - q z^\alpha) \prod_{\alpha \in R_t} (1 - t z^\alpha)} \right)\end{aligned}$$

for some  $c_\nu \in \pm q^{\mathbb{Z}} t^{\mathbb{Z}}$ .

# Haglund-Haiman-Loehr formula

Theorem (Haglund-Haiman-Loehr, 2005)










$$\tilde{H}_\mu(X; q, t) = \sum_D \left( \prod_{u \in D} q^{-\text{arm}(u)} t^{\text{leg}(u)+1} \right) \mathcal{G}_{\nu(\mu, D)}(X; q),$$

where

- the sum runs over all subsets  $D \subseteq \{(i, j) \in \mu \mid j > 1\}$ , and
- $\nu(\mu, D) = (\nu^{(1)}, \dots, \nu^{(k)})$  where  $k = \mu_1$  is the number of columns of  $\mu$ , and  $\nu^{(i)}$  is a ribbon of size  $\mu_i^*$ , i.e., box contents  $\{-1, -2, \dots, -\mu_i^*\}$ , and descent set  $\text{Des}(\nu^{(i)}) = \{-j \mid (i, j) \in D\}$ .

# Haglund-Haiman-Loehr formula example

$$\tilde{H}_\mu(X; q, t) = \sum_D \left( \prod_{u \in D} q^{-\text{arm}(u)} t^{\text{leg}(u)+1} \right) \mathcal{G}_{\nu(\mu, D)}(X; q)$$



## Putting it all together

- Take HHL formula  $\tilde{H}_\mu = \sum_D a_{\mu,D} \mathcal{G}_{\nu(\mu,D)}$  and apply  $\omega \nabla$ .



# Putting it all together

- Take HHL formula  $\tilde{H}_\mu = \sum_D a_{\mu,D} \mathcal{G}_{\nu(\mu,D)}$  and apply  $\omega \nabla$ .
- By construction, all the LLT Catalan animals  $H_{\nu(\mu,D)}$  appearing on the RHS will have the same root ideal data  $(R_q, R_t, R_{qt})$ .

# Putting it all together

- Take HHL formula  $\tilde{H}_\mu = \sum_D a_{\mu,D} \mathcal{G}_{\nu(\mu,D)}$  and apply  $\omega \nabla$ .
- By construction, all the LLT Catalan animals  $H_{\nu(\mu,D)}$  appearing on the RHS will have the same root ideal data  $(R_q, R_t, R_{qt})$ .
- Collect terms to get  $\prod_{\alpha_{ij} \in R_\mu \setminus \hat{R}_\mu} (1 - q^{\text{arm}(b_i)+1} t^{-\text{leg}(b_i)} z_i/z_j)$  factor.

$$\tilde{H}_\mu = \omega \sigma \left( z_1 \cdots z_n \frac{\prod_{\alpha_{ij} \in R_\mu \setminus \hat{R}_\mu} (1 - q^{\text{arm}(b_i)+1} t^{-\text{leg}(b_i)} z_i/z_j) \prod_{\alpha \in \hat{R}_\mu} (1 - qt z^\alpha)}{\prod_{\alpha \in R_+} (1 - qz^\alpha) \prod_{\alpha \in R_\mu} (1 - tz^\alpha)} \right).$$

# Outline

- ① Background on symmetric functions and Macdonald polynomials
- ② A new formula for Macdonald polynomials
- ③ **LLT polynomials in the elliptic Hall algebra**

# Elliptic Hall Algebra

Burban and Schiffmann studied a subalgebra  $\mathcal{E}$  of the Hall algebra of coherent sheaves on an elliptic curve over  $\mathbb{F}_p$ .

The *elliptic Hall algebra*  $\mathcal{E}$  is generated by subalgebras  $\Lambda(X^{a,b})$  isomorphic to the ring of symmetric functions  $\Lambda$  over  $\mathbb{k} = \mathbb{Q}(q, t)$ , one for each coprime pair  $(a, b) \in \mathbb{Z}^2$ , along with an additional central subalgebra.

# Shuffle algebra

Define a linear map

$$\sigma_{\Gamma}: \bigoplus_n \mathbb{k}(z_1, \dots, z_n) \rightarrow \bigoplus_n \mathbb{k}(z_1, \dots, z_n)^{S_n}$$

whose graded components  $\sigma_{\Gamma}^n$  are given by

# Shuffle algebra

Define a linear map

$$\sigma_{\Gamma}: \bigoplus_n \mathbb{k}(z_1, \dots, z_n) \rightarrow \bigoplus_n \mathbb{k}(z_1, \dots, z_n)^{S_n}$$

whose graded components  $\sigma_{\Gamma}^n$  are given by

$$\sigma_{\Gamma}^n: \mathbb{k}(z_1, \dots, z_n) \rightarrow \mathbb{k}(z_1, \dots, z_n)^{S_n}$$

$$\sigma_{\Gamma}^n(f) = \sum_{w \in S_n} w(f(z_1, \dots, z_n) \prod_{1 \leq i < j \leq n} \Gamma(z_i, z_j)),$$

$$\text{where } \Gamma(z_i, z_j) = \frac{1 - qtz_i/z_j}{(1 - z_j/z_i)(1 - qz_i/z_j)(1 - tz_i/z_j)}$$

# Shuffle algebra

Define a linear map

$$\sigma_{\Gamma}: \bigoplus_n \mathbb{k}(z_1, \dots, z_n) \rightarrow \bigoplus_n \mathbb{k}(z_1, \dots, z_n)^{S_n}$$

whose graded components  $\sigma_{\Gamma}^n$  are given by

$$\sigma_{\Gamma}^n: \mathbb{k}(z_1, \dots, z_n) \rightarrow \mathbb{k}(z_1, \dots, z_n)^{S_n}$$

$$\sigma_{\Gamma}^n(f) = \sum_{w \in S_n} w(f(z_1, \dots, z_n) \prod_{1 \leq i < j \leq n} \Gamma(z_i, z_j)),$$

$$\text{where } \Gamma(z_i, z_j) = \frac{1 - qtz_i/z_j}{(1 - z_j/z_i)(1 - qz_i/z_j)(1 - tz_i/z_j)}$$

The *shuffle algebra*  $\mathcal{S}_{\Gamma}$  is the image of  $\bigoplus_n \mathbb{k}[z_1^{\pm 1}, \dots, z_n^{\pm 1}]$  under the map  $\sigma_{\Gamma}$ , equipped with a variant of the concatenation product.

# Shuffle algebra

Define a linear map

$$\sigma_{\Gamma}: \bigoplus_n \mathbb{k}(z_1, \dots, z_n) \rightarrow \bigoplus_n \mathbb{k}(z_1, \dots, z_n)^{S_n}$$

whose graded components  $\sigma_{\Gamma}^n$  are given by

$$\sigma_{\Gamma}^n: \mathbb{k}(z_1, \dots, z_n) \rightarrow \mathbb{k}(z_1, \dots, z_n)^{S_n}$$

$$\sigma_{\Gamma}^n(f) = \sum_{w \in S_n} w(f(z_1, \dots, z_n) \prod_{1 \leq i < j \leq n} \Gamma(z_i, z_j)),$$

$$\text{where } \Gamma(z_i, z_j) = \frac{1 - qtz_i/z_j}{(1 - z_j/z_i)(1 - qz_i/z_j)(1 - tz_i/z_j)}$$

The *shuffle algebra*  $\mathcal{S}_{\Gamma}$  is the image of  $\bigoplus_n \mathbb{k}[z_1^{\pm 1}, \dots, z_n^{\pm 1}]$  under the map  $\sigma_{\Gamma}$ , equipped with a variant of the concatenation product.

Nice fact (up to some modifications of definitions)

Some Catalan animals are elements in  $\mathcal{S}_{\Gamma}$ . (“Tame Catalan animals”)



# Shuffle to elliptic Hall isomorphism

- The *right half-plane subalgebra*  $\mathcal{E}^+ \subseteq \mathcal{E}$  is generated by  $\Lambda(X^{a,b})$  for  $a > 0$ .

## Shuffle to elliptic Hall isomorphism

- The *right half-plane subalgebra*  $\mathcal{E}^+ \subseteq \mathcal{E}$  is generated by  $\Lambda(X^{a,b})$  for  $a > 0$ .
- $\mathcal{S}_\Gamma = \sigma_\Gamma \left( \bigoplus_n \mathbb{k}[z_1^{\pm 1}, \dots, z_n^{\pm 1}] \right)$  ( $\Gamma$ -symmetrized Laurent polynomials).

# Shuffle to elliptic Hall isomorphism

- The *right half-plane subalgebra*  $\mathcal{E}^+ \subseteq \mathcal{E}$  is generated by  $\Lambda(X^{a,b})$  for  $a > 0$ .
- $\mathcal{S}_\Gamma = \sigma_\Gamma \left( \bigoplus_n \mathbb{k}[z_1^{\pm 1}, \dots, z_n^{\pm 1}] \right)$  ( $\Gamma$ -symmetrized Laurent polynomials).

## Theorem (Schiffmann-Vasserot)

There is an algebra isomorphism  $\psi: \mathcal{S}_\Gamma \rightarrow \mathcal{E}^+$ .

## Elliptic Hall algebra action

Schiffmann-Vasserot and Feigin-Tsymbaliuk constructed an action of  $\mathcal{E}$  on  $\Lambda$ , where  $f(X^{0,1})$  acts by multiplication by  $f(X)$ .

# Elliptic Hall algebra action

Schiffmann-Vasserot and Feigin-Tsymbaliuk constructed an action of  $\mathcal{E}$  on  $\Lambda$ , where  $f(X^{0,1})$  acts by multiplication by  $f(X)$ .

## Proposition

*Conjugation by  $\nabla$  provides a symmetry of the action of  $\mathcal{E}$  on  $\Lambda$ ,*

$$\nabla f(X^{a,b}) \nabla^{-1} = f(X^{a+b,b}).$$

# Elliptic Hall algebra action

Schiffmann-Vasserot and Feigin-Tsymbaliuk constructed an action of  $\mathcal{E}$  on  $\Lambda$ , where  $f(X^{0,1})$  acts by multiplication by  $f(X)$ .

## Proposition

*Conjugation by  $\nabla$  provides a symmetry of the action of  $\mathcal{E}$  on  $\Lambda$ ,*

$$\nabla f(X^{a,b}) \nabla^{-1} = f(X^{a+b,b}).$$

## Corollary

$$f(X^{1,1}) \cdot 1 = \nabla f(X^{0,1}) \nabla^{-1} \cdot 1 = \nabla f.$$

# Elliptic Hall algebra action

Schiffmann-Vasserot and Feigin-Tsybaliuk constructed an action of  $\mathcal{E}$  on  $\Lambda$ , where  $f(X^{0,1})$  acts by multiplication by  $f(X)$ .

## Proposition

*Conjugation by  $\nabla$  provides a symmetry of the action of  $\mathcal{E}$  on  $\Lambda$ ,*

$$\nabla f(X^{a,b}) \nabla^{-1} = f(X^{a+b,b}).$$

## Corollary

$$f(X^{1,1}) \cdot 1 = \nabla f(X^{0,1}) \nabla^{-1} \cdot 1 = \nabla f.$$

## Theorem (Blasiak-Haiman-Morse-Pun-S.)

*Let  $H$  be a Catalan animal such that  $\psi(H) = f(X^{1,1})$ . Then*

$$\nabla f = \omega H.$$

# Shuffle to elliptic Hall summary

$$\begin{array}{ccc}
 & \mathcal{E} \curvearrowright \Lambda & f(X^{1,1}) \cdot 1 = \nabla f \\
 & \uparrow & \\
 \bigoplus_{\substack{a>0 \\ b \in \mathbb{Z} \\ (a,b)=1}} \Lambda(X^{a,b}) & \stackrel{\cong}{\underset{\text{v.sp.}}{\longrightarrow}} & \mathcal{E}^+ \\
 & \uparrow \psi \cong & \\
 \sigma_\Gamma \left( \bigoplus_n \mathbb{k}[z_1^{\pm 1}, \dots, z_n^{\pm 1}] \right) & \stackrel{\cong}{\underset{\text{v.sp.}}{\longrightarrow}} \mathcal{S}_\Gamma \ni H & \text{“tame” Catalanimal}
 \end{array}$$

Theorem (Blasiak-Haiman-Morse-Pun-S.)

$$\psi(H) = f(X^{1,1}) \implies f(X^{1,1}) \cdot 1 = \nabla f = \omega H.$$



# Proof of $\nabla \mathcal{G}_\nu$ formula

- ① LLT Catalananimals  $H_\nu$  are tame.
- ② LLT Catalananimals lie in  $\psi^{-1}(\Lambda(X^{1,1}))$ .
- ③ Describe coproduct  $\Delta$  on  $\mathcal{E}$  explicitly on tame Catalananimals and show  $\Delta H_\nu$  matches  $\Delta \mathcal{G}_\nu$ .
- ④ Conclude  $\psi(H_\nu) = c_\nu^{-1} \mathcal{G}_\nu(X^{1,1}) \in \mathcal{E}$ .
- ⑤ Apply previous theorem to conclude  $\nabla \mathcal{G}_\nu = c_\nu \omega H_\nu$

## A positivity conjecture

What can this formula tell us that other formulas for Macdonald polynomials do not?

# A positivity conjecture

What can this formula tell us that other formulas for Macdonald polynomials do not?

$$\tilde{H}_\mu^{(s)} := \omega \sigma \left( (z_1 \cdots z_n)^s \frac{\prod_{\alpha_{ij} \in R_\mu \setminus \hat{R}_\mu} (1 - q^{\text{arm}(b_i)+1} t^{-\text{leg}(b_i)} z_i / z_j) \prod_{\alpha \in \hat{R}_\mu} (1 - qt z^\alpha)}{\prod_{\alpha \in R_+} (1 - qz^\alpha) \prod_{\alpha \in R_\mu} (1 - tz^\alpha)} \right)$$

## Conjecture (Blasiak-Haiman-Morse-Pun-S.)

For any partition  $\mu$  and positive integer  $s$ , the symmetric function  $\tilde{H}_\mu^{(s)}$  is Schur positive. That is, the coefficients in

$$\tilde{H}_\mu^{(s)} = \sum_{\nu} K_{\nu, \mu}^{(s)}(q, t) s_\nu(X)$$

satisfy  $K_{\nu, \mu}^{(s)}(q, t) \in \mathbb{N}[q, t]$ .

# Thank you!

Blasiak, Jonah, Mark Haiman, Jennifer Morse, Anna Pun, and George H. Seelinger. 2021. *LLT Polynomials in the Schiffmann Algebra*, arXiv e-prints, arXiv:2112.07063.

———. 2023. *A Raising Operator Formula for Macdonald Polynomials*, arXiv e-prints, arXiv:2307.06517.

Burban, Igor and Olivier Schiffmann. 2012. *On the Hall algebra of an elliptic curve, I*, Duke Math. J. **161**, no. 7, 1171–1231, DOI 10.1215/00127094-1593263. MR2922373

Feigin, B. L. and Tsymbaliuk, A. I. 2011. *Equivariant K-theory of Hilbert Schemes via Shuffle Algebra*, Kyoto J. Math. **51**, no. 4, 831–854.

Garsia, Adriano M. and Mark Haiman. 1993. *A graded representation model for Macdonald's polynomials*, Proc. Nat. Acad. Sci. U.S.A. **90**, no. 8, 3607–3610, DOI 10.1073/pnas.90.8.3607. MR1214091

Haglund, J., M. Haiman, and N. Loehr. 2005. *A Combinatorial Formula for Macdonald Polynomials* **18**, no. 3, 735–761 (electronic).

Haiman, Mark. 2001. *Hilbert schemes, polygraphs and the Macdonald positivity conjecture*, J. Amer. Math. Soc. **14**, no. 4, 941–1006, DOI 10.1090/S0894-0347-01-00373-3. MR1839919

Lascoux, Alain, Bernard Leclerc, and Jean-Yves Thibon. 1995. *Ribbon tableaux, Hall-Littlewood functions and unipotent varieties*, Sémin. Lothar. Combin. **34**, Art. B34g, approx. 23. MR1399754

Negut, Andrei. 2014. *The shuffle algebra revisited*, Int. Math. Res. Not. IMRN **22**, 6242–6275, DOI 10.1093/imrn/rnt156. MR3283004

Schiffmann, Olivier and Vasserot, Eric. 2013. *The Elliptic Hall Algebra and the K-theory of the Hilbert Scheme of  $A_2$* , Duke Mathematical Journal **162**, no. 2, 279–366, DOI 10.1215/00127094-1961849.

Shimozono, Mark and Jerzy Weyman. 2000. *Graded Characters of Modules Supported in the Closure of a Nilpotent Conjugacy Class*, European Journal of Combinatorics **21**, no. 2, 257–288, DOI 10.1006/eujc.1999.0344.

Weyman, J. 1989. *The Equations of Conjugacy Classes of Nilpotent Matrices*, Inventiones mathematicae **98**, no. 2, 229–245, DOI 10.1007/BF01388851.

# Catalan animals in the shuffle algebra

For  $\lambda \in \mathbb{Z}^n$ ,

$$\begin{aligned}\sigma_{\Gamma}^n(\mathbf{z}^{\lambda}) &= \sum_{w \in S_n} w \left( \frac{\mathbf{z}^{\lambda} \prod_{\alpha \in R_+} (1 - qt\mathbf{z}^{\alpha})}{\prod_{\alpha \in R_+} ((1 - \mathbf{z}^{-\alpha})(1 - q\mathbf{z}^{\alpha})(1 - t\mathbf{z}^{\alpha}))} \right) \\ &= H(R_+, R_+, R_+, \lambda) \in \mathcal{S}_{\Gamma}.\end{aligned}$$

- Technicality: we have redefined

$\sigma(\mathbf{z}^{\gamma}) = \sum_{w \in S_n} \left( \frac{\mathbf{z}^{\gamma}}{\prod_{\alpha \in R_+} (1 - \mathbf{z}^{-\alpha})} \right) = \chi_{\gamma}$ , the irreducible  $\mathrm{GL}_n$  character.

- Let  $\mathrm{pol}_X$  send  $\chi_{\lambda} \mapsto s_{\lambda}$  if  $\lambda_n \geq 0$ , otherwise  $\chi_{\lambda} \mapsto 0$ .
- The  $\sigma$  from before is given by  $\sigma_{\mathrm{old}} = \mathrm{pol}_X \sigma_{\mathrm{new}}$ .

# Catalanimals in the Shuffle algebra

$\sigma_{\Gamma}^n(f)$  can lie in  $\mathcal{S}_{\Gamma}$  even when  $f$  is not a Laurent polynomial.

## Theorem (Negut)

*The following family of Catalanimals lie in the shuffle algebra:*

$$\sigma_{\Gamma}^n\left(\frac{z^{\lambda}}{\prod_{i=1}^{n-1}(1 - qtz_i/z_{i+1})}\right) = H(R_+, R_+, R'_+, \lambda) \in \mathcal{S}_{\Gamma},$$

where  $R'_+ = \{\alpha_{ij} \in R_+ \mid i + 1 < j\}$ .

# The wheel condition

- A symmetric Laurent polynomial  $g(\mathbf{z})$  satisfies the *wheel condition* if it vanishes whenever any three of the variables  $z_i, z_j, z_k$  are in the ratio  $(z_i : z_j : z_k) = (1 : q : qt) = (1 : t : qt)$ .
- Let  $\mathcal{S}_{\check{\Gamma}} \cong \mathcal{S}_{\Gamma}$  for  
$$\check{\Gamma}(z_i, z_j) = (1 - z_i/z_j)(1 - qz_j/z_i)(1 - tz_j/z_i)(1 - qtz_i/z_j).$$

## Theorem (Negut)




A symmetric Laurent polynomial  $g(z_1, \dots, z_n)$  belongs to  $\mathcal{S}_{\check{\Gamma}}$  if and only if it satisfies the wheel condition and vanishes whenever  $z_i = z_j$  for  $i \neq j$ .

# The wheel condition and tame Catalananimals

A Catalananimal  $H(R_q, R_t, R_{qt}, \lambda)$  is *tame* if

$$R_q + R_t \subseteq R_{qt},$$

where  $R_q + R_t = \{\alpha + \beta \mid \alpha \in R_q, \beta \in R_t\}$ .



The Catalananimals  $H(R_+, R_+, R'_+, \lambda)$  and the LLT Catalananimals are tame.

Using Negut's theorem, we show: Tame Catalananimals belong to the shuffle algebra  $\mathcal{S}_\Gamma$ .