

Flagged LLT polynomials and their applications

George H. Seelinger

ghseeli@umich.edu

joint work with Jonah Blasiak, Mark Haiman, Jennifer Morse, and Anna Pun

Center for Mathematics at the University of Notre Dame

Algebraic combinatorics and applications

6 July 2026

Symmetric polynomials and functions

- Symmetric polynomials $\mathbb{Q}[x_1, \dots, x_n]^{S_n}$
- Generators

$$e_r(x_1, \dots, x_n) = \sum_{1 \leq i_1 < i_2 < \dots < i_r \leq n} x_{i_1} \cdots x_{i_r}$$

$$e_1(x_1, x_2, x_3) = x_1 + x_2 + x_3$$

$$e_2(x_1, x_2, x_3) = x_1x_2 + x_1x_3 + x_2x_3$$

$$e_3(x_1, x_2, x_3) = x_1x_2x_3$$

Symmetric polynomials and functions

- Symmetric polynomials $\mathbb{Q}[x_1, \dots, x_n]^{S_n}$
- Generators

$$e_r(x_1, \dots, x_n) = \sum_{1 \leq i_1 < i_2 < \dots < i_r \leq n} x_{i_1} \cdots x_{i_r}$$

$$e_1(x_1, x_2, x_3) = x_1 + x_2 + x_3$$

$$e_2(x_1, x_2, x_3) = x_1x_2 + x_1x_3 + x_2x_3$$

$$e_3(x_1, x_2, x_3) = x_1x_2x_3$$

- $\Lambda = \mathbb{Q}[e_1, e_2, \dots]$

Symmetric polynomials and functions

- Symmetric polynomials $\mathbb{Q}[x_1, \dots, x_n]^{S_n}$
- Generators

$$e_r(x_1, \dots, x_n) = \sum_{1 \leq i_1 < i_2 < \dots < i_r \leq n} x_{i_1} \cdots x_{i_r}$$

$$e_1(x_1, x_2, x_3) = x_1 + x_2 + x_3$$

$$e_2(x_1, x_2, x_3) = x_1x_2 + x_1x_3 + x_2x_3$$

$$e_3(x_1, x_2, x_3) = x_1x_2x_3$$

- $\Lambda = \mathbb{Q}[e_1, e_2, \dots]$
- Degree d elements of $\Lambda = \mathbb{Q}$ -vector space of dimension $\dots$?

Symmetric polynomials and functions

- Symmetric polynomials $\mathbb{Q}[x_1, \dots, x_n]^{S_n}$
- Generators

$$e_r(x_1, \dots, x_n) = \sum_{1 \leq i_1 < i_2 < \dots < i_r \leq n} x_{i_1} \cdots x_{i_r}$$

$$e_1(x_1, x_2, x_3) = x_1 + x_2 + x_3$$

$$e_2(x_1, x_2, x_3) = x_1x_2 + x_1x_3 + x_2x_3$$

$$e_3(x_1, x_2, x_3) = x_1x_2x_3$$

- $\Lambda = \mathbb{Q}[e_1, e_2, \dots]$
- Degree d elements of $\Lambda = \mathbb{Q}$ -vector space of dimension $\dots$?
- Integer partitions of d .

Partitions

Definition

$n \in \mathbb{Z}_{>0}$, a *partition of n* is $\lambda = (\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_\ell > 0)$ such that $\lambda_1 + \lambda_2 + \cdots + \lambda_\ell = n$.

5 →

--	--	--	--	--

$4 + 1 \rightarrow$ 

$$3 + 2 \rightarrow \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}$$

$3 + 1 + 1 \rightarrow$ 

$$2 + 2 + 1 \rightarrow \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}$$

$$2 + 1 + 1 + 1 \rightarrow \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}$$

$$1 + 1 + 1 + 1 + 1 \rightarrow$$


Young Tableaux

Definition


Filling of partition diagram of λ with numbers such that

- 1 strictly increasing up columns
- 2 weakly increasing along rows

Collection is called $\text{SSYT}(\lambda)$.



For $\lambda = (2, 1)$,




Schur polynomials

Associate a polynomial to $\text{SSYT}(\lambda)$.

2		3		3		2		3		3		2		3	
1	1	1	1	2	2	1	2	1	3	2	3	1	3	1	2

Schur polynomials

Associate a polynomial to $\text{SSYT}(\lambda)$.



Schur polynomials

Associate a polynomial to $\text{SSYT}(\lambda)$.

$$\begin{array}{cccccccc} \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 2 & 2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 2 & 3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 2 \\ \hline \end{array} \\ \rightarrow & \begin{array}{|c|c|} \hline x_2 & \\ \hline x_1 & x_1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline x_3 & \\ \hline x_1 & x_1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline x_3 & \\ \hline x_2 & x_2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline x_2 & \\ \hline x_1 & x_2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline x_3 & \\ \hline x_1 & x_3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline x_3 & \\ \hline x_2 & x_3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline x_2 & \\ \hline x_1 & x_3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline x_3 & \\ \hline x_1 & x_2 \\ \hline \end{array} \end{array}$$

$$s_{(2,1)}(x_1, x_2, x_3) = x_1^2 x_2 + x_1^2 x_3 + x_2^2 x_3 + x_1 x_2^2 + x_1 x_3^2 + x_2 x_3^2 + 2x_1 x_2 x_3$$

Schur polynomials

Associate a polynomial to $\text{SSYT}(\lambda)$.

$$\begin{array}{cccccccc} \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 2 & 2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 2 & 3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 2 \\ \hline \end{array} \\ \rightarrow & \begin{array}{|c|c|} \hline x_2 & \\ \hline x_1 & x_1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline x_3 & \\ \hline x_1 & x_1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline x_3 & \\ \hline x_2 & x_2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline x_2 & \\ \hline x_1 & x_2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline x_3 & \\ \hline x_1 & x_3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline x_3 & \\ \hline x_2 & x_3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline x_2 & \\ \hline x_1 & x_3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline x_3 & \\ \hline x_1 & x_2 \\ \hline \end{array} \end{array}$$

$$s_{(2,1)}(x_1, x_2, x_3) = x_1^2 x_2 + x_1^2 x_3 + x_2^2 x_3 + x_1 x_2^2 + x_1 x_3^2 + x_2 x_3^2 + 2x_1 x_2 x_3$$

Definition

For λ a partition

$$s_\lambda = \sum_{T \in \text{SSYT}(\lambda)} x^T \text{ for } x^T = \prod_{i \in T} x_i$$

Schur polynomials

Associate a polynomial to $\text{SSYT}(\lambda)$.

$$\begin{array}{cccccccc} \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 2 & 2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 2 & 3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 2 \\ \hline \end{array} \\ \rightarrow & \begin{array}{|c|c|} \hline x_2 & \\ \hline x_1 & x_1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline x_3 & \\ \hline x_1 & x_1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline x_3 & \\ \hline x_2 & x_2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline x_2 & \\ \hline x_1 & x_2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline x_3 & \\ \hline x_1 & x_3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline x_3 & \\ \hline x_2 & x_3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline x_2 & \\ \hline x_1 & x_3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline x_3 & \\ \hline x_1 & x_2 \\ \hline \end{array} \end{array}$$

$$s_{(2,1)}(x_1, x_2, x_3) = x_1^2 x_2 + x_1^2 x_3 + x_2^2 x_3 + x_1 x_2^2 + x_1 x_3^2 + x_2 x_3^2 + 2x_1 x_2 x_3$$

Definition

For λ a partition

$$s_\lambda = \sum_{T \in \text{SSYT}(\lambda)} x^T \text{ for } x^T = \prod_{i \in T} x_i$$

- s_λ is a symmetric function.

Schur polynomials

Associate a polynomial to $\text{SSYT}(\lambda)$.

$$\begin{array}{cccccccc} \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 2 & 2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 2 & 3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 2 \\ \hline \end{array} \\ \rightarrow & \begin{array}{|c|c|} \hline x_2 & \\ \hline x_1 & x_1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline x_3 & \\ \hline x_1 & x_1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline x_3 & \\ \hline x_2 & x_2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline x_2 & \\ \hline x_1 & x_2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline x_3 & \\ \hline x_1 & x_3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline x_3 & \\ \hline x_2 & x_3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline x_2 & \\ \hline x_1 & x_3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline x_3 & \\ \hline x_1 & x_2 \\ \hline \end{array} \end{array}$$

$$s_{(2,1)}(x_1, x_2, x_3) = x_1^2 x_2 + x_1^2 x_3 + x_2^2 x_3 + x_1 x_2^2 + x_1 x_3^2 + x_2 x_3^2 + 2x_1 x_2 x_3$$

Definition

For λ a partition

$$s_\lambda = \sum_{T \in \text{SSYT}(\lambda)} x^T \text{ for } x^T = \prod_{i \in T} x_i$$

- s_λ is a symmetric function.
- $\{s_\lambda\}_\lambda$ forms a basis for Λ .

Representation theory and Schur functions

Irreducible representations of S_n are **also** labeled by partitions of n .

Representation theory and Schur functions

Irreducible representations of S_n are **also** labeled by partitions of n .

$$V_{\square\square\square} = \text{trivial}, \quad V_{\begin{smallmatrix}\square & \\ \square & \end{smallmatrix}} \cong \text{span}\{e_1 - e_2, e_2 - e_3\}, \quad V_{\begin{smallmatrix}\square \\ \square \\ \square \end{smallmatrix}} = \text{sign}$$

Representation theory and Schur functions

Irreducible representations of S_n are **also** labeled by partitions of n .

$$V_{\square\square\square} = \text{trivial}, \quad V_{\begin{smallmatrix}\square & \\ \square & \end{smallmatrix}} \cong \text{span}\{e_1 - e_2, e_2 - e_3\}, \quad V_{\begin{smallmatrix}\square \\ \square \\ \square \end{smallmatrix}} = \text{sign}$$

Frobenius characterstic, $\text{Frob}: \text{Rep}(S_n) \rightarrow \Lambda$, such that

Representation theory and Schur functions

Irreducible representations of S_n are **also** labeled by partitions of n .

$$V_{\square\square\square} = \text{trivial}, \quad V_{\begin{smallmatrix}\square & \\ \square & \end{smallmatrix}} \cong \text{span}\{e_1 - e_2, e_2 - e_3\}, \quad V_{\begin{smallmatrix}\square \\ \square \\ \square \end{smallmatrix}} = \text{sign}$$

Frobenius characterstic, $\text{Frob}: \text{Rep}(S_n) \rightarrow \Lambda$, such that

- Irreducible S_n -representation V_λ has $\text{Frob}(V_\lambda) = s_\lambda$

Representation theory and Schur functions

Irreducible representations of S_n are **also** labeled by partitions of n .

$$V_{\square\square\square} = \text{trivial}, \quad V_{\begin{smallmatrix} \square & \\ \square & \end{smallmatrix}} \cong \text{span}\{e_1 - e_2, e_2 - e_3\}, \quad V_{\begin{smallmatrix} \square \\ \square \\ \square \end{smallmatrix}} = \text{sign}$$

Frobenius characterstic, $\text{Frob}: \text{Rep}(S_n) \rightarrow \Lambda$, such that

- Irreducible S_n -representation V_λ has $\text{Frob}(V_\lambda) = s_\lambda$
- $U \cong V \oplus W \implies \text{Frob}(U) = \text{Frob}(V) + \text{Frob}(W)$

Representation theory and Schur functions

Irreducible representations of S_n are **also** labeled by partitions of n .

$$V_{\square\square\square} = \text{trivial}, \quad V_{\begin{smallmatrix} \square \\ \square \end{smallmatrix}} \cong \text{span}\{e_1 - e_2, e_2 - e_3\}, \quad V_{\begin{smallmatrix} \square \\ \square \\ \square \end{smallmatrix}} = \text{sign}$$

Frobenius characterstic, $\text{Frob}: \text{Rep}(S_n) \rightarrow \Lambda$, such that

- Irreducible S_n -representation V_λ has $\text{Frob}(V_\lambda) = s_\lambda$
- $U \cong V \oplus W \implies \text{Frob}(U) = \text{Frob}(V) + \text{Frob}(W)$
- $\text{Ind}_{S_m \times S_n}^{S_{m+n}}(V \times W) \mapsto \text{Frob}(V) \cdot \text{Frob}(W)$

Representation theory and Schur functions

Irreducible representations of S_n are **also** labeled by partitions of n .

$$V_{\square\square\square} = \text{trivial}, \quad V_{\begin{smallmatrix} \square \\ \square \end{smallmatrix}} \cong \text{span}\{e_1 - e_2, e_2 - e_3\}, \quad V_{\begin{smallmatrix} \square \\ \square \\ \square \end{smallmatrix}} = \text{sign}$$

Frobenius character map, $\text{Frob}: \text{Rep}(S_n) \rightarrow \Lambda$, such that

- Irreducible S_n -representation V_λ has $\text{Frob}(V_\lambda) = s_\lambda$
- $U \cong V \oplus W \implies \text{Frob}(U) = \text{Frob}(V) + \text{Frob}(W)$
- $\text{Ind}_{S_m \times S_n}^{S_{m+n}}(V \times W) \mapsto \text{Frob}(V) \cdot \text{Frob}(W)$
- $\text{Frob}(\text{Ind}_{S_1 \times S_1 \times S_1}^{S_3}(\mathbb{C} \times \mathbb{C} \times \mathbb{C})) = (s_{\square})^3 = s_{\square\square\square} + 2s_{\begin{smallmatrix} \square \\ \square \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ \square \end{smallmatrix}}$

Representation theory and Schur functions

Irreducible representations of S_n are **also** labeled by partitions of n .

$$V_{\square\square\square} = \text{trivial}, \quad V_{\begin{smallmatrix}\square & \\ \square & \end{smallmatrix}} \cong \text{span}\{e_1 - e_2, e_2 - e_3\}, \quad V_{\begin{smallmatrix}\square \\ \square \\ \square \end{smallmatrix}} = \text{sign}$$

Frobenius character map, $\text{Frob}: \text{Rep}(S_n) \rightarrow \Lambda$, such that

- Irreducible S_n -representation V_λ has $\text{Frob}(V_\lambda) = s_\lambda$
- $U \cong V \oplus W \implies \text{Frob}(U) = \text{Frob}(V) + \text{Frob}(W)$
- $\text{Ind}_{S_m \times S_n}^{S_{m+n}}(V \times W) \mapsto \text{Frob}(V) \cdot \text{Frob}(W)$
- $\text{Frob}(\text{Ind}_{S_1 \times S_1 \times S_1}^{S_3}(\mathbb{C} \times \mathbb{C} \times \mathbb{C})) = (s_{\square})^3 = s_{\square\square\square} + 2s_{\begin{smallmatrix}\square & \\ \square & \end{smallmatrix}} + s_{\begin{smallmatrix}\square \\ \square \\ \square \end{smallmatrix}}$

Hidden Guide: Schur Positivity

“Naturally occurring” symmetric functions which are non-negative (coefficients in $\mathbb{N}$) linear combinations in Schur polynomial basis are interesting since they could have representation-theoretic models.

Products of Schur polynomials

- Littlewood-Richardson rule: $s_\lambda s_\mu = \sum_\nu c_{\lambda\mu}^\nu s_\nu$ for $c_{\lambda\mu}^\nu \in \mathbb{N}$.

Products of Schur polynomials

- Littlewood-Richardson rule: $s_\lambda s_\mu = \sum_\nu c_{\lambda\mu}^\nu s_\nu$ for $c_{\lambda\mu}^\nu \in \mathbb{N}$.
- Skew Schur function: $s_{\nu/\lambda} = \sum_{T \in \text{SSYT}(\nu/\lambda)} x^T = \sum_\mu c_{\lambda\mu}^\nu s_\mu$.

Products of Schur polynomials

- Littlewood-Richardson rule: $s_\lambda s_\mu = \sum_\nu c_{\lambda\mu}^\nu s_\nu$ for $c_{\lambda\mu}^\nu \in \mathbb{N}$.
- Skew Schur function: $s_{\nu/\lambda} = \sum_{T \in \text{SSYT}(\nu/\lambda)} x^T = \sum_\mu c_{\lambda\mu}^\nu s_\mu$.

$$s_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} = x_1^3 x_2 + 2x_1^2 x_2^2 + \dots = s_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array}} + s_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}}$$

$$\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 1 & 1 \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 1 & 2 \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 2 & 2 \\ \hline 1 & 1 \\ \hline \end{array} \dots$$

Products of Schur polynomials

- Littlewood-Richardson rule: $s_\lambda s_\mu = \sum_\nu c_{\lambda\mu}^\nu s_\nu$ for $c_{\lambda\mu}^\nu \in \mathbb{N}$.
- Skew Schur function: $s_{\nu/\lambda} = \sum_{T \in \text{SSYT}(\nu/\lambda)} x^T = \sum_\mu c_{\lambda\mu}^\nu s_\mu$.

$$s_{\begin{smallmatrix} \square & \square \\ & \square \end{smallmatrix}} = x_1^3 x_2 + 2x_1^2 x_2^2 + \dots = s_{\begin{smallmatrix} \square & \square & \square \\ & \square & \square \end{smallmatrix}} + s_{\begin{smallmatrix} \square & \square \\ & \square \end{smallmatrix}}$$

$$\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 1 & 1 \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 1 & 2 \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 2 & 2 \\ \hline 1 & 1 \\ \hline \end{array} \dots$$

- Straightforwardly,

$$s_{\nu/\lambda} \cdot s_{\kappa/\mu} = \left(\sum_{T \in \text{SSYT}(\nu/\lambda)} x^T \right) \left(\sum_{S \in \text{SSYT}(\kappa/\mu)} x^S \right) = \sum_{\substack{T \in \text{SSYT}(\nu/\lambda) \\ S \in \text{SSYT}(\kappa/\mu)}} x^T x^S.$$

Products of Schur polynomials

- Littlewood-Richardson rule: $s_\lambda s_\mu = \sum_\nu c_{\lambda\mu}^\nu s_\nu$ for $c_{\lambda\mu}^\nu \in \mathbb{N}$.
- Skew Schur function: $s_{\nu/\lambda} = \sum_{T \in \text{SSYT}(\nu/\lambda)} x^T = \sum_\mu c_{\lambda\mu}^\nu s_\mu$.

$$s_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} = x_1^3 x_2 + 2x_1^2 x_2^2 + \dots = s_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array}} + s_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}}$$

$$\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 1 & 1 \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 1 & 2 \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 2 & 2 \\ \hline 1 & 1 \\ \hline \end{array} \dots$$

- Straightforwardly,

$$s_{\nu/\lambda} \cdot s_{\kappa/\mu} = \left(\sum_{T \in \text{SSYT}(\nu/\lambda)} x^T \right) \left(\sum_{S \in \text{SSYT}(\kappa/\mu)} x^S \right) = \sum_{\substack{T \in \text{SSYT}(\nu/\lambda) \\ S \in \text{SSYT}(\kappa/\mu)}} x^T x^S.$$

$$s_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} \cdot s_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} = x_1^5 x_2^3 + \dots$$

$$\left(\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 1 & 1 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 2 & 2 \\ \hline 1 & 1 \\ \hline \end{array} \right) \dots$$

Products of Schur polynomials

- Littlewood-Richardson rule: $s_\lambda s_\mu = \sum_\nu c_{\lambda\mu}^\nu s_\nu$ for $c_{\lambda\mu}^\nu \in \mathbb{N}$.
- Skew Schur function: $s_{\nu/\lambda} = \sum_{T \in \text{SSYT}(\nu/\lambda)} x^T = \sum_\mu c_{\lambda\mu}^\nu s_\mu$.

$$s_{\begin{smallmatrix} \square & \square \\ & \square \end{smallmatrix}} = x_1^3 x_2 + 2x_1^2 x_2^2 + \dots = s_{\begin{smallmatrix} \square & \square & \square \\ & & \square \end{smallmatrix}} + s_{\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}}$$

$$\begin{array}{|c|c|} \hline 1 & 2 \\ \hline & \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 1 & 2 \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 2 & 2 \\ \hline 1 & 1 \\ \hline \end{array} \quad \dots$$

- Straightforwardly,

$$s_{\nu/\lambda} \cdot s_{\kappa/\mu} = \left(\sum_{T \in \text{SSYT}(\nu/\lambda)} x^T \right) \left(\sum_{S \in \text{SSYT}(\kappa/\mu)} x^S \right) = \sum_{\substack{T \in \text{SSYT}(\nu/\lambda) \\ S \in \text{SSYT}(\kappa/\mu)}} x^T x^S.$$

$$s_{\begin{smallmatrix} \square & \square \\ & \square \end{smallmatrix}} \cdot s_{\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}} = x_1^5 x_2^3 + \dots$$


$$\left(\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 1 & 1 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 2 & 2 \\ \hline 1 & 1 \\ \hline \end{array} \right) \dots$$

- q -deformation?

Key Object: LLT Polynomials

Let $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$ be a tuple of skew shapes. (Skew shape = $\lambda \setminus \mu$)

$$\nu = \left(\begin{array}{|c|c|c|} \hline \square & \square & \\ \hline \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array} \right)$$



Key Object: LLT Polynomials

Let $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$ be a tuple of skew shapes. (Skew shape = $\lambda \setminus \mu$)

- The *content* of a box in row y , column x is $x - y$.

$$\nu = \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array} \right)$$

-4	-3	-2	-1	0	1
-3	-2	-1	0	1	2
-2	-1	0	1	2	3
-1	0	1	2	3	4
0	1	2	3	4	5

Key Object: LLT Polynomials

Let $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$ be a tuple of skew shapes. (Skew shape = $\lambda \setminus \mu$)

- The *content* of a box in row y , column x is $x - y$.
- *Reading order*: label boxes $b_1, \dots, b_n$ by scanning each diagonal from southwest to northeast, in order of increasing content.

$$\nu = \left(\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array} \right)$$

				b_3	b_6
				b_5	b_8
b_1	b_2				
	b_4	b_7			

Key Object: LLT Polynomials

Let $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$ be a tuple of skew shapes. (Skew shape $= \lambda \setminus \mu$)

- The *content* of a box in row y , column x is $x - y$.
- *Reading order*: label boxes $b_1, \dots, b_n$ by scanning each diagonal from southwest to northeast, in order of increasing content.
- A pair $(a, b) \in \nu$ is *attacking* if a precedes b in reading order and
 - $\text{content}(b) = \text{content}(a)$, or
 - $\text{content}(b) = \text{content}(a) + 1$ and $a \in \nu_{(i)}, b \in \nu_{(j)}$ with $i > j$.

$$\nu = \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right)$$

				b_3	b_6
				b_5	b_8
b_1	b_2				
	b_4	b_7			

Attacking pairs: $(b_2, b_3), (b_3, b_4), (b_4, b_5), (b_4, b_6), (b_5, b_7), (b_6, b_7), (b_7, b_8)$

Key Object: LLT Polynomials

Let $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$ be a tuple of skew shapes. (Skew shape $= \lambda \setminus \mu$)

- The *content* of a box in row y , column x is $x - y$.
- *Reading order*: label boxes $b_1, \dots, b_n$ by scanning each diagonal from southwest to northeast, in order of increasing content.
- A pair $(a, b) \in \nu$ is *attacking* if a precedes b in reading order and
 - $\text{content}(b) = \text{content}(a)$, or
 - $\text{content}(b) = \text{content}(a) + 1$ and $a \in \nu_{(i)}, b \in \nu_{(j)}$ with $i > j$.

$$\nu = \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right)$$

				b_3	b_6
				b_5	b_8
b_1	b_2				
	b_4	b_7			

Attacking pairs: $(b_2, b_3), (b_3, b_4), (b_4, b_5), (b_4, b_6), (b_5, b_7), (b_6, b_7), (b_7, b_8)$

Key Object: LLT Polynomials

Let $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$ be a tuple of skew shapes. (Skew shape $= \lambda \setminus \mu$)

- The *content* of a box in row y , column x is $x - y$.
- *Reading order*: label boxes $b_1, \dots, b_n$ by scanning each diagonal from southwest to northeast, in order of increasing content.
- A pair $(a, b) \in \nu$ is *attacking* if a precedes b in reading order and
 - $\text{content}(b) = \text{content}(a)$, or
 - $\text{content}(b) = \text{content}(a) + 1$ and $a \in \nu_{(i)}, b \in \nu_{(j)}$ with $i > j$.

$$\nu = \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right)$$

					b_3	b_6
					b_5	b_8
b_1	b_2					
	b_4	b_7				

Attacking pairs: (b_2, b_3) , (b_3, b_4) , (b_4, b_5) , (b_4, b_6) , (b_5, b_7) , (b_6, b_7) , (b_7, b_8)

Key Object: LLT Polynomials

Let $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$ be a tuple of skew shapes. (Skew shape $= \lambda \setminus \mu$)

- The *content* of a box in row y , column x is $x - y$.
- *Reading order*: label boxes $b_1, \dots, b_n$ by scanning each diagonal from southwest to northeast, in order of increasing content.
- A pair $(a, b) \in \nu$ is *attacking* if a precedes b in reading order and
 - $\text{content}(b) = \text{content}(a)$, or
 - $\text{content}(b) = \text{content}(a) + 1$ and $a \in \nu_{(i)}, b \in \nu_{(j)}$ with $i > j$.

$$\nu = \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right)$$

					b_3	b_6
					b_5	b_8
b_1	b_2					
	b_4	b_7				

Attacking pairs: $(b_2, b_3), (b_3, b_4), (b_4, b_5), (b_4, b_6), (b_5, b_7), (b_6, b_7), (b_7, b_8)$

Key Object: LLT Polynomials

Let $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$ be a tuple of skew shapes. (Skew shape $= \lambda \setminus \mu$)

- The *content* of a box in row y , column x is $x - y$.
- *Reading order*: label boxes $b_1, \dots, b_n$ by scanning each diagonal from southwest to northeast, in order of increasing content.
- A pair $(a, b) \in \nu$ is *attacking* if a precedes b in reading order and
 - $\text{content}(b) = \text{content}(a)$, or
 - $\text{content}(b) = \text{content}(a) + 1$ and $a \in \nu_{(i)}, b \in \nu_{(j)}$ with $i > j$.

$$\nu = \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right)$$

				b_3	b_6
				b_5	b_8
b_1	b_2				
	b_4	b_7			

Attacking pairs: $(b_2, b_3), (b_3, b_4), (b_4, b_5), (b_4, b_6), (b_5, b_7), (b_6, b_7), (b_7, b_8)$

Key Object: LLT Polynomials

Let $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$ be a tuple of skew shapes. (Skew shape $= \lambda \setminus \mu$)

- The *content* of a box in row y , column x is $x - y$.
- *Reading order*: label boxes $b_1, \dots, b_n$ by scanning each diagonal from southwest to northeast, in order of increasing content.
- A pair $(a, b) \in \nu$ is *attacking* if a precedes b in reading order and
 - $\text{content}(b) = \text{content}(a)$, or
 - $\text{content}(b) = \text{content}(a) + 1$ and $a \in \nu_{(i)}, b \in \nu_{(j)}$ with $i > j$.

$$\nu = \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right)$$

					b_3	b_6
					b_5	b_8
b_1	b_2					
	b_4	b_7				

Attacking pairs: $(b_2, b_3), (b_3, b_4), (b_4, b_5), (b_4, b_6), (b_5, b_7), (b_6, b_7), (b_7, b_8)$

Key Object: LLT Polynomials

Let $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$ be a tuple of skew shapes. (Skew shape $= \lambda \setminus \mu$)

- The *content* of a box in row y , column x is $x - y$.
- *Reading order*: label boxes $b_1, \dots, b_n$ by scanning each diagonal from southwest to northeast, in order of increasing content.
- A pair $(a, b) \in \nu$ is *attacking* if a precedes b in reading order and
 - $\text{content}(b) = \text{content}(a)$, or
 - $\text{content}(b) = \text{content}(a) + 1$ and $a \in \nu_{(i)}, b \in \nu_{(j)}$ with $i > j$.

$$\nu = \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right)$$

				b_3	b_6
				b_5	b_8
b_1	b_2				
	b_4	b_7			

Attacking pairs: $(b_2, b_3), (b_3, b_4), (b_4, b_5), (b_4, b_6), (b_5, b_7), (b_6, b_7), (b_7, b_8)$

Key Object: LLT Polynomials

Let $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$ be a tuple of skew shapes. (Skew shape $= \lambda \setminus \mu$)

- The *content* of a box in row y , column x is $x - y$.
- *Reading order*: label boxes $b_1, \dots, b_n$ by scanning each diagonal from southwest to northeast, in order of increasing content.
- A pair $(a, b) \in \nu$ is *attacking* if a precedes b in reading order and
 - $\text{content}(b) = \text{content}(a)$, or
 - $\text{content}(b) = \text{content}(a) + 1$ and $a \in \nu_{(i)}, b \in \nu_{(j)}$ with $i > j$.

$$\nu = \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right)$$

				b_3	b_6
				b_5	b_8
b_1	b_2				
	b_4	b_7			

Attacking pairs: $(b_2, b_3), (b_3, b_4), (b_4, b_5), (b_4, b_6), (b_5, b_7), (b_6, b_7), (b_7, b_8)$

LLT Polynomials

- A *semistandard tableau* on ν is a map $T: \nu \rightarrow \mathbb{Z}_+$ which restricts to a semistandard tableau on each $\nu_{(i)}$.

The *LLT polynomial* indexed by a tuple of skew shapes ν is

$$\mathcal{G}_\nu(x; q) = \sum_{T \in \text{SSYT}(\nu)} x^T,$$

$$x^T = \prod_{a \in \nu} x_{T(a)}.$$

$T =$

				5	6
				1	1
2	4				
	3	5			

$$x^T = x_1^2 x_2 x_3 x_4 x_5^2 x_6$$

LLT Polynomials

- A *semistandard tableau* on ν is a map $T: \nu \rightarrow \mathbb{Z}_+$ which restricts to a semistandard tableau on each $\nu_{(i)}$.
- An *attacking inversion* in T is an attacking pair (a, b) such that $T(a) > T(b)$.

The *LLT polynomial* indexed by a tuple of skew shapes ν is

$$\mathcal{G}_\nu(x; q) = \sum_{T \in \text{SSYT}(\nu)} q^{\text{inv}(T)} x^T,$$

where $\text{inv}(T)$ is the number of attacking inversions in T and $x^T = \prod_{a \in \nu} x_{T(a)}$.

$$T = \begin{array}{|c|c|c|c|c|c|} \hline & & & & 5 & 6 \\ \hline & & & & 1 & 1 \\ \hline & & & & & \\ \hline 2 & 4 & & & & \\ \hline & 3 & 5 & & & \\ \hline \end{array}$$

$$x^T = x_1^2 x_2 x_3 x_4 x_5^2 x_6$$

LLT Polynomials

- A *semistandard tableau* on ν is a map $T: \nu \rightarrow \mathbb{Z}_+$ which restricts to a semistandard tableau on each $\nu_{(i)}$.
- An *attacking inversion* in T is an attacking pair (a, b) such that $T(a) > T(b)$.

The *LLT polynomial* indexed by a tuple of skew shapes ν is

$$\mathcal{G}_\nu(x; q) = \sum_{T \in \text{SSYT}(\nu)} q^{\text{inv}(T)} x^T,$$

where $\text{inv}(T)$ is the number of attacking inversions in T and $x^T = \prod_{a \in \nu} x_{T(a)}$.

$T =$

				5	6
				1	1
2	4				
	3	5			

non-inversion

$$x^T = x_1^2 x_2 x_3 x_4 x_5^2 x_6$$

LLT Polynomials

- A *semistandard tableau* on ν is a map $T: \nu \rightarrow \mathbb{Z}_+$ which restricts to a semistandard tableau on each $\nu_{(i)}$.
- An *attacking inversion* in T is an attacking pair (a, b) such that $T(a) > T(b)$.

The *LLT polynomial* indexed by a tuple of skew shapes ν is

$$\mathcal{G}_\nu(x; q) = \sum_{T \in \text{SSYT}(\nu)} q^{\text{inv}(T)} x^T,$$

where $\text{inv}(T)$ is the number of attacking inversions in T and $x^T = \prod_{a \in \nu} x_{T(a)}$.

$T =$

				5	6
				1	1
2	4				
	3	5			

inversion

$$x^T = x_1^2 x_2 x_3 x_4 x_5^2 x_6$$

LLT Polynomials

- A *semistandard tableau* on ν is a map $T: \nu \rightarrow \mathbb{Z}_+$ which restricts to a semistandard tableau on each $\nu_{(i)}$.
- An *attacking inversion* in T is an attacking pair (a, b) such that $T(a) > T(b)$.

The *LLT polynomial* indexed by a tuple of skew shapes ν is

$$\mathcal{G}_\nu(x; q) = \sum_{T \in \text{SSYT}(\nu)} q^{\text{inv}(T)} x^T,$$

where $\text{inv}(T)$ is the number of attacking inversions in T and $x^T = \prod_{a \in \nu} x_{T(a)}$.

$T =$

				5	6	
				1	1	
2	4					
	3	5				

inversion

$$x^T = x_1^2 x_2 x_3 x_4 x_5^2 x_6$$

LLT Polynomials

- A *semistandard tableau* on ν is a map $T: \nu \rightarrow \mathbb{Z}_+$ which restricts to a semistandard tableau on each $\nu_{(i)}$.
- An *attacking inversion* in T is an attacking pair (a, b) such that $T(a) > T(b)$.

The *LLT polynomial* indexed by a tuple of skew shapes ν is

$$\mathcal{G}_\nu(x; q) = \sum_{T \in \text{SSYT}(\nu)} q^{\text{inv}(T)} x^T,$$

where $\text{inv}(T)$ is the number of attacking inversions in T and $x^T = \prod_{a \in \nu} x_{T(a)}$.

$T =$

				5	6
				1	1
2	4				
	3	5			

non-inversion

$$x^T = x_1^2 x_2 x_3 x_4 x_5^2 x_6$$

LLT Polynomials

- A *semistandard tableau* on ν is a map $T: \nu \rightarrow \mathbb{Z}_+$ which restricts to a semistandard tableau on each $\nu_{(i)}$.
- An *attacking inversion* in T is an attacking pair (a, b) such that $T(a) > T(b)$.

The *LLT polynomial* indexed by a tuple of skew shapes ν is

$$\mathcal{G}_\nu(x; q) = \sum_{T \in \text{SSYT}(\nu)} q^{\text{inv}(T)} x^T,$$

where $\text{inv}(T)$ is the number of attacking inversions in T and $x^T = \prod_{a \in \nu} x_{T(a)}$.

$T =$

				5	6
				1	1
2	4				
	3	5			

non-inversion

$$x^T = x_1^2 x_2 x_3 x_4 x_5^2 x_6$$

LLT Polynomials

- A *semistandard tableau* on ν is a map $T: \nu \rightarrow \mathbb{Z}_+$ which restricts to a semistandard tableau on each $\nu_{(i)}$.
- An *attacking inversion* in T is an attacking pair (a, b) such that $T(a) > T(b)$.

The *LLT polynomial* indexed by a tuple of skew shapes ν is

$$\mathcal{G}_\nu(x; q) = \sum_{T \in \text{SSYT}(\nu)} q^{\text{inv}(T)} x^T,$$

where $\text{inv}(T)$ is the number of attacking inversions in T and $x^T = \prod_{a \in \nu} x_{T(a)}$.

$T =$

				5	6	
				1	1	
2	4					
	3	5				

inversion

$$x^T = x_1^2 x_2 x_3 x_4 x_5^2 x_6$$

LLT Polynomials

- A *semistandard tableau* on ν is a map $T: \nu \rightarrow \mathbb{Z}_+$ which restricts to a semistandard tableau on each $\nu_{(i)}$.
- An *attacking inversion* in T is an attacking pair (a, b) such that $T(a) > T(b)$.

The *LLT polynomial* indexed by a tuple of skew shapes ν is

$$\mathcal{G}_\nu(x; q) = \sum_{T \in \text{SSYT}(\nu)} q^{\text{inv}(T)} x^T,$$

where $\text{inv}(T)$ is the number of attacking inversions in T and $x^T = \prod_{a \in \nu} x_{T(a)}$.

$T =$

				5	6	
				1	1	
2	4					
	3	5				

inversion

$$x^T = x_1^2 x_2 x_3 x_4 x_5^2 x_6$$

LLT Polynomials

- A *semistandard tableau* on ν is a map $T: \nu \rightarrow \mathbb{Z}_+$ which restricts to a semistandard tableau on each $\nu_{(i)}$.
- An *attacking inversion* in T is an attacking pair (a, b) such that $T(a) > T(b)$.

The *LLT polynomial* indexed by a tuple of skew shapes ν is

$$\mathcal{G}_\nu(x; q) = \sum_{T \in \text{SSYT}(\nu)} q^{\text{inv}(T)} x^T,$$

where $\text{inv}(T)$ is the number of attacking inversions in T and $x^T = \prod_{a \in \nu} x_{T(a)}$.

$$T = \begin{array}{|c|c|c|c|c|c|} \hline & & & & 5 & 6 \\ \hline & & & & 1 & 1 \\ \hline & & & & & \\ \hline 2 & 4 & & & & \\ \hline & 3 & 5 & & & \\ \hline \end{array}$$

$$\text{inv}(T) = 4, \quad x^T = x_1^2 x_2 x_3 x_4 x_5^2 x_6$$

LLT Polynomials $\mathcal{G}_\nu(X; q)$

- $\mathcal{G}_\nu(x; q)$ is a symmetric function

LLT Polynomials $\mathcal{G}_\nu(X; q)$

- $\mathcal{G}_\nu(x; q)$ is a symmetric function
- $\mathcal{G}_\nu(x; 1) = s_{\nu(1)} \cdots s_{\nu(r)}$

LLT Polynomials $\mathcal{G}_\nu(X; q)$

- $\mathcal{G}_\nu(x; q)$ is a symmetric function
- $\mathcal{G}_\nu(x; 1) = s_{\nu(1)} \cdots s_{\nu(r)}$
- $\mathcal{G}_\nu$ were originally defined by Lascoux, Leclerc, and Thibon to explore connections to Fock space representations of $U_q(\hat{\mathfrak{sl}}_r)$

LLT Polynomials $\mathcal{G}_\nu(X; q)$

- $\mathcal{G}_\nu(x; q)$ is a symmetric function
- $\mathcal{G}_\nu(x; 1) = s_{\nu(1)} \cdots s_{\nu(r)}$
- $\mathcal{G}_\nu$ were originally defined by Lascoux, Leclerc, and Thibon to explore connections to Fock space representations of $U_q(\hat{\mathfrak{sl}}_r)$
- When $\nu^{(i)}$ are partitions, the Schur-expansion coefficients are essentially parabolic Kazhdan-Luzstig polynomials.

LLT Polynomials $\mathcal{G}_\nu(X; q)$

- $\mathcal{G}_\nu(x; q)$ is a symmetric function
- $\mathcal{G}_\nu(x; 1) = s_{\nu(1)} \cdots s_{\nu(r)}$
- $\mathcal{G}_\nu$ were originally defined by Lascoux, Leclerc, and Thibon to explore connections to Fock space representations of $U_q(\hat{\mathfrak{sl}}_r)$
- When $\nu^{(i)}$ are partitions, the Schur-expansion coefficients are essentially parabolic Kazhdan-Luzstig polynomials.
- $\mathcal{G}_\nu$ is Schur-positive for any tuple of skew shapes ν [Grojnowski-Haiman, 2007].

LLT Polynomials $\mathcal{G}_\nu(X; q)$

- $\mathcal{G}_\nu(x; q)$ is a symmetric function
- $\mathcal{G}_\nu(x; 1) = s_{\nu(1)} \cdots s_{\nu(r)}$
- $\mathcal{G}_\nu$ were originally defined by Lascoux, Leclerc, and Thibon to explore connections to Fock space representations of $U_q(\hat{\mathfrak{sl}}_r)$
- When $\nu^{(i)}$ are partitions, the Schur-expansion coefficients are essentially parabolic Kazhdan-Luzstig polynomials.
- $\mathcal{G}_\nu$ is Schur-positive for any tuple of skew shapes ν [Grojnowski-Haiman, 2007].
- Thus, $\mathcal{G}_\nu$ are useful to show combinatorial quantities are Schur-positive!

Some notable occurrences of LLT Polynomials

- Haglund-Haiman-Loehr formula for Macdonald polynomials:

$$\tilde{H}_\mu(x; q, t) = t^{n(\mu)} \sum_R \left(\prod_{\substack{\square \\ u}} q^{a+1} t^l \right) \mathcal{G}_R(x; t^{-1}).$$

Some notable occurrences of LLT Polynomials

- Haglund-Haiman-Loehr formula for Macdonald polynomials:

$$\tilde{H}_{\mu}(x; q, t) = t^{n(\mu)} \sum_R \left(\prod_{\substack{\square \\ u}} q^{a+1} t^l \right) \mathcal{G}_R(x; t^{-1}).$$

- Shuffle theorem (and generalizations):

$$\nabla e_n = \sum_{\lambda \in DP_n} t^{\text{area}(\lambda)} q^{\text{dinv}(\lambda)} \mathcal{G}_{\nu(\lambda)}(x; q^{-1}).$$

Some notable occurrences of LLT Polynomials

- Haglund-Haiman-Loehr formula for Macdonald polynomials:

$$\tilde{H}_{\mu}(x; q, t) = t^{n(\mu)} \sum_R \left(\prod_{\substack{\square \\ u}} q^{a+1} t^l \right) \mathcal{G}_R(x; t^{-1}).$$

- Shuffle theorem (and generalizations):

$$\nabla e_n = \sum_{\lambda \in DP_n} t^{\text{area}(\lambda)} q^{\text{dinv}(\lambda)} \mathcal{G}_{\nu(\lambda)}(x; q^{-1}).$$

- For G = incomparability graph for natural unit interval order (encoded by Dyck path P), the t -chromatic symmetric polynomial is essentially a “signed” unicellular LLT polynomial (later), i.e.,

$$\chi_G(x; t) = (1 - t)^{-|V|} \mathcal{G}_{\nu(P)}[(1 - t)x; t].$$

Flagged Tableaux

- Fix partition shape $\lambda = (\lambda_1, \dots, \lambda_l)$ and *flag*
 $\mathbf{b} = (b_1 \leq \dots \leq b_l) \in \mathbb{N}^l$.

Flagged Tableaux

- Fix partition shape $\lambda = (\lambda_1, \dots, \lambda_l)$ and *flag* $\mathbf{b} = (b_1 \leq \dots \leq b_l) \in \mathbb{N}^l$.
- A *flagged semistandard Young tableau* of shape λ with flag $\mathbf{b}$ is a $T \in \text{SSYT}(\lambda)$ such that the entries of row i are bounded above by b_i .
- Denote the set of such tableaux via $\text{FT}(\lambda, \mathbf{b})$.
- E.g., $\lambda = (2, 1)$, $\mathbf{b} = (1, 3)$, $\text{FT}(\lambda, \mathbf{b}) =$

$$\begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 1 \\ \hline \end{array} \leq 1 \quad \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 1 \\ \hline \end{array} \leq 1$$

Flagged Schur functions (Lascoux-Schützenberger, Wachs)

- For partition λ and flag $\mathbf{b}$, the *flagged Schur function* is given by

$$s_{\lambda, \mathbf{b}}(x) = \sum_{T \in \text{FT}(\lambda, \mathbf{b})} x^T$$

Flagged Schur functions (Lascoux-Schützenberger, Wachs)

- For partition λ and flag $\mathbf{b}$, the *flagged Schur function* is given by

$$s_{\lambda, \mathbf{b}}(x) = \sum_{T \in \text{FT}(\lambda, \mathbf{b})} x^T$$

- Jacobi-Trudi: $s_{\lambda, \mathbf{b}}(x) = \det(h_{\lambda_i - i + j}(x_1, \dots, x_{b_i}))$

Flagged Schur functions (Lascoux-Schützenberger, Wachs)

- For partition λ and flag $\mathbf{b}$, the *flagged Schur function* is given by

$$s_{\lambda, \mathbf{b}}(x) = \sum_{T \in \text{FT}(\lambda, \mathbf{b})} x^T$$

- Jacobi-Trudi: $s_{\lambda, \mathbf{b}}(x) = \det(h_{\lambda_i - i + j}(x_1, \dots, x_{b_i}))$
- E.g., $s_{21, 13}(x_1, x_2, x_3) = x_1^2 x_2 + x_1^2 x_3 = h_2(x_1)h_1(x_1, x_2, x_3) - h_3(x_1)$

$$\begin{array}{|c|c|} \hline 2 & \leq 3 \\ \hline 1 & 1 \\ \hline \end{array} \leq 1 \quad \begin{array}{|c|c|} \hline 3 & \leq 3 \\ \hline 1 & 1 \\ \hline \end{array} \leq 1 \quad \det \begin{pmatrix} h_2(x_1) & h_3(x_1) \\ h_0(x_1, x_2, x_3) & h_1(x_1, x_2, x_3) \end{pmatrix}$$

Flagged Schur functions (Lascoux-Schützenberger, Wachs)

- For partition λ and flag $\mathbf{b}$, the *flagged Schur function* is given by

$$s_{\lambda, \mathbf{b}}(x) = \sum_{T \in \text{FT}(\lambda, \mathbf{b})} x^T$$

- Jacobi-Trudi: $s_{\lambda, \mathbf{b}}(x) = \det(h_{\lambda_i - i + j}(x_1, \dots, x_{b_i}))$
- E.g., $s_{21, 13}(x_1, x_2, x_3) = x_1^2 x_2 + x_1^2 x_3 = h_2(x_1)h_1(x_1, x_2, x_3) - h_3(x_1)$

$$\begin{array}{|c|c|} \hline 2 & \leq 3 \\ \hline 1 & 1 \\ \hline \end{array} \leq 1 \quad \begin{array}{|c|c|} \hline 3 & \leq 3 \\ \hline 1 & 1 \\ \hline \end{array} \leq 1 \quad \det \begin{pmatrix} h_2(x_1) & h_3(x_1) \\ h_0(x_1, x_2, x_3) & h_1(x_1, x_2, x_3) \end{pmatrix}$$

- Schubert $\mathfrak{S}_w(x) = s_{\lambda, \mathbf{b}}(x)$ for *vexillary* $w \in S_n$ (2143-avoiding).

Flagged Schur functions (Lascoux-Schützenberger, Wachs)

- For partition λ and flag $\mathbf{b}$, the *flagged Schur function* is given by

$$s_{\lambda, \mathbf{b}}(x) = \sum_{T \in \text{FT}(\lambda, \mathbf{b})} x^T$$

- Jacobi-Trudi: $s_{\lambda, \mathbf{b}}(x) = \det(h_{\lambda_i - i + j}(x_1, \dots, x_{b_i}))$
- E.g., $s_{21, 13}(x_1, x_2, x_3) = x_1^2 x_2 + x_1^2 x_3 = h_2(x_1)h_1(x_1, x_2, x_3) - h_3(x_1)$

$$\begin{array}{|c|c|} \hline 2 & \leq 3 \\ \hline 1 & 1 \\ \hline \end{array} \leq 1 \quad \begin{array}{|c|c|} \hline 3 & \leq 3 \\ \hline 1 & 1 \\ \hline \end{array} \leq 1 \quad \det \begin{pmatrix} h_2(x_1) & h_3(x_1) \\ h_0(x_1, x_2, x_3) & h_1(x_1, x_2, x_3) \end{pmatrix}$$

- Schubert $\mathfrak{S}_w(x) = s_{\lambda, \mathbf{b}}(x)$ for *vexillary* $w \in S_n$ (2143-avoiding).
- $s_{\lambda, \mathbf{b}}(x)$ are examples of *Demazure characters* (Reiner-Shimozono, 1995).

Demazure characters

The *Demazure operator* π_i acts on $f \in \mathbb{K}[x_1^{\pm 1}, \dots, x_N^{\pm 1}]$ by

$$\pi_i(f) = \frac{x_i f - x_{i+1} s_i(f)}{x_i - x_{i+1}}.$$

Demazure characters

The *Demazure operator* π_i acts on $f \in \mathbb{K}[x_1^{\pm 1}, \dots, x_N^{\pm 1}]$ by

$$\pi_i(f) = \frac{x_i f - x_{i+1} s_i(f)}{x_i - x_{i+1}}.$$

The *Demazure characters* or *key polynomials* are constructed from

- $\mathcal{D}_\lambda = x^\lambda := x_1^{\lambda_1} \cdots x_N^{\lambda_N}$ for partition λ .
- $\mathcal{D}_{s_i(\alpha)} = \pi_i \mathcal{D}_\alpha$ for $\alpha_i > \alpha_{i+1}$, for any $\alpha \in \mathbb{N}^N$.

Demazure characters

The *Demazure operator* π_i acts on $f \in \mathbb{k}[x_1^{\pm 1}, \dots, x_N^{\pm 1}]$ by

$$\pi_i(f) = \frac{x_i f - x_{i+1} s_i(f)}{x_i - x_{i+1}}.$$

The *Demazure characters* or *key polynomials* are constructed from

- $\mathcal{D}_\lambda = x^\lambda := x_1^{\lambda_1} \cdots x_N^{\lambda_N}$ for partition λ .
- $\mathcal{D}_{s_i(\alpha)} = \pi_i \mathcal{D}_\alpha$ for $\alpha_i > \alpha_{i+1}$, for any $\alpha \in \mathbb{N}^N$.

$$\mathcal{D}_{520} = x_1^5 x_2^2$$

$$\mathcal{D}_{250} = \pi_1 \mathcal{D}_{520} = \pi_1(x_1^5 x_2^2) = x_1^5 x_2^2 + x_1^4 x_2^3 + x_1^3 x_2^4 + x_1^2 x_2^5$$

$$\mathcal{D}_{205} = \pi_2 \mathcal{D}_{250} = \pi_2(x_1^5 x_2^2 + x_1^4 x_2^3 + x_1^3 x_2^4 + x_1^2 x_2^5)$$

Demazure characters

The *Demazure operator* π_i acts on $f \in \mathbb{k}[x_1^{\pm 1}, \dots, x_N^{\pm 1}]$ by

$$\pi_i(f) = \frac{x_i f - x_{i+1} s_i(f)}{x_i - x_{i+1}}.$$

The *Demazure characters* or *key polynomials* are constructed from

- $\mathcal{D}_\lambda = x^\lambda := x_1^{\lambda_1} \cdots x_N^{\lambda_N}$ for partition λ .
- $\mathcal{D}_{s_i(\alpha)} = \pi_i \mathcal{D}_\alpha$ for $\alpha_i > \alpha_{i+1}$, for any $\alpha \in \mathbb{N}^N$.

$$\mathcal{D}_{520} = x_1^5 x_2^2$$

$$\mathcal{D}_{250} = \pi_1 \mathcal{D}_{520} = \pi_1(x_1^5 x_2^2) = x_1^5 x_2^2 + x_1^4 x_2^3 + x_1^3 x_2^4 + x_1^2 x_2^5$$

$$\mathcal{D}_{205} = \pi_2 \mathcal{D}_{250} = \pi_2(x_1^5 x_2^2 + x_1^4 x_2^3 + x_1^3 x_2^4 + x_1^2 x_2^5)$$

- Demazure characters are characters of irreducible B -modules.

Demazure atoms

Demazure atoms are defined the same as keys but with $\hat{\pi}_i := \pi_i - 1$ in place of π_i :

- $\mathcal{A}_\lambda = x^\lambda$ for partition λ .
- $\mathcal{A}_{s_i(\alpha)} = \hat{\pi}_i \mathcal{A}_\alpha$ for $\alpha_i > \alpha_{i+1}$, for any $\alpha \in \mathbb{N}^N$.

Demazure atoms

Demazure atoms are defined the same as keys but with $\hat{\pi}_i := \pi_i - 1$ in place of π_i :

- $\mathcal{A}_\lambda = x^\lambda$ for partition λ .
- $\mathcal{A}_{s_i(\alpha)} = \hat{\pi}_i \mathcal{A}_\alpha$ for $\alpha_i > \alpha_{i+1}$, for any $\alpha \in \mathbb{N}^N$.
- Demazure atoms are related to Demazure characters by Bruhat order inclusion-exclusion:

$$\mathcal{D}_{520} = \mathcal{A}_{520}$$

$$\mathcal{D}_{250} = \mathcal{A}_{520} + \mathcal{A}_{250}$$

$$\mathcal{D}_{205} = \mathcal{A}_{520} + \mathcal{A}_{250} + \mathcal{A}_{502} + \mathcal{A}_{205}$$

$$\vdots$$

Recovering Symmetric Functions

- If $\mathbf{b} = (n, \dots, n)$ for $n = |\lambda|$, then $s_{\lambda, \mathbf{b}}(x) = s_{\lambda}(x_1, \dots, x_n)$.

Recovering Symmetric Functions

- If $\mathbf{b} = (n, \dots, n)$ for $n = |\lambda|$, then $s_{\lambda, \mathbf{b}}(x) = s_{\lambda}(x_1, \dots, x_n)$.
- For $w = s_{i_1} s_{i_2} \cdots s_{i_m} \in S_N$ reduced, $\pi_w := \pi_{i_1} \pi_{i_2} \cdots \pi_{i_m}$.

Recovering Symmetric Functions

- If $\mathbf{b} = (n, \dots, n)$ for $n = |\lambda|$, then $s_{\lambda, \mathbf{b}}(x) = s_{\lambda}(x_1, \dots, x_n)$.
- For $w = s_{i_1} s_{i_2} \cdots s_{i_m} \in S_N$ reduced, $\pi_w := \pi_{i_1} \pi_{i_2} \cdots \pi_{i_m}$.
- For *Weyl symmetrization operator* π_{w_0} , we have

Recovering Symmetric Functions

- If $\mathbf{b} = (n, \dots, n)$ for $n = |\lambda|$, then $s_{\lambda, \mathbf{b}}(x) = s_{\lambda}(x_1, \dots, x_n)$.
- For $w = s_{i_1} s_{i_2} \cdots s_{i_m} \in S_N$ reduced, $\pi_w := \pi_{i_1} \pi_{i_2} \cdots \pi_{i_m}$.
- For *Weyl symmetrization operator* π_{w_0} , we have

$$\pi_{w_0}(s_{\lambda, \mathbf{b}}(x)) = s_{\lambda}(x)$$

Recovering Symmetric Functions

- If $\mathbf{b} = (n, \dots, n)$ for $n = |\lambda|$, then $s_{\lambda, \mathbf{b}}(x) = s_{\lambda}(x_1, \dots, x_n)$.
- For $w = s_{i_1} s_{i_2} \cdots s_{i_m} \in S_N$ reduced, $\pi_w := \pi_{i_1} \pi_{i_2} \cdots \pi_{i_m}$.
- For *Weyl symmetrization operator* π_{w_0} , we have

$$\pi_{w_0}(s_{\lambda, \mathbf{b}}(x)) = s_{\lambda}(x), \quad \pi_{w_0}(\mathcal{D}_{\alpha}) = \mathcal{D}_{\text{sort}(\alpha)} = s_{\alpha+},$$


Recovering Symmetric Functions

- If $\mathbf{b} = (n, \dots, n)$ for $n = |\lambda|$, then $s_{\lambda, \mathbf{b}}(x) = s_{\lambda}(x_1, \dots, x_n)$.
- For $w = s_{i_1} s_{i_2} \cdots s_{i_m} \in S_N$ reduced, $\pi_w := \pi_{i_1} \pi_{i_2} \cdots \pi_{i_m}$.
- For *Weyl symmetrization operator* π_{w_0} , we have


$$\pi_{w_0}(s_{\lambda, \mathbf{b}}(x)) = s_{\lambda}(x), \quad \pi_{w_0}(\mathcal{D}_{\alpha}) = \mathcal{D}_{\text{sort}(\alpha)} = s_{\alpha+},$$

$$\pi_{w_0}(\mathcal{A}_{\alpha}) = \begin{cases} s_{\alpha} & \alpha \text{ a partition,} \\ 0 & \text{else.} \end{cases}$$

Summary of families



Summary of families



Question

Is there some “nice” family of polynomials that Weyl symmetrizes to LLT polynomials?

Flagged LLT Polynomials (Blasiak-Haiman-Morse-Pun-S.)

- Let $e_1, \dots, e_l$ be the row ends of ν , ordered in reverse reading order.
- Fix flag $\mathbf{b} = (b_1 \leq \dots \leq b_l)$.
- $\text{FT}(\nu, \mathbf{b})$ = set of semistandard tableaux T on ν satisfying $T(e_i) \leq b_i$.
- The *flagged LLT polynomial* indexed by ν and $\mathbf{b}$ is

$$\mathcal{G}_{\nu, \mathbf{b}}(x; t) = \sum_{T \in \text{FT}(\nu, \mathbf{b})} t^{\text{inv}(T)} x^T.$$

						e_3
						e_1
	e_4					
		e_2				

Flagged LLT Polynomials (Blasiak-Haiman-Morse-Pun-S.)

- Let $e_1, \dots, e_l$ be the row ends of ν , ordered in reverse reading order.
- Fix flag $\mathbf{b} = (b_1 \leq \dots \leq b_l)$.
- $\text{FT}(\nu, \mathbf{b})$ = set of semistandard tableaux T on ν satisfying $T(e_i) \leq b_i$.
- The *flagged LLT polynomial* indexed by ν and $\mathbf{b}$ is




$$\mathcal{G}_{\nu, \mathbf{b}}(x; t) = \sum_{T \in \text{FT}(\nu, \mathbf{b})} t^{\text{inv}(T)} x^T.$$

				2	3	≤ 3
				1	1	≤ 1
2	4					≤ 4
	1	2				≤ 2

Flagged LLT Polynomials

$$\nu = (\square\square, \square\square), \mathbf{b} = (1, 2)$$

T



$$\mathcal{G}_{r,\nu}(x; t) = x_1^4 + t x_1^3 x_2 + t^2 x_1^2 x_2^2$$

Flagged LLT Polynomials

- $\mathcal{G}_{\nu, \mathbf{b}}(x; t)$ Weyl symmetrizes to symmetric $\mathcal{G}_{\nu}(x; t)$.

Flagged LLT Polynomials

- $\mathcal{G}_{\nu, \mathbf{b}}(x; t)$ Weyl symmetrizes to symmetric $\mathcal{G}_{\nu}(x; t)$.
- For $\nu = (\lambda)$ a single shape, $\mathcal{G}_{\nu, \mathbf{b}}(x; t) = s_{\lambda, \mathbf{b}}(x)$.

Flagged LLT Polynomials

- $\mathcal{G}_{\nu, \mathbf{b}}(x; t)$ Weyl symmetrizes to symmetric $\mathcal{G}_{\nu}(x; t)$.
- For $\nu = (\lambda)$ a single shape, $\mathcal{G}_{\nu, \mathbf{b}}(x; t) = s_{\lambda, \mathbf{b}}(x)$.
- $\mathcal{G}_{\nu, \mathbf{b}}(x; t)$ also have an algebraic formula via “nonsymmetric Hall-Littlewood polynomials.”

Flagged LLT Polynomials

- $\mathcal{G}_{\nu, \mathbf{b}}(x; t)$ Weyl symmetrizes to symmetric $\mathcal{G}_{\nu}(x; t)$.
- For $\nu = (\lambda)$ a single shape, $\mathcal{G}_{\nu, \mathbf{b}}(x; t) = s_{\lambda, \mathbf{b}}(x)$.
- $\mathcal{G}_{\nu, \mathbf{b}}(x; t)$ also have an algebraic formula via “nonsymmetric Hall-Littlewood polynomials.”
- $\mathcal{G}_{\nu, \mathbf{b}}(x; t)$ are conjecturally positive in terms of Demazure atoms.

Signed flagged LLT polynomials

- Signed alphabet $\mathcal{A} = 1 < \bar{1} < 2 < \bar{2} \dots$

Signed flagged LLT polynomials

- Signed alphabet $\mathcal{A} = 1 < \bar{1} < 2 < \bar{2} \dots$
- $\text{FT}^\pm(\nu, \mathbf{b}) =$ fillings of ν from $\mathcal{A}$ satisfying
 - unbarred letters weakly increase in rows, strictly increase in columns.
 - barred letters strictly increase in rows, weakly increase in columns.
 - $T(e_i) \leq b_i$ for $i = 1, \dots, l$.

Signed flagged LLT polynomials

- Signed alphabet $\mathcal{A} = 1 < \bar{1} < 2 < \bar{2} \dots$
- $\text{FT}^\pm(\nu, \mathbf{b})$ = fillings of ν from $\mathcal{A}$ satisfying
 - unbarred letters weakly increase in rows, strictly increase in columns.
 - barred letters strictly increase in rows, weakly increase in columns.
 - $T(e_i) \leq b_i$ for $i = 1, \dots, l$.


The *signed flagged LLT polynomial* indexed by ν and flag $\mathbf{b}$ is

$$\mathcal{G}_{\nu, \mathbf{b}}^\pm(x; t) = \sum_{T \in \text{FT}^\pm(\nu, \mathbf{b})} t^{\text{inv}(T)} (-t)^{-\#\text{bar}(T)} x^{|T|},$$

where $|T|$ is the result of removing all bars from T .

Signed flagged LLT polynomials

$$\nu = (\square\square, \square\square), \mathbf{b} = (1, 2)$$



$$\begin{aligned} \mathcal{G}_{\nu,\mathbf{b}}^{\pm}(x;t) &= x_1^4 + tx_1^3x_2 + t^2x_1^2x_2^2 - x_1^4 - tx_1^3x_2 \\ &= t^2x_1^2x_2^2 \end{aligned}$$

Remark

Signed flagged LLTs are typically not positive in terms of Demazure atoms, nor are they generally even monomial positive.

Flagged plethysm

Define *flagged plethysm* $\Pi_{t,x}: \mathbb{k}[x_1, \dots, x_l] \rightarrow \mathbb{k}[x_1, \dots, x_l]$ to be the (over-determined) linear map

$$\Pi_{t,x} \mathcal{G}_{\nu, \mathbf{b}}^{\pm}(x; t^{-1}) = \mathcal{G}_{\nu, \mathbf{b}}(x; t^{-1}).$$

Flagged plethysm

Define *flagged plethysm* $\Pi_{t,x}: \mathbb{k}[x_1, \dots, x_l] \rightarrow \mathbb{k}[x_1, \dots, x_l]$ to be the (over-determined) linear map

$$\Pi_{t,x} \mathcal{G}_{\nu, \mathbf{b}}^{\pm}(x; t^{-1}) = \mathcal{G}_{\nu, \mathbf{b}}(x; t^{-1}).$$

Theorem (Blasiak-Haiman-Morse-Pun-S., 2025+)

$\Pi_{t,x}$ is well-defined.

Flagged plethysm

Define *flagged plethysm* $\Pi_{t,x}: \mathbb{k}[x_1, \dots, x_l] \rightarrow \mathbb{k}[x_1, \dots, x_l]$ to be the (over-determined) linear map

$$\Pi_{t,x} \mathcal{G}_{\nu, \mathbf{b}}^{\pm}(x; t^{-1}) = \mathcal{G}_{\nu, \mathbf{b}}(x; t^{-1}).$$

Theorem (Blasiak-Haiman-Morse-Pun-S., 2025+)

$\Pi_{t,x}$ is well-defined.

$\Pi_{t,x}$ is a “nonsymmetric analogue” of the plethystic map $f[X] \mapsto f[X/(1-t)]$ for symmetric function f .

Applications

- ① *Modified r -nonsymmetric Macdonald polynomials (or modern Macdonald polynomials) via a flagged HHL formula.*

Applications

- ① *Modified r -nonsymmetric Macdonald polynomials (or modern Macdonald polynomials)* via a flagged HHL formula.
- ② A “nonsymmetric shuffle theorem” expressed in terms of flagged LLT polynomials associated to flagged Dyck paths.

Applications

- ① *Modified r -nonsymmetric Macdonald polynomials (or modern Macdonald polynomials)* via a flagged HHL formula.
- ② A “nonsymmetric shuffle theorem” expressed in terms of flagged LLT polynomials associated to flagged Dyck paths.
- ③ Tewari-Wilson-Zhang define chromatic nonsymmetric polynomials associated to $d \times d$ Dyck paths starting with r north steps, $\chi_{\mathbf{b}, \pi}$. These are (essentially) examples of signed flagged LLT polynomials.

Macdonald polynomials

- Macdonald polynomials $P_\mu(x; q, t)$ form a basis for $\Lambda_{\mathbb{Q}(q,t)}(x)$.

Macdonald polynomials

- Macdonald polynomials $P_\mu(x; q, t)$ form a basis for $\Lambda_{\mathbb{Q}(q,t)}(x)$.
- Specializations give Hall-Littlewood polynomials, Schur polynomials, and many other famous bases.

Macdonald polynomials

- Macdonald polynomials $P_\mu(x; q, t)$ form a basis for $\Lambda_{\mathbb{Q}(q,t)}(x)$.
- Specializations give Hall-Littlewood polynomials, Schur polynomials, and many other famous bases.
- Integral form Macdonald polynomials $J_\mu(x; q, t) = c_\mu P_\mu(x; q, t)$ have coefficients in $\mathbb{Z}[q, t]$.

Macdonald polynomials

- Macdonald polynomials $P_\mu(x; q, t)$ form a basis for $\Lambda_{\mathbb{Q}(q,t)}(x)$.
- Specializations give Hall-Littlewood polynomials, Schur polynomials, and many other famous bases.
- Integral form Macdonald polynomials $J_\mu(x; q, t) = c_\mu P_\mu(x; q, t)$ have coefficients in $\mathbb{Z}[q, t]$.
- Macdonald provides a positivity conjecture about expanding J_μ into a funny basis.

Macdonald polynomials

- Macdonald polynomials $P_\mu(x; q, t)$ form a basis for $\Lambda_{\mathbb{Q}(q,t)}(x)$.
- Specializations give Hall-Littlewood polynomials, Schur polynomials, and many other famous bases.
- Integral form Macdonald polynomials $J_\mu(x; q, t) = c_\mu P_\mu(x; q, t)$ have coefficients in $\mathbb{Z}[q, t]$.
- Macdonald provides a positivity conjecture about expanding J_μ into a funny basis. What does it mean?

Modified Macdonald polynomials

- Plethystically modified Macdonald polynomials

$$H_\mu(x; q, t) = J_\mu[x/(1-t); q, t].$$

- Macdonald positivity: the H_μ are Schur positive.

Modified Macdonald polynomials

- Plethystically modified Macdonald polynomials

$$H_{\mu}(x; q, t) = J_{\mu}[x/(1-t); q, t].$$

- Macdonald positivity: the H_{μ} are Schur positive.
- $H_{\mu}(x; 1, 1) = (s_1)^n = h_{(1^n)}$.

Modified Macdonald polynomials

- Plethystically modified Macdonald polynomials

$$H_\mu(x; q, t) = J_\mu[x/(1-t); q, t].$$

- Macdonald positivity: the H_μ are Schur positive.
- $H_\mu(x; 1, 1) = (s_1)^n = h_{(1^n)}$.
- $t^{n(\mu)} H_\mu(x; q, t^{-1})$ = Frobenius series of the Garsia-Haiman module M_μ , a $\mathbb{Q}S_n$ -submodule of $\mathbb{Q}[x_1, \dots, x_n, y_1, \dots, y_n]$ of dimension $n!$.

Modified Macdonald polynomials

- Plethystically modified Macdonald polynomials

$$H_\mu(x; q, t) = J_\mu[x/(1-t); q, t].$$

- Macdonald positivity: the H_μ are Schur positive.
- $H_\mu(x; 1, 1) = (s_1)^n = h_{(1^n)}$.
- $t^{n(\mu)} H_\mu(x; q, t^{-1})$ = Frobenius series of the Garsia-Haiman module M_μ , a $\mathbb{Q}S_n$ -submodule of $\mathbb{Q}[x_1, \dots, x_n, y_1, \dots, y_n]$ of dimension $n!$.

$$H_{31} = ts_4 + (1 + qt + q^2t)s_{31} + (q + tq^2)s_{22} + (q + q^2 + q^3t)s_{211} + q^3s_{1111}$$


Haglund-Haiman-Loehr formula example

$$H_{\mu}(X; q, t) = t^{n(\mu)} \sum_R \left(\prod_{\substack{\square \\ u}} q^{a+1} t^l \right) \mathcal{G}_R(X; t^{-1})$$

Haglund-Haiman-Loehr formula example

$$H_{\mu}(X; q, t) = t^{n(\mu)} \sum_R \left(\prod_{\substack{u \\ \square}} q^{a+1} t^l \right) \mathcal{G}_R(X; t^{-1})$$


For $\mu = (3, 2)$, we get 8 LLT terms:



Haglund-Haiman-Loehr formula example

$$H_{\mu}(X; q, t) = t^{n(\mu)} \sum_R \left(\prod_{\substack{u \\ \square}} q^{a+1} t^l \right) \mathcal{G}_R(X; t^{-1})$$

For $\mu = (3, 2)$, we get 8 LLT terms:



Remark: Can permute the order of ribbons (but changes arm and leg statistics and LLTs.)

Nonsymmetric Macdonald polynomials

The Cherednik operators $Y_1, \dots, Y_N$ act on $\mathbb{Q}(q, t)[x_1^{\pm 1}, \dots, x_N^{\pm 1}]$.

$$T_i f = s_i f + (1 - t) x_i \frac{f - s_i f}{x_i - x_{i+1}}, \quad (\text{Demazure-Lusztig operators})$$

$$\Phi f = f(x_2, \dots, x_N, qx_1),$$

$$Y_i = t^{-i+1} T_{i-1} \cdots T_1 x_1 \Phi T_{N-1}^{-1} \cdots T_i^{-1}.$$

Definition

The *nonsymmetric Macdonald polynomials* $E_\alpha(x_1, \dots, x_N; q, t)$ are the joint eigenfunctions of the commuting operators $Y_1, \dots, Y_N$.

Nonsymmetric Macdonald polynomials

The Cherednik operators $Y_1, \dots, Y_N$ act on $\mathbb{Q}(q, t)[x_1^{\pm 1}, \dots, x_N^{\pm 1}]$.

$$T_i f = s_i f + (1 - t) x_i \frac{f - s_i f}{x_i - x_{i+1}}, \quad (\text{Demazure-Lusztig operators})$$

$$\Phi f = f(x_2, \dots, x_N, qx_1),$$

$$Y_i = t^{-i+1} T_{i-1} \cdots T_1 x_1 \Phi T_{N-1}^{-1} \cdots T_i^{-1}.$$

Definition

The *nonsymmetric Macdonald polynomials* $E_\alpha(x_1, \dots, x_N; q, t)$ are the joint eigenfunctions of the commuting operators $Y_1, \dots, Y_N$.

- $\{E_\alpha\}_{\alpha \in \mathbb{Z}^N}$ forms a basis for $\mathbb{Q}(q, t)[x_1^{\pm 1}, \dots, x_N^{\pm 1}]$.

Nonsymmetric Macdonald polynomials

The Cherednik operators $Y_1, \dots, Y_N$ act on $\mathbb{Q}(q, t)[x_1^{\pm 1}, \dots, x_N^{\pm 1}]$.

$$T_i f = s_i f + (1 - t) x_i \frac{f - s_i f}{x_i - x_{i+1}}, \quad (\text{Demazure-Lusztig operators})$$

$$\Phi f = f(x_2, \dots, x_N, qx_1),$$

$$Y_i = t^{-i+1} T_{i-1} \cdots T_1 x_1 \Phi T_{N-1}^{-1} \cdots T_i^{-1}.$$

Definition

The *nonsymmetric Macdonald polynomials* $E_\alpha(x_1, \dots, x_N; q, t)$ are the joint eigenfunctions of the commuting operators $Y_1, \dots, Y_N$.

- $\{E_\alpha\}_{\alpha \in \mathbb{Z}^N}$ forms a basis for $\mathbb{Q}(q, t)[x_1^{\pm 1}, \dots, x_N^{\pm 1}]$.
- Knop introduced *integral form nonsymmetric Macdonald polynomials* $\mathcal{E}_\alpha = c_\alpha E_\alpha$ with coefficients in $\mathbb{Z}[q, t]$.

Nonsymmetric Macdonald polynomials

The Cherednik operators $Y_1, \dots, Y_N$ act on $\mathbb{Q}(q, t)[x_1^{\pm 1}, \dots, x_N^{\pm 1}]$.

$$T_i f = s_i f + (1 - t) x_i \frac{f - s_i f}{x_i - x_{i+1}}, \quad (\text{Demazure-Lusztig operators})$$

$$\Phi f = f(x_2, \dots, x_N, qx_1),$$

$$Y_i = t^{-i+1} T_{i-1} \cdots T_1 x_1 \Phi T_{N-1}^{-1} \cdots T_i^{-1}.$$

Definition

The *nonsymmetric Macdonald polynomials* $E_\alpha(x_1, \dots, x_N; q, t)$ are the joint eigenfunctions of the commuting operators $Y_1, \dots, Y_N$.

- $\{E_\alpha\}_{\alpha \in \mathbb{Z}^N}$ forms a basis for $\mathbb{Q}(q, t)[x_1^{\pm 1}, \dots, x_N^{\pm 1}]$.
- Knop introduced *integral form nonsymmetric Macdonald polynomials* $\mathcal{E}_\alpha = c_\alpha E_\alpha$ with coefficients in $\mathbb{Z}[q, t]$.
- The $\mathcal{E}_\alpha$'s “Hecke symmetrize” to the integral form J_μ 's.

Nonsymmetric HHL formula example

$$\mathcal{E}_\alpha(x; q, t) = t^{n(\mu_+)} \sum_R \left(\prod_{\substack{u \\ \square}} q^{a+1} t^l \right) \mathcal{G}_{R, (1, 2, \dots, m)}^\pm(x; t^{-1}).$$

Nonsymmetric HHL formula example


$$\mathcal{E}_\alpha(x; q, t) = t^{n(\mu_+)} \sum_R \left(\prod_{\substack{\square \\ u}} q^{a+1} t^l \right) \mathcal{G}_{R, (1, 2, \dots, m)}^\pm(x; t^{-1}).$$

For $\alpha = (2, 0, 1)$, we get 8 *signed* flagged LLT terms:

Nonsymmetric HHL formula example

$$\mathcal{E}_\alpha(x; q, t) = t^{n(\mu_+)} \sum_R \left(\prod_{\substack{u \\ \square}} q^{a+1} t^l \right) \mathcal{G}_{R, (1, 2, \dots, m)}^\pm(x; t^{-1}).$$

For $\alpha = (2, 0, 1)$, we get 8 *signed* flagged LLT terms:



- Convention: bound boxes are always on the same content “diagonal.”
- Variation: bound boxes below shape bound shape strictly from below.

The Missing Corner

$$\begin{array}{ccc} \mathcal{E}_\alpha(x_1, \dots, x_N; q, t) & \xrightarrow{\quad} & ? \\ \text{Hecke sym.} \downarrow & & \downarrow \text{Weyl sym.} \\ J_\mu(x; q, t) & \xrightarrow{\text{Plethysm}} & H_\mu(x; q, t) \end{array}$$

Question

Is there a “nice” family of polynomials that Weyl symmetrizes to the modified Macdonald polynomials?

An important feature for “nice”? Positivity!

Toward positivity

- Knop (2007) formulated a positivity conjecture for a stable version of $\mathcal{E}_\alpha$ involving Kazhdan-Lusztig theory.
- Lapointe (2022) formulated another positivity conjecture for a stable version of $\mathcal{E}_\alpha$.
- Goodberry and Orr, Bechtloff Weising, and Bechtloff Weising and Orr consider stable versions of E_α .

Toward positivity

- Knop (2007) formulated a positivity conjecture for a stable version of $\mathcal{E}_\alpha$ involving Kazhdan-Lusztig theory.
- Lapointe (2022) formulated another positivity conjecture for a stable version of $\mathcal{E}_\alpha$.
- Goodberry and Orr, Bechtloff Weising, and Bechtloff Weising and Orr consider stable versions of E_α .
- We can understand stabilizations in terms of signed flagged LLT combinatorics.

Toward positivity

- Knop (2007) formulated a positivity conjecture for a stable version of $\mathcal{E}_\alpha$ involving Kazhdan-Lusztig theory.
- Lapointe (2022) formulated another positivity conjecture for a stable version of $\mathcal{E}_\alpha$.
- Goodberry and Orr, Bechtloff Weising, and Bechtloff Weising and Orr consider stable versions of E_α .
- We can understand stabilizations in terms of signed flagged LLT combinatorics.
- Upshot: look at stabilizations for “positivity.”

Filling in the missing corner

$$\mathcal{E}_\alpha(x_1, \dots, x_N; q, t) = \sum \text{signed flagged LLTs}$$

⋈ stabilize
↓

$$\mathcal{J}_{\eta|\lambda}(x; q, t) = \sum \text{signed flagged LLTs}$$

Hecke sym.
↓

$$J_{(\eta;\lambda)_+}(x; q, t) = \sum \text{signed LLTs}$$


→ ?

Weyl sym.
↓

$$H_{(\eta;\lambda)_+}(x; q, t) = \sum \text{LLTs}$$

Plethysm
→

Filling in the missing corner



Modern Macdonald polynomials

Definition (Blasiak-Haiman-Morse-Pun-S., 2025+)

The *modern Macdonald polynomial* indexed by $\eta \in \mathbb{N}^r$ and partition λ is

$$\text{nsH}_{\eta|\lambda}(x; q, t) = t^{n((\eta;\lambda)_+)} \sum_R \left(\prod_{\substack{u \\ \square}} q^{a+1} t^l \right) \mathcal{G}_{R,(1,2,\dots,r)}(x; t^{-1})$$

For $r = 2$, $\eta = (2, 1)$, $\lambda = \emptyset$, the flagged LLT terms are

$$\begin{array}{cc}
 \begin{array}{c} \square \square \\ \square \end{array} 1 & \begin{array}{c} \square \\ \square \end{array} 1 \\
 t & q t^2 \\
 \begin{array}{c} \square \\ 2 \end{array} & \begin{array}{c} \square \\ 2 \end{array}
 \end{array}$$

$$\text{nsH}_{21|\emptyset}(x_1, x_2, x_3; q, t) = t x_1^3 + x_1^2 x_2 + q t x_1^2 x_2 + q t x_1 x_2^2 + q t x_1^2 x_3 + q x_1 x_2 x_3$$

Modern Macdonald polynomials

- $nsH_{\eta|\lambda}(x; q, t)$ is monomial positive with a combinatorial formula.

Modern Macdonald polynomials

- $\text{nsH}_{\eta|\lambda}(x; q, t)$ is monomial positive with a combinatorial formula.
- $\text{nsH}_{\eta|\lambda}(x; q, t)$ Weyl symmetrizes to symmetric $H_{\beta_+}(x; q, t)$.

Modern Macdonald polynomials

- $\text{nsH}_{\eta|\lambda}(x; q, t)$ is monomial positive with a combinatorial formula.
- $\text{nsH}_{\eta|\lambda}(x; q, t)$ Weyl symmetrizes to symmetric $H_{\beta_+}(x; q, t)$.
- We conjecture $\text{nsH}_{\eta|\lambda}(x; q, t)$ is atom positive (would follow from flagged LLTs being atom positive).

Raising the flag

- Flagged LLT polynomials serve as a common combinatorial generalization of symmetric LLT polynomials and flagged Schur functions.

Raising the flag

- Flagged LLT polynomials serve as a common combinatorial generalization of symmetric LLT polynomials and flagged Schur functions.
- Flagged LLT polynomials and their signed variants provide a common framework for understanding various versions of (nonsymmetric) Macdonald polynomials.

Raising the flag

- Flagged LLT polynomials serve as a common combinatorial generalization of symmetric LLT polynomials and flagged Schur functions.
- Flagged LLT polynomials and their signed variants provide a common framework for understanding various versions of (nonsymmetric) Macdonald polynomials.
- Flagged LLT polynomials provide a tool to give nonsymmetric generalizations to the entire diagonal harmonics / shuffle theorem program.

Raising the flag

- Flagged LLT polynomials serve as a common combinatorial generalization of symmetric LLT polynomials and flagged Schur functions.
- Flagged LLT polynomials and their signed variants provide a common framework for understanding various versions of (nonsymmetric) Macdonald polynomials.
- Flagged LLT polynomials provide a tool to give nonsymmetric generalizations to the entire diagonal harmonics / shuffle theorem program.

Thank you!

Filling in the missing corner


- $\mathcal{P}(r) = \mathbb{Q}(q, t)[x_1, \dots, x_r] \otimes \Lambda_{\mathbb{Q}(q, t)}(x_{r+1}, \dots)$.
- Use a “stable” version of $\mathcal{E}_\alpha$, which we denote $\mathcal{J}_{\eta|\lambda}$.

$$\begin{array}{ccc}
 \mathcal{J}_{\eta|\lambda}(x; q, t) & \xrightarrow{\Pi_r} & \text{nsH}_{\eta|\lambda}(x; q, t) \\
 \downarrow \text{Hecke sym.} & & \downarrow \text{Weyl sym.} \\
 J_{(\eta; \lambda)_+}(x; q, t) & \xrightarrow{\text{Plethysm}} & H_{(\eta; \lambda)_+}(x; q, t)
 \end{array}$$

- $\mathcal{J}_{\eta|\lambda}(x; q, t)$ and $\text{nsH}_{\eta|\lambda}(x; q, t)$ for $(\eta|\lambda) \in \mathbb{N}^r \times \text{Par}$ form bases of $\mathcal{P}(r)$.
- $(\eta; \lambda)_+ =$ partition rearrangement of concatenating $(\eta; \lambda)$.
- Π_r is “ r -nonsymmetric plethysm”

$$\Pi_r(f(x_1, \dots, x_r) \otimes g(x)) = (\Pi_{t,x} f(x_1, \dots, x_r)) \otimes g[x/(1-t)] .$$

Arms and legs



$$a(u) = 2, l(u) = 3$$