

K -theoretic Catalan functions

George H. Seelinger (joint with J. Blasiak and J. Morse)

CAGE

ghs9ae@virginia.edu

6 February 2020

- Schubert calculus
- Catalan functions: a new approach to old problems
- K -theoretic Catalan functions

Overview of Schubert Calculus Combinatorics

Geometric problem

Find $c_{\lambda\mu}^\nu = \#$ of points in intersection of subvarieties in a variety X .

Overview of Schubert Calculus Combinatorics

Geometric problem

Find $c_{\lambda\mu}^\nu = \#$ of points in intersection of subvarieties in a variety X .



Cohomology

Schubert basis $\{\sigma_\lambda\}$ for $H^*(X)$ with property $\sigma_\lambda \cup \sigma_\mu = \sum_\nu c_{\lambda\mu}^\nu \sigma_\nu$

Overview of Schubert Calculus Combinatorics

Geometric problem

Find $c_{\lambda\mu}^\nu = \#$ of points in intersection of subvarieties in a variety X .



Cohomology

Schubert basis $\{\sigma_\lambda\}$ for $H^*(X)$ with property $\sigma_\lambda \cup \sigma_\mu = \sum_\nu c_{\lambda\mu}^\nu \sigma_\nu$



Representatives

Special basis of polynomials $\{f_\lambda\}$ such that $f_\lambda \cdot f_\mu = \sum_\nu c_{\lambda\mu}^\nu f_\nu$

Overview of Schubert Calculus Combinatorics (cont.)

Combinatorial study of $\{f_\lambda\}$ enlightens the geometry (and cohomology).

Goal

Identify $\{f_\lambda\}$ in explicit (simple) terms amenable to calculation and proofs.

Geometric problem

Find $c_{\lambda\mu}^\nu = \#$ of points in intersection of Schubert varieties $\{X_\lambda\}_{\lambda \subseteq (n^m)}$ in variety $X = \text{Gr}(m, n)$.

Classical Schubert Calculus

Geometric problem

Find $c_{\lambda\mu}^\nu = \#$ of points in intersection of Schubert varieties $\{X_\lambda\}_{\lambda \subseteq (n^m)}$ in variety $X = \text{Gr}(m, n)$.



Cohomology

Schubert basis $\{\sigma_\lambda\}_{\lambda \subseteq (n^m)}$ for $H^*(X)$ with property $\sigma_\lambda \cup \sigma_\mu = \sum_\nu c_{\lambda\mu}^\nu \sigma_\nu$

Classical Schubert Calculus

Geometric problem

Find $c_{\lambda\mu}^\nu = \#$ of points in intersection of Schubert varieties $\{X_\lambda\}_{\lambda \subseteq (n^m)}$ in variety $X = \text{Gr}(m, n)$.



Cohomology

Schubert basis $\{\sigma_\lambda\}_{\lambda \subseteq (n^m)}$ for $H^*(X)$ with property $\sigma_\lambda \cup \sigma_\mu = \sum_\nu c_{\lambda\mu}^\nu \sigma_\nu$



Representatives

Special basis of Schur polynomials $\{s_\lambda\}$ such that $s_\lambda \cdot s_\mu = \sum_\nu c_{\lambda\mu}^\nu s_\nu$ for Littlewood-Richardson coefficients $c_{\lambda\mu}^\nu$.

Next Step: Flag Variety

- $X = Fl_n(\mathbb{C}) = \{V_0 \subseteq V_1 \subseteq \cdots \subseteq V_n \mid \dim V_i = i\}$

Next Step: Flag Variety

- $X = Fl_n(\mathbb{C}) = \{V_0 \subseteq V_1 \subseteq \cdots \subseteq V_n \mid \dim V_i = i\}$
- Decomposes into Schubert varieties indexed by $w \in S_n$.

Next Step: Flag Variety

- $X = Fl_n(\mathbb{C}) = \{V_0 \subseteq V_1 \subseteq \cdots \subseteq V_n \mid \dim V_i = i\}$
- Decomposes into Schubert varieties indexed by $w \in S_n$.
- $H^*(Fl_n(\mathbb{C}))$ supported by Schubert polynomials $\mathfrak{S}_w \in \mathbb{Z}[x_1, \dots, x_n]$
(Not necessarily symmetric!)

Next Step: Flag Variety

- $X = Fl_n(\mathbb{C}) = \{V_0 \subseteq V_1 \subseteq \cdots \subseteq V_n \mid \dim V_i = i\}$
- Decomposes into Schubert varieties indexed by $w \in S_n$.
- $H^*(Fl_n(\mathbb{C}))$ supported by Schubert polynomials $\mathfrak{S}_w \in \mathbb{Z}[x_1, \dots, x_n]$
(Not necessarily symmetric!)

Open Problem

Structure constants $\mathfrak{S}_w \mathfrak{S}_u = c_{wu}^v \mathfrak{S}_v$ are combinatorially unknown.

There are many variations on classical Schubert calculus of the Grassmannian (Type A).

Schubert Calculus Variations

There are many variations on classical Schubert calculus of the Grassmannian (Type A).

Theory	f_λ
(Co)homology of Grassmannian	Schur functions
(Co)homology of flag variety	Schubert polynomials
Quantum cohomology of flag variety	Quantum Schuberts
(Co)homology of Types BCD Grassmannian	Schur- P and Q functions
(Co)homology of affine Grassmannian	(dual) k -Schur functions
K -theory of Grassmannian	Grothendieck polynomials
K -homology of affine Grassmannian	K - k -Schur functions

Schubert Calculus Variations

There are many variations on classical Schubert calculus of the Grassmannian (Type A).

Theory	f_λ
(Co)homology of Grassmannian	Schur functions
(Co)homology of flag variety	Schubert polynomials
Quantum cohomology of flag variety	Quantum Schuberts
(Co)homology of Types BCD Grassmannian	Schur- P and Q functions
(Co)homology of affine Grassmannian	(dual) k -Schur functions
K -theory of Grassmannian	Grothendieck polynomials
K -homology of affine Grassmannian	K - k -Schur functions

And many more!

- $QH^*(Fl_{k+1})$ quantum deformation of $H^*(Fl_{k+1})$.

Peterson Isomorphism

- $QH^*(Fl_{k+1})$ quantum deformation of $H^*(Fl_{k+1})$.
- Supported by quantum Schubert polynomials $\mathfrak{S}_w^Q$.

Peterson Isomorphism

- $QH^*(Fl_{k+1})$ quantum deformation of $H^*(Fl_{k+1})$.
- Supported by quantum Schubert polynomials $\mathfrak{S}_w^Q$.
- Peterson isomorphism

$$\Phi: QH^*(Fl_{k+1}) \rightarrow H_*(Gr_{SL_{k+1}})_{loc}$$

$$\mathfrak{S}_w^Q \mapsto \frac{s_\lambda^{(k)}}{\prod_{i \in Des(w)} \tau_i}$$

where $s_\lambda^{(k)}$ is a k -Schur function.

Peterson Isomorphism

- $QH^*(Fl_{k+1})$ quantum deformation of $H^*(Fl_{k+1})$.
- Supported by quantum Schubert polynomials $\mathfrak{S}_w^Q$.
- Peterson isomorphism

$$\Phi: QH^*(Fl_{k+1}) \rightarrow H_*(Gr_{SL_{k+1}})_{loc}$$

$$\mathfrak{S}_w^Q \mapsto \frac{s_\lambda^{(k)}}{\prod_{i \in Des(w)} \tau_i}$$

where $s_\lambda^{(k)}$ is a k -Schur function.

Upshot

Computations for (quantum) Schubert polynomials can be moved into symmetric functions.

k -Schur functions

- $s_{\lambda}^{(k)}$ for $\lambda_1 \leq k$ a basis for $\mathbb{Z}[s_1, s_2, \dots, s_k]$ (Lapointe et al., 2003).

k -Schur functions

- $s_{\lambda}^{(k)}$ for $\lambda_1 \leq k$ a basis for $\mathbb{Z}[s_1, s_2, \dots, s_k]$ (Lapointe et al., 2003).
- Schubert representatives for $H_*(Gr_{SL_{k+1}})$ (Lam, 2008).

k -Schur functions

- $s_{\lambda}^{(k)}$ for $\lambda_1 \leq k$ a basis for $\mathbb{Z}[s_1, s_2, \dots, s_k]$ (Lapointe et al., 2003).
- Schubert representatives for $H_*(Gr_{SL_{k+1}})$ (Lam, 2008).
- Has a tableaux formulation and Pieri rule: $s_1 s_{\lambda}^{(k)} = \sum_{\mu} a_{\lambda\mu} s_{\mu}^{(k)}$

k -Schur functions

- $s_{\lambda}^{(k)}$ for $\lambda_1 \leq k$ a basis for $\mathbb{Z}[s_1, s_2, \dots, s_k]$ (Lapointe et al., 2003).
- Schubert representatives for $H_*(Gr_{SL_{k+1}})$ (Lam, 2008).
- Has a tableaux formulation and Pieri rule: $s_1 s_{\lambda}^{(k)} = \sum_{\mu} a_{\lambda\mu} s_{\mu}^{(k)}$
- $s_{\lambda}^{(k)} = s_{\lambda}$ as $k \rightarrow \infty$.

k -Schur functions

- $s_{\lambda}^{(k)}$ for $\lambda_1 \leq k$ a basis for $\mathbb{Z}[s_1, s_2, \dots, s_k]$ (Lapointe et al., 2003).
- Schubert representatives for $H_*(Gr_{SL_{k+1}})$ (Lam, 2008).
- Has a tableaux formulation and Pieri rule: $s_1 s_{\lambda}^{(k)} = \sum_{\mu} a_{\lambda\mu} s_{\mu}^{(k)}$
- $s_{\lambda}^{(k)} = s_{\lambda}$ as $k \rightarrow \infty$.
- Branching with positive coefficients (Lam et al., 2010):

$$s_{\lambda}^{(2)} = s_{\lambda} + s_{\lambda} + s_{\lambda}$$

The diagram illustrates the branching of the 2-part Schur function $s_{\lambda}^{(2)}$ into three 3-part Schur functions $s_{\lambda}^{(3)}$. On the left, $s_{\lambda}^{(2)}$ is represented by a Young diagram with two rows of two boxes each. This is equal to the sum of three Young diagrams, each representing $s_{\lambda}^{(3)}$. The first diagram has two rows of two boxes. The second and third diagrams have a first row of two boxes and a second row of three boxes. Brackets are used to group the three diagrams on the right under the label $s_{\lambda}^{(3)}$.

k -Schur functions

- $s_{\lambda}^{(k)}$ for $\lambda_1 \leq k$ a basis for $\mathbb{Z}[s_1, s_2, \dots, s_k]$ (Lapointe et al., 2003).
- Schubert representatives for $H_*(Gr_{SL_{k+1}})$ (Lam, 2008).
- Has a tableaux formulation and Pieri rule: $s_1 s_{\lambda}^{(k)} = \sum_{\mu} a_{\lambda\mu} s_{\mu}^{(k)}$
- $s_{\lambda}^{(k)} = s_{\lambda}$ as $k \rightarrow \infty$.
- Branching with positive coefficients (Lam et al., 2010):

$$s_{\lambda}^{(2)} = s_{\lambda} + s_{\lambda} + s_{\lambda}$$

The diagram illustrates the branching rule for $k=2$. The left side shows $s_{\lambda}^{(2)}$ for $\lambda = (2, 2)$, represented by a 2x2 grid of boxes. The right side shows the sum of three terms, each representing a $s_{\lambda}^{(3)}$ for $\lambda = (2, 2)$. The first term is $s_{\lambda}^{(3)}$ for $\lambda = (2, 2)$, represented by a 2x2 grid of boxes. The second and third terms are $s_{\lambda}^{(3)}$ for $\lambda = (2, 2)$, represented by a 2x2 grid of boxes. The diagram uses curly braces to group the terms on the right side, indicating that they are all $s_{\lambda}^{(3)}$ for $\lambda = (2, 2)$.

- Has geometric interpretation.

k -Schur functions

- $s_{\lambda}^{(k)}$ for $\lambda_1 \leq k$ a basis for $\mathbb{Z}[s_1, s_2, \dots, s_k]$ (Lapointe et al., 2003).
- Schubert representatives for $H_*(Gr_{SL_{k+1}})$ (Lam, 2008).
- Has a tableaux formulation and Pieri rule: $s_1^r s_{\lambda}^{(k)} = \sum_{\mu} a_{\lambda\mu} s_{\mu}^{(k)}$
- $s_{\lambda}^{(k)} = s_{\lambda}$ as $k \rightarrow \infty$.
- Branching with positive coefficients (Lam et al., 2010):

$$s_{\lambda}^{(2)} = s_{\lambda}^{(3)} + s_{\mu}^{(3)} + s_{\nu}^{(3)}$$

- Has geometric interpretation.
- No combinatorial interpretation of branching coefficients.

k -Schur functions

- $s_{\lambda}^{(k)}$ for $\lambda_1 \leq k$ a basis for $\mathbb{Z}[s_1, s_2, \dots, s_k]$ (Lapointe et al., 2003).
- Schubert representatives for $H_*(Gr_{SL_{k+1}})$ (Lam, 2008).
- Has a tableaux formulation and Pieri rule: $s_1^r s_{\lambda}^{(k)} = \sum_{\mu} a_{\lambda\mu} s_{\mu}^{(k)}$
- $s_{\lambda}^{(k)} = s_{\lambda}$ as $k \rightarrow \infty$.
- Branching with positive coefficients (Lam et al., 2010):

$$s_{\lambda}^{(2)} = s_{\lambda} + s_{\mu} + s_{\nu}$$

The diagram shows the branching of the 2-partition $s_{(2)}^{(2)}$ into three 3-partitions. On the left is a 2x2 square labeled $s_{(2)}^{(2)}$. On the right is the sum of three partitions: a 2x2 square labeled $s_{(2)}^{(3)}$, a 2x1 rectangle labeled $s_{(1,1)}^{(3)}$, and a 1x3 horizontal row labeled $s_{(1,1,1)}^{(3)}$. Brackets group the three partitions on the right under the label $s_{(2)}^{(3)}$.

- Has geometric interpretation.
- No combinatorial interpretation of branching coefficients.
- Definition with t important for Macdonald polynomials.

k -Schur functions

- $s_{\lambda}^{(k)}$ for $\lambda_1 \leq k$ a basis for $\mathbb{Z}[s_1, s_2, \dots, s_k]$ (Lapointe et al., 2003).
- Schubert representatives for $H_*(Gr_{SL_{k+1}})$ (Lam, 2008).
- Has a tableaux formulation and Pieri rule: $s_1^r s_{\lambda}^{(k)} = \sum_{\mu} a_{\lambda\mu} s_{\mu}^{(k)}$
- $s_{\lambda}^{(k)} = s_{\lambda}$ as $k \rightarrow \infty$.
- Branching with positive coefficients (Lam et al., 2010):

$$s_{\lambda}^{(2)} = s_{\lambda}^{(3)} + s_{\lambda}^{(3)}$$

The diagram shows the branching of the Schur function $s_{(2)}^{(2)}$ into two components. On the left, $s_{(2)}^{(2)}$ is represented by a Young diagram with two rows of two boxes. This is equal to the sum of two terms. The first term is $s_{(2)}^{(3)}$, represented by a Young diagram with two rows of two boxes, with a brace underneath labeled $s_{(2)}^{(3)}$. The second term is $s_{(1,1)}^{(3)}$, represented by a Young diagram with two rows of one box each, with a brace underneath labeled $s_{(1,1)}^{(3)}$.

- Has geometric interpretation.
- No combinatorial interpretation of branching coefficients.
- Definition with t important for Macdonald polynomials.
- Many definitions. A new one makes proofs easier!

- Schubert calculus
- **Catalan functions: a new approach to old problems**
- K -theoretic Catalan functions

Raising Operators on Symmetric Functions

- Raising operators $R_{i,j}$ act on diagrams

$$R_{1,3} \left(\begin{array}{|c|c|c|} \hline \text{red} & & \\ \hline & & \\ \hline & & \\ \hline \end{array} \right) = \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \text{red} \\ \hline & & & \\ \hline \end{array} \quad R_{2,3} \left(\begin{array}{|c|} \hline \text{red} \\ \hline \\ \hline \\ \hline \end{array} \right) = \begin{array}{|c|c|} \hline & \text{red} \\ \hline & \\ \hline \end{array}$$

Raising Operators on Symmetric Functions

- Raising operators $R_{i,j}$ act on diagrams

$$R_{1,3} \left(\begin{array}{|c|c|c|} \hline \text{red} & & \\ \hline & & \\ \hline & & \\ \hline \end{array} \right) = \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \text{red} \\ \hline & & & \\ \hline \end{array} \quad R_{2,3} \left(\begin{array}{|c|} \hline \text{red} \\ \hline \\ \hline \\ \hline \end{array} \right) = \begin{array}{|c|c|} \hline & \text{red} \\ \hline & \\ \hline \end{array}$$

- Extend action to a symmetric function f_λ by $R_{i,j}(f_\lambda) = f_{\lambda + \epsilon_i - \epsilon_j}$.

Raising Operators on Symmetric Functions

- Raising operators $R_{i,j}$ act on diagrams

$$R_{1,3} \left(\begin{array}{|c|c|c|} \hline \text{red} & & \\ \hline & & \\ \hline & & \\ \hline \end{array} \right) = \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \text{red} \\ \hline & & & \\ \hline \end{array} \quad R_{2,3} \left(\begin{array}{|c|} \hline \text{red} \\ \hline \\ \hline \\ \hline \end{array} \right) = \begin{array}{|c|c|} \hline & \text{red} \\ \hline & \\ \hline \end{array}$$

- Extend action to a symmetric function f_λ by $R_{i,j}(f_\lambda) = f_{\lambda + \epsilon_i - \epsilon_j}$.
- For $h_\lambda = s_{\lambda_1} \cdots s_{\lambda_r}$, we have the *Jacobi-Trudi identity*

$$s_\lambda = \prod_{i < j} (1 - R_{ij}) h_\lambda$$

Raising Operators on Symmetric Functions

- Raising operators $R_{i,j}$ act on diagrams

$$R_{1,3} \left(\begin{array}{|c|c|c|c|} \hline \text{red} & & & \\ \hline & & & \\ \hline & & & \\ \hline \end{array} \right) = \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \text{red} \\ \hline & & & \\ \hline \end{array} \quad R_{2,3} \left(\begin{array}{|c|} \hline \text{red} \\ \hline \\ \hline \\ \hline \end{array} \right) = \begin{array}{|c|c|} \hline & \text{red} \\ \hline & \\ \hline & \\ \hline \end{array}$$

- Extend action to a symmetric function f_λ by $R_{i,j}(f_\lambda) = f_{\lambda+\epsilon_i-\epsilon_j}$.
- For $h_\lambda = s_{\lambda_1} \cdots s_{\lambda_r}$, we have the *Jacobi-Trudi identity*

$$s_\lambda = \prod_{i < j} (1 - R_{ij}) h_\lambda$$

$$s_{22} = (1 - R_{12}) h_{22} = h_{22} - h_{31}$$

$$s_{211} = (1 - R_{12})(1 - R_{23})(1 - R_{13}) h_{211}$$

$$= h_{211} - h_{301} - h_{220} - \text{red } h_{310} + \text{red } h_{310} + \underbrace{h_{32-1}}_{=0} + h_{400} - \underbrace{h_{41-1}}_{=0}$$

some terms cancel

Raising Operators on Symmetric Functions

Gives definition for Schur function indexed by any integer vector $\alpha \in \mathbb{Z}^\ell$.

Raising Operators on Symmetric Functions

Gives definition for Schur function indexed by any integer vector $\alpha \in \mathbb{Z}^\ell$.

Straightening:

$$s_\alpha = \prod_{i < j} (1 - R_{ij}) h_\alpha = \begin{cases} \pm s_\lambda & \text{for a partition } \lambda \\ 0 & \end{cases}$$

Raising Operators on Symmetric Functions

Gives definition for Schur function indexed by any integer vector $\alpha \in \mathbb{Z}^\ell$.

Straightening:

$$s_\alpha = \prod_{i < j} (1 - R_{ij}) h_\alpha = \begin{cases} \pm s_\lambda & \text{for a partition } \lambda \\ 0 & \end{cases}$$


For $\langle s_{1^r}^\perp s_\lambda, s_\mu \rangle = \langle s_\lambda, s_{1^r} s_\mu \rangle$,

$$s_{1^r}^\perp s_\lambda = \sum_{S \subseteq [1, \ell], |S|=r} s_{\lambda - \epsilon_S}$$

$$s_{1^2}^\perp s_{333} = s_{322} + s_{232} + s_{223}$$

Root Ideals


A root ideal Ψ of type $A_{\ell-1}$ positive roots: given by Dyck path (lattice path above diagonal).



$\Psi =$ Roots above Dyck path
 $\Delta_{\ell}^{+} \setminus \Psi =$ Non-roots below

Root Ideals

A root ideal Ψ of type $A_{\ell-1}$ positive roots: given by Dyck path (lattice path above diagonal).



Ψ = Roots above Dyck path
 $\Delta_{\ell}^+ \setminus \Psi$ = Non-roots below


Catalan Function (Chen, 2010; Panyushev, 2010; Blasiak et al., 2019)

For Ψ and $\gamma \in \mathbb{Z}^{\ell}$

$$H(\Psi; \gamma)(x) = \prod_{(i,j) \in \Delta_{\ell}^+ \setminus \Psi} (1 - R_{ij}) h_{\gamma}(x)$$

Root Ideals

A root ideal Ψ of type $A_{\ell-1}$ positive roots: given by Dyck path (lattice path above diagonal).



Ψ = Roots above Dyck path
 $\Delta_{\ell}^{+} \setminus \Psi$ = Non-roots below

Catalan Function (Chen, 2010; Panyushev, 2010; Blasiak et al., 2019)


For Ψ and $\gamma \in \mathbb{Z}^{\ell}$

$$H(\Psi; \gamma)(x) = \prod_{(i,j) \in \Delta_{\ell}^{+} \setminus \Psi} (1 - R_{ij}) h_{\gamma}(x)$$

- $\Psi = \emptyset \implies H(\emptyset; \gamma) = s_{\gamma}$

Root Ideals

A root ideal Ψ of type $A_{\ell-1}$ positive roots: given by Dyck path (lattice path above diagonal).



Ψ = Roots above Dyck path
 $\Delta_{\ell}^{+} \setminus \Psi$ = Non-roots below

Catalan Function (Chen, 2010; Panyushev, 2010; Blasiak et al., 2019)

For Ψ and $\gamma \in \mathbb{Z}^{\ell}$

$$H(\Psi; \gamma)(x) = \prod_{(i,j) \in \Delta_{\ell}^{+} \setminus \Psi} (1 - R_{ij}) h_{\gamma}(x)$$

- $\Psi = \emptyset \implies H(\emptyset; \gamma) = s_{\gamma}$
- $\Psi = \text{all roots} \implies H(\Psi; \gamma) = h_{\gamma}$

k -Schur root ideal for λ

$$\begin{aligned}\psi = \Delta^k(\lambda) &= \{(i, j) : j > k - \lambda_i\} \\ &= \text{root ideal with } k - \lambda_i \text{ non-roots in row } i\end{aligned}$$

k -Schur root ideal for λ

$$\begin{aligned}\psi = \Delta^k(\lambda) &= \{(i, j) : j > k - \lambda_i\} \\ &= \text{root ideal with } k - \lambda_i \text{ non-roots in row } i\end{aligned}$$

$$\Delta^4(3, 3, 2, 2, 1, 1) =$$

3					
	3				
		2			
			2		
				1	
					1

$\leftarrow$ row i has $4 - \lambda_i$ non-roots

Catalan functions

k -Schur root ideal for λ

$$\begin{aligned}\psi = \Delta^k(\lambda) &= \{(i, j) : j > k - \lambda_i\} \\ &= \text{root ideal with } k - \lambda_i \text{ non-roots in row } i\end{aligned}$$

$$\Delta^4(3, 3, 2, 2, 1, 1) =$$

3					
	3				
		2			
			2		
				1	
					1

$\leftarrow$ row i has $4 - \lambda_i$ non-roots

k -Schur is a Catalan function (Blasiak et al., 2019).

For partition λ with $\lambda_1 \leq k$,

$$s_{\lambda}^{(k)} = H(\Delta^k(\lambda); \lambda).$$

Key ingredient of branching proof

Dual vertical Pieri rule: $s_{1^r}^\perp s_\lambda^{(k)} = \sum_\mu a_{\lambda\mu} s_\mu^{(k)}$ for $\langle s_{1^r}^\perp f, g \rangle = \langle f, s_{1^r} g \rangle$.

Key ingredient of branching proof

Dual vertical Pieri rule: $s_{1^r}^\perp s_\lambda^{(k)} = \sum_\mu a_{\lambda\mu} s_\mu^{(k)}$ for $\langle s_{1^r}^\perp f, g \rangle = \langle f, s_{1^r} g \rangle$.

Shift Invariance (Blasiak et al., 2019)

For partition λ of length ℓ with $\lambda_1 \leq k$,

$$s_{1^\ell}^\perp s_{\lambda+1^\ell}^{(k+1)} = s_\lambda^{(k)}.$$

Key ingredient of branching proof

Dual vertical Pieri rule: $s_{1^r}^\perp s_\lambda^{(k)} = \sum_\mu a_{\lambda\mu} s_\mu^{(k)}$ for $\langle s_{1^r}^\perp f, g \rangle = \langle f, s_{1^r} g \rangle$.

Shift Invariance (Blasiak et al., 2019)

For partition λ of length ℓ with $\lambda_1 \leq k$,

$$s_{1^\ell}^\perp s_{\lambda+1^\ell}^{(k+1)} = s_\lambda^{(k)}.$$

Proof: $k - \lambda_i = (k + 1) - (\lambda_i + 1)$

Key ingredient of branching proof

Dual vertical Pieri rule: $s_{1^r}^\perp s_\lambda^{(k)} = \sum_\mu a_{\lambda\mu} s_\mu^{(k)}$ for $\langle s_{1^r}^\perp f, g \rangle = \langle f, s_{1^r} g \rangle$.

Shift Invariance (Blasiak et al., 2019)

For partition λ of length ℓ with $\lambda_1 \leq k$,

$$s_{1^\ell}^\perp s_{\lambda+1^\ell}^{(k+1)} = s_\lambda^{(k)}.$$

Proof: $k - \lambda_i = (k + 1) - (\lambda_i + 1)$

$$\Delta^4(3, 3, 2, 2, 1, 1) =$$

3					
	3				
		2			
			2		
				1	
					1

$$\Delta^5(4, 4, 3, 3, 2, 2) =$$

4					
	4				
		3			
			3		
				2	
					2

Key ingredient of branching proof


Dual vertical Pieri rule: $s_{1^r}^\perp s_\lambda^{(k)} = \sum_\mu a_{\lambda\mu} s_\mu^{(k)}$ for $\langle s_{1^r}^\perp f, g \rangle = \langle f, s_{1^r} g \rangle$.

Shift Invariance (Blasiak et al., 2019)

For partition λ of length ℓ with $\lambda_1 \leq k$,

$$s_{1^\ell}^\perp s_{\lambda+1^\ell}^{(k+1)} = s_\lambda^{(k)}.$$

Proof: $k - \lambda_i = (k + 1) - (\lambda_i + 1)$

$$\Delta^4(3, 3, 2, 2, 1, 1) =$$


$$\Delta^5(4, 4, 3, 3, 2, 2) =$$


Pieri:

$$s_{1^\ell}^\perp s_{\lambda+1^\ell}^{(k+1)} = \sum_\mu a_{\lambda+1^\ell, \mu} s_\mu^{(k+1)}$$

Key ingredient of branching proof


Dual vertical Pieri rule: $s_{1^r}^\perp s_\lambda^{(k)} = \sum_\mu a_{\lambda\mu} s_\mu^{(k)}$ for $\langle s_{1^r}^\perp f, g \rangle = \langle f, s_{1^r} g \rangle$.


Shift Invariance (Blasiak et al., 2019)

For partition λ of length ℓ with $\lambda_1 \leq k$,

$$s_{1^\ell}^\perp s_{\lambda+1^\ell}^{(k+1)} = s_\lambda^{(k)}.$$

Proof: $k - \lambda_i = (k + 1) - (\lambda_i + 1)$

$$\Delta^4(3, 3, 2, 2, 1, 1) =$$


$$\Delta^5(4, 4, 3, 3, 2, 2) =$$


Branching is a special case of Pieri:

$$s_\lambda^{(k)} = s_{1^\ell}^\perp s_{\lambda+1^\ell}^{(k+1)} = \sum_\mu a_{\lambda+1^\ell, \mu} s_\mu^{(k+1)}$$

- Schubert calculus
- Catalan functions: a new approach to old problems
- ***K*-theoretic Catalan functions**

Dual Grothendieck polynomials

- Inhomogeneous basis: $g_\lambda = s_\lambda + \text{lower degree terms}$.

Dual Grothendieck polynomials

- Inhomogeneous basis: $g_\lambda = s_\lambda +$ lower degree terms.
- Satisfies Pieri rule on “set-valued strips”

Dual Grothendieck polynomials

- Inhomogeneous basis: $g_\lambda = s_\lambda + \text{lower degree terms}$.
- Satisfies Pieri rule on “set-valued strips”

$$g_{1^2}g_{3,2} = g_{43} + g_{421} + g_{331} - g_{42} - g_{33} - 2g_{321} + g_{31}$$



Dual Grothendieck polynomials

- Inhomogeneous basis: $g_\lambda = s_\lambda + \text{lower degree terms}$.
- Satisfies Pieri rule on “set-valued strips”

$$g_{1^2}g_{3,2} = g_{43} + g_{421} + g_{331} - g_{42} - g_{33} - 2g_{321} + g_{31}$$



- $g_\lambda = \prod_{i < j} (1 - R_{ij}) k_\lambda$ for k_λ and inhomogeneous analogue of h_λ .

Dual Grothendieck polynomials

- Inhomogeneous basis: $g_\lambda = s_\lambda + \text{lower degree terms}$.
- Satisfies Pieri rule on “set-valued strips”

$$g_{1^2}g_{3,2} = g_{43} + g_{421} + g_{331} - g_{42} - g_{33} - 2g_{321} + g_{31}$$



- $g_\lambda = \prod_{i < j} (1 - R_{ij}) k_\lambda$ for k_λ and inhomogeneous analogue of h_λ .
- Dual to Grothendieck polynomials G_λ : Schubert representatives for $K^*(Gr(m, n))$

K - k -Schur functions

- Inhomogeneous basis: $g_{\lambda}^{(k)} = s_{\lambda}^{(k)} + \text{lower degree terms}$

K - k -Schur functions

- Inhomogeneous basis: $g_{\lambda}^{(k)} = s_{\lambda}^{(k)} + \text{lower degree terms}$
- Satisfies Pieri rule on “affine set-valued strips”

K - k -Schur functions

- Inhomogeneous basis: $g_{\lambda}^{(k)} = s_{\lambda}^{(k)} + \text{lower degree terms}$
- Satisfies Pieri rule on “affine set-valued strips”

$$g_1 g_{211}^{(2)} = g_{2111}^{(2)} - 2g_{211}^{(2)}$$

2-bounded partitions $\leftrightarrow$ 3-cores

K - k -Schur functions

- Inhomogeneous basis: $g_{\lambda}^{(k)} = s_{\lambda}^{(k)} + \text{lower degree terms}$
- Satisfies Pieri rule on “affine set-valued strips”

$$g_1 g_{211}^{(2)} = g_{2111}^{(2)} - 2g_{211}^{(2)}$$

2-bounded partitions $\leftrightarrow$ 3-cores

The diagram illustrates the Pieri rule for K -Schur functions. It shows the product of a 1-strip (represented by a 5x5 grid of colored dots) and a 2-bounded partition (represented by a 5x5 grid of colored dots). The result is the difference of two 2-bounded partitions (represented by 5x5 grids of colored dots). The partitions are labeled as 2-bounded partitions $\leftrightarrow$ 3-cores.

- Conjecture: $g_{\lambda}^{(k)}$ have positive branching into $g_{\mu}^{(k+1)}$ (Lam et al., 2010; Morse, 2011).

K - k -Schur functions

- Inhomogeneous basis: $g_{\lambda}^{(k)} = s_{\lambda}^{(k)} + \text{lower degree terms}$
- Satisfies Pieri rule on “affine set-valued strips”

$$g_1 g_{211}^{(2)} = g_{2111}^{(2)} - 2g_{211}^{(2)}$$

2-bounded partitions $\leftrightarrow$ 3-cores

- Conjecture: $g_{\lambda}^{(k)}$ have positive branching into $g_{\mu}^{(k+1)}$ (Lam et al., 2010; Morse, 2011).

Problem

No direct formula for $g_{\lambda}^{(k)}$

An Extra Ingredient: Lowering Operators

Lowering Operators $L_j(f_\lambda) = f_{\lambda - \epsilon_j}$

$$L_3 \left(\begin{array}{|c|c|c|c|} \hline \text{red} & & & \\ \hline & & & \\ \hline & & & \\ \hline \end{array} \right) = \begin{array}{|c|c|c|} \hline & & \\ \hline & & \\ \hline \end{array}, \quad L_1 \left(\begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline & & \text{red} & \\ \hline \end{array} \right) = \begin{array}{|c|c|} \hline & \\ \hline & \\ \hline & \\ \hline \end{array}$$

K -theoretic Catalan function

Let $\Psi, \mathcal{L} \subseteq \Delta_\ell^+$ be order ideals of positive roots and $\gamma \in \mathbb{Z}^\ell$, then

$$K(\Psi; \mathcal{L}; \gamma) := \prod_{(i,j) \in \mathcal{L}} (1 - L_j) \prod_{(i,j) \in \Delta_\ell^+ \setminus \Psi} (1 - R_{ij}) k_\gamma$$

Affine K -Theory Representatives with Raising Operators

K -theoretic Catalan function

Let $\Psi, \mathcal{L} \subseteq \Delta_\ell^+$ be order ideals of positive roots and $\gamma \in \mathbb{Z}^\ell$, then

$$K(\Psi; \mathcal{L}; \gamma) := \prod_{(i,j) \in \mathcal{L}} (1 - L_j) \prod_{(i,j) \in \Delta_\ell^+ \setminus \Psi} (1 - R_{ij}) k_\gamma$$

Example

non-roots of Ψ , roots of $\mathcal{L}$

	(12)	(13)	(14)	(15)
		(23)	(24)	(25)
			(34)	(35)
				(45)

$$\begin{aligned} K(\Psi; \mathcal{L}; 54332) \\ = (1 - L_4)^2 (1 - L_5)^2 (1 - R_{12}) (1 - R_{34}) (1 - R_{45}) k_{54332} \end{aligned}$$

Answer (Blasiak-Morse-S., 2020)

Answer (Blasiak-Morse-S., 2020)


For K -homology of affine Grassmannian, $g_{\lambda}^{(k)} = K(\Delta^k(\lambda); \Delta^{k+1}(\lambda); \lambda)$ since this family satisfies the Pieri rule.

Affine K -Theory Representatives with Raising Operators

Answer (Blasiak-Morse-S., 2020)

For K -homology of affine Grassmannian, $g_{\lambda}^{(k)} = K(\Delta^k(\lambda); \Delta^{k+1}(\lambda); \lambda)$ since this family satisfies the Pieri rule.

Example



$$\Delta_9^+ / \Delta^4(332111111), \Delta^5(332111111)$$

Theorem (Blasiak-Morse-S., 2020)

Theorem (Blasiak-Morse-S., 2020)

The $g_{\lambda}^{(k)}$ are “shift invariant”, i.e. for $\ell = \ell(\lambda)$

$$G_{1^{\ell}}^{\perp} g_{\lambda+1^{\ell}}^{(k+1)} = g_{\lambda}^{(k)}$$

Theorem (Blasiak-Morse-S., 2020)

The $g_\lambda^{(k)}$ are “shift invariant”, i.e. for $\ell = \ell(\lambda)$

$$G_{1^\ell}^\perp g_{\lambda+1^\ell}^{(k+1)} = g_\lambda^{(k)}$$

Theorem (Blasiak-Morse-S., 2020)

The branching coefficients in

$$g_\lambda^{(k)} = \sum_{\mu} a_{\lambda\mu} g_\mu^{(k+1)}$$

satisfy $(-1)^{|\lambda|-|\mu|} a_{\lambda\mu} \in \mathbb{Z}_{\geq 0}$.

For $G_{\lambda}^{(k)}$ an affine Grothendieck polynomial (dual to $g_{\lambda}^{(k)}$),

For $G_\lambda^{(k)}$ an affine Grothendieck polynomial (dual to $g_\lambda^{(k)}$),

- ① Combinatorially describe dual Pieri rule:

$$G_{1^r}^\perp g_\lambda^{(k)} = \sum_\mu ?? g_\mu^{(k)} \iff G_{1^r} G_\mu^{(k)} = \sum_\lambda ?? G_\lambda^{(k)}, \quad 1 \leq r \leq k.$$

For $G_\lambda^{(k)}$ an affine Grothendieck polynomial (dual to $g_\lambda^{(k)}$),

- 1 Combinatorially describe dual Pieri rule:

$$G_{1^r}^\perp g_\lambda^{(k)} = \sum_\mu ?? g_\mu^{(k)} \iff G_{1^r} G_\mu^{(k)} = \sum_\lambda ?? G_\lambda^{(k)}, \quad 1 \leq r \leq k.$$

- 2 Combinatorially describe branching coefficients: $g_\lambda^{(k)} = \sum_\mu ?? g_\mu^{(k+1)}$.

For $G_\lambda^{(k)}$ an affine Grothendieck polynomial (dual to $g_\lambda^{(k)}$),

- 1 Combinatorially describe dual Pieri rule:

$$G_{1^r}^\perp g_\lambda^{(k)} = \sum_\mu ?? g_\mu^{(k)} \iff G_{1^r} G_\mu^{(k)} = \sum_\lambda ?? G_\lambda^{(k)}, \quad 1 \leq r \leq k.$$

- 2 Combinatorially describe branching coefficients: $g_\lambda^{(k)} = \sum_\mu ?? g_\mu^{(k+1)}$.

- 3 Combinatorially describe $g_\lambda^{(k)} = \sum_\mu ?? s_\mu^{(k)}$.

Thank you!

- Anderson, David, Linda Chen, and Hsian-Hua Tseng. 2017. *On the quantum K -ring of the flag manifold*, preprint. arXiv: 1711.08414.
- Blasiak, Jonah, Jennifer Morse, Anna Pun, and Daniel Summers. 2019. *Catalan Functions and k -Schur Positivity*, J. Amer. Math. Soc. **32**, no. 4, 921–963.
- Chen, Li-Chung. 2010. *Skew-linked partitions and a representation theoretic model for k -Schur functions*, Ph.D. thesis.
- Ikedo, Takeshi, Shinsuke Iwao, and Toshiaki Maeno. 2018. *Peterson Isomorphism in K -theory and Relativistic Toda Lattice*, preprint. arXiv: 1703.08664.
- Lam, Thomas. 2008. *Schubert polynomials for the affine Grassmannian*, J. Amer. Math. Soc. **21**, no. 1, 259–281.
- Lam, Thomas, Luc Lapointe, Jennifer Morse, and Mark Shimozono. 2010. *Affine insertion and Pieri rules for the affine Grassmannian*, Mem. Amer. Math. Soc. **208**, no. 977.
- Lam, Thomas, Anne Schilling, and Mark Shimozono. 2010. *K -theory Schubert calculus of the affine Grassmannian*, Compositio Math. **146**, 811–852.
- Lapointe, Luc, Alain Lascoux, and Jennifer Morse. 2003. *Tableau atoms and a new Macdonald positivity conjecture*, Duke Mathematical Journal **116**, no. 1, 103–146.
- Morse, Jennifer. 2011. *Combinatorics of the K -theory of affine Grassmannians*, Advances in Mathematics.
- Panyushev, Dmitri I. 2010. *Generalised Kostka-Foulkes polynomials and cohomology of line bundles on homogeneous vector bundles*, Selecta Math. (N.S.) **16**, no. 2, 315–342.