

Catalan numbers, shuffle theorems, and Macdonald polynomials

George H. Seelinger

joint work with J. Blasiak, M. Haiman, J. Morse, and A. Pun


ghseeli@umich.edu

SMRI: Modern Perspectives in Representation Theory


22 May 2025

Algebraic quantity = Combinatorial generating function


Overview



Overview



Overview



- ① **Background on symmetric functions and Macdonald polynomials**
- ② Shuffle theorems, combinatorics, and LLT polynomials
- ③ A new formula for Macdonald polynomials

Symmetric polynomials and functions

- Symmetric polynomials $\mathbb{Q}[z_1, \dots, z_n]^{S_n}$
- Generators

$$e_r(z_1, \dots, z_n) = \sum_{1 \leq i_1 < i_2 < \dots < i_r \leq n} z_{i_1} \cdots z_{i_r}$$

$$e_1(z_1, z_2, z_3) = z_1 + z_2 + z_3$$

$$e_2(z_1, z_2, z_3) = z_1 z_2 + z_1 z_3 + z_2 z_3$$

$$e_3(z_1, z_2, z_3) = z_1 z_2 z_3$$

Symmetric polynomials and functions

- Symmetric polynomials $\mathbb{Q}[z_1, \dots, z_n]^{S_n}$
- Generators

$$e_r(z_1, \dots, z_n) = \sum_{1 \leq i_1 < i_2 < \dots < i_r \leq n} z_{i_1} \cdots z_{i_r}$$

$$e_1(z_1, z_2, z_3) = z_1 + z_2 + z_3$$

$$e_2(z_1, z_2, z_3) = z_1 z_2 + z_1 z_3 + z_2 z_3$$

$$e_3(z_1, z_2, z_3) = z_1 z_2 z_3$$

- $\Lambda = \mathbb{Q}[e_1, e_2, \dots]$

Symmetric polynomials and functions

- Symmetric polynomials $\mathbb{Q}[z_1, \dots, z_n]^{S_n}$
- Generators

$$e_r(z_1, \dots, z_n) = \sum_{1 \leq i_1 < i_2 < \dots < i_r \leq n} z_{i_1} \cdots z_{i_r}$$

$$e_1(z_1, z_2, z_3) = z_1 + z_2 + z_3$$

$$e_2(z_1, z_2, z_3) = z_1 z_2 + z_1 z_3 + z_2 z_3$$

$$e_3(z_1, z_2, z_3) = z_1 z_2 z_3$$

- $\Lambda = \mathbb{Q}[e_1, e_2, \dots]$
- Degree d elements of $\Lambda = \mathbb{Q}$ -vector space of dimension. . . ?

Symmetric polynomials and functions

- Symmetric polynomials $\mathbb{Q}[z_1, \dots, z_n]^{S_n}$
- Generators

$$e_r(z_1, \dots, z_n) = \sum_{1 \leq i_1 < i_2 < \dots < i_r \leq n} z_{i_1} \cdots z_{i_r}$$

$$e_1(z_1, z_2, z_3) = z_1 + z_2 + z_3$$

$$e_2(z_1, z_2, z_3) = z_1 z_2 + z_1 z_3 + z_2 z_3$$

$$e_3(z_1, z_2, z_3) = z_1 z_2 z_3$$

- $\Lambda = \mathbb{Q}[e_1, e_2, \dots]$
- Degree d elements of $\Lambda = \mathbb{Q}$ -vector space of dimension $\dots$?
- Integer partitions of d .

Partitions

Definition

$n \in \mathbb{Z}_{>0}$, a *partition of n* is $\lambda = (\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_\ell > 0)$ such that $\lambda_1 + \lambda_2 + \cdots + \lambda_\ell = n$.

Partitions

Definition

$n \in \mathbb{Z}_{>0}$, a *partition* of n is $\lambda = (\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_\ell > 0)$ such that $\lambda_1 + \lambda_2 + \cdots + \lambda_\ell = n$.

5 $\rightarrow$

--	--	--	--	--

$$4 + 1 \rightarrow \begin{array}{|c|c|c|c|} \hline \square & & & \\ \hline \square & \square & \square & \square \\ \hline \end{array}$$

$3 + 2 \rightarrow$ 

$$3 + 1 + 1 \rightarrow \begin{array}{|c|c|c|} \hline \square & & \\ \hline \square & & \\ \hline \square & \square & \square \\ \hline \end{array}$$

$$2 + 2 + 1 \rightarrow \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}$$

$$2 + 1 + 1 + 1 \rightarrow \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}$$

$$1 + 1 + 1 + 1 + 1 \rightarrow$$

Young Tableaux

Definition

Filling of partition diagram of λ with numbers such that

Young Tableaux

Definition

Filling of partition diagram of λ with numbers such that

- 1 strictly increasing up columns

Young Tableaux

Definition

Filling of partition diagram of λ with numbers such that

- 1 strictly increasing up columns
- 2 weakly increasing along rows

Young Tableaux

Definition

Filling of partition diagram of λ with numbers such that

- ① strictly increasing up columns
- ② weakly increasing along rows

Collection is called $\text{SSYT}(\lambda)$.


Young Tableaux

Definition


Filling of partition diagram of λ with numbers such that

- 1 strictly increasing up columns
- 2 weakly increasing along rows

Collection is called $\text{SSYT}(\lambda)$.



For $\lambda = (2, 1)$,



Polynomials from tableaux

Associate a polynomial to $\text{SSYT}(\lambda)$.

Polynomials from tableaux

Associate a polynomial to $\text{SSYT}(\lambda)$.

2		3		3		2		3		3		2		3	
1	1	1	1	2	2	1	2	1	3	2	3	1	3	1	2

Polynomials from tableaux

Associate a polynomial to $\text{SSYT}(\lambda)$.

$$\begin{array}{cccccccc} \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 2 & 2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 2 & 3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 2 \\ \hline \end{array} \\ \rightarrow & \begin{array}{|c|c|} \hline z_2 & \\ \hline z_1 & z_1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_3 & \\ \hline z_1 & z_1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_3 & \\ \hline z_2 & z_2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_2 & \\ \hline z_1 & z_2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_3 & \\ \hline z_1 & z_3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_3 & \\ \hline z_2 & z_3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_2 & \\ \hline z_1 & z_3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_3 & \\ \hline z_1 & z_2 \\ \hline \end{array} \end{array}$$

Polynomials from tableaux

Associate a polynomial to $\text{SSYT}(\lambda)$.

$$\begin{array}{cccccccc} \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 2 & 2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 2 & 3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 2 \\ \hline \end{array} \\ \rightarrow & \begin{array}{|c|c|} \hline z_2 & \\ \hline z_1 & z_1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_3 & \\ \hline z_1 & z_1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_3 & \\ \hline z_2 & z_2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_2 & \\ \hline z_1 & z_2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_3 & \\ \hline z_1 & z_3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_3 & \\ \hline z_2 & z_3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_2 & \\ \hline z_1 & z_3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_3 & \\ \hline z_1 & z_2 \\ \hline \end{array} \end{array}$$

$$s_{(2,1)}(z_1, z_2, z_3) = z_1^2 z_2 + z_1^2 z_3 + z_2^2 z_3 + z_1 z_2^2 + z_1 z_3^2 + z_2 z_3^2 + 2z_1 z_2 z_3$$

Polynomials from tableaux

Associate a polynomial to $\text{SSYT}(\lambda)$.

$$\begin{array}{cccccccc} \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 2 & 2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 2 & 3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 2 \\ \hline \end{array} \\ \rightarrow & \begin{array}{|c|c|} \hline z_2 & \\ \hline z_1 & z_1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_3 & \\ \hline z_1 & z_1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_3 & \\ \hline z_2 & z_2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_2 & \\ \hline z_1 & z_2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_3 & \\ \hline z_1 & z_3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_3 & \\ \hline z_2 & z_3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_2 & \\ \hline z_1 & z_3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_3 & \\ \hline z_1 & z_2 \\ \hline \end{array} \end{array}$$

$$s_{(2,1)}(z_1, z_2, z_3) = z_1^2 z_2 + z_1^2 z_3 + z_2^2 z_3 + z_1 z_2^2 + z_1 z_3^2 + z_2 z_3^2 + 2z_1 z_2 z_3$$

Definition

For λ a partition

$$s_\lambda = \sum_{T \in \text{SSYT}(\lambda)} \mathbf{z}^T \text{ for } \mathbf{z}^T = \prod_{i \in T} z_i$$

Polynomials from tableaux

Associate a polynomial to $\text{SSYT}(\lambda)$.

$$\begin{array}{cccccccc} \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 2 & 2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 2 & 3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 2 \\ \hline \end{array} \\ \rightarrow & \begin{array}{|c|c|} \hline z_2 & \\ \hline z_1 & z_1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_3 & \\ \hline z_1 & z_1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_3 & \\ \hline z_2 & z_2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_2 & \\ \hline z_1 & z_2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_3 & \\ \hline z_1 & z_3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_3 & \\ \hline z_2 & z_3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_2 & \\ \hline z_1 & z_3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_3 & \\ \hline z_1 & z_2 \\ \hline \end{array} \end{array}$$

$$s_{(2,1)}(z_1, z_2, z_3) = z_1^2 z_2 + z_1^2 z_3 + z_2^2 z_3 + z_1 z_2^2 + z_1 z_3^2 + z_2 z_3^2 + 2z_1 z_2 z_3$$

Definition

For λ a partition

$$s_\lambda = \sum_{T \in \text{SSYT}(\lambda)} \mathbf{z}^T \text{ for } \mathbf{z}^T = \prod_{i \in T} z_i$$

- s_λ is a symmetric function.

Polynomials from tableaux

Associate a polynomial to $\text{SSYT}(\lambda)$.

$$\begin{array}{cccccccc} \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 2 & 2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 2 & 3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 2 \\ \hline \end{array} \\ \rightarrow & \begin{array}{|c|c|} \hline z_2 & \\ \hline z_1 & z_1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_3 & \\ \hline z_1 & z_1 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_3 & \\ \hline z_2 & z_2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_2 & \\ \hline z_1 & z_2 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_3 & \\ \hline z_1 & z_3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_3 & \\ \hline z_2 & z_3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_2 & \\ \hline z_1 & z_3 \\ \hline \end{array}, & \begin{array}{|c|c|} \hline z_3 & \\ \hline z_1 & z_2 \\ \hline \end{array} \end{array}$$

$$s_{(2,1)}(z_1, z_2, z_3) = z_1^2 z_2 + z_1^2 z_3 + z_2^2 z_3 + z_1 z_2^2 + z_1 z_3^2 + z_2 z_3^2 + 2z_1 z_2 z_3$$

Definition

For λ a partition

$$s_\lambda = \sum_{T \in \text{SSYT}(\lambda)} \mathbf{z}^T \text{ for } \mathbf{z}^T = \prod_{i \in T} z_i$$

- s_λ is a symmetric function.
- $\{s_\lambda\}_\lambda$ forms a basis for Λ .

Representation theory and Schur functions

Irreducible representations of S_n are **also** labeled by partitions of n .

Representation theory and Schur functions

Irreducible representations of S_n are **also** labeled by partitions of n .

$$V_{\square\square\square} = \text{trivial}, \quad V_{\begin{smallmatrix}\square & \\ \square & \end{smallmatrix}} \cong \text{span}\{e_1 - e_2, e_2 - e_3\}, \quad V_{\begin{smallmatrix}\square \\ \square \\ \square \end{smallmatrix}} = \text{sign}$$

Representation theory and Schur functions

Irreducible representations of S_n are **also** labeled by partitions of n .

$$V_{\square\square\square} = \text{trivial}, \quad V_{\begin{smallmatrix} \square \\ \square \end{smallmatrix}} \cong \text{span}\{e_1 - e_2, e_2 - e_3\}, \quad V_{\begin{smallmatrix} \square \\ \square \\ \square \end{smallmatrix}} = \text{sign}$$

Frobenius characterstic, $\text{Frob}: \text{Rep}(S_n) \rightarrow \Lambda$, such that

Representation theory and Schur functions

Irreducible representations of S_n are **also** labeled by partitions of n .

$$V_{\square\square\square} = \text{trivial}, \quad V_{\begin{smallmatrix}\square & \\ \square & \end{smallmatrix}} \cong \text{span}\{e_1 - e_2, e_2 - e_3\}, \quad V_{\begin{smallmatrix}\square \\ \square \\ \square \end{smallmatrix}} = \text{sign}$$

Frobenius characterstic, $\text{Frob}: \text{Rep}(S_n) \rightarrow \Lambda$, such that

- Irreducible S_n -representation V_λ has $\text{Frob}(V_\lambda) = s_\lambda$

Representation theory and Schur functions

Irreducible representations of S_n are **also** labeled by partitions of n .

$$V_{\square\square\square} = \text{trivial}, \quad V_{\begin{smallmatrix} \square \\ \square \end{smallmatrix}} \cong \text{span}\{e_1 - e_2, e_2 - e_3\}, \quad V_{\begin{smallmatrix} \square \\ \square \\ \square \end{smallmatrix}} = \text{sign}$$

Frobenius characterstic, $\text{Frob}: \text{Rep}(S_n) \rightarrow \Lambda$, such that

- Irreducible S_n -representation V_λ has $\text{Frob}(V_\lambda) = s_\lambda$
- $U \cong V \oplus W \implies \text{Frob}(U) = \text{Frob}(V) + \text{Frob}(W)$

Representation theory and Schur functions

Irreducible representations of S_n are **also** labeled by partitions of n .

$$V_{\square\square\square} = \text{trivial}, \quad V_{\begin{smallmatrix} \square \\ \square \end{smallmatrix}} \cong \text{span}\{e_1 - e_2, e_2 - e_3\}, \quad V_{\begin{smallmatrix} \square \\ \square \\ \square \end{smallmatrix}} = \text{sign}$$

Frobenius characterstic, $\text{Frob}: \text{Rep}(S_n) \rightarrow \Lambda$, such that

- Irreducible S_n -representation V_λ has $\text{Frob}(V_\lambda) = s_\lambda$
- $U \cong V \oplus W \implies \text{Frob}(U) = \text{Frob}(V) + \text{Frob}(W)$
- $\text{Ind}_{S_m \times S_n}^{S_{m+n}}(V \times W) \mapsto \text{Frob}(V) \cdot \text{Frob}(W)$

Representation theory and Schur functions

Irreducible representations of S_n are **also** labeled by partitions of n .

$$V_{\square\square\square} = \text{trivial}, \quad V_{\begin{smallmatrix} \square \\ \square \end{smallmatrix}} \cong \text{span}\{e_1 - e_2, e_2 - e_3\}, \quad V_{\begin{smallmatrix} \square \\ \square \\ \square \end{smallmatrix}} = \text{sign}$$

Frobenius characterstic, $\text{Frob}: \text{Rep}(S_n) \rightarrow \Lambda$, such that

- Irreducible S_n -representation V_λ has $\text{Frob}(V_\lambda) = s_\lambda$
- $U \cong V \oplus W \implies \text{Frob}(U) = \text{Frob}(V) + \text{Frob}(W)$
- $\text{Ind}_{S_m \times S_n}^{S_{m+n}}(V \times W) \mapsto \text{Frob}(V) \cdot \text{Frob}(W)$
- $\text{Frob}(\text{Ind}_{S_1 \times S_1 \times S_1}^{S_3}(\mathbb{C} \times \mathbb{C} \times \mathbb{C})) = (s_{\square})^3 = s_{\square\square\square} + 2s_{\begin{smallmatrix} \square \\ \square \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ \square \end{smallmatrix}}$

Representation theory and Schur functions

Irreducible representations of S_n are **also** labeled by partitions of n .

$$V_{\square\square\square} = \text{trivial}, \quad V_{\begin{smallmatrix} \square \\ \square \end{smallmatrix}} \cong \text{span}\{e_1 - e_2, e_2 - e_3\}, \quad V_{\begin{smallmatrix} \square \\ \square \\ \square \end{smallmatrix}} = \text{sign}$$

Frobenius character map, $\text{Frob}: \text{Rep}(S_n) \rightarrow \Lambda$, such that

- Irreducible S_n -representation V_λ has $\text{Frob}(V_\lambda) = s_\lambda$
- $U \cong V \oplus W \implies \text{Frob}(U) = \text{Frob}(V) + \text{Frob}(W)$
- $\text{Ind}_{S_m \times S_n}^{S_{m+n}}(V \times W) \mapsto \text{Frob}(V) \cdot \text{Frob}(W)$
- $\text{Frob}(\text{Ind}_{S_1 \times S_1 \times S_1}^{S_3}(\mathbb{C} \times \mathbb{C} \times \mathbb{C})) = (s_{\square})^3 = s_{\square\square\square} + 2s_{\begin{smallmatrix} \square \\ \square \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ \square \end{smallmatrix}}$

Hidden Guide: Schur Positivity

“Naturally occurring” symmetric functions which are non-negative (coefficients in $\mathbb{N}$) linear combinations in Schur polynomial basis are interesting since they could have representation-theoretic models.

A Graded Example

$$\Delta = \det \begin{vmatrix} x_1^2 & x_1 & 1 \\ x_2^2 & x_2 & 1 \\ x_3^2 & x_3 & 1 \end{vmatrix} = x_1^2(x_2 - x_3) - x_2^2(x_1 - x_3) + x_3^2(x_1 - x_2)$$

$$S_3 \curvearrowright \text{span}\{\Delta\} \text{ via } \sigma.\Delta = \text{sgn}(\sigma)\Delta.$$

A Graded Example

$$\Delta = \det \begin{vmatrix} x_1^2 & x_1 & 1 \\ x_2^2 & x_2 & 1 \\ x_3^2 & x_3 & 1 \end{vmatrix} = x_1^2(x_2 - x_3) - x_2^2(x_1 - x_3) + x_3^2(x_1 - x_2)$$

$S_3 \curvearrowright \text{span}\{\Delta\}$ via $\sigma.\Delta = \text{sgn}(\sigma)\Delta$.

Consider $S_3 \curvearrowright M = \text{all partial derivatives of } \Delta$.

$$\begin{array}{c} \text{span}\{\Delta\} \\ \downarrow \partial_{x_i} \\ \text{span}\{2x_1(x_2 - x_3) - x_2^2 + x_3^2, 2x_2(x_3 - x_1) - x_3^2 + x_1^2\} \\ \downarrow \partial_{x_i} \\ \text{span}\{x_3 - x_1, x_2 - x_3\} \\ \downarrow \partial_{x_i} \\ \text{span}\{1\} \end{array}$$

A Graded Example

$$\begin{array}{ccc}
 \text{span}\{\Delta\} & & \cong V_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}} \\
 \downarrow \partial_{x_i} & & \\
 \text{span}\{2x_1(x_2 - x_3) - x_2^2 + x_3^2, 2x_2(x_3 - x_1) - x_3^2 + x_1^2\} & & \cong V_{\begin{array}{|c|} \hline \square \square \\ \hline \square \square \\ \hline \end{array}} \\
 \downarrow \partial_{x_i} & & \\
 \text{span}\{x_3 - x_1, x_2 - x_3\} & & \cong V_{\begin{array}{|c|} \hline \square \square \\ \hline \square \square \\ \hline \end{array}} \\
 \downarrow \partial_{x_i} & & \\
 \text{span}\{1\} & & \cong V_{\begin{array}{|c|} \hline \square \square \square \\ \hline \end{array}}
 \end{array}$$

A Graded Example

$$\begin{array}{ccc}
 \text{span}\{\Delta\} & & \cong V_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}} \\
 \downarrow \partial_{x_i} & & \\
 \text{span}\{2x_1(x_2 - x_3) - x_2^2 + x_3^2, 2x_2(x_3 - x_1) - x_3^2 + x_1^2\} & & \cong V_{\begin{array}{|c|} \hline \square \\ \hline \square \square \\ \hline \end{array}} \\
 \downarrow \partial_{x_i} & & \\
 \text{span}\{x_3 - x_1, x_2 - x_3\} & & \cong V_{\begin{array}{|c|} \hline \square \\ \hline \square \square \\ \hline \end{array}} \\
 \downarrow \partial_{x_i} & & \\
 \text{span}\{1\} & & \cong V_{\begin{array}{|c|} \hline \square \\ \hline \square \square \square \\ \hline \end{array}}
 \end{array}$$

- $\text{Frob}(M) = s_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}} + 2s_{\begin{array}{|c|} \hline \square \\ \hline \square \square \\ \hline \end{array}} + s_{\begin{array}{|c|} \hline \square \square \square \\ \hline \end{array}}$

A Graded Example

$$\begin{array}{ccc}
 \text{span}\{\Delta\} & & \cong V_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}} \\
 \downarrow \partial_{x_i} & & \\
 \text{span}\{2x_1(x_2 - x_3) - x_2^2 + x_3^2, 2x_2(x_3 - x_1) - x_3^2 + x_1^2\} & & \cong V_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} \\
 \downarrow \partial_{x_i} & & \\
 \text{span}\{x_3 - x_1, x_2 - x_3\} & & \cong V_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}} \\
 \downarrow \partial_{x_i} & & \\
 \text{span}\{1\} & & \cong V_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}}
 \end{array}$$

- $\text{Frob}(M) = s_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}} + 2s_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} + s_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}}$
- Problem: what about grading?

Getting More Information

$$\begin{array}{llll} \text{span}\{\Delta\} & & \cong V_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}} & \text{deg} = 3 \\ \downarrow \partial_{x_i} & & & \\ \text{span}\{2x_1(x_2 - x_3) - x_2^2 + x_3^2, 2x_2(x_3 - x_1) - x_3^2 + x_1^2\} & & \cong V_{\begin{array}{|c|} \hline \square \\ \hline \square \square \\ \hline \end{array}} & \text{deg} = 2 \\ \downarrow \partial_{x_i} & & & \\ \text{span}\{x_3 - x_1, x_2 - x_3\} & & \cong V_{\begin{array}{|c|} \hline \square \\ \hline \square \square \\ \hline \end{array}} & \text{deg} = 1 \\ \downarrow \partial_{x_i} & & & \\ \text{span}\{1\} & & \cong V_{\begin{array}{|c|} \hline \square \square \square \\ \hline \end{array}} & \text{deg} = 0 \end{array}$$

Getting More Information

$$\begin{array}{lll}
 \text{span}\{\Delta\} & \cong V_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}} & \text{deg} = 3 \\
 \downarrow \partial_{x_i} & & \\
 \text{span}\{2x_1(x_2 - x_3) - x_2^2 + x_3^2, 2x_2(x_3 - x_1) - x_3^2 + x_1^2\} & \cong V_{\begin{array}{|c|} \hline \square \\ \hline \square \square \\ \hline \end{array}} & \text{deg} = 2 \\
 \downarrow \partial_{x_i} & & \\
 \text{span}\{x_3 - x_1, x_2 - x_3\} & \cong V_{\begin{array}{|c|} \hline \square \\ \hline \square \square \\ \hline \end{array}} & \text{deg} = 1 \\
 \downarrow \partial_{x_i} & & \\
 \text{span}\{1\} & \cong V_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}} & \text{deg} = 0
 \end{array}$$

- $(V_\lambda \text{ in degree } d) \mapsto q^d s_\lambda$

Getting More Information

$$\begin{array}{llll}
 \text{span}\{\Delta\} & & \cong V_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}} & \text{deg} = 3 \\
 \downarrow \partial_{x_i} & & & \\
 \text{span}\{2x_1(x_2 - x_3) - x_2^2 + x_3^2, 2x_2(x_3 - x_1) - x_3^2 + x_1^2\} & & \cong V_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} & \text{deg} = 2 \\
 \downarrow \partial_{x_i} & & & \\
 \text{span}\{x_3 - x_1, x_2 - x_3\} & & \cong V_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} & \text{deg} = 1 \\
 \downarrow \partial_{x_i} & & & \\
 \text{span}\{1\} & & \cong V_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}} & \text{deg} = 0
 \end{array}$$

- $(V_\lambda \text{ in degree } d) \mapsto q^d s_\lambda$
- $\text{Frob}(M) = q^3 s_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}} + q^2 s_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} + q^1 s_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} + q^0 s_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}}.$

Getting More Information

$$\begin{array}{lll}
 \text{span}\{\Delta\} & \cong V_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}} & \text{deg} = 3 \\
 \downarrow \partial_{x_i} & & \\
 \text{span}\{2x_1(x_2 - x_3) - x_2^2 + x_3^2, 2x_2(x_3 - x_1) - x_3^2 + x_1^2\} & \cong V_{\begin{array}{|c|} \hline \square & \square \\ \hline \end{array}} & \text{deg} = 2 \\
 \downarrow \partial_{x_i} & & \\
 \text{span}\{x_3 - x_1, x_2 - x_3\} & \cong V_{\begin{array}{|c|} \hline \square & \square \\ \hline \end{array}} & \text{deg} = 1 \\
 \downarrow \partial_{x_i} & & \\
 \text{span}\{1\} & \cong V_{\begin{array}{|c|} \hline \square & \square & \square \\ \hline \end{array}} & \text{deg} = 0
 \end{array}$$

- $(V_\lambda \text{ in degree } d) \mapsto q^d s_\lambda$
- $\text{Frob}(M) = q^3 s_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}} + q^2 s_{\begin{array}{|c|} \hline \square & \square \\ \hline \end{array}} + q^1 s_{\begin{array}{|c|} \hline \square & \square \\ \hline \end{array}} + q^0 s_{\begin{array}{|c|} \hline \square & \square & \square \\ \hline \end{array}}.$
- Hall-Littlewood polynomial $H_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}}(X; q).$

Getting More Information

$$\begin{array}{rcl}
 \text{span}\{\Delta\} & \cong V_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}} & \text{deg} = 3 \\
 \downarrow \partial_{x_i} & & \\
 \text{span}\{2x_1(x_2 - x_3) - x_2^2 + x_3^2, 2x_2(x_3 - x_1) - x_3^2 + x_1^2\} & \cong V_{\begin{array}{|c|} \hline \square \\ \hline \square \square \\ \hline \end{array}} & \text{deg} = 2 \\
 \downarrow \partial_{x_i} & & \\
 \text{span}\{x_3 - x_1, x_2 - x_3\} & \cong V_{\begin{array}{|c|} \hline \square \\ \hline \square \square \\ \hline \end{array}} & \text{deg} = 1 \\
 \downarrow \partial_{x_i} & & \\
 \text{span}\{1\} & \cong V_{\begin{array}{|c|} \hline \square \\ \hline \square \square \\ \hline \square \square \square \\ \hline \end{array}} & \text{deg} = 0
 \end{array}$$

- $(V_\lambda \text{ in degree } d) \mapsto q^d s_\lambda$
- $\text{Frob}(M) = q^3 s_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}} + q^2 s_{\begin{array}{|c|} \hline \square \\ \hline \square \square \\ \hline \end{array}} + q^1 s_{\begin{array}{|c|} \hline \square \\ \hline \square \square \\ \hline \end{array}} + q^0 s_{\begin{array}{|c|} \hline \square \square \square \\ \hline \end{array}}.$
- Hall-Littlewood polynomial $H_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}}(X; q).$
- Remark: $M \cong \mathbb{Z}[x_1, x_2, x_3]/\text{Sym}^+ \cong H^*(Fl_3)$ as **graded** S_3 -representations.

A Problem

- In 1988, Macdonald introduces one basis of symmetric polynomials to rule them all!

A Problem

- In 1988, Macdonald introduces one basis of symmetric polynomials to rule them all!
- Coefficients in $\mathbb{Q}(q, t)$, specializations give Hall-Littlewood polynomials, Schur polynomials, and many other famous bases.

A Problem

- In 1988, Macdonald introduces one basis of symmetric polynomials to rule them all!
- Coefficients in $\mathbb{Q}(q, t)$, specializations give Hall-Littlewood polynomials, Schur polynomials, and many other famous bases.
- Defined by orthogonality and triangularity under a certain inner-product.

A Problem

- In 1988, Macdonald introduces one basis of symmetric polynomials to rule them all!
- Coefficients in $\mathbb{Q}(q, t)$, specializations give Hall-Littlewood polynomials, Schur polynomials, and many other famous bases.
- Defined by orthogonality and triangularity under a certain inner-product.
- Garsia modifies these polynomials so

$$\tilde{H}_\lambda(X; q, t) = \sum_{\mu} \tilde{K}(q, t) s_{\mu} \text{ conjecturally satisfies } \tilde{K}(q, t) \in \mathbb{N}[q, t]$$

A Problem

- In 1988, Macdonald introduces one basis of symmetric polynomials to rule them all!
- Coefficients in $\mathbb{Q}(q, t)$, specializations give Hall-Littlewood polynomials, Schur polynomials, and many other famous bases.
- Defined by orthogonality and triangularity under a certain inner-product.
- Garsia modifies these polynomials so

$$\tilde{H}_\lambda(X; q, t) = \sum_{\mu} \tilde{K}(q, t) s_{\mu} \text{ conjecturally satisfies } \tilde{K}(q, t) \in \mathbb{N}[q, t]$$

- $\tilde{H}_\lambda(X; 1, 1) = s_1^{|\lambda|}$.

A Problem

- In 1988, Macdonald introduces one basis of symmetric polynomials to rule them all!
- Coefficients in $\mathbb{Q}(q, t)$, specializations give Hall-Littlewood polynomials, Schur polynomials, and many other famous bases.
- Defined by orthogonality and triangularity under a certain inner-product.
- Garsia modifies these polynomials so

$$\tilde{H}_\lambda(X; q, t) = \sum_{\mu} \tilde{K}(q, t) s_{\mu} \text{ conjecturally satisfies } \tilde{K}(q, t) \in \mathbb{N}[q, t]$$

- $\tilde{H}_\lambda(X; 1, 1) = s_1^{|\lambda|}$.
- Does there exist a family of S_n -regular representations whose bigraded Frobenius characteristics equal $\tilde{H}_\lambda(X; q, t)$?

Garsia-Haiman modules

- $S_n \curvearrowright \mathbb{Q}[x_1, \dots, x_n, y_1, \dots, y_n]$ with $\sigma(x_i) = x_{\sigma(i)}$, $\sigma(y_j) = y_{\sigma(j)}$.

Garsia-Haiman modules

- $S_n \curvearrowright \mathbb{Q}[x_1, \dots, x_n, y_1, \dots, y_n]$ with $\sigma(x_i) = x_{\sigma(i)}$, $\sigma(y_j) = y_{\sigma(j)}$.
- Garsia-Haiman (1993): $M_\mu = \text{span of partial derivatives of}$
 $\Delta_\mu = \det_{(i,j) \in \mu, k \in [n]} (x_k^{i-1} y_k^{j-1})$

Garsia-Haiman modules

- $S_n \curvearrowright \mathbb{Q}[x_1, \dots, x_n, y_1, \dots, y_n]$ with $\sigma(x_i) = x_{\sigma(i)}$, $\sigma(y_j) = y_{\sigma(j)}$.
- Garsia-Haiman (1993): $M_\mu = \text{span of partial derivatives of}$
 $\Delta_\mu = \det_{(i,j) \in \mu, k \in [n]} (x_k^{i-1} y_k^{j-1})$

$$\Delta_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} = \det \begin{vmatrix} 1 & y_1 & x_1 \\ 1 & y_2 & x_2 \\ 1 & y_3 & x_3 \end{vmatrix} = x_3 y_2 - y_3 x_2 - y_1 x_3 + y_1 x_2 + y_3 x_1 - y_2 x_1$$

Garsia-Haiman modules

- $S_n \curvearrowright \mathbb{Q}[x_1, \dots, x_n, y_1, \dots, y_n]$ with $\sigma(x_i) = x_{\sigma(i)}$, $\sigma(y_j) = y_{\sigma(j)}$.
- Garsia-Haiman (1993): $M_\mu = \text{span of partial derivatives of } \Delta_\mu = \det_{(i,j) \in \mu, k \in [n]} (x_k^{i-1} y_k^{j-1})$

$$\Delta_{\begin{smallmatrix} \square & \\ \square & \end{smallmatrix}} = \det \begin{vmatrix} 1 & y_1 & x_1 \\ 1 & y_2 & x_2 \\ 1 & y_3 & x_3 \end{vmatrix} = x_3 y_2 - y_3 x_2 - y_1 x_3 + y_1 x_2 + y_3 x_1 - y_2 x_1$$

$$M_{2,1} =$$

$$\underbrace{\text{span}\{\Delta_{2,1}\}}_{\text{deg}=(1,1)} \oplus \underbrace{\text{span}\{y_3 - y_1, y_1 - y_2\}}_{\text{deg}=(0,1)} \oplus \underbrace{\text{span}\{x_3 - x_1, x_1 - x_2\}}_{\text{deg}=(1,0)} \oplus \underbrace{\text{span}\{1\}}_{\text{deg}=(0,0)}$$

Garsia-Haiman modules

- $S_n \curvearrowright \mathbb{Q}[x_1, \dots, x_n, y_1, \dots, y_n]$ with $\sigma(x_i) = x_{\sigma(i)}$, $\sigma(y_j) = y_{\sigma(j)}$.
- Garsia-Haiman (1993): $M_\mu = \text{span of partial derivatives of } \Delta_\mu = \det_{(i,j) \in \mu, k \in [n]} (x_k^{i-1} y_k^{j-1})$

$$\Delta_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} = \det \begin{vmatrix} 1 & y_1 & x_1 \\ 1 & y_2 & x_2 \\ 1 & y_3 & x_3 \end{vmatrix} = x_3 y_2 - y_3 x_2 - y_1 x_3 + y_1 x_2 + y_3 x_1 - y_2 x_1$$

$$M_{2,1} =$$

$$\underbrace{\text{span}\{\Delta_{2,1}\}}_{\text{deg}=(1,1)} \oplus \underbrace{\text{span}\{y_3 - y_1, y_1 - y_2\}}_{\text{deg}=(0,1)} \oplus \underbrace{\text{span}\{x_3 - x_1, x_1 - x_2\}}_{\text{deg}=(1,0)} \oplus \underbrace{\text{span}\{1\}}_{\text{deg}=(0,0)}$$

Irreducible S_n -representation V_λ with bidegree $(a, b) \mapsto q^a t^b s_\lambda$

Garsia-Haiman modules

- $S_n \curvearrowright \mathbb{Q}[x_1, \dots, x_n, y_1, \dots, y_n]$ with $\sigma(x_i) = x_{\sigma(i)}$, $\sigma(y_j) = y_{\sigma(j)}$.
- Garsia-Haiman (1993): $M_\mu = \text{span of partial derivatives of } \Delta_\mu = \det_{(i,j) \in \mu, k \in [n]} (x_k^{i-1} y_k^{j-1})$

$$\Delta_{\begin{smallmatrix} \square & \\ \square & \end{smallmatrix}} = \det \begin{vmatrix} 1 & y_1 & x_1 \\ 1 & y_2 & x_2 \\ 1 & y_3 & x_3 \end{vmatrix} = x_3 y_2 - y_3 x_2 - y_1 x_3 + y_1 x_2 + y_3 x_1 - y_2 x_1$$

$$M_{2,1} =$$

$$\underbrace{\text{span}\{\Delta_{2,1}\}}_{\text{deg}=(1,1)} \oplus \underbrace{\text{span}\{y_3 - y_1, y_1 - y_2\}}_{\text{deg}=(0,1)} \oplus \underbrace{\text{span}\{x_3 - x_1, x_1 - x_2\}}_{\text{deg}=(1,0)} \oplus \underbrace{\text{span}\{1\}}_{\text{deg}=(0,0)}$$

Irreducible S_n -representation V_λ with bidegree $(a, b) \mapsto q^a t^b s_\lambda$

$$\tilde{H}_{\begin{smallmatrix} \square & \\ \square & \end{smallmatrix}} = q^1 t^1 s_{\begin{smallmatrix} \square \\ \square \\ \square \end{smallmatrix}} + t^1 s_{\begin{smallmatrix} \square & \\ \square & \end{smallmatrix}} + q^1 s_{\begin{smallmatrix} \square & \\ \square & \end{smallmatrix}} + s_{\begin{smallmatrix} \square & \square & \square \end{smallmatrix}}$$

Garsia-Haiman modules

Theorem (Haiman, 2001)

The Garsia-Haiman module M_λ has bigraded Frobenius characteristic given by $\tilde{H}_\lambda(X; q, t)$

Theorem (Haiman, 2001)

The Garsia-Haiman module M_λ has bigraded Frobenius characteristic given by $\tilde{H}_\lambda(X; q, t)$

- Proved via connection to the Hilbert Scheme $\text{Hilb}^n(\mathbb{C}^2)$.

Garsia-Haiman modules

Theorem (Haiman, 2001)

The Garsia-Haiman module M_λ has bigraded Frobenius characteristic given by $\tilde{H}_\lambda(X; q, t)$

- Proved via connection to the Hilbert Scheme $\text{Hilb}^n(\mathbb{C}^2)$.

Corollary

$\tilde{H}_\lambda(X; q, t) = \sum_\mu \tilde{K}_{\lambda\mu}(q, t) s_\mu$ satisfies $\tilde{K}_{\lambda\mu}(q, t) \in \mathbb{N}[q, t]$.

Garsia-Haiman modules

Theorem (Haiman, 2001)

The Garsia-Haiman module M_λ has bigraded Frobenius characteristic given by $\tilde{H}_\lambda(X; q, t)$

- Proved via connection to the Hilbert Scheme $\text{Hilb}^n(\mathbb{C}^2)$.

Corollary

$\tilde{H}_\lambda(X; q, t) = \sum_\mu \tilde{K}_{\lambda\mu}(q, t) s_\mu$ satisfies $\tilde{K}_{\lambda\mu}(q, t) \in \mathbb{N}[q, t]$.

- No combinatorial description of $\tilde{K}_{\lambda\mu}(q, t)$.

Symmetric functions, representation theory, and combinatorics

Symmetric function	Representation theory	Combinatorics
s_λ	Irreducible V_λ	SSYT(λ)
$\tilde{H}_\lambda(X; q, t)$	Garsia-Haiman M_λ	Later

Observation

All of these Garsia-Haiman modules are contained in the module of diagonal harmonics:

$$DH_n = \text{span}\{f \in \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n] \mid \left(\sum_{j=1}^n \partial_{x_j}^r \partial_{y_j}^s \right) f = 0, \forall r+s > 0\}$$

Garsia-Haiman modules

Observation

All of these Garsia-Haiman modules are contained in the module of diagonal harmonics:

$$DH_n = \text{span}\{f \in \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n] \mid \left(\sum_{j=1}^n \partial_{x_j}^r \partial_{y_j}^s \right) f = 0, \forall r+s > 0\}$$

Question

What symmetric function is the bigraded Frobenius characteristic of DH_n ?

Frobenius characteristic of DH_3

Frobenius characteristic of DH_3

$$= \frac{t^3 \tilde{H}_{1,1,1}}{-qt^2 + t^3 + q^2 - qt} - \frac{(q^2 t + qt^2 + qt) \tilde{H}_{2,1}}{-q^2 t^2 + q^3 + t^3 - qt} - \frac{q^3 \tilde{H}_3}{-q^3 + q^2 t + qt - t^2}$$

Frobenius characteristic of DH_3

$$= \frac{t^3 \tilde{H}_{1,1,1}}{-qt^2+t^3+q^2-qt} - \frac{(q^2t+qt^2+qt)\tilde{H}_{2,1}}{-q^2t^2+q^3+t^3-qt} - \frac{q^3\tilde{H}_3}{-q^3+q^2t+qt-t^2}$$

Compare to

$$e_3 = \frac{\tilde{H}_{1,1,1}}{-qt^2+t^3+q^2-qt} - \frac{(q+t+1)\tilde{H}_{2,1}}{-q^2t^2+q^3+t^3-qt} - \frac{\tilde{H}_3}{-q^3+q^2t+qt-t^2}$$

Frobenius characteristic of DH_3

$$= \frac{t^3 \tilde{H}_{1,1,1}}{-qt^2 + t^3 + q^2 - qt} - \frac{(q^2 t + qt^2 + qt) \tilde{H}_{2,1}}{-q^2 t^2 + q^3 + t^3 - qt} - \frac{q^3 \tilde{H}_3}{-q^3 + q^2 t + qt - t^2}$$

Compare to

$$e_3 = \frac{\tilde{H}_{1,1,1}}{-qt^2 + t^3 + q^2 - qt} - \frac{(q+t+1) \tilde{H}_{2,1}}{-q^2 t^2 + q^3 + t^3 - qt} - \frac{\tilde{H}_3}{-q^3 + q^2 t + qt - t^2}$$

Operator ∇

$$\nabla \tilde{H}_\lambda(X; q, t) = q^{n(\lambda)} t^{n(\lambda^*)} \tilde{H}_\lambda(X; q, t),$$

where $n(\lambda) = \sum_i (i-1)\lambda_i$ and λ^* is the transpose partition to λ .

Frobenius characteristic of DH_3

$$= \frac{t^3 \tilde{H}_{1,1,1}}{-qt^2 + t^3 + q^2 - qt} - \frac{(q^2 t + qt^2 + qt) \tilde{H}_{2,1}}{-q^2 t^2 + q^3 + t^3 - qt} - \frac{q^3 \tilde{H}_3}{-q^3 + q^2 t + qt - t^2}$$

Compare to

$$e_3 = \frac{\tilde{H}_{1,1,1}}{-qt^2 + t^3 + q^2 - qt} - \frac{(q+t+1) \tilde{H}_{2,1}}{-q^2 t^2 + q^3 + t^3 - qt} - \frac{\tilde{H}_3}{-q^3 + q^2 t + qt - t^2}$$

Operator ∇

$$\nabla \tilde{H}_\lambda(X; q, t) = q^{n(\lambda)} t^{n(\lambda^*)} \tilde{H}_\lambda(X; q, t),$$

where $n(\lambda) = \sum_i (i-1)\lambda_i$ and λ^* is the transpose partition to λ .

Theorem (Haiman, 2002)

The bigraded Frobenius characteristic of DH_n is given by ∇e_n .

Symmetric functions, representation theory, and combinatorics

Symmetric function	Representation theory	Combinatorics
s_λ	Irreducible V_λ	SSYT(λ)
$\tilde{H}_\lambda(X; q, t)$	Garsia-Haiman M_λ	Later
∇e_n	DH_n	Now: Shuffle theorem

Outline

- ① Background on symmetric functions and Macdonald polynomials
- ② **Shuffle theorems, combinatorics, and LLT polynomials**
- ③ A new formula for Macdonald polynomials

A Combinatorial Connection: Shuffle Theorem

Theorem (Carlsson-Mellit, 2018)

$$\nabla e_k = \sum_{\lambda} (q, t \text{ monomial})(LLT \text{ polynomial})$$

- Summation over all k -by- k Dyck paths.

A Combinatorial Connection: Shuffle Theorem

Theorem (Carlsson-Mellit, 2018)

$$\nabla e_k = \sum_{\lambda} t^{\text{area}(\lambda)} q^{\text{dinv}(\lambda)} (LLT \text{ polynomial})$$

- Summation over all k -by- k Dyck paths.
- $\text{area}(\lambda)$ and $\text{dinv}(\lambda)$ statistics of Dyck paths.

A Combinatorial Connection: Shuffle Theorem

Theorem (Carlsson-Mellit, 2018)

$$\nabla e_k = \sum_{\lambda} t^{\text{area}(\lambda)} q^{\text{dinv}(\lambda)} \omega \mathcal{G}_{\nu(\lambda)}(X; q^{-1})$$

- Summation over all k -by- k Dyck paths.
- $\text{area}(\lambda)$ and $\text{dinv}(\lambda)$ statistics of Dyck paths.
- $\mathcal{G}_{\nu(\lambda)}(X; q)$ a symmetric LLT polynomial indexed by a tuple of offset (skew) rows.

A Combinatorial Connection: Shuffle Theorem

Theorem (Carlsson-Mellit, 2018)

$$\nabla e_k = \sum_{\lambda} t^{\text{area}(\lambda)} q^{\text{dinv}(\lambda)} \omega \mathcal{G}_{\nu(\lambda)}(X; q^{-1})$$

- Summation over all k -by- k Dyck paths.
- $\text{area}(\lambda)$ and $\text{dinv}(\lambda)$ statistics of Dyck paths.
- $\mathcal{G}_{\nu(\lambda)}(X; q)$ a symmetric LLT polynomial indexed by a tuple of offset (skew) rows.
- ω a standard involution of symmetric polynomials.

A Combinatorial Connection: Shuffle Theorem

Theorem (Carlsson-Mellit, 2018)


$$\nabla e_k = \sum_{\lambda} t^{\text{area}(\lambda)} q^{\text{dinv}(\lambda)} \omega \mathcal{G}_{\nu(\lambda)}(X; q^{-1})$$

- Summation over all k -by- k Dyck paths.
- $\text{area}(\lambda)$ and $\text{dinv}(\lambda)$ statistics of Dyck paths.
- $\mathcal{G}_{\nu(\lambda)}(X; q)$ a symmetric LLT polynomial indexed by a tuple of offset (skew) rows.
- ω a standard involution of symmetric polynomials.
- Conjectured by (Haiman-Haglund-Loehr-Remmel-Ulyanov, 2002).

Dyck paths

Dyck paths


A Dyck path λ is a south-east lattice path lying below the line segment from $(0, k)$ to $(k, 0)$.



Dyck paths

Dyck paths

A Dyck path λ is a south-east lattice path lying below the line segment from $(0, k)$ to $(k, 0)$.




- $\text{area}(\lambda) = \text{number of squares above } \lambda \text{ but below the path } \delta \text{ of alternating S-E steps.}$

Dyck paths

Dyck paths

A Dyck path λ is a south-east lattice path lying below the line segment from $(0, k)$ to $(k, 0)$.




- $\text{area}(\lambda)$ = number of squares above λ but below the path δ of alternating S-E steps.
- E.g., above $\text{area}(\lambda) = 10$.

Dyck paths

Dyck paths

A Dyck path λ is a south-east lattice path lying below the line segment from $(0, k)$ to $(k, 0)$.




- $\text{area}(\lambda)$ = number of squares above λ but below the path δ of alternating S-E steps.
- E.g., above $\text{area}(\lambda) = 10$.
- Catalan-number many Dyck paths for fixed k .

Dyck paths

Dyck paths


A Dyck path λ is a south-east lattice path lying below the line segment from $(0, k)$ to $(k, 0)$.




- $\text{area}(\lambda) =$ number of squares above λ but below the path δ of alternating S-E steps.
- E.g., above $\text{area}(\lambda) = 10$.
- Catalan-number many Dyck paths for fixed k . $(1, 2, 5, 14, 42, \dots)$

dinv

$\text{dinv}(\lambda) = \#$ of balanced hooks in diagram below λ .



$\text{dinv}(\lambda) = \#$ of balanced hooks in diagram below λ .




Balanced hook is given by a cell below λ satisfying

$$\frac{\ell}{a+1} < 1 - \epsilon < \frac{\ell+1}{a}, \quad \epsilon \text{ small.}$$

Key Object: LLT Polynomials

Let $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$ be a tuple of skew shapes. (Skew shape = $\lambda \setminus \mu$)

$$\nu = \left(\begin{array}{|c|c|c|} \hline \square & \square & \\ \hline \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array} \right)$$



Key Object: LLT Polynomials

Let $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$ be a tuple of skew shapes. (Skew shape = $\lambda \setminus \mu$)

- The *content* of a box in row y , column x is $x - y$.

$$\nu = \left(\begin{array}{|c|c|c|} \hline \square & \square & \\ \hline \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array} \right)$$

-4	-3	-2	-1	0	1
-3	-2	-1	0	1	2
-2	-1	0	1	2	3
-1	0	1	2	3	4
0	1	2	3	4	5

Key Object: LLT Polynomials

Let $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$ be a tuple of skew shapes. (Skew shape = $\lambda \setminus \mu$)

- The *content* of a box in row y , column x is $x - y$.
- *Reading order*: label boxes $b_1, \dots, b_n$ by scanning each diagonal from southwest to northeast, in order of increasing content.

$$\nu = \left(\begin{array}{|c|c|c|} \hline \square & \square & \\ \hline \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array} \right)$$

				b_3	b_6
				b_5	b_8
b_1	b_2				
	b_4	b_7			

Key Object: LLT Polynomials

Let $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$ be a tuple of skew shapes. (Skew shape $= \lambda \setminus \mu$)

- The *content* of a box in row y , column x is $x - y$.
- *Reading order*: label boxes $b_1, \dots, b_n$ by scanning each diagonal from southwest to northeast, in order of increasing content.
- A pair $(a, b) \in \nu$ is *attacking* if a precedes b in reading order and
 - $\text{content}(b) = \text{content}(a)$, or
 - $\text{content}(b) = \text{content}(a) + 1$ and $a \in \nu_{(i)}, b \in \nu_{(j)}$ with $i > j$.

$$\nu = \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right)$$

				b_3	b_6
				b_5	b_8
b_1	b_2				
	b_4	b_7			

Attacking pairs: $(b_2, b_3), (b_3, b_4), (b_4, b_5), (b_4, b_6), (b_5, b_7), (b_6, b_7), (b_7, b_8)$

Key Object: LLT Polynomials

Let $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$ be a tuple of skew shapes. (Skew shape $= \lambda \setminus \mu$)

- The *content* of a box in row y , column x is $x - y$.
- *Reading order*: label boxes $b_1, \dots, b_n$ by scanning each diagonal from southwest to northeast, in order of increasing content.
- A pair $(a, b) \in \nu$ is *attacking* if a precedes b in reading order and
 - $\text{content}(b) = \text{content}(a)$, or
 - $\text{content}(b) = \text{content}(a) + 1$ and $a \in \nu_{(i)}, b \in \nu_{(j)}$ with $i > j$.

$$\nu = \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right)$$

				b_3	b_6
				b_5	b_8
b_1	b_2				
	b_4	b_7			

Attacking pairs: $(b_2, b_3), (b_3, b_4), (b_4, b_5), (b_4, b_6), (b_5, b_7), (b_6, b_7), (b_7, b_8)$

Key Object: LLT Polynomials

Let $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$ be a tuple of skew shapes. (Skew shape $= \lambda \setminus \mu$)

- The *content* of a box in row y , column x is $x - y$.
- *Reading order*: label boxes $b_1, \dots, b_n$ by scanning each diagonal from southwest to northeast, in order of increasing content.
- A pair $(a, b) \in \nu$ is *attacking* if a precedes b in reading order and
 - $\text{content}(b) = \text{content}(a)$, or
 - $\text{content}(b) = \text{content}(a) + 1$ and $a \in \nu_{(i)}, b \in \nu_{(j)}$ with $i > j$.

$$\nu = \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right)$$

				b_3	b_6
				b_5	b_8
b_1	b_2				
	b_4	b_7			

Attacking pairs: (b_2, b_3) , (b_3, b_4) , (b_4, b_5) , (b_4, b_6) , (b_5, b_7) , (b_6, b_7) , (b_7, b_8)

Key Object: LLT Polynomials

Let $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$ be a tuple of skew shapes. (Skew shape $= \lambda \setminus \mu$)

- The *content* of a box in row y , column x is $x - y$.
- *Reading order*: label boxes $b_1, \dots, b_n$ by scanning each diagonal from southwest to northeast, in order of increasing content.
- A pair $(a, b) \in \nu$ is *attacking* if a precedes b in reading order and
 - $\text{content}(b) = \text{content}(a)$, or
 - $\text{content}(b) = \text{content}(a) + 1$ and $a \in \nu_{(i)}, b \in \nu_{(j)}$ with $i > j$.

$$\nu = \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right)$$

					b_3	b_6
					b_5	b_8
b_1	b_2					
	b_4	b_7				

Attacking pairs: $(b_2, b_3), (b_3, b_4), (b_4, b_5), (b_4, b_6), (b_5, b_7), (b_6, b_7), (b_7, b_8)$

Key Object: LLT Polynomials

Let $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$ be a tuple of skew shapes. (Skew shape $= \lambda \setminus \mu$)

- The *content* of a box in row y , column x is $x - y$.
- *Reading order*: label boxes $b_1, \dots, b_n$ by scanning each diagonal from southwest to northeast, in order of increasing content.
- A pair $(a, b) \in \nu$ is *attacking* if a precedes b in reading order and
 - $\text{content}(b) = \text{content}(a)$, or
 - $\text{content}(b) = \text{content}(a) + 1$ and $a \in \nu_{(i)}, b \in \nu_{(j)}$ with $i > j$.

$$\nu = \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right)$$

				b_3	b_6
				b_5	b_8
b_1	b_2				
	b_4	b_7			

Attacking pairs: $(b_2, b_3), (b_3, b_4), (b_4, b_5), (b_4, b_6), (b_5, b_7), (b_6, b_7), (b_7, b_8)$

Key Object: LLT Polynomials

Let $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$ be a tuple of skew shapes. (Skew shape $= \lambda \setminus \mu$)

- The *content* of a box in row y , column x is $x - y$.
- *Reading order*: label boxes $b_1, \dots, b_n$ by scanning each diagonal from southwest to northeast, in order of increasing content.
- A pair $(a, b) \in \nu$ is *attacking* if a precedes b in reading order and
 - $\text{content}(b) = \text{content}(a)$, or
 - $\text{content}(b) = \text{content}(a) + 1$ and $a \in \nu_{(i)}, b \in \nu_{(j)}$ with $i > j$.

$$\nu = \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right)$$

					b_3	b_6
					b_5	b_8
b_1	b_2					
	b_4	b_7				

Attacking pairs: $(b_2, b_3), (b_3, b_4), (b_4, b_5), (b_4, b_6), (b_5, b_7), (b_6, b_7), (b_7, b_8)$

Key Object: LLT Polynomials

Let $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$ be a tuple of skew shapes. (Skew shape $= \lambda \setminus \mu$)

- The *content* of a box in row y , column x is $x - y$.
- *Reading order*: label boxes $b_1, \dots, b_n$ by scanning each diagonal from southwest to northeast, in order of increasing content.
- A pair $(a, b) \in \nu$ is *attacking* if a precedes b in reading order and
 - $\text{content}(b) = \text{content}(a)$, or
 - $\text{content}(b) = \text{content}(a) + 1$ and $a \in \nu_{(i)}, b \in \nu_{(j)}$ with $i > j$.

$$\nu = \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right)$$

				b_3	b_6
				b_5	b_8
b_1	b_2				
	b_4	b_7			

Attacking pairs: $(b_2, b_3), (b_3, b_4), (b_4, b_5), (b_4, b_6), (b_5, b_7), (b_6, b_7), (b_7, b_8)$

Key Object: LLT Polynomials

Let $\nu = (\nu_{(1)}, \dots, \nu_{(k)})$ be a tuple of skew shapes. (Skew shape $= \lambda \setminus \mu$)

- The *content* of a box in row y , column x is $x - y$.
- *Reading order*: label boxes $b_1, \dots, b_n$ by scanning each diagonal from southwest to northeast, in order of increasing content.
- A pair $(a, b) \in \nu$ is *attacking* if a precedes b in reading order and
 - $\text{content}(b) = \text{content}(a)$, or
 - $\text{content}(b) = \text{content}(a) + 1$ and $a \in \nu_{(i)}, b \in \nu_{(j)}$ with $i > j$.

$$\nu = \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right)$$

				b_3	b_6
				b_5	b_8
b_1	b_2				
	b_4	b_7			

Attacking pairs: $(b_2, b_3), (b_3, b_4), (b_4, b_5), (b_4, b_6), (b_5, b_7), (b_6, b_7), (b_7, b_8)$

LLT Polynomials

- A *semistandard tableau* on ν is a map $T: \nu \rightarrow \mathbb{Z}_+$ which restricts to a semistandard tableau on each $\nu_{(i)}$.

The *LLT polynomial* indexed by a tuple of skew shapes ν is

$$\mathcal{G}_\nu(\mathbf{z}; q) = \sum_{T \in \text{SSYT}(\nu)} \mathbf{z}^T,$$

$$\mathbf{z}^T = \prod_{a \in \nu} z_{T(a)}.$$

$T =$

				5	6
				1	1
2	4				
	3	5			

$$\mathbf{z}^T = z_1^2 z_2 z_3 z_4 z_5^2 z_6$$

LLT Polynomials

- A *semistandard tableau* on ν is a map $T: \nu \rightarrow \mathbb{Z}_+$ which restricts to a semistandard tableau on each $\nu_{(i)}$.
- An *attacking inversion* in T is an attacking pair (a, b) such that $T(a) > T(b)$.

The *LLT polynomial* indexed by a tuple of skew shapes ν is

$$\mathcal{G}_\nu(\mathbf{z}; q) = \sum_{T \in \text{SSYT}(\nu)} q^{\text{inv}(T)} \mathbf{z}^T,$$

where $\text{inv}(T)$ is the number of attacking inversions in T and $\mathbf{z}^T = \prod_{a \in \nu} z_{T(a)}$.

$$T = \begin{array}{|c|c|c|c|c|c|} \hline & & & & 5 & 6 \\ \hline & & & & 1 & 1 \\ \hline & & & & & \\ \hline 2 & 4 & & & & \\ \hline & 3 & 5 & & & \\ \hline \end{array}$$

$$\mathbf{z}^T = z_1^2 z_2 z_3 z_4 z_5^2 z_6$$

LLT Polynomials

- A *semistandard tableau* on ν is a map $T: \nu \rightarrow \mathbb{Z}_+$ which restricts to a semistandard tableau on each $\nu_{(i)}$.
- An *attacking inversion* in T is an attacking pair (a, b) such that $T(a) > T(b)$.

The *LLT polynomial* indexed by a tuple of skew shapes ν is

$$\mathcal{G}_\nu(\mathbf{z}; q) = \sum_{T \in \text{SSYT}(\nu)} q^{\text{inv}(T)} \mathbf{z}^T,$$

where $\text{inv}(T)$ is the number of attacking inversions in T and $\mathbf{z}^T = \prod_{a \in \nu} z_{T(a)}$.

$T =$

				5	6
				1	1
2	4				
	3	5			

non-inversion

$$\mathbf{z}^T = z_1^2 z_2 z_3 z_4 z_5^2 z_6$$

LLT Polynomials

- A *semistandard tableau* on ν is a map $T: \nu \rightarrow \mathbb{Z}_+$ which restricts to a semistandard tableau on each $\nu_{(i)}$.
- An *attacking inversion* in T is an attacking pair (a, b) such that $T(a) > T(b)$.

The *LLT polynomial* indexed by a tuple of skew shapes ν is

$$\mathcal{G}_\nu(\mathbf{z}; q) = \sum_{T \in \text{SSYT}(\nu)} q^{\text{inv}(T)} \mathbf{z}^T,$$

where $\text{inv}(T)$ is the number of attacking inversions in T and $\mathbf{z}^T = \prod_{a \in \nu} z_{T(a)}$.

$T =$

				5	6	
				1	1	
2	4					
	3	5				

inversion

$$\mathbf{z}^T = z_1^2 z_2 z_3 z_4 z_5^2 z_6$$

LLT Polynomials

- A *semistandard tableau* on ν is a map $T: \nu \rightarrow \mathbb{Z}_+$ which restricts to a semistandard tableau on each $\nu_{(i)}$.
- An *attacking inversion* in T is an attacking pair (a, b) such that $T(a) > T(b)$.

The *LLT polynomial* indexed by a tuple of skew shapes ν is

$$\mathcal{G}_\nu(\mathbf{z}; q) = \sum_{T \in \text{SSYT}(\nu)} q^{\text{inv}(T)} \mathbf{z}^T,$$

where $\text{inv}(T)$ is the number of attacking inversions in T and $\mathbf{z}^T = \prod_{a \in \nu} z_{T(a)}$.

$T =$

				5	6	
				1	1	
2	4					
	3	5				

inversion

$$\mathbf{z}^T = z_1^2 z_2 z_3 z_4 z_5^2 z_6$$

LLT Polynomials

- A *semistandard tableau* on ν is a map $T: \nu \rightarrow \mathbb{Z}_+$ which restricts to a semistandard tableau on each $\nu_{(i)}$.
- An *attacking inversion* in T is an attacking pair (a, b) such that $T(a) > T(b)$.

The *LLT polynomial* indexed by a tuple of skew shapes ν is

$$\mathcal{G}_\nu(\mathbf{z}; q) = \sum_{T \in \text{SSYT}(\nu)} q^{\text{inv}(T)} \mathbf{z}^T,$$

where $\text{inv}(T)$ is the number of attacking inversions in T and $\mathbf{z}^T = \prod_{a \in \nu} z_{T(a)}$.

$T =$

				5	6
				1	1
2	4				
	3	5			

non-inversion

$$\mathbf{z}^T = z_1^2 z_2 z_3 z_4 z_5^2 z_6$$

LLT Polynomials

- A *semistandard tableau* on ν is a map $T: \nu \rightarrow \mathbb{Z}_+$ which restricts to a semistandard tableau on each $\nu_{(i)}$.
- An *attacking inversion* in T is an attacking pair (a, b) such that $T(a) > T(b)$.

The *LLT polynomial* indexed by a tuple of skew shapes ν is

$$\mathcal{G}_\nu(\mathbf{z}; q) = \sum_{T \in \text{SSYT}(\nu)} q^{\text{inv}(T)} \mathbf{z}^T,$$

where $\text{inv}(T)$ is the number of attacking inversions in T and $\mathbf{z}^T = \prod_{a \in \nu} z_{T(a)}$.

$T =$

				5	6
				1	1
2	4				
	3	5			

non-inversion

$$\mathbf{z}^T = z_1^2 z_2 z_3 z_4 z_5^2 z_6$$

LLT Polynomials

- A *semistandard tableau* on ν is a map $T: \nu \rightarrow \mathbb{Z}_+$ which restricts to a semistandard tableau on each $\nu_{(i)}$.
- An *attacking inversion* in T is an attacking pair (a, b) such that $T(a) > T(b)$.

The *LLT polynomial* indexed by a tuple of skew shapes ν is

$$\mathcal{G}_\nu(\mathbf{z}; q) = \sum_{T \in \text{SSYT}(\nu)} q^{\text{inv}(T)} \mathbf{z}^T,$$

where $\text{inv}(T)$ is the number of attacking inversions in T and $\mathbf{z}^T = \prod_{a \in \nu} z_{T(a)}$.

$$T = \begin{array}{|c|c|c|c|c|c|} \hline & & & & 5 & 6 \\ \hline & & & & 1 & 1 \\ \hline & & & & & \\ \hline 2 & 4 & & & & \\ \hline & 3 & 5 & & & \\ \hline \end{array} \quad \text{inversion}$$

$$\mathbf{z}^T = z_1^2 z_2 z_3 z_4 z_5^2 z_6$$

LLT Polynomials

- A *semistandard tableau* on ν is a map $T: \nu \rightarrow \mathbb{Z}_+$ which restricts to a semistandard tableau on each $\nu_{(i)}$.
- An *attacking inversion* in T is an attacking pair (a, b) such that $T(a) > T(b)$.

The *LLT polynomial* indexed by a tuple of skew shapes ν is

$$\mathcal{G}_\nu(\mathbf{z}; q) = \sum_{T \in \text{SSYT}(\nu)} q^{\text{inv}(T)} \mathbf{z}^T,$$

where $\text{inv}(T)$ is the number of attacking inversions in T and $\mathbf{z}^T = \prod_{a \in \nu} z_{T(a)}$.

$$T = \begin{array}{|c|c|c|c|c|c|} \hline & & & & 5 & 6 \\ \hline & & & & 1 & 1 \\ \hline & & & & & \\ \hline 2 & 4 & & & & \\ \hline & 3 & 5 & & & \\ \hline \end{array} \quad \text{inversion}$$

$$\mathbf{z}^T = z_1^2 z_2 z_3 z_4 z_5^2 z_6$$

LLT Polynomials

- A *semistandard tableau* on ν is a map $T: \nu \rightarrow \mathbb{Z}_+$ which restricts to a semistandard tableau on each $\nu_{(i)}$.
- An *attacking inversion* in T is an attacking pair (a, b) such that $T(a) > T(b)$.

The *LLT polynomial* indexed by a tuple of skew shapes ν is

$$\mathcal{G}_\nu(\mathbf{z}; q) = \sum_{T \in \text{SSYT}(\nu)} q^{\text{inv}(T)} \mathbf{z}^T,$$

where $\text{inv}(T)$ is the number of attacking inversions in T and $\mathbf{z}^T = \prod_{a \in \nu} z_{T(a)}$.

$$T = \begin{array}{|c|c|c|c|c|c|} \hline & & & & 5 & 6 \\ \hline & & & & 1 & 1 \\ \hline & & & & & \\ \hline 2 & 4 & & & & \\ \hline & 3 & 5 & & & \\ \hline \end{array}$$

$$\text{inv}(T) = 4, \quad \mathbf{z}^T = z_1^2 z_2 z_3 z_4 z_5^2 z_6$$

LLT Polynomials $\mathcal{G}_\nu(X; q)$

- $\mathcal{G}_\nu(X; q)$ is a symmetric function

LLT Polynomials $\mathcal{G}_\nu(X; q)$

- $\mathcal{G}_\nu(X; q)$ is a symmetric function
- $\mathcal{G}_\nu(X; 1) = s_{\nu(1)} \cdots s_{\nu(r)}$

LLT Polynomials $\mathcal{G}_\nu(X; q)$

- $\mathcal{G}_\nu(X; q)$ is a symmetric function
- $\mathcal{G}_\nu(X; 1) = s_{\nu(1)} \cdots s_{\nu(r)}$
- $\mathcal{G}_\nu$ were originally defined by Lascoux, Leclerc, and Thibon to explore connections to Fock space representations of $U_q(\hat{\mathfrak{sl}}_r)$

LLT Polynomials $\mathcal{G}_\nu(X; q)$

- $\mathcal{G}_\nu(X; q)$ is a symmetric function
- $\mathcal{G}_\nu(X; 1) = s_{\nu(1)} \cdots s_{\nu(r)}$
- $\mathcal{G}_\nu$ were originally defined by Lascoux, Leclerc, and Thibon to explore connections to Fock space representations of $U_q(\hat{\mathfrak{sl}}_r)$
- $\mathcal{G}_\nu$ is Schur-positive for any tuple of skew shapes ν [Grojnowski-Haiman, 2007].

Example ∇e_3

$$\lambda \rightarrow q^{\text{dinv}(\lambda)} t^{\text{area}(\lambda)} \rightarrow q^{\text{dinv}(\lambda)} t^{\text{area}(\lambda)} \omega \mathcal{G}_{\nu(\lambda)}(X; q^{-1})$$

Example ∇e_3

$$\lambda \quad q^{\text{dinv}(\lambda)} t^{\text{area}(\lambda)} \quad q^{\text{dinv}(\lambda)} t^{\text{area}(\lambda)} \omega \mathcal{G}_{\nu(\lambda)}(X; q^{-1})$$



Example ∇e_3

$$\lambda \quad q^{\text{dinv}(\lambda)} t^{\text{area}(\lambda)} \quad q^{\text{dinv}(\lambda)} t^{\text{area}(\lambda)} \omega \mathcal{G}_{\nu(\lambda)}(X; q^{-1})$$



$$q^3$$



$$q^2 t$$



$$qt$$



$$qt^2$$



$$t^3$$

Example ∇e_3

λ	$q^{\text{dinv}(\lambda)} t^{\text{area}(\lambda)}$	$q^{\text{dinv}(\lambda)} t^{\text{area}(\lambda)} \omega \mathcal{G}_{\nu(\lambda)}(X; q^{-1})$
	q^3	$s_3 + qs_{2,1} + q^2 s_{2,1} + q^3 s_{1,1,1}$
	$q^2 t$	$qts_{2,1} + q^2 ts_{1,1,1}$
	qt	$ts_{2,1} + qts_{1,1,1}$
	qt^2	$t^2 s_{2,1} + qt^2 s_{1,1,1}$
	t^3	$t^3 s_{1,1,1}$

Example ∇e_3

λ	$q^{\text{dinv}(\lambda)} t^{\text{area}(\lambda)}$	$q^{\text{dinv}(\lambda)} t^{\text{area}(\lambda)} \omega \mathcal{G}_{\nu(\lambda)}(X; q^{-1})$
	q^3	$s_3 + qs_{2,1} + q^2 s_{2,1} + q^3 s_{1,1,1}$
	$q^2 t$	$qts_{2,1} + q^2 ts_{1,1,1}$
	qt	$ts_{2,1} + qts_{1,1,1}$
	qt^2	$t^2 s_{2,1} + qt^2 s_{1,1,1}$
	t^3	$t^3 s_{1,1,1}$

- Entire quantity is q, t -symmetric

Example ∇e_3

λ	$q^{\text{dinv}(\lambda)} t^{\text{area}(\lambda)}$	$q^{\text{dinv}(\lambda)} t^{\text{area}(\lambda)} \omega \mathcal{G}_{\nu(\lambda)}(X; q^{-1})$
	q^3	$s_3 + qs_{2,1} + q^2 s_{2,1} + q^3 s_{1,1,1}$
	$q^2 t$	$qts_{2,1} + q^2 ts_{1,1,1}$
	qt	$ts_{2,1} + qts_{1,1,1}$
	qt^2	$t^2 s_{2,1} + qt^2 s_{1,1,1}$
	t^3	$t^3 s_{1,1,1}$

- Entire quantity is q, t -symmetric
- Coefficient of $s_{1,1,1}$ in sum is a “ (q, t) -Catalan number” $(q^3 + q^2 t + qt + qt^2 + t^3)$.

Generalizing Shuffle Theorem

When a problem is too difficult, try generalizing!

Generalizing Shuffle Theorem

When a problem is too difficult, try generalizing!

Algebraic Expression

Combinatorial Expression

$$\nabla e_k = \sum q, t\text{-weighted Dyck paths}$$

Generalizing Shuffle Theorem

When a problem is too difficult, try generalizing!

Algebraic Expression

Combinatorial Expression

$$\nabla e_k = \sum q, t\text{-weighted Dyck paths}$$

What generalizes ∇e_k ?

Elliptic Hall Algebra

Algebra $\mathcal{E} \curvearrowright \Lambda =$ symmetric polynomials

Elliptic Hall Algebra

Algebra $\mathcal{E} \curvearrowright \Lambda =$ symmetric polynomials

$\mathcal{E}$ comes from algebraic geometry

Elliptic Hall Algebra

Algebra $\mathcal{E} \curvearrowright \Lambda =$ symmetric polynomials

$\mathcal{E}$ comes from algebraic geometry


$$\mathcal{E} \cong_{\text{v.sp.}} \begin{matrix} \text{central} \\ \text{subalgebra} \end{matrix} \oplus \bigoplus_{m,n \text{ coprime}} \Lambda^{(m,n)}$$

Elliptic Hall Algebra

Algebra $\mathcal{E} \curvearrowright \Lambda =$ symmetric polynomials

$\mathcal{E}$ comes from algebraic geometry

$$\mathcal{E} \cong_{\text{v.sp.}} \text{central subalgebra} \oplus \bigoplus_{m,n \text{ coprime}} \Lambda^{(m,n)}$$




Elliptic Hall Algebra

Algebra $\mathcal{E} \curvearrowright \Lambda =$ symmetric polynomials

$\mathcal{E}$ comes from algebraic geometry

$$\mathcal{E} \cong_{\text{v.sp.}} \text{central subalgebra} \bigoplus_{m,n \text{ coprime}} \bigoplus \Lambda^{(m,n)}$$



Each $\Lambda^{(m,n)} \cong$ symmetric polynomials

Elliptic Hall Algebra


Algebra $\mathcal{E} \curvearrowright \Lambda =$ symmetric polynomials

$\mathcal{E}$ comes from algebraic geometry

$$\mathcal{E} \cong_{\text{v.sp.}} \text{central subalgebra} \bigoplus_{m,n \text{ coprime}} \bigoplus \Lambda^{(m,n)}$$

Each $\Lambda^{(m,n)} \cong$ symmetric polynomials

- For $f^{(m,n)} \in \Lambda^{(m,n)}$, $\nabla f^{(m,n)} \nabla^{-1} = f^{(m+n,n)}$



Elliptic Hall Algebra


Algebra $\mathcal{E} \curvearrowright \Lambda =$ symmetric polynomials

$\mathcal{E}$ comes from algebraic geometry

$$\mathcal{E} \cong \text{central v.sp. subalgebra} \oplus \bigoplus_{m,n \text{ coprime}} \Lambda^{(m,n)}$$

Each $\Lambda^{(m,n)} \cong$ symmetric polynomials

- For $f^{(m,n)} \in \Lambda^{(m,n)}$, $\nabla f^{(m,n)} \nabla^{-1} = f^{(m+n,n)}$
- Algebraic side of Shuffle Theorem = $e_k^{(1,1)} \in \Lambda^{(1,1)}$ acting on $1 \in \Lambda$.




Elliptic Hall Algebra

Algebra $\mathcal{E} \curvearrowright \Lambda =$ symmetric polynomials

$\mathcal{E}$ comes from algebraic geometry

$$\mathcal{E} \cong_{\text{v.sp. subalgebra}} \text{central} \bigoplus_{m,n \text{ coprime}} \bigoplus \Lambda^{(m,n)}$$




- For $f^{(m,n)} \in \Lambda^{(m,n)}$, $\nabla f^{(m,n)} \nabla^{-1} = f^{(m+n,n)}$
- Algebraic side of Shuffle Theorem = $e_k^{(1,1)} \in \Lambda^{(1,1)}$ acting on $1 \in \Lambda$.
- $e_k^{(1,1)} \cdot 1 = \nabla e_k^{(0,1)} \nabla^{-1} \cdot 1 = \nabla e_k$.

Elliptic Hall Algebra

Algebra $\mathcal{E} \curvearrowright \Lambda =$ symmetric polynomials

$\mathcal{E}$ comes from algebraic geometry

$$\mathcal{E} \cong_{\text{v.sp. subalgebra}} \text{central} \bigoplus_{m,n \text{ coprime}} \bigoplus \Lambda^{(m,n)}$$



- For $f^{(m,n)} \in \Lambda^{(m,n)}$, $\nabla f^{(m,n)} \nabla^{-1} = f^{(m+n,n)}$
- Algebraic side of Shuffle Theorem = $e_k^{(1,1)} \in \Lambda^{(1,1)}$ acting on $1 \in \Lambda$.
- $e_k^{(1,1)} \cdot 1 = \nabla e_k^{(0,1)} \nabla^{-1} \cdot 1 = \nabla e_k$.
- Can be difficult to work with in general. Can we make it more explicit?

Root ideals

$R_+ = \{\alpha_{ij} \mid 1 \leq i < j \leq n\}$ denotes the set of positive roots for GL_n , where $\alpha_{ij} = \epsilon_i - \epsilon_j$.

	(12)	(13)	(14)	(15)
		(23)	(24)	(25)
			(34)	(35)
				(45)

Root ideals

$R_+ = \{\alpha_{ij} \mid 1 \leq i < j \leq n\}$ denotes the set of positive roots for GL_n , where $\alpha_{ij} = \epsilon_i - \epsilon_j$.

	(12)	(13)	(14)	(15)
		(23)	(24)	(25)
			(34)	(35)
				(45)

A root ideal $\Psi \subseteq R_+$ is an upper order ideal of positive roots.

	(12)	(13)	(14)	(15)
		(23)	(24)	(25)
			(34)	(35)
				(45)

$\Psi = \text{Roots above Dyck path}$

Weyl symmetrization

Define the *Weyl symmetrization operator*

$\sigma: \mathbb{Q}[z_1^{\pm 1}, \dots, z_n^{\pm 1}] \rightarrow \mathbb{Q}[z_1^{\pm 1}, \dots, z_n^{\pm 1}]^{S_n}$ by

$$f(\mathbf{z}) \mapsto \sum_{w \in S_n} w \left(\frac{f(\mathbf{z})}{\prod_{i < j} (1 - z_j/z_i)} \right).$$

Weyl symmetrization

Define the *Weyl symmetrization operator*

$\sigma: \mathbb{Q}[z_1^{\pm 1}, \dots, z_n^{\pm 1}] \rightarrow \mathbb{Q}[z_1^{\pm 1}, \dots, z_n^{\pm 1}]^{S_n}$ by

$$f(\mathbf{z}) \mapsto \sum_{w \in S_n} w \left(\frac{f(\mathbf{z})}{\prod_{i < j} (1 - z_j/z_i)} \right).$$

$\sigma(z_1^{\gamma_1} \cdots z_n^{\gamma_n}) = \pm$ irreducible GL_n character or 0.

Weyl symmetrization

Define the *Weyl symmetrization operator*

$\sigma: \mathbb{Q}[z_1^{\pm 1}, \dots, z_n^{\pm 1}] \rightarrow \mathbb{Q}[z_1^{\pm 1}, \dots, z_n^{\pm 1}]^{S_n}$ by

$$f(\mathbf{z}) \mapsto \sum_{w \in S_n} w \left(\frac{f(\mathbf{z})}{\prod_{i < j} (1 - z_j/z_i)} \right).$$

$\sigma(z_1^{\gamma_1} \cdots z_n^{\gamma_n}) = \pm$ irreducible GL_n character or 0.

Example

$$\sigma(\mathbf{z}^{111} + \mathbf{z}^{201} + \mathbf{z}^{210} + \mathbf{z}^{3-11}) = \chi_{111} + 0 + \chi_{210} - \chi_{300}$$

Weyl symmetrization

Define the *Weyl symmetrization operator*

$\sigma: \mathbb{Q}[z_1^{\pm 1}, \dots, z_n^{\pm 1}] \rightarrow \mathbb{Q}[z_1^{\pm 1}, \dots, z_n^{\pm 1}]^{S_n}$ by

$$f(\mathbf{z}) \mapsto \sum_{w \in S_n} w \left(\frac{f(\mathbf{z})}{\prod_{i < j} (1 - z_j/z_i)} \right).$$

$\sigma(z_1^{\gamma_1} \cdots z_n^{\gamma_n}) = \pm$ irreducible GL_n character or 0.

Example

$$\sigma(\mathbf{z}^{111} + \mathbf{z}^{201} + \mathbf{z}^{210} + \mathbf{z}^{3-11}) = \chi_{111} + 0 + \chi_{210} - \chi_{300}$$

- For λ a partition, $\chi_\lambda(\mathbf{z}) = s_\lambda(\mathbf{z})$.

Weyl symmetrization

Define the *Weyl symmetrization operator*

$\sigma: \mathbb{Q}[z_1^{\pm 1}, \dots, z_n^{\pm 1}] \rightarrow \mathbb{Q}[z_1^{\pm 1}, \dots, z_n^{\pm 1}]^{S_n}$ by

$$f(\mathbf{z}) \mapsto \sum_{w \in S_n} w \left(\frac{f(\mathbf{z})}{\prod_{i < j} (1 - z_j/z_i)} \right).$$

$\sigma(z_1^{\gamma_1} \cdots z_n^{\gamma_n}) = \pm$ irreducible GL_n character or 0.

Example

$$\sigma(\mathbf{z}^{111} + \mathbf{z}^{201} + \mathbf{z}^{210} + \mathbf{z}^{3-11}) = \chi_{111} + 0 + \chi_{210} - \chi_{300}$$

- For λ a partition, $\chi_\lambda(\mathbf{z}) = s_\lambda(\mathbf{z})$.
- Define $\text{pol}_\chi \chi_\lambda(\mathbf{z}) = s_\lambda$ if $\lambda_i \geq 0$, otherwise 0.

Catalanimals

Definition

The *Catalanimal* indexed by $R_q, R_t, R_{qt} \subseteq R_+$ and $\lambda \in \mathbb{Z}^n$ is

Catalan animals

Definition

The *Catalan animal* indexed by $R_q, R_t, R_{qt} \subseteq R_+$ and $\lambda \in \mathbb{Z}^n$ is

$$H(R_q, R_t, R_{qt}, \lambda) = \text{pol}_X \sigma \left(\frac{z^\lambda \prod_{\alpha \in R_{qt}} (1 - qt z^\alpha)}{\prod_{\alpha \in R_q} (1 - q z^\alpha) \prod_{\alpha \in R_t} (1 - t z^\alpha)} \right),$$

where $z^{\alpha_{ij}} = z_i/z_j$ and $(1 - tz_i/z_j)^{-1} = 1 + tz_i/z_j + t^2 z_i^2/z_j^2 + \dots$.

Catalanimals

Definition

The *Catalanimal* indexed by $R_q, R_t, R_{qt} \subseteq R_+$ and $\lambda \in \mathbb{Z}^n$ is

$$H(R_q, R_t, R_{qt}, \lambda) = \text{pol}_X \sigma \left(\frac{z^\lambda \prod_{\alpha \in R_{qt}} (1 - qt z^\alpha)}{\prod_{\alpha \in R_q} (1 - q z^\alpha) \prod_{\alpha \in R_t} (1 - t z^\alpha)} \right),$$

where $z^{\alpha_{ij}} = z_i/z_j$ and $(1 - tz_i/z_j)^{-1} = 1 + tz_i/z_j + t^2 z_i^2/z_j^2 + \dots$.

With $n = 3$, $R_+ =$

$$H(R_+, R_+, \{\alpha_{13}\}, (111)) =$$

Definition

The *Catalanimal* indexed by $R_q, R_t, R_{qt} \subseteq R_+$ and $\lambda \in \mathbb{Z}^n$ is

$$H(R_q, R_t, R_{qt}, \lambda) = \text{pol}_X \sigma \left(\frac{z^\lambda \prod_{\alpha \in R_{qt}} (1 - qt z^\alpha)}{\prod_{\alpha \in R_q} (1 - q z^\alpha) \prod_{\alpha \in R_t} (1 - t z^\alpha)} \right),$$

where $z^{\alpha_{ij}} = z_i/z_j$ and $(1 - tz_i/z_j)^{-1} = 1 + tz_i/z_j + t^2 z_i^2/z_j^2 + \dots$.

With $n = 3$, $R_+ =$

$$\begin{aligned} H(R_+, R_+, \{\alpha_{13}\}, (111)) &= \text{pol}_X \sigma \left(\frac{z^{111} (1 - qt z_1/z_3)}{\prod_{1 \leq i < j \leq 3} (1 - q z_i/z_j) (1 - t z_i/z_j)} \right) \\ &= s_{111} + (q + t + q^2 + qt + t^2) s_{21} + (qt + q^3 + q^2 t + qt^2 + t^3) s_3 \\ &= \omega \nabla e_3. \end{aligned}$$

Why?

Let $R_+ = \{\alpha_{ij} \mid 1 \leq i < j \leq l\}$ and $R_+^0 = \{\alpha_{ij} \in R_+ \mid i + 1 < j\}$.

Why?

Let $R_+ = \{\alpha_{ij} \mid 1 \leq i < j \leq l\}$ and $R_+^0 = \{\alpha_{ij} \in R_+ \mid i+1 < j\}$.

Proposition

For $(m, n) \in \mathbb{Z}_+^2$ coprime,

$$e_k^{(m,n)} \cdot 1 = H(R_+, R_+, R_+^0, \mathbf{b})$$

for

Why?

Let $R_+ = \{\alpha_{ij} \mid 1 \leq i < j \leq l\}$ and $R_+^0 = \{\alpha_{ij} \in R_+ \mid i+1 < j\}$.


Proposition

For $(m, n) \in \mathbb{Z}_+^2$ coprime,

$$e_k^{(m,n)} \cdot 1 = H(R_+, R_+, R_+^0, \mathbf{b})$$

for $\mathbf{b} = (b_0, \dots, b_{km-1})$ satisfying $b_i =$ the number of south steps on vertical line $x = i$ of highest lattice path under line $y + \frac{n}{m}x = n$.

δ = highest Dyck path.



δ


$$\mathbf{b} = (1, 1, 0, 1, 0, 1, 0, 1, 0, 1, 0)$$

Results

Manipulating Catalanimal $\implies$ a proof of the Rational Shuffle Theorem + a generalization.

Theorem (Blasiak-Haiman-Morse-Pun-S., 2023)

Given $r, s \in \mathbb{R}_{>0}$ such that $p = s/r$ irrational, take $\mathbf{b} = (b_1, \dots, b_l) \in \mathbb{Z}^l$ to be the south step sequence of highest path δ under the line $y + px = s$.




Results

Manipulating Catalanimal $\implies$ a proof of the Rational Shuffle Theorem + a generalization.

Theorem (Blasiak-Haiman-Morse-Pun-S., 2023)

Given $r, s \in \mathbb{R}_{>0}$ such that $p = s/r$ irrational, take $\mathbf{b} = (b_1, \dots, b_l) \in \mathbb{Z}^l$ to be the south step sequence of highest path δ under the line $y + px = s$.

$$H(R_+, R_+, R_+^0, \mathbf{b})$$




Results

Manipulating Catalanimal $\implies$ a proof of the Rational Shuffle Theorem + a generalization.

Theorem (Blasiak-Haiman-Morse-Pun-S., 2023)

Given $r, s \in \mathbb{R}_{>0}$ such that $p = s/r$ irrational, take $\mathbf{b} = (b_1, \dots, b_l) \in \mathbb{Z}^l$ to be the south step sequence of highest path δ under the line $y + px = s$.

$$H(R_+, R_+, R_+^0, \mathbf{b}) =$$



Results


Manipulating Catalanimal $\implies$ a proof of the Rational Shuffle Theorem + a generalization.

Theorem (Blasiak-Haiman-Morse-Pun-S., 2023)

Given $r, s \in \mathbb{R}_{>0}$ such that $p = s/r$ irrational, take $\mathbf{b} = (b_1, \dots, b_l) \in \mathbb{Z}^l$ to be the south step sequence of highest path δ under the line $y + px = s$.

$$H(R_+, R_+, R_+^0, \mathbf{b}) = \sum_{\lambda} \omega \mathcal{G}_{\nu(\lambda)}(X; q^{-1})$$

where summation is over all lattice paths under the line $y + px = s$,



Results


Manipulating Catalan animal $\implies$ a proof of the Rational Shuffle Theorem + a generalization.

Theorem (Blasiak-Haiman-Morse-Pun-S., 2023)

Given $r, s \in \mathbb{R}_{>0}$ such that $p = s/r$ irrational, take $\mathbf{b} = (b_1, \dots, b_l) \in \mathbb{Z}^l$ to be the south step sequence of highest path δ under the line $y + px = s$.

$$H(R_+, R_+, R_+^0, \mathbf{b}) = \sum_{\lambda} t^{\text{area}(\lambda)} q^{\text{div}_p(\lambda)} \omega \mathcal{G}_{\nu(\lambda)}(X; q^{-1})$$

where summation is over all lattice paths under the line $y + px = s$,



$\text{area}(\lambda)$ as before

$\text{div}_p(\lambda) = \#p\text{-balanced hooks}$

$$\frac{\ell}{a+1} < p < \frac{\ell+1}{a}$$

A Question

Why stop at $e_k^{(m,n)}$?

A Question

Why stop at $e_k^{(m,n)}$?

For which symmetric functions f can we find a Catalan animal such that $f^{(m,n)} \cdot 1 = \text{a Catalan animal}$?

A Question

Why stop at $e_k^{(m,n)}$?

For which symmetric functions f can we find a Catalan animal such that $f^{(m,n)} \cdot 1 = \text{a Catalan animal}$?

Answer: for f equal to any LLT polynomial!

A Question

Why stop at $e_k^{(m,n)}$?

For which symmetric functions f can we find a Catalan animal such that $f^{(m,n)} \cdot 1 = \text{a Catalan animal}$?

Answer: for f equal to any LLT polynomial!

Special case: $\mathcal{G}_\nu^{(1,1)} \cdot 1 = \nabla \mathcal{G}_\nu(X; q)$.

LLT Catalanimals

For a tuple of skew shapes ν , the *LLT Catalanimal* $H_\nu = H(R_q, R_t, R_{qt}, \lambda)$ is determined by

- $R_+ \supseteq R_q \supseteq R_t \supseteq R_{qt}$,

LLT Catalananimals

For a tuple of skew shapes ν , the *LLT Catalananimal* $H_\nu = H(R_q, R_t, R_{qt}, \lambda)$ is determined by

- $R_+ \supseteq R_q \supseteq R_t \supseteq R_{qt}$,
- $R_+ \setminus R_q$ = pairs of boxes in the same diagonal in the same shape,
- $R_q \setminus R_t$ = the attacking pairs,
- $R_t \setminus R_{qt}$ = pairs going between adjacent diagonals,

LLT Catalananimals

For a tuple of skew shapes ν , the *LLT Catalananimal* $H_\nu = H(R_q, R_t, R_{qt}, \lambda)$ is determined by


- $R_+ \supseteq R_q \supseteq R_t \supseteq R_{qt}$,
- $R_+ \setminus R_q$ = pairs of boxes in the same diagonal in the same shape,
- $R_q \setminus R_t$ = the attacking pairs,
- $R_t \setminus R_{qt}$ = pairs going between adjacent diagonals,
- λ : fill each diagonal D of ν with $1 + \chi(D \text{ contains a row start}) - \chi(D \text{ contains a row end})$.
Listing this filling in reading order gives λ .

LLT Catalanimals


- $R_+ \setminus R_q =$ pairs of boxes in the same diagonal in the same shape,
- $R_q \setminus R_t =$ the attacking pairs,
- $R_t \setminus R_{qt} =$ pairs going between adjacent diagonals,
- $R_{qt} =$ all other pairs,

λ : fill each diagonal D of ν with

$1 + \chi(D \text{ contains a row start}) - \chi(D \text{ contains a row end})$.



ν




LLT Catalanimals


- $R_+ \setminus R_q =$ pairs of boxes in the same diagonal in the same shape,
- $R_q \setminus R_t =$ the attacking pairs,
- $R_t \setminus R_{qt} =$ pairs going between adjacent diagonals,
- $R_{qt} =$ all other pairs,

λ : fill each diagonal D of ν with

$1 + \chi(D \text{ contains a row start}) - \chi(D \text{ contains a row end})$.



λ , as a filling of ν



Theorem (Blasiak-Haiman-Morse-Pun-S., 2024)

Let ν be a tuple of skew shapes and let $H_\nu = H(R_q, R_t, R_{qt}, \lambda)$ be the associated LLT Catalanimal. Then

$$\begin{aligned}\nabla \mathcal{G}_\nu(X; q) &= c_\nu \omega H_\nu \\ &= c_\nu \omega \sigma \left(\frac{z^\lambda \prod_{\alpha \in R_{qt}} (1 - qt z^\alpha)}{\prod_{\alpha \in R_q} (1 - q z^\alpha) \prod_{\alpha \in R_t} (1 - t z^\alpha)} \right)\end{aligned}$$

for some $c_\nu \in \pm q^{\mathbb{Z}} t^{\mathbb{Z}}$.

What about Macdonald polynomials?!

- Remember $\nabla \tilde{H}_\mu = q^{n(\mu)} t^{n(\mu^*)} \tilde{H}_\mu$.

What about Macdonald polynomials?!

- Remember $\nabla \tilde{H}_\mu = q^{n(\mu)} t^{n(\mu^*)} \tilde{H}_\mu$.
- We have a formula for $\nabla \mathcal{G}_\nu$.

What about Macdonald polynomials?!

- Remember $\nabla \tilde{H}_\mu = q^{n(\mu)} t^{n(\mu^*)} \tilde{H}_\mu$.
- We have a formula for $\nabla \mathcal{G}_\nu$.
- Does there exist formula $\tilde{H}_\mu = \sum_\nu a_{\mu\nu}(q, t) \mathcal{G}_\nu$?

What about Macdonald polynomials?!

- Remember $\nabla \tilde{H}_\mu = q^{n(\mu)} t^{n(\mu^*)} \tilde{H}_\mu$.
- We have a formula for $\nabla \mathcal{G}_\nu$.
- Does there exist formula $\tilde{H}_\mu = \sum_\nu a_{\mu\nu}(q, t) \mathcal{G}_\nu$? Yes!

Outline










- ① Background on symmetric functions and Macdonald polynomials
- ② Shuffle theorems, combinatorics, and LLT polynomials
- ③ **A new formula for Macdonald polynomials**

Haglund-Haiman-Loehr formula example

$$\tilde{H}_\mu(X; q, t) = \sum_D \left(\prod_{u \in D} q^{-\text{arm}(u)} t^{\text{leg}(u)+1} \right) \mathcal{G}_{\nu(\mu, D)}(X; q)$$

Haglund-Haiman-Loehr formula example

$$\tilde{H}_\mu(X; q, t) = \sum_D \left(\prod_{u \in D} q^{-\text{arm}(u)} t^{\text{leg}(u)+1} \right) \mathcal{G}_{\nu(\mu, D)}(X; q)$$



Putting it all together

- Take HHL formula $\tilde{H}_\mu = \sum_D a_{\mu,D} \mathcal{G}_{\nu(\mu,D)}$ and apply $\omega \nabla$.

Putting it all together

- Take HHL formula $\tilde{H}_\mu = \sum_D a_{\mu,D} \mathcal{G}_{\nu(\mu,D)}$ and apply $\omega \nabla$.
- By construction, all the LLT Catalan animals $H_{\nu(\mu,D)}$ appearing on the RHS will have the same root ideal data (R_q, R_t, R_{qt}) .

Putting it all together

- Take HHL formula $\tilde{H}_\mu = \sum_D a_{\mu,D} \mathcal{G}_{\nu(\mu,D)}$ and apply $\omega \nabla$.
- By construction, all the LLT Catalan animals $H_{\nu(\mu,D)}$ appearing on the RHS will have the same root ideal data (R_q, R_t, R_{qt}) .
- Collect terms to get $\prod_{(b_i, b_j) \in V(\mu)} (1 - q^{\text{arm}(b_i)+1} t^{-\text{leg}(b_i)} z_i/z_j)$ factor for $V(\mu)$ the set of vertical dominoes (b_i, b_j) in μ .

$$\tilde{H}_\mu = \omega \text{pol}_X \sigma \left(z_1 \cdots z_n \frac{\prod_{\substack{\alpha_{ij} \in V(\mu)}} (1 - q^{\text{arm}(b_i)+1} t^{-\text{leg}(b_i)} z_i/z_j) \prod_{\alpha \in \hat{R}_\mu} (1 - qtz^\alpha)}{\prod_{\alpha \in R_+} (1 - qz^\alpha) \prod_{\alpha \in R_\mu} (1 - tz^\alpha)} \right).$$

The root ideal R_μ

b_1		
b_2	b_3	
b_4	b_5	b_6
b_7	b_8	b_9

row reading order

$$b_1 \prec b_2 \prec \cdots \prec b_n$$


$$R_\mu := \{\alpha_{ij} \in R_+ \mid \text{south}(b_i) \preceq b_j\},$$

$$\widehat{R}_\mu := \{\alpha_{ij} \in R_+ \mid \text{south}(b_i) \prec b_j\},$$

$$R_\mu \setminus \widehat{R}_\mu \leftrightarrow V(\mu) = \text{vertical dominoes in } \mu$$

Example:

$$R_{3321} =$$



The root ideal R_μ

b_1		
b_2	b_3	
b_4	b_5	b_6
b_7	b_8	b_9


row reading order
 $b_1 \prec b_2 \prec \cdots \prec b_n$

$$R_\mu := \{\alpha_{ij} \in R_+ \mid \text{south}(b_i) \preceq b_j\},$$

$$\hat{R}_\mu := \{\alpha_{ij} \in R_+ \mid \text{south}(b_i) \prec b_j\},$$

$$R_\mu \setminus \hat{R}_\mu \leftrightarrow V(\mu) = \text{vertical dominoes in } \mu$$


Example:




Remark

$$\tilde{H}_\mu(X; 0, t) = \omega \operatorname{pol}_X \sigma \left(\frac{z_1 \cdots z_n}{\prod_{\alpha \in R_\mu} (1 - t z^\alpha)} \right)$$

Example




partition $\mu = 22211$



Example

$1 - q^{\frac{z_1}{z_2}}$	
$1 - qt^{-1} \frac{z_2}{z_3}$	
$1 - q^2 t^{-2} \frac{z_3}{z_5}$	$1 - q^{\frac{z_4}{z_6}}$
$1 - q^2 t^{-3} \frac{z_5}{z_7}$	$1 - qt^{-1} \frac{z_6}{z_8}$

numerator factors $1 - q^{\text{arm}+1} t^{-\text{leg}} z_i/z_j$



● $R_\mu \setminus \hat{R}_\mu$ (t factors)

■ $\hat{R}_\mu$ (t and qt factors)

A positivity conjecture

What can this formula tell us that other formulas for Macdonald polynomials do not?

A positivity conjecture

What can this formula tell us that other formulas for Macdonald polynomials do not?

$$\tilde{H}_\mu^{(s)} = \omega \operatorname{pol}_X \sigma \left((z_1 \cdots z_n)^s \frac{\prod_{\alpha_{ij} \in R_\mu \setminus \hat{R}_\mu} (1 - q^{\operatorname{arm}(b_i)+1} t^{-\operatorname{leg}(b_i)} z_i / z_j) \prod_{\alpha \in \hat{R}_\mu} (1 - q t z^\alpha)}{\prod_{\alpha \in R_+} (1 - q z^\alpha) \prod_{\alpha \in R_\mu} (1 - t z^\alpha)} \right)$$

Conjecture (Blasiak-Haiman-Morse-Pun-S., 2025)

For any partition μ and positive integer s , the symmetric function $\tilde{H}_\mu^{(s)}$ is Schur positive. That is, the coefficients in

$$\tilde{H}_\mu^{(s)} = \sum_{\nu} K_{\nu, \mu}^{(s)}(q, t) s_\nu$$

satisfy $K_{\nu, \mu}^{(s)}(q, t) \in \mathbb{N}[q, t]$.

Symmetric functions, representation theory, and combinatorics

Symmetric function	Representation theory	Combinatorics
s_λ	Irreducible V_λ	SSYT(λ)
$\tilde{H}_\lambda(X; q, t)$	Garsia-Haiman M_λ	HHL
∇e_n	DH_n	Shuffle theorem
$H(R_+, R_+, R_+^0, \mathbf{b})$	??	Generalized shuffle theorem
$\tilde{H}_\lambda^{(s)}(X; q, t)$	??	??

Thank you!

- Blasiak, Jonah, Mark Haiman, Jennifer Morse, Anna Pun, and George H. Seelinger. 2023/ed. *A Shuffle Theorem for Paths Under Any Line*, Forum of Mathematics, Pi **11**, e5, DOI 10.1017/fmp.2023.4.
- . 2024. *LLT Polynomials in the Schiffmann Algebra*, Journal für die reine und angewandte Mathematik (Crelles Journal) **811**, 93–133, DOI 10.1515/crelle-2024-0012.
- . 2025. *A Raising Operator Formula for Macdonald Polynomials*, Forum of Math, Sigma **13**, e47, DOI 10.1017/fms.2025.8.
- Burban, Igor and Olivier Schiffmann. 2012. *On the Hall algebra of an elliptic curve, I*, Duke Math. J. **161**, no. 7, 1171–1231, DOI 10.1215/00127094-1593263. MR2922373
- Carlsson, Erik and Mellit, Anton. 2018. *A Proof of the Shuffle Conjecture* **31**, no. 3, 661–697, DOI 10.1090/jams/893.
- Feigin, B. L. and Tsymbaliuk, A. I. 2011. *Equivariant K-theory of Hilbert Schemes via Shuffle Algebra*, Kyoto J. Math. **51**, no. 4, 831–854.
- Garsia, Adriano M. and Mark Haiman. 1993. *A graded representation model for Macdonald's polynomials*, Proc. Nat. Acad. Sci. U.S.A. **90**, no. 8, 3607–3610, DOI 10.1073/pnas.90.8.3607. MR1214091
- Haglund, J., M. Haiman, and N. Loehr. 2005. *A Combinatorial Formula for Macdonald Polynomials* **18**, no. 3, 735–761 (electronic).
- Haglund, J. and Haiman, M. and Loehr. 2005. *A combinatorial formula for the character of the diagonal coinvariants*, Duke Math. J. **126**, no. 2, 195–232, DOI 10.1215/S0012-7094-04-12621-1.
- Haiman, Mark. 2001. *Hilbert schemes, polygraphs and the Macdonald positivity conjecture*, J. Amer. Math. Soc. **14**, no. 4, 941–1006, DOI 10.1090/S0894-0347-01-00373-3. MR1839919
- . 2002. *Vanishing theorems and character formulas for the Hilbert scheme of points in the plane*, Invent. Math. **149**, no. 2, 371–407, DOI 10.1007/s002220200219. MR1918676
- Lascoux, Alain, Bernard Leclerc, and Jean-Yves Thibon. 1995. *Ribbon tableaux, Hall-Littlewood functions and unipotent varieties*, Sémin. Lothar. Combin. **34**, Art. B34g, approx. 23. MR1399754
- Mellit, Anton. 2021. *Toric Braids and (m,n) -Parking Functions*, Duke Math. J. **170**, no. 18, 4123–4169, DOI 10.1215/00127094-2021-0011.
- Negut, Andrei. 2014. *The shuffle algebra revisited*, Int. Math. Res. Not. IMRN **22**, 6242–6275, DOI 10.1093/imrn/rnt156. MR3283004
- Schiffmann, Olivier and Vasserot, Eric. 2013. *The Elliptic Hall Algebra and the K-theory of the Hilbert Scheme of A_2* , Duke Mathematical Journal **162**, no. 2, 279–366, DOI 10.1215/00127094-1961849.

Schiffmann to Shuffle

- Shuffle algebra S given by the image of Laurent polynomials $\phi \in \mathbb{Q}(q, t)[x_1^{\pm 1}, \dots, x_l^{\pm 1}]$ via map

$$H_{q,t}: \phi \mapsto \sum_{w \in S_l} w \left(\frac{\phi \prod_{i < j} (1 - qtx_i/x_j)}{\prod_{i < j} ((1 - x_j/x_i)(1 - qx_i/x_j)(1 - tx_i/x_j))} \right)$$

- (Schiffmann-Vasserot, 2013) There exists isomorphism $\psi: S \rightarrow \mathcal{E}^+$.
- (Negut, 2014) gives well-defined

$$D_{b_1, \dots, b_l} = \psi \left(H_{q,t} \left(\frac{x_1^{b_1} \cdots x_l^{b_l}}{\prod_{i=1}^{l-1} (1 - qtx_i/x_{i+1})} \right) \right)$$

Key Relationship (BHMPs, 2023)

For $\zeta = \psi(\phi) \in \mathcal{E}^+$,

$$\omega(\zeta \cdot 1) = \omega \operatorname{pol}_X H_{q,t}(\phi).$$

Cauchy Identity

- (Twisted) non-symmetric Hall-Littlewood polynomials $E_{\lambda}^{\sigma}(x; q)$ defined via Demazure-Lusztig operators.

$$T_i = qs_i + (1 - q) \frac{s_i - 1}{1 - x_{i+1}/x_i}$$

- Dual basis F_{λ}^{σ} .

Cauchy identity

$$\frac{\prod_{i < j} (1 - q t x_i y_j)}{\prod_{i \leq j} (1 - t x_i y_j)} = \sum_{\mathbf{a} \geq 0} t^{|\mathbf{a}|} E_{\mathbf{a}}^{\sigma}(x_1, \dots, x_l; q^{-1}) F_{\mathbf{a}}^{\sigma}(y_1, \dots, y_l; q),$$

- $\mathcal{L}_{\beta/\alpha} = H_q(w_0(F_{\beta}^{\sigma^{-1}}(x; q) \overline{E_{\alpha}^{\sigma^{-1}}(x; q)}))$