Math 115 Worksheet Section 3.2

Warm-up question

 $(e^x)' = \underline{\qquad} (a^x)' = \underline{\qquad} (\sin(x))' = \underline{\qquad} (\cos(x))' = \underline{\qquad}$

Problem 1. An animal population is given by $P(t) = 300(1.044)^t$ where t is the number of years since the study of the population began. Find P'(5) and provide a practical interpretation of your result.

Problem 2. Find the quadratic polynomial $g(x) = ax^2 + bx + c$ which best fits the function $f(x) = e^x$ at x = 0, in the sense that

$$g(0) = f(0)$$
 and $g'(0) = f'(0)$ and $g''(0) = f''(0)$

Using a computer or calculator, sketch the graphs of f and g on the same axes. What do you notice?

Problem 3. Using the equation of the tangent line to the graph of e^x at x=0, show that

$$e^x > 1 + x$$

for all values of x. A sketch may be helpful.

Problem 4. Find the 50^{th} derivative of $y = \cos(x)$.

Problem 5. Find the tangent lines to $f(x) = \sin(x)$ at $x = \frac{4\pi}{3}$ and at $x = \frac{10\pi}{3}$. Graph them, along with $\sin(x)$. What do you notice? Express the second as a transformation of the first.

Problem 6. Are the following statements true or false? Give an explanation for your answer.

- (a) If f(x) is increasing, then f'(x) is increasing.
- (b) There is no function such that f'(x) = f(x) for all x besides the constant function f(x) = 0.
- (c) There is no function such that f'(x) = -f(x) for all x besides the constant function f(x) = 0.
- (d) There is no function such that f''(x) = -f(x) for all x besides the constant function f(x) = 0.
- (e) If f(x) is defined for all x, then f'(x) is defined for all x.

Problem 7. For what value(s) of a are $y = a^x$ and y = x + 1 tangent at x = 0?

Problem 8. (Winter 2017 Exam 3) A Math 115 coordinator is trying to create functions with certain properties in order to test students' understanding of various calculus concepts. She wants a function f(x) of the form

$$f(x) = \begin{cases} ax^2 + ax + be^x & \text{for } x < 0\\ a + 2\cos(x) & \text{for } x \ge 0 \end{cases}$$

where a and b are constants. Find all value(s) of a and b for which f(x) be differentiable at x = 0. Show enough work to justify your answer.

Problem 9. Explain for which values of a > 0 the function a^x is increasing and for which values it is decreasing. Is this consistent with your formula for $(a^x)'$?

Problem 10. Give an example of:

- (a) An exponential function for which the derivative is always negative.
- (b) A function f such that f'''(x) = f(x).

Problem 11. In 2009, the population of Mexico was 111 million and growing 1.13% annually, while the population of the US was 307 million and growing 0.975% annually. If we measure growth rates in people/year, which population was growing faster in 2009? Note $\ln(1.0113) \approx 0.01124$ and $\ln(1.00975) \approx 0.009702$.