

HW due Friday, 1/14

$$1.1: 14, 20, 30$$

$$1.2: 10, 44, 45 \text{ (use calculator for 45)}$$

HW due Friday, 1/21

$$1.3: 4, 48ab, 58$$

$$2.3: 2, 4, 8$$

Last time Matrix Operations and linear combinations

Putting it all together

Then Linear system of equations with augmented matrix

$(A; \vec{b})$ can be written $A\vec{x} = \vec{b}$ in matrix form,

Eq $\begin{cases} x_1 + x_2 = 5 \\ x_1 - x_2 = 1 \end{cases}$ Augmented matrix $\left(\begin{array}{cc|c} 1 & 1 & 5 \\ 1 & -1 & 1 \end{array} \right)$

Matrix form: $\begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 5 \\ 1 \end{pmatrix}$

Is $\begin{pmatrix} 5 \\ 1 \end{pmatrix}$ a linear combination of $\left\{ \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \end{pmatrix} \right\}$?

In other words, is there an x_1, x_2 such that

$$x_1 \begin{pmatrix} 1 \\ 1 \end{pmatrix} + x_2 \begin{pmatrix} 1 \\ -1 \end{pmatrix} = \begin{pmatrix} 5 \\ 1 \end{pmatrix}$$

Same as solving linear system!

$$\left(\begin{array}{cc|c} 1 & 1 & 5 \\ 1 & -1 & 1 \end{array} \right) \xrightarrow{R_2 \leftrightarrow R_1} \left(\begin{array}{cc|c} 1 & 1 & 5 \\ 0 & -2 & 1 \end{array} \right) \xrightarrow{R_2 \leftarrow -\frac{1}{2}R_2} \left(\begin{array}{cc|c} 1 & 1 & 5 \\ 0 & 1 & 2 \end{array} \right) \xrightarrow{R_1 \leftarrow R_1 - R_2} \left(\begin{array}{cc|c} 1 & 0 & 3 \\ 0 & 1 & 2 \end{array} \right)$$

$$x_1 = 3, x_2 = 2$$

$$\text{So, } 3 \begin{pmatrix} 1 \\ 1 \end{pmatrix} + 2 \begin{pmatrix} 1 \\ -1 \end{pmatrix} = \begin{pmatrix} 5 \\ 1 \end{pmatrix}.$$

Matrix Products

Def B - $n \times p$ matrix

A - $p \times m$ matrix

then BA is the $n \times m$ matrix given by

$$BA = B \begin{pmatrix} 1 & 1 & 1 \\ \bar{v}_1 & \bar{v}_2 & \dots & \bar{v}_m \\ 1 & 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 \\ B\bar{v}_1 & B\bar{v}_2 & \dots & B\bar{v}_m \\ 1 & 1 & 1 \end{pmatrix}$$

$$(n \times p)(p \times m) = (n \times m)$$

Warning The number of cols of B = The number of rows of A ,
Otherwise this is undefined!

Eg a) $\begin{pmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{pmatrix} \begin{pmatrix} 2 & 0 & 1 \\ 0 & 2 & 1 \end{pmatrix} = \begin{pmatrix} 2 & 8 & 5 \\ 4 & 10 & 7 \\ 6 & 12 & 9 \end{pmatrix}$

$$\begin{pmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{pmatrix} \begin{pmatrix} 2 \\ 0 \\ 1 \end{pmatrix} = 2 \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} + 0 \begin{pmatrix} 4 \\ 5 \\ 6 \end{pmatrix} = \begin{pmatrix} 2 \\ 4 \\ 6 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{pmatrix} \begin{pmatrix} 0 \\ 2 \end{pmatrix} = 0 \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} + 2 \begin{pmatrix} 4 \\ 5 \\ 6 \end{pmatrix} = \begin{pmatrix} 8 \\ 10 \\ 12 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = 1 \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} + 1 \begin{pmatrix} 4 \\ 5 \\ 6 \end{pmatrix} = \begin{pmatrix} 5 \\ 7 \\ 9 \end{pmatrix}$$

b) $\begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} \begin{pmatrix} 5 & 6 \\ 7 & 8 \end{pmatrix} = \begin{pmatrix} 17 & 22 \\ 43 & 50 \end{pmatrix}$

$$\begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} \begin{pmatrix} 5 \\ 7 \end{pmatrix} = 5 \begin{pmatrix} 1 \\ 3 \end{pmatrix} + 7 \begin{pmatrix} 2 \\ 4 \end{pmatrix} = \begin{pmatrix} 19 \\ 43 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} \begin{pmatrix} 6 \\ 8 \end{pmatrix} = 6 \begin{pmatrix} 1 \\ 3 \end{pmatrix} + 8 \begin{pmatrix} 2 \\ 4 \end{pmatrix} = \begin{pmatrix} 22 \\ 50 \end{pmatrix}$$

What about $\underbrace{\begin{pmatrix} 5 & 6 \\ 7 & 8 \end{pmatrix}}_{\text{Check}} \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$?

$$\begin{pmatrix} 23 & 34 \\ 31 & 46 \end{pmatrix} \neq \begin{pmatrix} 19 & 22 \\ 43 & 50 \end{pmatrix} = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} \begin{pmatrix} 5 & 6 \\ 7 & 8 \end{pmatrix}$$

Warning! BA and AB are not always equal!

What is the i, j -entry of BA ?

i, j -entry of BA = i^{th} component of $B\vec{v}_j = B \begin{pmatrix} a_{ij} \\ \vdots \\ a_{pj} \end{pmatrix}$

$$= (b_{i1} \dots b_{ip}) \cdot \begin{pmatrix} a_{ij} \\ \vdots \\ a_{pj} \end{pmatrix} = \sum_{k=1}^p b_{ik} a_{kj}$$

Thm Let B be an $n \times p$ matrix and A be a $p \times m$ matrix.

Then, the i, j -entry of BA is $\sum_{k=1}^p b_{ik} a_{kj}$.

Trick $\begin{pmatrix} 5 & 6 \\ 7 & 8 \end{pmatrix}$

$$\begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} \begin{pmatrix} 19 & 22 \\ 43 & 50 \end{pmatrix}$$

$$\vec{e}_i = \begin{pmatrix} 0 \\ \vdots \\ 1 \\ \vdots \\ 0 \end{pmatrix} \leftarrow i^{\text{th}} \text{ component}$$

Some Matrix Product Rules

Thm a) For A an $n \times m$ matrix,

$$A I_m \checkmark = A = I_n A$$

$$A \cdot \begin{pmatrix} \vec{e}_1 & \cdots & \vec{e}_n \end{pmatrix} = \begin{pmatrix} 1 & & 1 \\ A\vec{e}_1 & \cdots & A\vec{e}_n \end{pmatrix} \stackrel{(*)}{=} \begin{pmatrix} a_{11} & \cdots & a_{1m} \\ \vdots & \cdots & \vdots \\ a_{m1} & \cdots & a_{mm} \end{pmatrix} = A \checkmark$$

$I_k = k \times k$ identity matrix

$$= \begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & & \vdots \\ \vdots & & \ddots & 0 \\ 0 & \cdots & 0 & 1 \end{pmatrix} \left. \begin{array}{l} \text{k rows} \\ \text{k columns} \end{array} \right\}$$

$$(*) A\vec{e}_i = \begin{pmatrix} a_{1i} \\ \vdots \\ a_{ni} \end{pmatrix}$$

$$= \begin{pmatrix} 1 & 1 & \cdots & 1 \\ \vec{e}_1 & \vec{e}_2 & \cdots & \vec{e}_n \\ 1 & 1 & & 1 \end{pmatrix}$$

b) $(AB)C = A(BC)$, so we can write ABC .

Let $C = \begin{pmatrix} \frac{1}{v_1} & \dots & \frac{1}{v_m} \\ 1 & \dots & 1 \end{pmatrix}$, then $(AB)C = \begin{pmatrix} \frac{1}{(AB)\vec{v}_1} & \dots & \frac{1}{(AB)\vec{v}_m} \\ 1 & \dots & 1 \end{pmatrix} \Rightarrow (*)$

and $A(BC) = \begin{pmatrix} \frac{1}{A(B\vec{v}_1)} & \dots & \frac{1}{A(B\vec{v}_m)} \\ 1 & \dots & 1 \end{pmatrix} \Rightarrow$

(*) Check $\underbrace{(AB)\vec{v}_i}_{\text{in}} = A(\vec{B}\vec{v}_i)$ for $\vec{v}_i = \begin{pmatrix} c_{1i} \\ \vdots \\ c_{pi} \end{pmatrix}$ ✓

$$AB = A \begin{pmatrix} \frac{1}{\vec{u}_1} & \dots & \frac{1}{\vec{u}_p} \\ 1 & \dots & 1 \end{pmatrix} = \begin{pmatrix} 1 & \dots & 1 \\ A\vec{u}_1 & \dots & A\vec{u}_p \\ 1 & \dots & 1 \end{pmatrix}$$

so, $(AB)\vec{v}_i = c_{1i}A\vec{u}_1 + \dots + c_{pi}A\vec{u}_p \Rightarrow$
 and $A(\vec{B}\vec{v}_i) = A(c_{1i}\vec{u}_1 + \dots + c_{pi}\vec{u}_p) \Rightarrow \square$

c) If A and B are $n \times p$ matrices and C and D are $p \times m$ matrices,
 then

$$A(C+D) = AC + AD$$

$$(A+B)C = AC + BC$$

d) If A is an $n \times p$ matrix, B is a $p \times m$ matrix,
 and k is a scalar,

$$(kA)B = A(kB) = k(AB)$$

Why multiply matrices?



A $n \times m$ matrix
 $\vec{v} \in \mathbb{R}^n \longmapsto A\vec{v} \in \mathbb{R}^m$