

Written HW 10 (due 9/8)

7.1: 4, 6, 12, 16, 18

7.2: 8, 12, 18, 38

7.3: 8, 10, 24

Quiz on 7.1-7.3 4/11

Written HW 11 (due 9/15)

7.5: 14, 20

Will post Midterm 2 grades today

- Adding +15 Upoints to scores
- Grade on Final can replace one of two midterm grades if final score is better

Complex Eigenvalues (7.5)

Some matrices have no real eigenvalues:

Rotation matrix $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \rightsquigarrow \det \begin{pmatrix} -\lambda & -1 \\ 1 & -\lambda \end{pmatrix} = \lambda^2 - (-1) = \lambda^2 + 1$

\uparrow
has no real solutions

$$\begin{aligned}\lambda^2 + 1 &= 0 \\ \Rightarrow \lambda^2 &= -1\end{aligned}$$

However, over complex numbers, \mathbb{C} , $\lambda = \pm i$ are solutions ($i^2 = -1$)

A complex number z is expressed as $a+bi$ for a, b real numbers and i is the "imaginary root" satisfying $i^2 = -1$

Eg $i, -i, 2+3i, -5-5i, 4-1.5i, 7, -4$ are all in \mathbb{C}

Facts a) $(a+bi) + (c+di) = (a+c) + (b+d)i$

b) $(a+bi)(c+di) = ac + bci + adi + bdi^2$
 $= (ac - bd) + (bc + ad)i$

λ just a generic variable here.

Fundamental Theorem of Algebra Any polynomial $p(\lambda)$ of degree n

can be written as a product of n linear factors over \mathbb{C} :

$$p(\lambda) = k(\lambda - \lambda_1)(\lambda - \lambda_2) \cdots (\lambda - \lambda_n) \quad \text{for complex numbers } \lambda_1, \dots, \lambda_n, k.$$

$$\text{Eq } x^2 + 1 = (x+i)(x-i)$$

Eg Eigenvectors of $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ (will have complex entries)

$$\lambda = i \quad \ker(A - iI_2) = \ker \begin{pmatrix} -i & -1 \\ 1 & -i \end{pmatrix}$$

$$\text{Solve } x_1 \begin{pmatrix} -i \\ 1 \end{pmatrix} + x_2 \begin{pmatrix} -1 \\ -i \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\begin{aligned} x_1 &= i : & \begin{pmatrix} 1 \\ i \end{pmatrix} + x_2 \begin{pmatrix} -1 \\ -i \end{pmatrix} &= \begin{pmatrix} 0 \\ 0 \end{pmatrix} \\ (-i)i &= -i^2 \\ &= -(-1) \\ &= 1 & \Rightarrow x_2 &= 1 \end{aligned}$$

$$\text{So, } E_i = \text{span} \left\{ \begin{pmatrix} i \\ 1 \end{pmatrix} \right\} \text{ and, similarly } E_{-i} = \text{span} \left\{ \begin{pmatrix} -i \\ 1 \end{pmatrix} \right\}.$$

Eg Find all complex eigenvalues of

$$\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 5 & -7 & 3 \end{pmatrix}. \quad \text{tr}A = 3 \quad \det A = 5$$

$$\begin{aligned} \rightarrow \det \begin{pmatrix} -\lambda & 1 & 0 \\ 0 & -\lambda & 1 \\ 5 & -7 & 3-\lambda \end{pmatrix} &= (-\lambda)^2(3-\lambda) + 5 + 0 - 0 - (-7)(1)(-\lambda) - 0 \\ &= (-\lambda)^3 + 3\lambda^2 - 7\lambda + 5 \\ &= -\lambda^3 + 3\lambda^2 - 7\lambda + 5 \end{aligned}$$

Recall 3×3 matrix must have at least 1 real eigenvalue.

$$\begin{aligned} &= -(\lambda-1)(\lambda^2 - 2\lambda + 5) \\ &\quad \uparrow \quad \frac{2 \pm \sqrt{4-20}}{2} \\ &= \frac{2 \pm \sqrt{-16}}{2} \\ &= \frac{2 \pm 4i}{2} = 1 \pm 2i \end{aligned}$$

So, eigenvalues are $\lambda = 1, 1+2i, 1-2i$.

Thm If A is a 2×2 real matrix with eigenvalues $a \pm bi$, then

a) A is similar to $B = \begin{pmatrix} a & -b \\ b & a \end{pmatrix}$ (rotation-scaling) and

b) If $\vec{v} + i\vec{w}$ is an eigenvector for A ($\vec{v}, \vec{w} \in \mathbb{R}^2$), then

$$S^{-1}AS = B \quad \text{where} \quad S = \begin{pmatrix} 1 & 1 \\ \vec{w} & \vec{v} \end{pmatrix}. \quad \underbrace{\text{Note Entries of } S \text{ are real.}}$$

Eg For $A = \begin{pmatrix} 3 & -2 \\ 4 & -1 \end{pmatrix}$, find an invertible matrix S such that $S^{-1}AS$ is a rotation-scaling matrix.

$$\det \begin{pmatrix} 3-\lambda & -2 \\ 4 & -1-\lambda \end{pmatrix} = 5 - 2\lambda + \lambda^2$$

$$\lambda = \frac{2 \pm \sqrt{4-20}}{2}$$

$$= 1 \pm 2i \quad \Rightarrow \quad S^{-1}AS = \begin{pmatrix} 1 & -2 \\ 2 & 1 \end{pmatrix}$$

$$E_{1+2i} = \ker(A - (1+2i)I_2) = \ker \begin{pmatrix} 2-2i & -2 \\ 4 & -2-2i \end{pmatrix}$$

Useful trick $(a+bi)(a-bi) = a^2 + abi - abi - b^2i^2$
 $= a^2 + b^2$ ← real number

$$\text{Solve } x_1 \begin{pmatrix} 2-2i \\ 4 \end{pmatrix} + x_2 \begin{pmatrix} -2 \\ -2-2i \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\text{Note } (-2+2i) \begin{pmatrix} -2 \\ -2-2i \end{pmatrix} = \begin{pmatrix} 4-4i \\ 8 \end{pmatrix}$$

$$\text{So } x_2 = (-2+2i) \rightarrow \text{Solve } x_1 \begin{pmatrix} 2-2i \\ 4 \end{pmatrix} + \begin{pmatrix} 4-4i \\ 8 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\text{So } x_1 = -2$$

$$\Rightarrow E_{1+2i} = \text{span} \left\{ \begin{pmatrix} -2 \\ -2+2i \end{pmatrix} \right\} \Rightarrow S = \begin{pmatrix} 0 & -2 \\ 2 & -2 \end{pmatrix}$$

$$\underbrace{\begin{pmatrix} -2 \\ -2 \end{pmatrix}}_{\vec{v}''} + i \underbrace{\begin{pmatrix} 0 \\ 2 \end{pmatrix}}_{\vec{w}''}$$

$$\rightsquigarrow \begin{pmatrix} 0 & -2 \\ 2 & -2 \end{pmatrix}^{-1} \begin{pmatrix} 3 & -2 \\ 4 & -1 \end{pmatrix} \begin{pmatrix} 0 & -2 \\ 2 & -2 \end{pmatrix} = \begin{pmatrix} 1 & -2 \\ 2 & 1 \end{pmatrix}$$

Thm For $n \times n$ matrix A with complex eigenvalues $\lambda_1, \dots, \lambda_n$,
listed with algebraic multiplicity,

$$\operatorname{tr} A = \lambda_1 + \lambda_2 + \dots + \lambda_n$$

$$\det A = \lambda_1 \lambda_2 \dots \lambda_n$$

Eq $A = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ $\operatorname{tr} A = 0 = i + (-i)$ $\lambda_1 = i, \lambda_2 = -i$.

Rank Complex numbers \mathbb{C} form a linear space of dimension 2.

For instance, $B = \{1, i\}$ is a basis of \mathbb{C} (when scalars = \mathbb{R})

$$\rightsquigarrow [a+bi]_B = \begin{pmatrix} a \\ b \end{pmatrix}.$$