

Lars B. Wahlbin

# Superconvergence in Galerkin Finite Element Methods

1605



Springer

# Lecture Notes in Mathematics

1605

Editors:

A. Dold, Heidelberg

F. Takens, Groningen

**Springer**

*Berlin*

*Heidelberg*

*New York*

*Barcelona*

*Budapest*

*Hong Kong*

*London*

*Milan*

*Paris*

*Tokyo*

Lars B. Wahlbin

# Superconvergence in Galerkin Finite Element Methods



Springer

**Author**

Lars B. Wahlbin  
Department of Mathematics  
White Hall  
Cornell University  
Ithaca, NY 14853, USA  
E-mail: [wahlbin@math.cornell.edu](mailto:wahlbin@math.cornell.edu)

Library of Congress Cataloging-in-Publication Data. Wahlbin, Lars B., 1945–  
Superconvergence in Galerkin finite element methods/Lars B. Wahlbin. p.cm. –  
(Lecture notes in mathematics; 1605) Includes bibliographical references (p. -) and  
index. ISBN 3-540-60011-6 (acid-free) 1. Differential equation, Elliptic – Numerical  
solutions. 2. Convergence. 3. Galerkin methods. I. Title. II. Series: Lecture notes  
in mathematics (Springer-Verlag); 1605.

QA3. L28 no. 1605

[QA377]  
510 s--dc20  
[515'.353]

Mathematics Subject Classification (1991): 65N30, 65N15

ISBN 3-540-60011-6 Springer-Verlag Berlin Heidelberg New York

This work is subject to copyright. All rights are reserved, whether the whole or part  
of the material is concerned, specifically the rights of translation, reprinting, re-use  
of illustrations, recitation, broadcasting, reproduction on microfilms or in any other  
way, and storage in data banks. Duplication of this publication or parts thereof is  
permitted only under the provisions of the German Copyright Law of September 9,  
1965, in its current version, and permission for use must always be obtained from  
Springer-Verlag. Violations are liable for prosecution under the German Copyright  
Law.

© Springer-Verlag Berlin Heidelberg 1995  
Printed in Germany

SPIN: 10130328      46/3142-543210 - Printed on acid-free paper

## Preface.

These notes are from a graduate seminar at Cornell in Spring 1994. They are devoted mainly to basic concepts of superconvergence in second-order time-independent elliptic problems.

A brief chapter-by-chapter description is as follows: Chapter 1 considers one-dimensional problems and is intended to get us moving quickly into the subject matter of superconvergence. (The results in Sections 1.8 and 1.10 are new.) Some standard results and techniques used there are then expounded on in Chapters 2 and 3. Chapter 4 gives a few selected results about superconvergence in  $L_2$ -projections in any number of space dimensions. In Chapter 5 we elucidate local maximum-norm error estimates in second order elliptic partial differential equations and the techniques used in proving them, without aiming for complete detail. Theorems 5.5.1 and 5.5.2 are basic technical results; they will be used over and over again in the rest of the notes.

In Chapters 6 through 12 we treat a variety of topics in superconvergence for second order elliptic problems. Some are old and established, some are very recent and not yet published. Some of the earlier contributions have benefitted from later sharpening of tools, in particular with respect to local maximum-norm estimates.

In Chapter 6 we consider tensor-product elements. Using ideas of [Douglas, Dupont and Wheeler 1974b] we show that in some situations one-dimensional superconvergence results automatically translate to several dimensions. Chapter 7, “Superconvergence by local symmetry”, presents recent fundamental results from [Schatz, Sloan and Wahlbin 1994]. Chapter 8 treats difference quotients for approximating derivatives of any order on translation invariant meshes. Here we follow the basic ideas of [Nitsche and Schatz 1974, Section 6]. In Chapter 9 we briefly comment on how, in many cases, results about superconvergence in linear problems automatically carry over to nonlinear problems. The essential idea is from [Douglas and Dupont 1975], which in turn is essentially the quadratic convergence of Newton’s method. Chapter 10 is concerned with superconvergence on curved meshes which come about via isoparametric mappings of straight-lined meshes; it is based on [Cayco, Schatz and Wahlbin 1994]. Chapter 11 reverts back to the seventies. It is mainly concerned with the  $K$ -operator of [Bramble and Schatz 1974]; the presentation follows [Thomée 1977]. We give an application to boundary integral equations, [Tran 1993]. Also, we briefly mention a method for obtaining higher order accuracy in outflow derivatives, [Douglas, Dupont and Wheeler 1974a], and an averaging method of [Louis 1979]. Finally, in Chapter 12, we review the computational investigation of [Babuška, Strouboulis, Upadhyay and Gangaraj 1993] and comment on it in light of the theories of Chapters 6 and 7.

Previous treatises of superconvergence are [Chen 1982a] and [Zhu and Lin 1989]. Both are in Chinese and, although there is a description of Zhu and Lin’s book in Mathematical Reviews, it is hard for me to judge how they compare with the present account. The use of local maximum-norm estimates seems common to all three. *For what appears to be a major difference of approach, see Remark 7.4.i.*

Surveys of superconvergence with a more limited scope have appeared in [Křížek and Neittaanmäki 1987a] and [Wahlbin 1991, Chapter VII].

Let me next list some topics that are *not* included in these notes. The first is superconvergence in collocation finite element methods for differential equations. For

this topic I have not even included references; the reader is referred to [Křížek and Neittaanmäki 1987a]. The second omitted topic is superconvergence in boundary integral methods or in integral equations (except for Section 11.5). Third, extrapolation methods, involving computation on two or more meshes, and fourth, the use of superconvergence in construction of smooth stress fields and the related use of superconvergence for a posteriori error estimation and adaptive refinement. For the second through fourth topics I have included a fair number of references so that the interested reader may, by glancing through the list of references, easily gain an inroad to the literature on the subjects. Finally, I have not considered superconvergence in Galerkin finite element methods for time-dependent problems. Here, though, I have included all references that I know of.

The references are likewise “complete” with respect to the mathematical literature for the main topics treated. Of course, many of these references touch only briefly on superconvergence. In preparing the references I have used MathSci ( $\simeq$  Mathematical Reviews since 1972 on line) which does not systematically cover the vast engineering literature. The number of references given is more than it makes sense to actually refer to in the text, unless one resorts to plain listing, which I have not done. I hope that nevertheless some readers will find the list of references useful. (As an example, if a reader interested in early history wants to find references to papers on superconvergence in finite element methods before 1970, our list gives: [Stricklin 1966], [Filho 1968], [Stricklin 1968], [Oganesyan and Rukhovetz 1969] and [Tong 1969].)

Basic discoveries continue to be made at present. Furthermore, the theory of superconvergence is very immature in carrying results up close to boundaries (or, internal lines of discontinuity). Today, such investigations are carried out on a case-by-case basis, and, of course, not all results hold all the way up to boundaries (see Section 1.7). Most results that are proven up to boundaries in several dimension pertain to axes-parallel parallelepipeds, or, locally, to straight boundaries. For these and other reasons I have decided to offer these notes essentially as they were written week-by-week during the seminar, rather than rework them into “text-book” form; such a textbook would most likely be out-of-date when it appears. The reader may be warned that, reflecting my lecturing style, proofs often appear before theorems; indeed, the “theorem” may be only an informal statement. Also, true to the principle that repetition is the mother of studies, there is a fair amount of such. E.g., symmetry considerations are first met with in two-point boundary value problems, then in  $L_2$ -projections and finally in multidimensional elliptic problems; difference quotients first occur in two-point boundary value problems and later in many dimensions; and, tensor product elements are considered as well for  $L_2$ -projections as for elliptic problems.

Let me penultimately remark that there are three concepts which may be confused with superconvergence. “Supraconvergence” is a concept in finite-difference theory for irregular meshes, cf. [Kreiss, Manteuffel, Swartz, Wendroff and White 1986], [Manteuffel and White 1986], [Heinrich 1987, p. 107], and cf. also [Bramble 1970]. “Superapproximation” will be explained in these notes and “superlinear convergence” occurs in the theory of iterative methods, cf. e.g. [Ortega and Rheinboldt 1979, pp. 285 and 291].

Finally, I thank Arletta Havlik for her superior typing, Al Schatz for many stimulating discussions, Stig Larsson for timely help and the National Science Foundation for financial support.

Ithaca, January 1995  
Lars B. Wahlbin

## Table of Contents.

Chapter 1. Some one-dimensional superconvergence results.	1
1.1. Introduction	1
1.2. Nodal superconvergence for function values in continuous elements ( $\mu = 0$ ).	3
1.3. Reduction to a model problem.	4
1.4. Existence of superconvergence points in general.	8
1.5. Superconvergence for interior points of mesh-intervals for continuous elements ( $\mu = 0$ ).	8
1.6. Superconvergence in derivatives at points about which the meshes are locally symmetric ( $r$ even).	10
1.7. Necessity of staying $C_1 h \ln 1/h$ away from the boundary for superconvergence in the uniform mesh case ( $\mu = 1$ ).	13
1.8. Finding all superconvergent points for function values and derivatives in the case of a locally uniform mesh for $\mu = 1$ and $r$ even (with a remark about smoothest cubics).	14
1.9. Superconvergence in function values at points about which the meshes are locally symmetric ( $r$ odd).	16
1.10. Finding all superconvergent points for function values and derivatives in the case of a locally uniform mesh for $\mu = 1$ and $r$ odd.	19
1.11. First order difference quotients of $u_h$ as superconvergent approximations to $u'$ on locally uniform meshes.	22
1.12. Two examples of superconvergence by “iteration”.	25
1.13. A graphical illustration of superconvergence.	26
Chapter 2. Remarks about some of the tools used in Chapter 1.	28
2.1. Inverse estimates.	28
2.2. On approximation theory, and duality.	30
2.3. Superapproximation.	32
2.4. A typical combination of inverse estimates and approximation theory used in Chapter 1.	35
Chapter 3. Local and global properties of $L_2$ -projections.	36
3.1. Assumptions.	36
3.2. Estimates for $L_2$ -projections.	38
Chapter 4. Introduction to several space dimensions: some results about superconvergence in $L_2$ -projections.	42
4.1. Negative norm estimates and existence of general superconvergence points for function values.	42
4.2. Superconvergence in $L_2$ -projections on $n$ -dimensional tensor product spaces.	43
4.3. Superconvergence by symmetry in $L_2$ -projections.	44

Chapter 5. Second order elliptic boundary value problems in any number of space dimensions: preliminary considerations on local and global estimates and presentation of the main technical tools for showing superconvergence.	48
5.1. Introduction.	48
5.2. Existence of superconvergence points in general: an example (also an example of a multi-dimensional duality argument).	49
5.3. General comments on local a priori error estimates.	52
5.4. General comments on $L_\infty$ estimates.	58
5.5. The main technical tools for proving superconvergence in second order elliptic problems in several space dimensions.	62
Chapter 6. Superconvergence in tensor-product elements.	65
6.1. Introduction.	65
6.2. Superconvergence in derivatives for the case of the Laplacian.	65
6.3. Negative norm estimates for $u - u_h$ : Examples.	69
6.4. Superconvergence in derivatives for the case of (6.1.1.).	70
6.5. Superconvergence in function values for the Laplacian and $r \geq 3$ .	72
Chapter 7. Superconvergence by local symmetry.	74
7.1. Introduction.	74
7.2. The case of a symmetric form with constant coefficients.	74
7.3. The general case of (5.1.3) with variable smooth coefficients.	78
7.4. Historical remarks.	79
Chapter 8. Superconvergence for difference quotients on translation invariant meshes.	84
8.1. Introduction.	84
8.2. Constant coefficient operators and unit separation, $d \simeq 1$ .	86
8.3. Constant coefficient operators and general separation $d$ .	89
8.4. Variable coefficients.	89
Chapter 9. On superconvergence in nonlinear problems.	93
Chapter 10. Superconvergence in isoparametric mappings of translation invariant meshes: an example.	98
10.1. Introduction.	98
10.2. Superconvergence in difference quotients for first derivatives.	101
Chapter 11. Superconvergence by averaging: mainly, the $K$ -operator.	107
11.1. Introduction	107
11.2. Preliminaries on Fourier transforms and multipliers.	107
11.3. The $K$ -operator in general.	111
11.4. The $K$ -operator applied to finite element approximations in second order elliptic problems.	115
11.5. Boundary integral equations and the $K$ -operator: an example.	116
11.6. Remarks, including some other averaging methods.	121
11.7. A superconvergent “global” averaging technique for function values.	123

Chapter 12. A computational investigation of superconvergence for first derivatives in the plane.	125
12.1. Introduction.	125
12.2. Proof of (12.1.5), (12.1.6) and precise definition of the principal error term $\psi$ .	128
12.3. Results of computational studies, with comments.	132
References.	136
Subject index.	165

## Chapter 1. Some one-dimensional superconvergence results.

We shall move briskly into the subject of superconvergence. In Chapters 2 and 3 we shall then backtrack and elucidate some of the techniques and results used in this chapter.

### 1.1. Introduction.

Consider the following two-point boundary value problem: Find  $u = u(x)$  such that

$$(1.1.1) \quad \begin{cases} -(a_2(x)u')' - (a_1(x)u)' + a_0(x)u = f & \text{in } I = (0, 1), \\ u(0) = u(1) = 0. \end{cases}$$

We shall assume throughout this chapter that the coefficients  $a_i$  and the right hand side  $f$  are as smooth as necessary on  $\bar{I}$  for our analyses to carry through. Furthermore, we demand that

$$(1.1.2) \quad a_2(x) \geq a_2 > 0, \text{ for all } x \in \bar{I}.$$

The weak formulation of (1.1.1) is to find  $u \in \overset{\circ}{H}^1(I)$  such that

$$(1.1.3) \quad A(u, \chi) \equiv \int_I (a_2 u' \chi' + a_1 u \chi' + a_0 u \chi) dx = \int_I f \chi dx, \text{ for all } \chi \in \overset{\circ}{H}^1(I).$$

Let  $T_h$ ,  $0 < h < 1/2$ , be a sequence of subdivisions of  $\bar{I}$ ,

$$(1.1.4) \quad T_h = \{x_i\}_{i=0}^{N(h)}, \quad 0 = x_0 < x_1 < \dots < x_N = 1,$$

$$I_i = (x_i, x_{i+1}), \quad \bar{I} = \bigcup_{i=0}^{N(h)-1} \bar{I}_i.$$

Let  $h_i = x_{i+1} - x_i$ . With, perhaps, a slight abuse of notation, we shall let  $h = \max_i h_i$ . With integers  $0 \leq \mu < r - 1$ , we set

$$(1.1.5) \quad \overset{\circ}{S}_h = \overset{\circ}{S}_h^{\mu, r} = \{\chi(x) : \chi \in C^\mu(I) \cap C^0(\bar{I}), \chi(0) = \chi(1) = 0, \\ \chi|_{I_i} \in \Pi_{r-1}(I_i)\}$$

where  $\Pi_{r-1}(I_i)$  denotes the polynomials of degree  $\leq r-1$  on  $I_i$ . When no confusion can arise we use the shorter notation  $\overset{\circ}{S}_h$ . Note that  $\overset{\circ}{S}_h \subseteq \overset{\circ}{H}^1(I)$  since  $\mu \geq 0$ .

Well known examples are:

$\mu = 0, r = 2$  : piecewise linears

$\mu = 1, r = 4$  : Hermite cubics

$\mu = r - 2$  : smoothest splines.

The finite element solution is sought as follows: Find  $u_h \in \overset{\circ}{S}_h$  such that

$$(1.1.6) \quad A(u_h, \chi) = (f, \chi), \text{ for all } \chi \in \overset{\circ}{S}_h.$$

(We shall not take into account numerical integration in (1.1.6).)

In our analyses below we shall need to use various results from the literature. These can be found with varying degrees of generality, in particular with respect to dependence on the distribution of mesh sizes  $h_i$  in a subdivision. To keep things

simple in this chapter we make the following explicit assumption of quasi-uniformity of meshes, except in the case of continuous elements ( $\mu = 0$ ) when no such restriction is made.

(1.1.7)      If  $\mu \geq 1$ , then there exists a positive constant  $C_{QU}$  independent of  $h$  such that in any subdivision  $\tau_h$ , we have  $h \leq C_{QU} \min_i h_i$ .

We shall assume that (1.1.3) has a unique solution  $u$  for any  $f$  in  $L_2(I)$ , say. Following [Schatz 1974], cf. also [Hildebrandt and Weinholtz 1964] and [Schatz and Wang 1994], we then know that there exists  $h_0 > 0$  such that for  $h \leq h_0$ , given any  $f \in L_2(I)$  there is a unique solution  $u_h$  to (1.1.6). Furthermore, under our general conditions on smooth enough data, we have the following estimates for the error  $e = u - u_h$ :

$$(1.1.8) \quad \|e\|_{L_2(I)} + h\|e\|_{H^1(I)} \leq Ch^r \|u\|_{H^r(I)}$$

and

$$(1.1.9) \quad \|e\|_{H^{-s}(I)} \leq Ch^{r+s} \|u\|_{H^r(I)}, \text{ for } s \leq r - 2.$$

Here

$$(1.1.10) \quad \|v\|_{H^{-s}(I)} = \sup_{\substack{w \in H^s(I) \\ \|w\|_{H^s(I)}=1}} (v, w).$$

Correspondingly we have in maximum-norm,

$$(1.1.11) \quad \|e\|_{L_\infty(I)} + h\|e\|_{W_\infty^1(I)} \leq Ch^r \|u\|_{W_\infty^r(I)},$$

and

$$(1.1.12) \quad \|e\|_{W_\infty^{-s}(I)} \leq Ch^{r+s} \|u\|_{W_\infty^r(I)}, \text{ for } s \leq r - 2,$$

where now

$$(1.1.13) \quad \|v\|_{W_\infty^{-s}(I)} = \sup_{\substack{w \in W_1^s(I) \\ \|w\|_{W_1^s(I)}=1}} (v, w).$$

The constants  $C$  occurring are independent of  $h$  and  $u$ . They do depend on the coefficients  $a_i$  and on  $\mu$  and  $r$ ; in the case of  $\mu \geq 1$  they may also depend on  $C_{QU}$  in (1.1.7).

The  $\overset{\circ}{H}{}^1$ -estimate in (1.1.8) can be found in this generality in [Schatz 1974], after use of standard approximation theory. The  $L_2$ -estimate and the negative norm estimates (1.1.9) follow by standard duality arguments, cf. (2.2.4) below. The  $L_\infty$ -estimate (1.1.11) for  $\mu = 0$  is in [Wheeler, M. F. 1973]. The case of general  $\mu$ , with the quasi-uniformity condition (1.1.7), is in [Douglas, Dupont and Wahlbin 1975]. From this the  $W_\infty^1$ -estimate follows in the quasi-uniform case. The  $W_\infty^1$ -estimate in the case  $\mu = 0$ , although “well-known folklore”, we have not been able to locate without restrictions on the meshes. We will therefore give a proof, in Remark 1.3.2 below. The negative norm estimates in (1.1.12) follow in a standard duality fashion from (1.1.11).

It is not hard to convince one–self that the powers of  $h$  occurring in (1.1.11) are the best possible in general. E.g.,

$$\begin{aligned}
 (1.1.14) \quad & \min_{\chi \in \Pi_{r-1}(0,h)} \|x^r - \chi\|_{L_\infty(0,h)} \\
 &= h^r \min_{\chi \in \Pi_{r-1}(0,h)} \left\| \left(\frac{x}{h}\right)^r - \frac{1}{h^r} \chi(x) \right\|_{L_\infty(0,h)} \\
 &= h^r \min_{\chi \in \Pi_{r-1}(0,h)} \left\| \xi^r - \frac{1}{h^r} \chi(h\xi) \right\|_{L_\infty(0,1)} \\
 &= h^r \min_{\tilde{\chi} \in \Pi_{r-1}(0,1)} \|\xi^r - \tilde{\chi}(\xi)\|_{L_\infty(0,1)} = Ch^r.
 \end{aligned}$$

A superconvergent “point” for function values of order  $\sigma$  is now a family of points  $\xi = \xi(h)$  such that

$$(1.1.15) \quad |e(\xi)| \leq Ch^{r+\sigma},$$

where  $\sigma > 0$  and  $C = C(u, a_2, a_1, a_0)$  (and possibly also depending on  $G_{QU}$  in (1.1.7)). Similarly,  $\eta = \eta(h)$  is superconvergent of order  $\sigma$  for first derivatives if

$$(1.1.16) \quad |e'(\eta)| \leq Ch^{r-1+\sigma},$$

with  $\sigma > 0$ .

In principle, we could talk about superconvergence points for a particular problem or for a particular solution. Generally what we have in mind, though, is some class of problems with (locally) smooth coefficients and solutions, and we then wish to determine points which are superconvergent for the whole class.

Furthermore, we point out that what is described in (1.1.15) and (1.1.16) is only so-called “natural” superconvergence. That is,  $u_h$  (or  $u'_h$ ) is simply evaluated at the point  $\xi$  (or  $\eta$ ) and then compared with  $u(\xi)$  (or  $u'(\eta)$ ). In these notes we shall also see many examples of superconvergence involving postprocessing of  $u_h$  (trivial, or not so trivial). In fact, in terms of implementation on a computer, many postprocessing methods are simpler to implement than is evaluating  $u_h$  at a (non-node) point.

## 1.2. Nodal superconvergence for function values in continuous elements ( $\mu = 0$ ).

We shall give the argument of [Douglas and Dupont 1974]. Let  $G(\xi; \cdot)$  be the Green’s function for (1.1.1), cf. e.g. [Birkhoff and Rota 1969, Theorem 10, p.52–53], so that

$$(1.2.1) \quad e(\xi) = A(e, G(\xi; \cdot)).$$

From (1.1.3) and (1.1.6),  $A(e, \chi) = 0$ , for  $\chi \in \overset{\circ}{S}_h$  and thus

$$(1.2.2) \quad e(\xi) = A(e, G(\xi; \cdot) - \chi), \text{ for any } \chi \in \overset{\circ}{S}_h.$$

Let now  $\xi = x_i$ , a meshpoint. Since  $G(\xi; \cdot)$  is continuous and (uniformly in  $\xi$ ) smooth on both sides of  $x = \xi$ , and since  $\mu = 0$ , we have from standard approximation theory that, for a suitable  $\chi$ ,

$$(1.2.3) \quad \|G(\xi; \cdot) - \chi\|_{H^1(I)} \leq Ch^{r-1}.$$

Since by (1.1.8) also  $\|e\|_{H^1(I)} \leq Ch^{r-1}\|u\|_{H^r(I)}$  we thus obtain from (1.2.2),

$$(1.2.4) \quad |e(x_i)| \leq Ch^{2r-2}\|u\|_{H^r(I)},$$

for  $x_i$  a meshpoint. Thus, provided  $r \geq 3$  (i.e., continuous piecewise quadratics or higher elements are used), we have superconvergence at the knots. There is no restriction on the geometries of the meshes allowed.

We state the above as a theorem.

**Theorem 1.2.1.** *Under the assumptions of Section 1.1, for  $\mu = 0$  (with no mesh restrictions),*

$$(1.2.5) \quad |(u - u_h)(x_i)| \leq Ch^{2r-2}\|u\|_{H^r(I)}.$$

Let us mention that [Douglas and Dupont 1974] gives an explicit example (with  $a_2 \in C^{r-1}(I)$  only, though) showing that the power  $h^{2r-2}$  is sharp.

### 1.3. Reduction to a model problem.

In Sections 1.5–1.11 we shall investigate superconvergence of, typically, order one (i.e.,  $\sigma = 1$  in (1.1.15) or (1.1.16)). It turns out to be convenient to reduce the investigations to the case  $a_2 \equiv 1$ ,  $a_1 = a_0 \equiv 0$ . The argument follows [Wahlbin 1992] which in turn was based on [Douglas, Dupont and Wahlbin 1975]. Let thus

$$(1.3.1) \quad A(u - u_h, \chi) = 0, \text{ for } \chi \in \overset{\circ}{S}_h,$$

(cf., (1.1.3) and (1.1.6)) and let  $\tilde{u}_h \in \overset{\circ}{S}_h$  be another approximation to  $u$  given by

$$(1.3.2) \quad (u' - \tilde{u}'_h, \chi') = 0, \text{ for } \chi \in \overset{\circ}{S}_h.$$

Then, with  $\theta = \tilde{u}_h - u_h \in \overset{\circ}{S}_h$ ,

$$\begin{aligned} (1.3.3) \quad (\theta', \chi') &= ((\tilde{u}_h - u)', \chi') - ((u_h - u)', \chi') \\ &= -\left(a_2(u_h - u)', \frac{1}{a_2}\chi'\right) \\ &= -\left(a_2(u_h - u)', \left(\frac{1}{a_2}\chi\right)'\right) + \left(u_h - u, \left(\frac{a'_2}{a_2}\chi\right)'\right) \\ &= -\left(a_2(u_h - u)', \left(\frac{1}{a_2}\chi\right)' - \psi'\right) + \left(a_1(u_h - u), \psi'\right) \\ &\quad + (a_0(u_h - u), \psi) + \left(u_h - u, \left(\frac{a'_2}{a_2}\chi\right)'\right), \text{ for } \chi, \psi \in \overset{\circ}{S}_h. \end{aligned}$$

Let us now consider  $\left(\frac{1}{a_2}\chi\right)' - \psi'$ , and let us treat in detail the continuous case ( $\mu = 0$ ). Let then  $\psi$  be the natural (Lagrange) interpolant (at  $x_i$ ,  $x_{i+1}$ ; and the appropriate number of equispaced points in the interior of  $I_i$ ) to  $\frac{1}{a_2}\chi$ . By standard approximation theory,

$$(1.3.4) \quad \left\| \left(\frac{1}{a_2}\chi\right)' - \psi' \right\|_{L_1(I_i)} \leq Ch_i^{r-1} \left\| \left(\frac{1}{a_2}\chi\right)^{(r)} \right\|_{L_1(I_i)},$$

and since  $\chi^{(r)} \equiv 0$  we have by Leibniz' formula and inverse estimates (cf. Section 2.1),

$$(1.3.5) \quad \left\| \left( \frac{1}{a_2} \chi \right)' - \psi' \right\|_{L_1(I_i)} \leq C h_i \|\chi\|_{W_1^1(I_i)},$$

where  $C$  depends on  $a_2$ . Thus, summing over all intervals,

$$(1.3.6) \quad \left\| \left( \frac{1}{a_2} \chi \right)' - \psi' \right\|_{L_1(I)} \leq C h \|\chi\|_{W_1^1(I)}.$$

This result is a special case of “superapproximation”, cf. [Nitsche and Schatz 1974]. Employing a quasi-interpolant à la [deBoor and Fix 1973] (see e.g. [Wahlbin 1991, Lemma 3.2, p.367]), (1.3.6) is true for any  $\overset{\circ}{S}_h$  considered by us, cf. (1.1.7). We shall give some detail on this in Section 2.3 below. From (1.3.3) we now have using also (1.1.11),

$$(1.3.7) \quad |(\theta', \chi')| \leq C h^r \|u\|_{W_\infty^r(I)} \|\chi\|_{W_1^1(I)}.$$

We proceed to estimate  $\|\theta'\|_{L_\infty(I)}$ . We have

$$(1.3.8) \quad \|\theta'\|_{L_\infty(I)} = \sup_{\|\psi\|_{L_1(I)}=1} (\theta', \psi).$$

Let

$$(1.3.9) \quad \$_h = \$_h^{\mu-1, r-1} = \left\{ \chi(x) : \chi \in C^{\mu-1}(I); \chi|_{I_i} \in \Pi_{r-2} \right\},$$

and let  $P$  denote the  $L_2$ -projection into  $\$_h$ . For each  $\psi$  occurring in (1.3.8) we have

$$(1.3.10) \quad (\theta', \psi) = (\theta', P\psi)$$

since  $\theta' \in \$_h$ . Now set

$$(1.3.11) \quad \phi(x) = \int_0^x P\psi(y) dy - x MV(P\psi)$$

where  $MV(P\psi) = \int_0^1 P\psi dy$ . Then  $\phi \in \overset{\circ}{S}_h$ , and  $\phi' = P\psi - MV(P\psi)$ . Since  $(\theta', 1) = 0$  we have from (1.3.10) and (1.3.7)

$$(1.3.12) \quad |(\theta', \psi)| = |(\theta', \phi')| \leq C h^r \|u\|_{W_\infty^r(I)} \|\phi\|_{W_1^1(I)}.$$

Since  $\|\phi\|_{W_1^1(I)} \leq C \|P\psi\|_{L_1(I)}$  and the  $L_2$ -projection into  $\$_h$  is bounded in  $L_1$  (easy if  $\mu = 0$  so that  $P$  is completely local; for the cases  $\mu \geq 1$  see [Douglas, Dupont and Wahlbin 1975] or [Wahlbin 1991, Lemma 3.5] or, Theorem 3.2.3 below) we thus have  $\|\phi\|_{W_1^1(I)} \leq C \|\psi\|_{L_1(I)} = C$ . Hence

$$(1.3.13) \quad \|\theta'\|_{L_\infty(I)} \leq C h^r \|u\|_{W_\infty^r(I)}.$$

We state this result as a theorem.

**Theorem 1.3.1.** *Let  $u_h$  be the projection into  $\overset{\circ}{S}_h$  of  $u$  based on the form  $A$ , which satisfies the assumptions of Section 1.1. Let  $\tilde{u}_h \in \overset{\circ}{S}_h$  be given by  $(\tilde{u}'_h - u', \chi') = 0$ , for  $\chi \in \overset{\circ}{S}_h$ . For  $\mu \geq 1$  assume quasiuniformity as in (1.1.7). Then*

$$(1.3.14) \quad \|(u_h - \tilde{u}_h)'\|_{L_\infty(I)} \leq C h^r \|u\|_{W_\infty^r(I)}.$$

We shall next prove an analogue of this for function values.

**Theorem 1.3.2.** *With assumptions as in Theorem 1.3.1 and, in addition,  $r \geq 3$ , we have*

$$(1.3.15) \quad \|u_h - \tilde{u}_h\|_{L_\infty(I)} \leq Ch^{r+1} \|u\|_{W_\infty^r(I)}.$$

Proof: In the case  $\mu = 0$  we have from Theorem 1.2.1 that  $\theta(x_i) \equiv (u_h - \tilde{u}_h)(x_i) = 0(h^{2r-2}) \leq Ch^{r+1}$ . For  $x \in I_i$ ,  $\theta(x) = \theta(x_i) + \int_{x_i}^x \theta'(y) dy$  and (1.3.15) follows in this case from Theorem 1.3.1.

For the general case of  $\mu \geq 1$  we write

$$(1.3.16) \quad \|\theta\|_{L_\infty(I)} = \sup_{\|v\|_{L_1(I)}=1} (\theta, v).$$

For each such  $v$ , let  $-w'' = v$  in  $I$ ,  $w(0) = w(1) = 0$ . Then

$$(1.3.17) \quad (\theta, v) = (\theta', w') = (\theta', P(w'))$$

where  $P$  is the  $L_2$ -projection into  $\$_h = \$_h^{\mu-1,r-1}$ . Since  $MV(P(w')) = MV(w') = 0$ ,  $Q = \int_0^x P(w') \in \overset{\circ}{S}_h$  with  $Q' = P(w')$  so that from (1.3.3), with  $e = u_h - u$ ,

$$\begin{aligned} (1.3.18) \quad (\theta, v) &= (\theta', Q') \\ &= -\left(a_2 e', \left(\frac{1}{a_2} Q\right)' - \psi'\right) + (a_1 e, \psi') + (a_0 e, \psi) \\ &\quad + \left(e, \left(\frac{a'_2}{a_2} Q\right)'\right) \\ &\equiv I_1 + I_2 + I_3 + I_4. \end{aligned}$$

Integrating by parts, and using (1.1.11) and superapproximation,

$$\begin{aligned} (1.3.19) \quad |I_1| &= \left| \left( e, \left( a_2 \left( \frac{1}{a_2} Q \right)' - \psi' \right)' \right) \right| \\ &\leq C \|e\|_{L_\infty(I)} \left\| \left( \frac{1}{a_2} Q \right)' - \psi' \right\|_{W_1^1(I)} \\ &\leq Ch^{r+1} \|Q\|_{W_1^2(I)}. \end{aligned}$$

Similarly, using the  $W_\infty^{-1}$  error estimate of (1.1.12),

$$(1.3.20) \quad |I_2 + I_3 + I_4| \leq Ch^{r+1} \|Q\|_{W_1^2(I)}.$$

Under our quasi-uniformly assumption (1.1.7) it is easy to see that, since  $P$  is stable in  $L_1$ , it is also stable in  $W_1^1(I)$ . Thus  $\|Q\|_{W_1^2(I)} \leq C \|w\|_{W_1^2(I)} \leq C \|v\|_{L_1} = C$ . It follows from this and (1.3.16)–(1.3.18) that

$$(1.3.21) \quad \|\theta\|_{L_\infty(I)} \leq Ch^{r+1}.$$

This completes the proof of the theorem.  $\square$

Remark 1.3.1. In various special cases, better results can be had. E.g., if  $r \geq 3$ , if  $a_2 \equiv 1$  and  $a_1 \equiv 0$ , then (1.3.14) in Theorem 1.3.1 may be replaced by

$$\|(u_h - \tilde{u}_h)'\|_{L_\infty(I)} \leq Ch^{r+1} \|u\|_{W_\infty^r(I)}.$$

This is easily seen from the proof. Similarly we then have  $Ch^{r+2}$  in Theorem 1.3.2 for  $r \geq 4$ .  $\square$

Remark 1.3.2. In the case of  $\mu = 0$  we will now briefly point out how the  $W_\infty^1$  error estimate in (1.1.11) follows without any mesh restrictions. With  $e = u_h - u$  we have as in (1.3.3)

$$\begin{aligned} (\theta', \chi') &= -\left(a_2 e', \frac{1}{a_2} \chi'\right) = -\left(a_2 e', \left(\frac{1}{a_2} \chi\right)'\right) + \left(e', \frac{a'_2}{a_2} \chi\right) \\ &= -\left(a_2 e', \left(\frac{1}{a_2} \chi\right)' - \psi'\right) + (a_1 e, \psi') + (a_0 e, \psi) + \left(e', \frac{a'_2}{a_2} \chi\right) \\ &\equiv I_1 + I_2 + I_3 + I_4, \text{ for } \chi, \psi \in \overset{\circ}{S}_h. \end{aligned}$$

Using the  $H^1$  estimate (1.1.10),

$$|I_1| \leq Ch^{r-1} \|u\|_{H^r(I)} \left\| \left(\frac{1}{a_2} \chi\right)' - \psi' \right\|_{L_2(I)}.$$

By the  $L_2$  analogue of the superapproximation result (1.3.6), and using inverse estimates, cf. Section 2.1 below (recall that  $\mu = 0$  now),

$$\left\| \left(\frac{1}{a_2} \chi\right)' \right\|_{L_2(I_i)}^2 \leq C^2 h_i^2 \|\chi\|_{W_2^1(I_i)}^2 \leq C^2 h_i \|\chi\|_{W_1^1(I_i)}^2 \leq C^2 h_i \|\chi\|_{W_1^1(I)}^2.$$

Thus,

$$\left\| \left(\frac{1}{a_2} \chi\right)' - \psi' \right\|_{L_2(I)}^2 \leq C^2 \left( \sum_i h_i \right) \|\chi\|_{W_1^1(I)}^2 \leq C^2 \|\chi\|_{W_1^1(I)}^2,$$

and hence,

$$|I_1| \leq Ch^{r-1} \|u\|_{H^r(I)} \|\chi\|_{W_1^1(I)}.$$

Similarly, integrating by parts and using also a Sobolev inequality in one space dimension,

$$\begin{aligned} |I_2| &= |-(a_1 e)', \psi| \leq C \|e\|_{H^1(I)} \|\psi\|_{L_2(I)} \leq Ch^{r-1} \|u\|_{H^r(I)} \|\psi\|_{L_2(I)} \\ &\leq Ch^{r-1} \|u\|_{H^r(I)} \|\psi\|_{W_1^1(I)} \leq Ch^{r-1} \|u\|_{H^r(I)} \|\chi\|_{W_1^1(I)}. \end{aligned}$$

One treats the two remaining terms  $I_3$  and  $I_4$  in a similar manner and arrives at

$$|(\theta', \chi')| \leq Ch^{r-1} \|u\|_{H^r(I)} \|\chi\|_{W_1^1(I)}, \text{ for } \chi \in \overset{\circ}{S}_h.$$

Note that we have only used the  $H^1$ -error estimates for  $u_h - u$  in obtaining this result. These are well-known to hold for  $\mu = 0$  without mesh restrictions by use of Lagrange interpolation on each mesh interval (cf. [Schatz 1974]). As in (1.3.8)–(1.3.13) this leads to

$$\|\theta'\|_{L_\infty} \leq Ch^{r-1} \|u\|_{H^r(I)} \leq Ch^{r-1} \|u\|_{W_\infty^r(I)}.$$

Thus, to conclude the  $W_\infty^1$ -error estimate, it now suffices to consider the error in  $(u - \tilde{u}_h)'$ . But  $\tilde{u}'_h$  is the  $L_2$ -projection of  $u'$  into the now discontinuous space  $\mathbb{S}_h = \mathbb{S}_h^{-1,r-1}$  and as such  $\tilde{u}'_h$  is completely determined on each  $I_i$  by  $u'|_{I_i}$ . It follows easily that this  $L_2$ -projection is bounded in  $L_\infty$  without any mesh restrictions. So,  $\|(u - \tilde{u}_h)'\|_{L_\infty(I)} \leq C \min_{\chi \in \mathbb{S}_h} \|u' - \chi\|_{L_\infty(I)} \leq Ch^{r-1} \|u\|_{W_\infty^r(I)}$ . Thus the  $W_\infty^1$  error estimate of (1.1.11) obtains without any mesh restriction in the case  $\mu = 0$ .  $\square$

#### 1.4. Existence of superconvergence points in general.

In this section we shall consider only  $\tilde{u}_h \in \overset{\circ}{S}_h$  given by

$$(1.4.1) \quad ((u - \tilde{u}_h)', \chi') = 0, \text{ for } \chi \in \overset{\circ}{S}_h.$$

No mesh restrictions at all will be involved in this section, even for  $\mu \geq 1$ . The results may be combined with those of Section 1.3 which, however, do involve mesh restrictions for  $\mu \geq 1$ .

Our first result concerns derivatives. Let  $[\sigma]^+$  denote the smallest integer  $\geq \sigma$ .

**Theorem 1.4.1.** *Let  $k_d = [\frac{r-1}{r-1-\mu}]^+$ , and let  $\mathcal{J}_i = (x_i, x_{i+k_d})$ , for any  $i = 0, 1, \dots, N - k_d$ . There exists a point  $\eta_i \in \mathcal{J}_i$  such that  $(u - \tilde{u}_h)'(\eta_i) = 0$ .*

Proof: By [deBoor 1978, pp.108-] we may construct  $B_i \in \$^{\mu-1,r-1}$ , cf. (1.3.9), with support in  $\overline{\mathcal{J}}_i$  and  $B_i(x) > 0$  on  $\mathcal{J}_i$  (a  $B$ -spline basis function). Let now

$$(1.4.1) \quad \psi_i(x) = \int_0^x B_i(y) ds - x MV(B_i)$$

which belongs to  $\overset{\circ}{S}_h$ . Then

$$(1.4.2) \quad \int_{\mathcal{J}_i} (u - \tilde{u}_h)' B_i = ((u - \tilde{u}_h)', \psi_i' + MV(B_i)) = ((u - \tilde{u}_h)', \psi_i') = 0.$$

The theorem follows since  $B_i > 0$  on  $\mathcal{J}_i$  (note that if  $\mu = 0$ ,  $\mathcal{J}_i = I_i$ ).  $\square$

A corresponding result for function values is as follows.

**Theorem 1.4.2. a)** *For  $\mu = 0$ ,  $(u - \tilde{u}_h)(x_i) = 0$ , for all meshpoints  $x_i$ .*

*b) For  $r \geq 3$ , let  $k_f = [\frac{r-2}{r-3-\max(-1,\mu-2)}]^+$  and  $\mathcal{J}_i = (x_i, x_{i+k_f})$ .*

*There exists a point  $\xi_i \in \mathcal{J}_i$  such that  $(u - \tilde{u}_h)(\xi_i) = 0$ .*

Proof: a) follows as in Section 1.2 by noting that now  $G(x_i; \cdot) \in \overset{\circ}{S}_h$ . For b), as in the proof of Theorem 1.4.1, we may find  $B_i \in \$^{\mu-2,r-2}$  such that  $B_i$  is supported in  $\overline{\mathcal{J}}_i$  and positive in the interior. Letting  $-\psi'' = B_i$  on  $I$ ,  $\psi(0) = \psi(1) = 0$  we have  $\psi \in \overset{\circ}{S}_h$  and so

$$(1.4.3) \quad \int_{\mathcal{J}_i} (u - \tilde{u}_h) B_i = ((u - \tilde{u}_h)', \psi') = 0. \quad \square$$

Of course, Theorems 1.4.1 and 1.4.2 give no information as how to nail down what these points may be. They do, however, give us a hunting license.

#### 1.5. Superconvergence for interior points of mesh-intervals for continuous elements ( $\mu = 0$ ).

For the results in this section we refer to [Chen 1979], [Lesaint and Zlamal 1979] and also to [Bakker 1982, 1984], where many results were rediscovered but also extended to  $2m^{\text{th}}$  order problems. We point out that no mesh restrictions will be imposed.

We shall first show superconvergence for derivatives at certain points. For a mesh interval  $I_i$ ,  $L_{i,k}(x)$  shall denote the  $k^{\text{th}}$  Legendre polynomial normalized to  $I_i$ , i.e.,

if  $I_i = (x_i, x_{i+1})$ , then  $L_{i,k}(x) = L_k(2(x - (x_i + x_{i+1})/2)/h_i)$  so that  $x_i$  corresponds to  $-1$ , the midpoint to  $0$  and  $x_{i+1}$  to  $1$ .

**Theorem 1.5.1.** *For  $\mu = 0$ ,*

$$(1.5.1) \quad |(u - u_h)'(\eta_i)| \leq Ch^r \left( \|u\|_{W_\infty^r(I)} + \|u\|_{W_\infty^{r+1}(I_i)} \right)$$

where  $\eta_i$  is any zero of  $L_{i,r-1}$ .

We note that  $L_{i,r-1}$  has  $r-1$  zeroes inside  $I_i$  and that  $u'_h$  is a polynomial of degree  $r-2$ . The number of superconvergence points found is thus “maximal”. For, if we had  $r$  stable points  $\eta_j$  on each  $I_i$  where  $(u - u_h)'(\eta_j) = 0(h^r)$ , then we would have  $\|(u - u_h)'\|_{L_\infty(I)} = 0(h^r)$  which is not true in general.

Proof: By Theorem 1.3.1 it suffices to consider  $\tilde{u}_h$  instead of  $u_h$ . Now with  $P$  the  $L_2$ -projection into  $\$_h = \$_h^{-1,r-1}$ , i.e., discontinuous piecewise polynomials of degree  $\leq r-2$ , we have  $\tilde{u}'_h = Pu'$ , or, that  $\tilde{e}' \equiv \tilde{u}'_h - u'$  satisfies

$$(1.5.2) \quad (\tilde{e}', \psi) = 0, \text{ for all } \psi \in \$_h,$$

or,

$$(1.5.3) \quad \int_{I_i} \tilde{e}' \psi = 0, \text{ for all } \psi \in \Pi_{r-2}(I_i).$$

Taylor-expanding  $\tilde{e}'$  around the midpoint  $(x_i + x_{i+1})/2$  and writing the expansion in terms of the Legendre polynomials, we have

$$(1.5.4) \quad \tilde{e}'(x) = c_0 L_{i,0}(x) + c_1 L_{i,1}(x) + \cdots + c_{r-1} L_{i,r-1}(x) + 0(h^r),$$

where the  $0(h^r)$  term now involves  $(r+1)^{\text{th}}$  derivatives of  $u$  on  $I_i$ . By (1.5.3),  $c_0, c_1, \dots, c_{r-2}$  are all  $0(h^r)$ , so that

$$(1.5.5) \quad \tilde{e}'(x) = c_{r-1} L_{i,r-1}(x) + 0(h^r).$$

The desired result follows. □

The corresponding superconvergence result for function values is as follows.

**Theorem 1.5.2.** *For  $\mu = 0$ ,  $r \geq 3$ ,*

$$(1.5.6) \quad (u - u_h)(\xi_i) \leq Ch^{r+1} \left( \|u\|_{W_\infty^r(I)} + \|u\|_{W_\infty^{r+1}(I_i)} \right)$$

where  $\xi_i$  is any zero of  $L'_{i,r-1}$  (or, a mesh point, where we have  $0(h^{2r-2})$ ).

Again, the number of superconvergence points found is maximal.

Proof: From Theorem 1.3.2 it is again enough to consider  $\tilde{u}_h$ . By (1.5.5) and Theorem 1.2.1,

$$(1.5.7) \quad \begin{aligned} \tilde{e}(x) &= \tilde{e}(x_i) + \int_{x_i}^x \tilde{e}'(y) dy \\ &= 0(h^{2r-2}) + c_{r-1} \int_{x_i}^x L_{i,r-1}(y) dy + 0(h^{r+1}). \end{aligned}$$

Since  $(r-1)rL_{r-1} = -((1-x^2)L'_{r-1})'$  by Legendre's differential equation, the result follows. □

### 1.6. Superconvergence in derivatives at points about which the meshes are locally symmetric ( $r$ even).

In this section we shall consider general  $\mu$  and since Theorem 1.5.1 gives a maximally possible number of superconvergence points when  $\mu = 0$ , identifiable a priori to boot, we have in mind here  $\mu \geq 1$ . Our arguments in this section follow [Wahlbin 1992] and are restricted to one-dimensional problems. In return, we then obtain more sharply delineated results than in several dimensions, in particular with respect to mesh-geometry outside an immediate neighborhood of the point of interest and with respect to how close to the boundary of  $I$  the results are valid. By Theorem 1.3.1 it turns out to be sufficient to treat only  $\tilde{u}_h \in \overset{\circ}{S}_h$  given by  $((u - \tilde{u}_h)', \chi') = 0$ , for  $\chi \in \overset{\circ}{S}_h$ ; this is because only superconvergence of order 1 will be at issue. The quasi-uniformity condition (1.1.7) is assumed throughout.

Let  $w = u'$  and  $w_h = \tilde{u}'_h \in \$_h = \$^{\mu-1, r-1}$ ; then as we have seen before,

$$(1.6.1) \quad w_h = P(w)$$

where  $P$  is the  $L_2$ -projection into  $\$_h$ . Fixing a point  $\eta$ , there exists  $\delta = \delta_h^\eta \in \$_h$  such that

$$(1.6.2) \quad \chi(\eta) = (\chi, \delta), \text{ for all } \chi \in \$_h.$$

We shall refer to  $\delta$  as a “discrete delta-function” centered at  $\eta$ . In case  $\mu = 0$ ,  $\delta$  is of course a function with support on  $I_i \ni \eta$ . For  $\mu \geq 1$  this is no longer true;  $\delta_h^\eta$  suffers some influence from outside  $I_i$ . However such influences are small. In fact we have:

**Lemma 1.6.1.** *Assume (1.1.7). There exist constants  $C$  and  $c > 0$ , independent of  $h$  and  $\eta$ , such that*

$$(1.6.3) \quad |\delta_h^\eta(x)| \leq \frac{C}{h} e^{-c|x-\eta|/h}.$$

Such an exponential decay property was noted in [Descloux 1972] and [Douglas, Dupont and Wahlbin 1975]. Lemma 1.6.1 is taken straight from [Wahlbin 1991, Lemma 3.4]. It will also be proven below in Theorem 3.2.2.

Let now  $\widehat{w}_s(\cdot; \eta)$  denote the Taylor polynomial of degree  $s$  for  $w$  centered at  $\eta$  and consider in particular

$$(1.6.4) \quad \widehat{w}_{r-1}(x; \eta) = \widehat{w}_{r-2}(x; \eta) + w^{(r-1)}(\eta)(x - \eta)^{r-1}/(r-1)!.$$

Then since  $\widehat{w}_{r-2}(\cdot; \eta) \in \$_h$  and  $w_h = P(w)$ ,

$$\begin{aligned} (1.6.5) \quad w(\eta) - w_h(\eta) &= \widehat{w}_{r-2}(\eta; \eta) - w_h(\eta) \\ &= (\widehat{w}_{r-2} - w_h, \delta) = (\widehat{w}_{r-2} - w, \delta) \\ &= (\widehat{w}_{r-1} - w^{(r-1)}(\eta)(x - \eta)^{r-1}/(r-1)! - w, \delta) \\ &= (\widehat{w}_{r-1} - w, \delta) - \frac{w^{(r-1)}(\eta)}{(r-1)!} ((x - \eta)^{r-1}, \delta). \end{aligned}$$

Here

$$(1.6.6) \quad |(\widehat{w}_{r-1} - w)(x)| \leq \frac{1}{r!} |x - \eta|^r \|w\|_{W_\infty^r(I)} \leq \frac{1}{r!} |x - \eta|^r \|u\|_{W_\infty^{r+1}(I)}$$

and using (1.6.3) we thus find that

$$(1.6.7) \quad \begin{aligned} & |(\widehat{w}_{r-1} - w, \delta)| \\ & \leq C \left( \int_0^1 |x - \eta|^r h^{-1} \exp(-c|x - \eta|/h) dx \right) \|u\|_{W_\infty^{r+2}(I)} \\ & \leq Ch^r \|u\|_{W_\infty^{r+1}(I)}. \end{aligned}$$

Thus, in total, from (1.6.5)–(1.6.7),

$$(1.6.8) \quad |(u' - \tilde{u}'_h)(\eta)| \leq Ch^r \|u\|_{W_\infty^{r+1}(I)} + \frac{|u^{(r)}(\eta)|}{(r-1)!} |((x-\eta)^{r-1}, \delta_h^\eta(x))|.$$

In general there is no reason to expect the last term on the right in (1.6.8) to be less in order than  $O(h^{r-1})$  (which it clearly is from (1.6.3), cf. the calculation in (1.6.7)). However, if the mesh happened to be symmetric around  $\eta$  we might expect  $\delta_h^\eta$  to be evenly symmetric around  $\eta$ , at least “locally” (Occam’s razor). If then  $r$  is even so that  $(x - \eta)^{r-1}$  is oddly symmetric around  $\eta$ , maybe the term will be of higher order due to cancellation.

Let us formalize “mesh-symmetry around  $\eta$ ” as follows: For a sufficiently large constant  $C_1$ , with  $\mathcal{J} = \mathcal{J}(h, \eta, C_1) = \{x \in R : |x - \eta| \leq C_1 h \ln 1/h\}$ , we assume that  $\mathcal{J} \subseteq I$  and also, if  $\chi(x) \in \$_h(\mathcal{J}) = \$_h|_{\mathcal{J}}$ , then

$$(1.6.9) \quad \bar{\chi}(x) \equiv \chi(\eta - (x - \eta)) \in \$_h(\mathcal{J}).$$

In words, if we reflect (locally) through  $\eta$  we stay in  $\$_h$ .

Sharpening the razor, we shall prove the following:

**Theorem 1.6.1.** *Assume the quasi-uniformity condition (1.1.7), that  $r$  is even and that  $\mu \geq 1$ . There exist constants  $C_1$  and  $C_2$ , independent of  $u$ ,  $h$  and  $\eta$ , such that if  $\eta$  ( $= \eta(h)$ ) is such that the mesh is symmetric around  $\eta$  with constant  $C_1$  (i.e., in a symmetric  $C_1 h \ln 1/h$  neighborhood inside  $I$ ), then for  $u_h$  the solution of (1.1.6),*

$$(1.6.10) \quad |(u - u_h)'(\eta)| \leq C_2 h^r \|u\|_{W_\infty^{r+1}(I)}.$$

Remark 1.6.1. The reader may wonder about the restriction  $\mu \geq 1$ . Note that if  $\mu = 0$  and  $\eta$  a mesh point (which is symmetric) then  $u'_h$  is not well defined. This can be gotten around by taking limits of averages of left and right derivatives. However, the case  $\mu = 0$  is so thoroughly analyzed in Section 1.5 that we shall wait to give the argument until the multi-dimensional case, where it is more urgent.  $\square$

Proof of Theorem 1.6.1: As already noted, by Theorem 1.3.1 it suffices to consider  $u - \tilde{u}_h$ , and by (1.6.8) we need only consider the last term there. Introducing the notation  $(f, g)_{I'} = \int_{I'} fg$  for  $I'$  a set in  $I$  we first have using (1.6.3),

$$(1.6.11) \quad \begin{aligned} & |((x - \eta)^{r-1}, \delta)_{I \setminus \mathcal{J}}| \\ & \leq C \int_{|x - \eta| \geq C_1 h \ln 1/h} |x - \eta|^{r-1} h^{-1} \exp(-c|x - \eta|/h) dx \\ & \leq Ch^{r-1} \int_{C_1 \ln 1/h}^\infty y^{r-1} e^{-cy} dy \\ & \leq Ch^{r-1} e^{-(c/2)C_1 \ln 1/h} \leq Ch^r, \end{aligned}$$

provided  $C_1 c/2 \geq 1$ . Furthermore, since  $(x - \eta)^{r-1}$  is odd around  $\eta$ ,

$$(1.6.12) \quad ((x - \eta)^{r-1}, \delta)_{\mathcal{J}} = ((x - \eta)^{r-1}, \delta_{odd})_{\mathcal{J}}$$

where

$$(1.6.13) \quad f_{odd} = (f(x) - \bar{f}(x))/2 = (f(x) - f(2\eta - x))/2, \text{ for } x \in \mathcal{J},$$

and  $f = f_{odd} + f_{even}$ . We shall need an extension operator  $E$  from  $\$h(\mathcal{J})$  to  $\$h = S_h(I)$ . Assuming (as we may possibly after a slight enlargement) that  $\mathcal{J}$  is a union of mesh intervals, a simple such operator  $E$  is constructed in [Douglas, Dupont and Wahlbin 1975, Lemma 2.1] in such a fashion that

$$(1.6.14) \quad \|E\chi\|_{L_p(I \setminus \mathcal{J})} \leq C\|\chi\|_{L_p(\mathcal{J}')}, \text{ for } \chi \in \$h(\mathcal{J}), 1 \leq p \leq \infty.$$

Here  $\mathcal{J}'$  denotes the union of the two mesh intervals at the very ends of  $\mathcal{J}$ . Now since  $(\delta_{odd}, \delta_{even})_{\mathcal{J}} = 0$ ,

$$(1.6.15) \quad \begin{aligned} \|\delta_{odd}\|_{L_2(\mathcal{J})}^2 &= (\delta_{odd}, \delta_{odd})_{\mathcal{J}} = (\delta_{odd}, \delta)_{\mathcal{J}} \\ &= (E\delta_{odd}, \delta) - (E\delta_{odd}, \delta)_{I \setminus \mathcal{J}}. \end{aligned}$$

Since  $(E\delta_{odd}, \delta) = \delta_{odd}(\eta) = 0$  (here we used that  $E\delta_{odd} \in \$h(I)$ , i.e., that  $\delta_{odd} \in \$h(\mathcal{J})$ , i.e., the symmetry of the mesh for the one and only time!) and using (1.6.3) twice,

$$(1.6.16) \quad |(E\delta_{odd}, \delta)_{I \setminus \mathcal{J}}| \leq Ch^{2k} \|\delta_{odd}\|_{L_1} \leq Ch^{2k},$$

for any given  $k$ , provided  $C_1 = C_1(k)$  is taken large enough. It follows from (1.6.15) that

$$(1.6.17) \quad \|\delta_{odd}\|_{L_2(\mathcal{J})} \leq Ch^k$$

and hence from (1.6.12) that

$$(1.6.18) \quad |((x - \eta)^{r-1}, \delta)_{\mathcal{J}}| \leq Ch^r$$

(with  $C_1$  further enlarged if necessary). This and (1.6.11) shows that

$$(1.6.19) \quad |((x - \eta)^{r-1}, \delta)| \leq Ch^r$$

and, as noted, the theorem now follows from (1.6.8).  $\square$

Remark 1.6.2. It is easily seen from (1.6.7) that the full  $\|u\|_{W_\infty^{r+1}(I)}$  norm may be replaced by  $\|u\|_{W_\infty^{r+1}(\mathcal{J})} + \|u\|_{W_\infty(I)}$ . In fact, further localization with respect to smoothness demands is possible but we shall wait with that argument until the multi-dimensional case so as not to overburden the present exposition.  $\square$

A particular example of symmetry points  $\eta$  which satisfy (1.6.9) occurs in a uniform mesh,  $h_i \equiv h$  for all  $i$ . Then both meshpoints and midpoints of mesh-intervals are symmetric, *provided* they stay away from the boundary by a distance  $C_1 h \ln 1/h$ .

**Corollary 1.6.2.** *Let  $r$  be even and the meshes uniform. Then for  $\eta$  a meshpoint or midpoint with*

$$(1.6.20) \quad \text{dist}(\eta, \{0, 1\}) \geq C_1 h \ln 1/h,$$

*we have*

$$(1.6.21) \quad |(u - u_h)'(\eta)| \leq C_2 h^r \|u\|_{W_\infty^{r+1}(I)}.$$

Perhaps the most interesting aspect of this result is that the condition (1.6.20) is necessary in general, as we shall see in the next section. Intuitively, the discrete delta-function, which now suffers from outside influences, loses its symmetry when the point  $\eta$  at which it is centered approaches the boundary.

### 1.7. Necessity of staying $C_1 h \ln 1/h$ away from the boundary for superconvergence in the uniform mesh case ( $\mu = 1$ ).

We place ourselves in the situation of Corollary 1.6.2, i.e., uniform meshes, and we take the particular “lowest” case of Hermite cubics, i.e.,  $r = 4$ ,  $\mu = 1$ . If our original problem is  $-u'' = -3x^2$ ,  $u(0) = u(1) = 0$ , with solution  $u(x) = (x^4 - x)/4$ , we thus look at the  $L_2$ -projection  $P(x^3 - 1/4)$  of  $u'$  into  $\$_h = \$_h^{0,3}(I)$ , the continuous quadratics. Since constants are clearly reproduced by  $P$ , we consider the error in

$$(1.7.1) \quad x^3 - P(x^3)$$

at mesh and midpoints. Notice that now  $u^{(r+1)} = u^{(5)} \equiv 0$  which perhaps makes the counterexample even more convincing. Again, our arguments are taken from [Wahlbin 1992, Section 3]; indeed, here we shall merely describe them and refer to this reference for further details.

There, the error of meshpoints and midpoints was explicitly calculated for  $x^3 - P(x^3)$ , modulo terms which are of order  $e^{-cN}$ ,  $c > 0$  constant,  $N = 1/h$ . This was done by using different basis functions for  $\$_h$  when regarded as a trial or test-space, respectively; the system of equations to be solved was then reduced to easily analyzable three-term recurrence relations.

Let, in this section,  $e = x^3 - P(x^3)$  and

$$(1.7.2) \quad e_{2i} = e(x_i), \quad e_{2i+1} = e(x_{i+1/2}) \equiv e((x_i + x_{i+1})/2), \quad i = 0, N-1.$$

From [Wahlbin 1992], by explicit calculation and a simple algebraic manipulation,

$$(1.7.3) \quad -e_{2i-1} + 6e_{2i+1} - e_{2i+3} = 0, \quad i = 1, 2, \dots, N-2$$

and

$$(1.7.4) \quad -e_{2i-2} + 6e_{2i} - e_{2i+2} = 0, \quad i = 1, 2, \dots, N-1,$$

as well as

$$(1.7.5) \quad e_0 + 2e_1 = -\frac{h^3}{20}.$$

From (1.7.3)–(1.7.5), with  $\rho = 3 - \sqrt{8}$  (considering points in the left half of  $I$ ),

$$(1.7.6) \quad (u - \tilde{u}_h)'(\eta_i) = \begin{cases} A_1 h^3 \rho^i + E_{i,1}, & \text{if } \eta_i = x_i, \\ A_2 h^3 \rho^i + E_{i,2}, & \text{if } \eta_i = x_{i+1/2} \end{cases}$$

where  $A_1 \simeq -0.071$ ,  $A_2 \simeq 0.010$  (and  $\rho \simeq 0.172$ ). Here, for  $i \leq [N/2]$ ,  $|E_{i,j}| \leq e^{-cN}$ , with  $c > 0$ .

Thus, if we want convergence as  $Ch^{3+\sigma}$  for some  $\sigma > 0$  we need (for  $i \leq [N/2]$ )  $\rho^i \leq Ch^\sigma$  or,  $i \geq (\ln C + \sigma \ln 1/h)/(\ln 1/\rho)$ , i.e.,

$$\text{dist}(\eta_i, 0) \simeq ih \geq C\sigma h(\ln 1/h)/(\ln 1/\rho).$$

(Of course, in general the presence of a non-zero fifth derivative of  $u$  precludes superconvergence to extra order  $> 1$  in the case of Hermite cubics.)

### 1.8. Finding all superconvergent points for function values and derivatives in the case of a locally uniform mesh for $\mu = 1$ and $r$ even (with a remark about smoothest cubics).

We place ourselves in a situation with  $r$  even and locally uniform meshes. Then for  $\mu \geq 1$ , meshpoints and midpoints are superconvergent for derivatives to order 1, provided the patches of uniform meshes, of extent  $C_1 h \ln 1/h$  themselves, stay  $Ch \ln 1/h$  away from the boundary of  $I$ . In this section we shall investigate the consequences which this has for superconvergence in both function-values and derivatives, mainly in the case of  $\mu = 1$ . In fact, we shall find all such superconvergence points.

We take the case of  $\tilde{u}_h$ , i.e.,  $a_2 \equiv 1$ ,  $a_1 \equiv a_0 \equiv 0$ , cf. Section 1.3. Let us work on an interval  $I_i = [x_i, x_{i+1}]$  where we now know that  $e \equiv u - \tilde{u}_h$  satisfies (with  $x_{i+1/2} = (x_i + x_{i+1})/2$ )

$$(1.8.1) \quad e'(x_i), e'(x_{i+1/2}) \text{ and } e'(x_{i+1}) = O(h^r).$$

Let

$$(1.8.2) \quad \varphi_k = \begin{cases} x^k & \text{on } I_i, \\ 0 & \text{on } I \setminus \bar{I}_i, \end{cases} \quad k = 0, 1, \dots, r-3,$$

(note that now  $r \geq 4$  since  $\mu \geq 1$  and  $r$  is even). With  $w_k$  given by

$$(1.8.3) \quad \begin{aligned} -w_k'' &= \varphi_k, \quad x \in I, \\ w_k(0) &= w_k(1) = 0, \end{aligned}$$

we have the relations

$$(1.8.4) \quad \int_{I_i} e \varphi_k = \int_I e' w'_k.$$

Since  $\mu = 1$  and  $k \leq r-3$ , we find that  $w_k \in \overset{\circ}{S}_h$  and so

$$(1.8.5) \quad \int_{I_i} e x^k = 0, \quad k = 0, 1, \dots, r-3.$$

If we now Taylor-expand  $e$  on  $I_i$  around  $x_{i+1/2}$  to order  $r$  and write the expansion in terms of (normalized) Legendre polynomials  $L_{i,k} = L_k(2(x - x_{i+1/2})/h_i)$  where  $L_k(1) = 1$ , we have

$$(1.8.6) \quad e(x) = c_0 L_{i,0} + c_1 L_{i,1} + \dots + c_{r-3} L_{i,r-3} + c_{r-2} L_{i,r-2} + c_{r-1} L_{i,r-1} + c_r L_{i,r} + R,$$

where

$$(1.8.7) \quad R = R(x, x_{i+1/2}) = \int_{x_{i+1/2}}^x \frac{(x-t)^r}{r!} u^{(r+1)}(t) dt = 0(h^{r+1}) \|u\|_{W_\infty^{r+1}(I_i)}.$$

Due to (1.8.5),  $c_0, c_1, \dots, c_{r-3}$  are all  $0(h^{r+1})$ . Hence, since  $R' = 0(h^r)$  and  $L'_{i,k} = \frac{1}{h_i} L'_k$ ,

$$(1.8.8) \quad e'(x) = c_{r-2} L'_{i,r-2} + c_{r-1} L'_{i,r-1} + c_r L'_{i,r} + 0(h^r).$$

Thus, from (1.8.1), since  $L'_{i,r-2}$  and  $L'_{i,r}$  are odd about  $x_{i+1/2}$ , and  $L'_{i,r-1}$  is even,

$$(1.8.9) \quad e'(x_{i+1/2}) = c_{r-1} L'_{i,r-1}(x_{i+1/2}) + 0(h^r) = 0(h^r)$$

so that since  $L'_{i,r-1}(x_{i+1/2}) = \frac{1}{h_i} L'_{r-1}(0) \neq 0$ , we have

$$(1.8.10) \quad c_{r-1} = 0(h^{r+1}).$$

Therefore

$$(1.8.11) \quad e'(x) = c_{r-2} L'_{i,r-2} + c_r L'_{i,r} + 0(h^r).$$

We next plug in the fact that  $e'(x_i), e'(x_{i+1}) = 0(h^r)$  from (1.8.1):

$$(1.8.12) \quad \begin{aligned} e'(x_i) &= c_{r-2} \frac{1}{h_i} L'_{r-2}(-1) + \frac{c_r}{h_i} L'_r(-1) = 0(h^r), \\ e'(x_{i+1}) &= c_{r-2} \frac{1}{h_i} L'_{r-2}(-1) + \frac{c_r}{h_i} L'_r(1) = 0(h^r), \end{aligned}$$

from which we obtain

$$(1.8.13) \quad c_r = -c_{r-2} \frac{L'_{r-2}(1)}{L'_r(1)} + 0(h^{r+1}).$$

Hence, from (1.8.6) and the above

$$(1.8.14) \quad \begin{aligned} e(x) &= c_{r-2} \left[ L_{i,r-2}(x) - \frac{L'_{r-2}(1)}{L'_r(1)} L_{i,r}(x) \right] + 0(h^{r+1}) \\ &= c_{r-2} Q(x) + 0(h^{r+1}) \end{aligned}$$

where

$$(1.8.15) \quad Q(x) = L_{i,r-2}(x) - \frac{(r-2)(r-1)}{r(r+1)} L_{i,r}(x)$$

if we use the common ( $L_k(1) = 1$ ) normalization of the Legendre polynomials. One may use Rodrigues' formula and elementary manipulations to see that (normalized to  $(-1, 1)$ )

$$(1.8.16) \quad Q(x) = \text{const.} \frac{d^{r-2}}{dx^{r-2}} \left[ (x^2 - 1)^{r-2} \left( x^2 - \frac{r+2}{r-2} \right) \right]$$

so that repeated application of Rolle's theorem shows that  $Q(x)$  has exactly  $r-2$  roots inside  $I_i$  (and two roots outside).

Using now also Theorem 1.3.2 (note that  $r \geq 4$  since  $r$  is even and  $\mu \geq 1$ ) we thus have the following:

**Theorem 1.8.1.** Let  $r$  be even,  $\mu = 1$  and, with  $e = u - u_h$ ,  $u_h$  the solution to (1.1.6),  $e'(x_i)$ ,  $e'(x_{i+1/2})$  and  $e'(x_{i+1}) = 0(h^r)$  on  $I_i$ . Then

$$(1.8.17) \quad |e(\xi)| = 0(h^{r+1})$$

for  $\xi$  any of the  $r - 2$  roots in  $\bar{I}_i$  of  $Q(x)$  given in (1.8.15), and only then in general. In particular, this is so if the mesh is uniform in a symmetric  $C_1 h \ln 1/h$  neighborhood of  $I_i$  which is inside  $I$ , with  $C_1$  large enough.

In contradistinction to the case  $\mu = 0$  (see Theorem 1.5.2) we now don't have the "maximum" possible number of superconvergence points on  $\bar{I}_i$ , namely  $r$ , but merely  $r - 2$  of them. (The two extra roots occur outside the interval  $\bar{I}_i$ ; in principle there would be no problem in using them too.)

Example 1.8.1. In the case of Hermite cubics ( $r = 4$ ,  $\mu = 1$ ), the two superconvergent points for function values are given by

$$(1.8.18) \quad \xi = x_{i+1/2} \pm \left( \frac{1}{4} - \frac{1}{\sqrt{30}} \right)^{1/2} h \simeq x_{i+1/2} \pm (.2596648112\dots)h. \quad \square$$

One may now go back and differentiate (1.8.14) to deduce that  $e'(x)$  is superconvergent to order 1 for points  $\eta \in \bar{I}_i$  such that

$$(1.8.19) \quad Q'(\eta) = L'_{i,r-2}(\eta) - \frac{L'_{r-2}(1)}{L'_r(1)} L'_{i,r}(\eta) = 0.$$

There are of course always such roots at  $x_i$ ,  $x_{i+1/2}$  and  $x_{i+1}$ ; the corresponding superconvergence points are now "rediscovered". However, apart from these,  $Q'$  has  $r - 4$  additional roots inside  $I_i$  which we didn't know about before. Indeed, this gives a "full" complement of  $r - 1$  superconvergence points on  $\bar{I}_i$  for first derivatives.

**Theorem 1.8.2.** Let  $r$  be even,  $\mu = 1$ , and with  $e = u - u_h$ ,  $u_h$  the solution to (1.1.6),  $e'(x_i)$ ,  $e'(x_{i+1/2})$  and  $e'(x_{i+1}) = 0(h^r)$  on  $I_i$ . Then

$$(1.8.20) \quad |e'(\eta)| = 0(h^r)$$

where  $\eta$  is any of the roots of  $Q'$  in  $\bar{I}_i$ . In particular, this is so if the mesh is uniform in a symmetric  $C_1 h \ln 1/h$  neighborhood of  $I_i$  which is inside  $I$ , with  $C_1$  large enough.

Example 1.8.2. One may treat cases of other  $\mu$  in ad hoc manners. For example, in the case of smoothest cubics ( $\mu = 2$ ,  $r = 4$ ) one may consider two adjoining equal length intervals  $\bar{I}_i \cup \bar{I}_{i+1}$  and use that  $e' = 0(h^4)$  at five points: the meshpoints and the midpoints; that  $e$  is  $C^2$  across  $x_{i+1}$ ; and also the fact that (if  $a_2 \equiv 1$ ,  $a_1 = a_0 \equiv 0$ )  $\int_{\bar{I}_i \cup \bar{I}_{i+1}} e B_i dx = 0$  where  $B_i$  is the piecewise linear "tent" function;  $B_i \equiv 0$  outside  $I \setminus (\bar{I}_i \cup \bar{I}_{i+1})$ ,  $B_i(x_{i+1}) = 1$  and  $B_i \in C^0(I)$ . This follows since, if  $-w'' = B_i$ ,  $w(0) = w(1) = 0$ , then  $w \in \overset{\circ}{S}_h$ ; it corresponds to the relations (1.8.5). The two superconvergent points for function values turn out to be the very same as for Hermite cubics (Example 1.8.1, (1.8.18)).  $\square$

## 1.9. Superconvergence in function values at points about which the meshes are locally symmetric ( $r$ odd).

This section is analogous to the investigation for even  $r$  in Section 1.6.

**Theorem 1.9.1.** Assume the quasi-uniformity condition (1.1.7), that  $r$  is odd and that  $\mu \geq 1$ . There exist constants  $C_1$  and  $C_2$ , independent of  $u$ ,  $h$  and  $\xi$  such that if  $\xi$  ( $= \xi(h)$ ) is such that the mesh is symmetric around  $\xi$  with constant  $C_1$  (i.e., symmetric in a  $C_1 h \ln 1/h$  neighborhood, cf. definition (1.6.9)), then for  $u_h$  the solution of (1.1.6),

$$(1.9.1) \quad |(u - u_h)(\xi)| \leq C_2 h^{r+1} \|u\|_{W_\infty^{r+1}(I)}.$$

Proof: Since  $\mu \geq 1$  and hence  $r \geq 3$ , by Theorem 1.3.2 it suffices to treat  $\tilde{u}_h$  given by  $((u - \tilde{u}_h)', \chi') = 0$ , for  $\chi \in \mathbb{S}_h$ . Let  $\xi$  be a point in  $I$  and

$$(1.9.2) \quad \varphi = \varphi(x; \xi) = \frac{1}{2}(\chi_{[0, \xi]} - \chi_{[\xi, 1]}),$$

half the difference of the characteristic functions of  $[0, \xi]$  and  $[\xi, 1]$ , respectively. Then

$$(1.9.3) \quad (u - \tilde{u}_h)(\xi) = ((u - \tilde{u}_h)', \varphi),$$

or, with  $w = u'$  since then  $\tilde{u}'_h = P(u')$ , the  $L_2$ -projection into  $\mathbb{S}_h = \mathbb{S}_h^{\mu-1, r-1}$ ,

$$(1.9.4) \quad \begin{aligned} (u - \tilde{u}_h)(\xi) &= (w - P(w), \varphi) = (w - P(w), \varphi - P(\varphi)) \\ &= (w - \chi, \varphi - P(\varphi)), \text{ for any } \chi \in \mathbb{S}_h. \end{aligned}$$

Letting  $\hat{w}_s = \hat{w}_s(x; \xi)$  be the Taylor polynomial of degree  $s$  centered at  $\xi$  we have taking  $\chi = \hat{w}_{r-2} \in \mathbb{S}_h$  in (1.9.4),

$$(1.9.5) \quad (u - u_h)(\xi) = (w - \hat{w}_{r-1}, \varphi - P(\varphi)) - \frac{w^{(r)}(\xi)}{(r-1)!}((x - \xi)^{r-1}, \varphi - P(\varphi)).$$

We now need a suitable analogue of the exponential decay properties for the discrete delta-function, given in Lemma 1.6.1. In fact, based on this, we have the following:

**Lemma 1.9.1.** There exist  $C$  and  $c > 0$  such that with  $\varphi$  as above,

$$(1.9.6) \quad |(\varphi - P(\varphi))(x)| \leq Ce^{-c|x-\xi|/h}.$$

Proof: From [Wahlbin 1991, Theorem 3.3] or Theorem 3.2.4 below we have with  $A_d = A_d(x) = \{y : |x - y| \leq d\} \cap I$ , for  $d \geq c_0 h$  ( $c_0$  large enough),

$$(1.9.7) \quad \begin{aligned} |(\varphi - P(\varphi))(x)| &\leq C \min_{\chi \in \mathbb{S}_h} \|\varphi - \chi\|_{L_\infty(A_d)} + Cd^{-1/2} e^{-cd/h} \|\varphi - P(\varphi)\|_{L_2(A_d)}, \end{aligned}$$

with  $c > 0$ . If  $A_d$  doesn't include  $\xi$ , then  $\varphi \equiv 1/2$  or  $-1/2$  so that the first term on the right of (1.9.7) is zero. Furthermore, we have already remarked that  $P$  is stable in any  $L_p$ -space, cf. Theorem 3.2.3 below. Since then

$$(1.9.8) \quad \|\varphi - P(\varphi)\|_{L_2(A_d)} \leq Cd^{1/2} \|\varphi - P(\varphi)\|_{L_\infty(A_d)} \leq Cd^{1/2}$$

we obtain the lemma.  $\square$

Continuing now with the proof of Theorem 1.9.1, from (1.9.6) we have for the first term in (1.9.5),

$$(1.9.9) \quad \begin{aligned} |(w - \widehat{w}_{r-1}, \varphi - P(\varphi))| &\leq C\|u\|_{W_\infty^{r+1}(I)} \int_0^1 |x - \xi|^r e^{-c|x-\xi|/h} dx \\ &\leq Ch^{r+1}\|u\|_{W_\infty^{r+1}(I)}, \end{aligned}$$

so that (corresponding to (1.6.8)),

$$(1.9.10) \quad |(u - u_h)(\xi)| \leq Ch^{r+1}\|u\|_{W_\infty^{r+1}(I)} + \frac{|u^{(r)}(\xi)|}{(r-1)!}|((x - \xi)^{r-1}, \varphi - P(\varphi))|.$$

Let now  $\mathcal{J} = \mathcal{J}(h, \xi, C_1)$  be a symmetric  $C_1 h \ln 1/h$  neighborhood about  $\xi$ , assumed inside  $I$ . Again by (1.9.6), and with the notation  $(f, g)_\mathcal{J} = \int_{\mathcal{J}} f g$ ,

$$(1.9.11) \quad |((x - \xi)^{r-1}, \varphi - P(\varphi))_{I \setminus \mathcal{J}}| \leq Ch^k, \text{ for any } k = k(C_1).$$

It remains to estimate

$$(1.9.12) \quad \begin{aligned} ((x - \xi)^{r-1}, \varphi - P(\varphi))_\mathcal{J} &= ((x - \xi)^{r-1}, \varphi_{even} - (P(\varphi))_{even})_\mathcal{J} \\ &= -((x - \xi)^{r-1}, (P(\varphi))_{even})_\mathcal{J} \end{aligned}$$

where we used that  $(r-1)$  is now even and  $\varphi$  odd about  $\xi$ , and where  $f_{even}$  denotes the even part of  $f$  (on  $\mathcal{J}$ ). Further, again by trivial odd-even considerations,

$$(1.9.13) \quad \begin{aligned} \|(P(\varphi))_{even}\|_{L_2(\mathcal{J})}^2 &= ((P(\varphi))_{even}, (P(\varphi))_{even})_\mathcal{J} \\ &= ((P(\varphi))_{even}, P(\varphi))_\mathcal{J} = ((P(\varphi))_{even}, P(\varphi) - \varphi)_\mathcal{J}. \end{aligned}$$

Using then for the first and only time in this proof the assumption that the mesh is symmetric about  $\xi$  (on  $\mathcal{J}$ ), cf. (1.6.9), we have  $(P(\varphi))_{even} \in \$h(\mathcal{J})$ . Employing the extension operator  $E$  from  $\$h(\mathcal{J})$  to  $\$h(I)$ , cf. (1.6.14),

$$(1.9.14) \quad \begin{aligned} \|(P(\varphi))_{even}\|_{L_2(\mathcal{J})}^2 &= (E((P(\varphi))_{even}), P(\varphi) - \varphi)_\mathcal{J} \\ &= (E((P(\varphi))_{even}), P(\varphi) - \varphi) - (E((P(\varphi))_{even}), P(\varphi) - \varphi)_{I \setminus \mathcal{J}} \\ &= -(E((P(\varphi))_{even}), P(\varphi) - \varphi)_{I \setminus \mathcal{J}} \\ &\leq Ch^{2k}, \end{aligned}$$

for any  $k$ , for  $C_1$  large enough, where we again used (1.9.6). The theorem obtains from this, reported into (1.9.12), and then used together with (1.9.11) in (1.9.10).  $\square$

Remark 1.9.1. From (1.9.9) it is easy to replace  $\|u\|_{W_\infty^{r+1}(I)}$  in (1.9.1) by  $\|u\|_{W_\infty^{r+1}(\mathcal{J})} + \|u\|_{W_\infty(I)}$ .  $\square$

Corresponding to Corollary 1.6.2 we now have the following.

**Corollary 1.9.2.** *Let  $r$  be odd and the meshes uniform. Then for  $\xi$  a meshpoint or midpoint with*

$$(1.9.15) \quad \text{dist}(\xi, \{0, 1\}) \geq C_1 h \ln 1/h,$$

*we have*

$$(1.9.16) \quad |(u - u_h)(\xi)| \leq C_2 h^{r+1} \|u\|_{W_\infty^{r+1}(I)}.$$

### 1.10. Finding all superconvergent points for function values and derivatives in the case of a locally uniform mesh for $\mu = 1$ and $r$ odd.

Corresponding to Section 1.8 we consider locally uniform meshes so that for  $r$  odd, meshpoints and midpoints are superconvergent for function values, locally. Let  $e = u - \tilde{u}_h$ ,  $\tilde{u}_h$  the projection with respect to the simple form  $(e', \chi')$ . Taking  $\mu = 1$ , as in (1.8.5) we have for any  $I_i$ ,

$$(1.10.1) \quad \int_{I_i} ex^k = 0, \text{ for } k = 0, 1, \dots, r-3.$$

Thus, using Taylor expansion about the midpoint  $x_{i+1/2}$  to polynomial degree  $r$ , and writing this in terms of the (normalized) Legendre polynomials.

$$(1.10.2) \quad e(x) = c_{r-2} L_{i,r-2}(x) + c_{r-1} L_{i,r-1}(x) + c_r L_{i,r}(x) + O(h^{r+1}).$$

Since  $r$  is odd,  $e(x_{i+1/2}) = O(h^{r+1})$  means that  $c_{r-1} = O(h^{r+1})$  and so, (if  $L_k$  normalized to  $L_k(1) = 1$ )

$$(1.10.3) \quad e(x_{i+1}) = (c_{r-2} + c_r) + O(h^{r+1});$$

the assumption  $e(x_{i+1}) = O(h^{r+1})$  then gives

$$(1.10.4) \quad e(x) = c_{r-2}(L_{i,r-2}(x) - L_{i,r}(x)) + O(h^{r+1}).$$

(Use of  $e(x_i) = O(h^{r+1})$  gives nothing new.)

Now, [Gradshteyn and Ryzhik 1979, 8.914, p.1026],

$$(1.10.5) \quad L_{r-2}(x) - L_r(x) = \text{const}(r)(1-x^2)L'_{r-1}(x)$$

and we thus have the following (which corresponds in a sense rather to Theorem 1.8.2 than Theorem 1.8.1).

**Theorem 1.10.1.** *Let  $r$  be odd,  $\mu = 1$  and, with  $e = u - u_h$ ,  $u_h$  the solution to (1.1.6), and  $e(x_i)$ ,  $e(x_{i+1/2})$  and  $e(x_{i+1}) = O(h^{r+1})$  on  $I_i$ . Then*

$$(1.10.6) \quad |e(\xi)| = O(h^{r+1})$$

where  $\xi$  is any root of  $L'_{i,r-1}$  on  $I_i$  (or, the given meshpoints and midpoints). In particular, this is so if the mesh is uniform in a symmetric  $C_1 h \ln 1/h$  neighborhood of  $I_i$  which is inside  $I$ , with  $C_1$  large enough.

Now,  $L_{r-1}$  has  $r-1$  roots on  $(-1, 1)$  and hence  $L'_{r-1}$  has  $r-2$  roots. One of those is the midpoint. Thus, the number of superconvergence points on  $\bar{I}_i$  for function values is now  $r$ , a full complement; this is in contrast to the situation for  $r$  even for function values. We also note that the points in Theorem 1.10.1 are exactly those occurring for  $\mu = 0$  (without mesh restrictions or staying away from the boundary), see Theorem 1.5.2.

Finally, differentiating (1.10.4), (1.10.5) and using Legendre's differential equation,  $((1 - x^2)L'_{r-1})' = \text{const.}L_{r-1}$ , we have:

**Theorem 1.10.2.** *Let  $r$  be odd,  $\mu = 1$  and  $e(x_i)$ ,  $e(x_{i+1/2})$  and  $e(x_{i+1}) = O(h^{r+1})$  on  $I_i$ . Then*

$$(1.10.7) \quad |e'(\eta)| = O(h^r)$$

*where  $\eta$  is any root of  $L_{i,r-1}$  on  $I_i$ . In particular, this is so if the mesh is uniform in a symmetric  $C_1 h \ln 1/h$  neighborhood of  $I_i$  which is inside  $I$ , with  $C_1$  large enough.*

Thus we have exactly the same superconvergence points for derivatives as for  $\mu = 0$ , see Theorem 1.5.1.

The following table summarizes our superconvergence results in Sections 1.2 and 1.5–1.10.

$\mu : r$	Function Values	First Derivative
$\mu = 0 : \text{Any } r$ Completely General Meshes.	a) $O(h^{2r-2})$ at Meshpoints. b) $O(h^{r+1})$ at Zeroes of $L'_{r-1}(x)$ . (For $r \geq 3$ , Total of $r$ Points, “Maximal #”.)	$O(h^r)$ at Zeroes of $L_{r-1}(x)$ . (Total of $r - 1$ Points, “Maximal #”.)
$\mu = 1 : r$ Even Meshes Uniform in a $C_1 h \ln 1/h$ neighborhood of the Point (and similarly, away from the boundary $\partial I$ ).	$O(h^{r+1})$ at the $r - 2$ Zeroes of $Q(x) := L_{r-2}(x) - \frac{L'_{r-2}(1)}{L'_r(1)} L_r(x).$ (Total of $r - 2$ Points, <b>NOT</b> “Maximal #”.)	$O(h^r)$ at Mesh- and Midpoints, and at the Additional $r - 4$ Zeroes of $Q'(x)$ . (Total of $r - 1$ Points, “Maximal #”.)
$\mu = 1 : r$ Odd Meshes as for $\mu = 1$ , $r$ Even, above.	$O(h^{r+1})$ at Mesh- and Midpoints, and at Zeroes of $L'_{r-1}(x)$ . (Cf. $\mu = 0$ , any $r (\geq 3)$ above.) ("Maximal #".)	$O(h^r)$ at Zeroes of $L_{r-1}(x)$ . (Cf. $\mu = 0$ , any $r$ above.) ("Maximal #".)
Smoothest Cubics $\mu = 2 : r = 4$ Meshes as immediately above.	$O(h^{r+1})$ at Two Points, same as for Hermite Cubics ( $\mu = 1$ , $r = 4$ ) above. <b>(NOT</b> “Maximal #”.)	$O(h^r)$ at Mesh- and Midpoints. ("Maximal #".)
$\mu \geq 1 : r$ Even General Meshes Symmetric About the Point in a $C_1 h \ln 1/h$ -neighborhood (and similarly away from the boundary $\partial I$ ).		$O(h^r)$ at the Point.
$\mu \geq 1 : r$ Odd General Meshes: Same as for $\mu \geq 1$ , $r$ Even, immediately above.	$O(h^{r+1})$ at the Point.	

### 1.11. First order difference quotients of $u_h$ as superconvergent approximations to $u'$ on locally uniform meshes.

This section is an exposé of some of the ideas of [Nitsche and Schatz 1974, Section 6]; that work was in turn inspired by [Thomée and Westergren 1968] in finite difference approximations. In the present situation we shall rely on the results of Section 1.3 and thus treat only the case of first derivatives. Similar results for higher order derivatives will be treated in the multidimensional situation; they would not follow from the techniques of this section for variable coefficients  $a_2$ ,  $a_1$  and  $a_0$ .

On the other hand, the present restricted technique, viz., utilizing that  $u'_h \simeq P(u')$  to the hilt, gives somewhat sharper results with respect to how close to the boundary of  $I$  the result is valid and the size of domains on which the mesh is assumed uniform. Let us also remark that if approximations to  $u$  and  $u'$  of a certain order are available, then in our present case, a same order approximation to  $u''$  obtains readily from the differential equation (1.1.1) itself.

We shall thus consider  $\tilde{u}_h \in \overset{\circ}{S}_h$  given by

$$(1.11.1) \quad ((u - \tilde{u}_h)', \chi') = 0, \text{ for } \chi \in \overset{\circ}{S}_h = \overset{\circ}{S}_h^{\mu, r}.$$

With  $P$  the  $L_2$ -projection into  $\$_h = \$_h^{\mu-1, r-1}$  and  $w = u'$ , we then have  $w_h = \tilde{u}'_h = P(w) \in \$_h$ , i.e.,

$$(1.11.2) \quad (w - w_h, \chi) = 0, \text{ for } \chi \in \$_h.$$

For  $\mathcal{J} \subseteq I$  an interval, let  $\$_h^{comp}(\mathcal{J})$  denote the functions in  $\$_h(\mathcal{J})$  which have compact support in  $\mathcal{J}$ . For technical reasons we shall need to operate with four intervals inside each other, all centered about the same point.  $\mathcal{J}_1$  is the basic interval on which we wish to prove our final estimate.  $\mathcal{J}_2 = \{x \in R : dist(x, \mathcal{J}_1) \leq Kh\}$  where  $K$  is to be given below, (1.11.5) (and, if necessary, slightly enlarged later).  $\mathcal{J}_3 = \{x \in R : dist(x, \mathcal{J}_2) \leq C_1 h \ln 1/h\}$  where  $C_1$  will be taken large enough. Finally,  $\mathcal{J}_4 = \{x \in R : dist(x, \mathcal{J}_3) \leq Kh\}$ . Note that  $\mathcal{J}_1 \subseteq \mathcal{J}_2 \subseteq \mathcal{J}_3 \subseteq \mathcal{J}_4$ . It is assumed that

$$(1.11.3) \quad \mathcal{J}_4 \subseteq I \text{ and that the mesh is uniform on } \mathcal{J}_4; \quad h_i \equiv h \text{ there.}$$




Figure 1.11.1

With  $T_h f(x) = f(x + h)$ , translation by  $h$ , we have

$$(1.11.4) \quad (T_h(w - w_h), \chi) = (w - w_h, T_{-h}\chi) = 0, \text{ for } \chi \in \$_h^{comp}(\mathcal{J}_3),$$

since then  $T_{-h}\chi \in \$_h$ .

Let now

$$(1.11.5) \quad \partial_h = \sum_{i=-K}^K c_i(h) T_{ih}$$

denote a difference quotient operator which is supposed to be an approximation to  $\partial/\partial x$ . Specifically we assume that

$$(1.11.6) \quad |\partial_h f(x)| \leq C \|f\|_{W_\infty^1(A_{Kh}(x))}$$

where  $A_{Kh}(x) = \{y : |x - y| \leq Kh\}$ , and also that

$$(1.11.7) \quad |(\partial_h f - f')(x)| \leq Ch^s \|f\|_{W_\infty^{s+1}(A_{Kh}(x))}, \text{ for } s = r - 1 \text{ and } r.$$

Now, as in (1.11.4),

$$(1.11.8) \quad (\partial_h(w - w_h), \chi) = 0, \text{ for } \chi \in \$_h^{comp}(\mathcal{J}_3).$$

Since  $\partial_h w_h \in \$_h(\mathcal{J}_3)$  we may utilize the following result on local behavior in ‘‘local  $L_2$ -projections’’, see [Wahlbin 1991, Theorem 3.3] or Theorem 3.2.4 below. In fact, we have already used it in proving Lemma 1.9.1, see (1.9.7).

**Lemma 1.11.1.** *There exist three constants  $C$ ,  $c_0$ , and  $c > 0$  such that the following holds. For any  $v$  and  $v_h \in \$_h(\mathcal{J})$  such that*

$$(1.11.9) \quad (v - v_h, \chi) = 0, \text{ for all } \chi \in \$_h^{comp}(\mathcal{J}),$$

*if  $d \geq c_0 h$  with  $A_d = A_d(x) = \{y : |x - y| \leq d\} \subseteq \mathcal{J}$ ,*

$$(1.11.10) \quad |(v - v_h)(x)| \leq C \min_{\chi \in \$_h(\mathcal{J})} \|v - \chi\|_{L_\infty(A_d)} + Cd^{-1/2} e^{-cd/h} \|v - v_h\|_{L_2(A_d)}.$$

We clearly have

$$(1.11.11) \quad d^{-1/2} \|v - v_h\|_{L_2(A_d)} \leq C \|v - v_h\|_{L_\infty(A_d)}.$$

We apply Lemma 1.11.1 to  $v = \partial_h w = \partial_h u'$  and  $v_h = \partial_h w_h = \partial_h P(w)$  for  $x \in \mathcal{J}_2$ , and we may then take  $d = C_1 h \ln 1/h$  with  $A_d(x) \subseteq \mathcal{J}_3$ . Since  $P$  is  $L_p$ -stable and hence  $W_p^1$ -stable on our quasi-uniform mesh, by (1.11.6) we find that

$$(1.11.12) \quad \begin{aligned} \|\partial_h(w - w_h)\|_{L_\infty(A_d)} &\leq \|\partial_h(w - w_h)\|_{L_\infty(\mathcal{J}_3)} \\ &\leq C \|w - w_h\|_{W_\infty^1(I)} \leq C \|u\|_{W_\infty^2(I)}. \end{aligned}$$

Then for  $x \in \mathcal{J}_2$ , using (1.11.10) and (1.11.11),

$$(1.11.13) \quad \begin{aligned} |(\partial_h w - \partial_h w_h)(x)| &\leq C \min_{\chi \in \$_h(\mathcal{J}_3)} \|\partial_h w - \chi\|_{L_\infty(\mathcal{J}_3)} + Ch^k \|u\|_{W_\infty^2(I)}, \\ &\text{for } x \in \mathcal{J}_2, \end{aligned}$$

for any  $k = k(C_1)$  provided  $C_1$  is large enough. Finally, using (1.11.7) with  $s = r - 1$  and standard approximation theory for a suitable  $\chi$ ,

$$(1.11.14) \quad \begin{aligned} \|\partial_h w - \chi\|_{L_\infty(\mathcal{J}_3)} &\leq \|\partial_h w - w'\|_{L_\infty(\mathcal{J}_3)} + \|w' - \chi\|_{L_\infty(\mathcal{J}_3)} \\ &\leq Ch^{r-1} \|w\|_{W_\infty^r(I)} \leq Ch^{r-1} \|u\|_{W_\infty^{r+1}(I)}. \end{aligned}$$

Thus

$$(1.11.15) \quad \|\partial_h(u' - \tilde{u}'_h)\|_{L_\infty(\mathcal{J}_2)} \leq Ch^{r-1} \|u\|_{W_\infty^{r+1}(I)}.$$

To convert this to a result about  $\partial_h(u - \tilde{u}_h)$ , we first consider the case  $r \geq 3$ . Let  $x \in \mathcal{J}_1$ . Following the technique of Section 1.4, we introduce  $B \in \$h^{\mu-2,r-2}$ , a nonnegative function with small compact support containing  $x$  and of length  $M(\mu, r)h$ . With  $\partial_h^* = \sum_{i=-K}^K c_i(h)T_{-ih}$  we have  $(\partial_h\chi, \psi) = (x, \partial_h^*\psi)$  for  $\psi$  with compact support in  $\mathcal{J}_3$ . After enlarging  $K$  (in the definition of  $\mathcal{J}_2$ ) if necessary, we have  $\partial_h^*B \in \$h^{comp}(\mathcal{J}_2)$ . Defining  $\varphi$  by  $-\varphi'' = \partial_h^*B$  on  $I$ ,  $\varphi(0) = \varphi(1) = 0$ , we see  $\varphi \in \overset{\circ}{S}_h(I)$  so that

$$(1.11.16) \quad \int_{supp B} \partial_h(u - \tilde{u}_h)B = (u - \tilde{u}_h, \partial_h^*B) = ((u - \tilde{u}_h)', \varphi') = 0.$$

Thus if we consider a point  $x \in \mathcal{J}_1$ ,  $\partial_h(u - \tilde{u}_h)$  vanishes at a point  $\xi = \xi(x)$  within a distance  $hM(\mu, r)$  of  $x$ . Again enlarging  $K$  in the definition of  $\mathcal{J}_2$  if necessary, we may assume that  $\xi(x) \in \mathcal{J}_2$ . Hence

$$(1.11.17) \quad \partial_h(u - \tilde{u}_h)(x) = \int_\xi^x (\partial_h(u - \tilde{u}_h))' = \int_\xi^x \partial_h(u' - \tilde{u}'_h)$$

so that by (1.11.15),

$$(1.11.18) \quad |\partial_h(u - \tilde{u}_h)(x)| \leq Ch^r \|u\|_{W_\infty^{r+1}(I)}, \text{ for } r \geq 3, x \in \mathcal{J}_1.$$

Using (1.11.7) with  $s = r$ , finally, for  $r \geq 3$ ,

$$(1.11.19) \quad \|u' - \partial_h \tilde{u}_h\|_{L_\infty(\mathcal{J}_1)} \leq Ch^r \|u\|_{W_\infty^{r+1}(I)}.$$

For the case  $r = 2$ , i.e., the piecewise linear case, we know in our present setting ( $a_2 \equiv 1, a_1 \equiv a_0 \equiv 0$ ) that  $(u - \tilde{u}_h)(x_i) = 0$  at meshpoints. Consequently, also  $\partial_h(u - \tilde{u}_h)(x_i) = 0$  (for  $x_i \in \mathcal{J}_1$ ). The result (1.11.19) follows as before.

Finally, we now combine (1.11.19), derived for the case  $a_2 \equiv 1, a_1 \equiv a_0 \equiv 0$ , with Theorem 1.3.1. By (1.11.6) and that theorem, with  $\theta$  the difference between our present simple projection and that based on the general form  $A(\cdot, \cdot)$ , we find that

$$(1.11.20) \quad \|\partial_h \theta\|_{L_\infty(\mathcal{J}_1)} \leq C \|\theta\|_{W_\infty^1(I)} \leq Ch^r.$$

Noting that  $C_1 h \ln 1/h >> Kh$  for  $h$  small we thus have upon using (1.11.20) in (1.11.19):

**Theorem 1.11.1.** *There exist constants  $C_1$  and  $C_2$  such that the following holds. Let  $\mathcal{J}_1$  be any interval in  $I$  such that  $\mathcal{J}_4 = \{x \in R : dist(x, \mathcal{J}_1) \leq C_1 h \ln 1/h\} \subseteq I$  and the mesh is uniform on  $\mathcal{J}_4$  with meshsize  $h$ . Let  $\partial_h$  be a difference operator, based on translations by  $h$ , approximating  $\frac{\partial}{\partial x}$  in the sense of (1.11.6) and (1.11.7). Let  $u_h$  be the solution to (1.1.6). Then*

$$(1.11.21) \quad \|u' - \partial_h u_h\|_{L_\infty(\mathcal{J}_1)} \leq C_2 h^r \|u\|_{W_\infty^{r+1}(I)}.$$

In other words, unless  $x \in \mathcal{J}_1$  happens to be a superconvergent point for derivatives,  $\partial_h u_h(x)$  is an asymptotically better approximation to  $u'(x)$  than  $u'_h(x)$ .

### 1.12. Two examples of superconvergence by “iteration”.

The general structure is as follows:  $Q$  is a quantity of interest,  $Q = Q(u)$ , which may be given by two different “expressions”,  $Q = Q_1(u)$  as well as  $Q = Q_2(u)$ . Using  $Q_{1,h} := Q_1(u_h)$  or  $Q_{2,h} := Q_2(u_h)$  as an approximation may give approximations of different order.

1.12.a. This example is due to [Wheeler, J. A. 1973], cf. also [Douglas, Dupont and Wheeler 1974a]. The quantity of interest is the “outflow” derivative  $u'(1)$ , say. Using  $Q_{1,h} = u'_h(1)$  as an approximation we have no reason to expect better rate of convergence than

$$(1.12.1) \quad |u'(1) - u'_h(1)| \leq Ch^{r-1}.$$

Now, multiplying the equation (1.1.1) by  $x$  and integrating by parts over  $I$  a few times, we have using that  $u(0) = u(1) = 0$ ,

$$(1.12.2) \quad u'(1) = Q_2(u) = \frac{1}{a_2(1)} \int_0^1 (-fx - ua'_2 + a_1u + a_0ux)dx.$$

Define then

$$(1.12.3) \quad Q_{2,h} := \frac{1}{a_2(1)} \int_0^1 (-fx - u_h a'_2 + a_1 u_h + a_0 u_h x)dx.$$

Leaving numerical quadrature out of the picture (easily handled for smooth data) we have with  $e = u - u_h$ ,

$$(1.12.4) \quad u'(1) - Q_{2,h} = \frac{1}{a_2(1)} \int_0^1 (-ea'_2 + a_1 e + a_0 ex)dx.$$

Thus,

$$(1.12.5) \quad |u'(1) - Q_{2,h}| \leq C\|e\|_{W_1^{-s}(I)} \equiv \sup_{\substack{\varphi \in W_\infty^s \\ \|\varphi\|_{W_\infty^{-s}(I)}=1}} (e, \varphi).$$

A standard duality argument then establishes that, for smooth data, taking  $s = r - 2$ ,

$$(1.12.6) \quad |u'(1) - Q_{2,h}| \leq Ch^{2r-2},$$

an asymptotically considerable improvement over (1.12.1).

1.12.b. This iteration method was first used and analyzed in connection with Fredholm integral equations of the second kind, cf. [Nyström 1930] (where the motivation for its use was somewhat different than here), [Sloan, Burn and Datyner 1975] and [Sloan 1976]. The analogue for two-point boundary value problems presented here is in [Volk 1986], where also numerical illustrations can be found.

First, if necessary by fiddling around in (1.1.1), let us assume that the highest order coefficient  $a_2 \equiv 1$ . then

$$(1.12.7) \quad -u'' = f + (a_1 u)' - a_0 u, \quad u(0) = u(1) = 0.$$

At issue is to approximate  $u(x)$  itself at any point; of course we could let  $Q_{1,h}(x) = u_h(x)$  do it. However, if somewhat laboriously, the operator on the left of (1.12.7),

$-d^2/dx^2$  with boundary conditions, can be inverted by integration: I.e., if  $v$  such that

$$(1.12.8) \quad -v'' = g \text{ on } I, \quad v(0) = v(1) = 0$$

is desired, then

$$(1.12.9) \quad v(x) = \int_0^x \int_0^y g(z) dz dy - x \int_0^1 \int_0^y g(z) dz dy.$$

(Again, numerical quadrature will not be taken into account here.) Thus we replace  $u$  by  $u_h$  on the right hand side of (1.12.7) and consequently define

$$(1.12.10) \quad u_*(x) = Q_{2,h}(x) := v(x)$$

where  $v(x)$  is as in (1.12.9) with

$$(1.12.11) \quad g = f + (a_1 u_h)' - a_0 u_h.$$

Then

$$(1.12.12) \quad \|u - u_*\|_{L_\infty(I)} = \sup_{\|v\|_{L_1}=1} (u - u_*, v).$$

Letting  $-\psi'' = v$  on  $I$ ,  $\psi(0) = \psi(1) = 0$ , and letting  $e = u - u_h$ ,

$$\begin{aligned} (1.12.13) \quad (u - u_*, v) &= ((u - u_*)', \psi') = -((u - u_*)'', \psi) \\ &= ((a_1 e)', \psi) - (a_0 e, \psi) = -(a_1 e, \psi') - (a_0 e, \psi) \\ &\leq C(a_1) \|e\|_{W_\infty^{-1}(I)} + C(a_0) \|e\|_{W_\infty^{-2}(I)}. \end{aligned}$$

Thus,

$$(1.12.14) \quad \|u - u_*\|_{L_\infty(I)} \leq \begin{cases} Ch^{r+1}, & \text{if } r \geq 3, \\ Ch^{r+2}, & \text{if } a_1 \equiv 0 \text{ and } r \geq 4. \end{cases}$$

(The reader may wish to compare this to Remark 1.3.1.)

### 1.13. A graphical illustration of superconvergence.

We consider the equation

$$(1.13.1) \quad -u'' + u = f \text{ in } I = (0, 1),$$

with the boundary conditions

$$(1.13.2) \quad u'(0) = u'(1) = 0.$$

We shall give one example from [Dunlap, Li, Taylor, Warren and Wahlbin 1995].

Let us first describe the general approach. Working with a known function and having in mind  $\mu \geq 1$ , let the meshes be uniform of size  $h$  and  $\tilde{I} \subset\subset I$  an interior subinterval of  $I$ , cf. Section 1.7. Introduce testpoints  $t_{h,i,j} \in I_i$ ,  $j = 0, \dots, J$ , by taking  $J + 1$  uniformly spaced points on each mesh interval  $\tilde{I}_i$ . The error for the  $j^{\text{th}}$  set of testpoints with respect to  $\tilde{I}$  (and the given function and problem) is then

$$(1.13.3) \quad \tilde{e}_{h,j} = \max_{t_{h,i,j} \in \tilde{I}} |(u - u_h)(t_{h,i,j})|.$$

The observed rate of convergence  $\tilde{r}_{h,j}$  at the  $j^{\text{th}}$  set of testpoints is now naturally defined as

$$(1.13.4) \quad \tilde{r}_{h,j} = \log_2 \left( \frac{\tilde{e}_{2h,j}}{\tilde{e}_{h,j}} \right).$$

(A similar computation may of course be done for the error in derivatives.)

For the graph below we have  $u(x) = \sin(5x) - \frac{(5\cos 5 - 5)}{2}x^2 - 5x$ ,  $r = 4$ ,  $\mu = 1$ ,  $\tilde{I} = [0.2, 0.8]$ ,  $h = 1/50$  and  $J = 1000$ . The graph gives  $\tilde{r}_{h,j}$  versus  $j$ , cf. (1.8.18).




Figure 1.13.1

## Chapter 2. Remarks about some of the tools used in Chapter 1.

A seasoned Finite Elementer need not bother with this chapter.

We shall give sketchy generalities about certain standard tools of the trade such as inverse estimates, duality estimates and superapproximation. We shall not, however, give much detail concerning basic approximation theory; doing so would swell this chapter to unmanageable proportions. As for the local properties of  $L_2$ -projections, so heavily used in Chapter 1, we shall derive them in Chapter 3. In fact, they will be given there in any number of space dimensions.

### 2.1. Inverse estimates.

Let  $\mathcal{J}$  be an interval  $(a, b)$  of length  $|\mathcal{J}| = (b - a)$  and consider  $\Pi_s(\mathcal{J})$ , the polynomials of degree  $\leq s$  on  $\mathcal{J}$ . We have the following:

**Theorem 2.1.1.** *There exists a constant  $C(s)$  such that for any  $p$  and  $q$  with  $1 \leq p \leq \infty$ ,  $1 \leq q \leq \infty$ ,*

$$(2.1.1) \quad \|\chi\|_{L_p(\mathcal{J})} \leq C(s)|\mathcal{J}|^{-(\frac{1}{q} - \frac{1}{p})}\|\chi\|_{L_q(\mathcal{J})}, \text{ for } \chi \in \Pi_s(\mathcal{J}).$$

Proof: We treat the case  $p < \infty$ . By Hölder's inequality we have

$$(2.1.2) \quad \|\chi\|_{L_p(\mathcal{J})} \leq |\mathcal{J}|^{1/p}\|\chi\|_{L_\infty(\mathcal{J})}.$$

With the transformation  $\tilde{x} = (x - a)/|\mathcal{J}|$  from  $\mathcal{J}$  to  $I = [0, 1]$ , and with  $\tilde{\chi}(\tilde{x}) = \chi(x)$ ,

$$(2.1.3) \quad \|\chi(x)\|_{L_\infty(\mathcal{J})} = \|\tilde{\chi}(\tilde{x})\|_{L_\infty(I)}.$$

Now,  $\tilde{\chi} \in \Pi_s(I)$  and hence by the fact that any norms on a finite dimensional space are equivalent,

$$(2.1.4) \quad \|\tilde{\chi}\|_{L_\infty(I)} \leq C(s)\|\tilde{\chi}\|_{L_1(I)}.$$

By transformation of variables and Hölder's inequality with  $1/q + 1/q' = 1$ ,

$$(2.1.5) \quad \begin{aligned} \|\tilde{\chi}\|_{L_1(I)} &= \int_I |\chi(x)| \frac{dx}{|\mathcal{J}|} \leq \frac{1}{|\mathcal{J}|} \|\chi\|_{L_q(\mathcal{J})} |\mathcal{J}|^{1/q'} \\ &= |\mathcal{J}|^{-1/q} \|\chi\|_{L_q(\mathcal{J})}. \end{aligned}$$

The desired result follows from (2.1.2)–(2.1.5).  $\square$

Let next  $\tau$  be a quasi-uniform partition of  $I = [0, 1]$  with  $h = \max_i h_i$ , and  $h \leq C_{QU} \min_i h_i$ , cf. (1.1.7). Let  $S_h = S_h^{\mu, s+1}(I)$ .

**Theorem 2.1.2.** *There exists a constant  $C(s)$  such that for  $p > q$ ,*

$$(2.1.6) \quad \|\chi\|_{L_p(I)} \leq C(s) \left( \frac{h}{C_{QU}} \right)^{-(\frac{1}{q} - \frac{1}{p})} \|\chi\|_{L_q(I)}, \text{ for } \chi \in S_h.$$

Proof: We have, for  $q < p < \infty$ , using Theorem 2.1.1,

$$\begin{aligned}
(2.1.7) \quad \|\chi\|_{L_p(I)}^p &= \sum_{i=0}^{N-1} \|\chi\|_{L_p(I_i)}^p \\
&\leq C(s)^p \sum_{i=0}^{N-1} |h_i|^{(1-p/q)} \|\chi\|_{L_q(I_i)}^p \\
&\leq C(s)^p (\min_i h_i)^{1-p/q} \sum_{i=0}^{N-1} \|\chi\|_{L_q(I_i)}^p \\
&\leq C(s)^p \left(\frac{h}{C_{QU}}\right)^{1-p/q} \sum_{i=0}^{N-1} \|\chi\|_{L_q(I_i)}^p.
\end{aligned}$$

For an array  $z = \{z_i\}_{i=1}^{N-1}$  we have  $\|z\|_{\ell_p} \leq \|z\|_{\ell_q}$ , for  $p > q$ . Thus, from (2.1.7),

$$(2.1.8) \quad \|\chi\|_{L_p(I)}^p \leq C(s)^p \left(\frac{h}{C_{QU}}\right)^{1-p/q} \left(\sum_{i=0}^{N-1} \|\chi\|_{L_q(I_i)}^q\right)^{p/q}.$$

Taking  $p^{\text{th}}$  roots concludes the argument. The case  $p = \infty$  is similar.  $\square$

The estimate (2.1.6) is a typical case of an “inverse estimate”. It is a property that does not hold for “general” functions, but only for some restricted class. Inverse estimates do not only come into play when going from a high  $L_p$ -norm to a lower one, but also in bounding higher derivatives by lower ones.

**Theorem 2.1.3.** *There exists a constant  $C(s)$  such that for any  $1 \leq p \leq \infty$  and any integers  $k > \ell$ ,*

$$(2.1.9) \quad \|\chi^{(k)}\|_{L_p(\mathcal{J})} \leq C(s)|\mathcal{J}|^{-(k-\ell)} \|\chi^{(\ell)}\|_{L_p(\mathcal{J})}, \text{ for } \chi \in \Pi_s(\mathcal{J}).$$

Proof: If  $k > s$  there is nothing to prove. Since  $\chi^{(k)} = \frac{\delta^{k-\ell}}{\partial x^{k-\ell}}(\chi^{(\ell)})$  it is enough to consider  $\ell = 0$ . Now, by Hölder’s inequality,

$$(2.1.10) \quad \|\chi^{(k)}\|_{L_p(\mathcal{J})} \leq |\mathcal{J}|^{1/p} \|\chi^{(k)}\|_{L_\infty(\mathcal{J})}.$$

Letting  $\tilde{x} = \frac{(x-a)}{|\mathcal{J}|}$ ,  $\tilde{\chi}(\tilde{x}) = \chi(x)$ , we have  $\frac{d^k}{dx^k} \chi(x) = \frac{1}{|\mathcal{J}|^k} \frac{d^k}{d\tilde{x}^k} \tilde{\chi}(\tilde{x})$ , so that

$$(2.1.11) \quad \|\chi^{(k)}\|_{L_p(\mathcal{J})} \leq |\mathcal{J}|^{1/p-k} \left\| \frac{d^k}{d\tilde{x}^k} \tilde{\chi} \right\|_{L_\infty(I)}.$$

and again appealing to the equivalence of any two norms on a finite dimensional space,

$$\begin{aligned}
(2.1.12) \quad \|\chi^{(k)}\|_{L_p(\mathcal{J})} &\leq C(s)|\mathcal{J}|^{1/p-k} C_1(s) \|\tilde{\chi}\|_{L_\infty(I)} \\
&= C(s) C_1(s) |\mathcal{J}|^{1/p-k} \|\chi\|_{L_\infty(\mathcal{J})} \\
&\leq C(s) C_1(s) |\mathcal{J}|^{-k} \|\chi\|_{L_p(\mathcal{J})}
\end{aligned}$$

where we used Theorem 2.1.1 in the last step. This proves the theorem.  $\square$

Corresponding to Theorem 2.1.2 we now have in the quasi-uniform situation:

**Theorem 2.1.4.** *There exists a constant  $C(s)$  s.t. for  $k > \ell$ ,  $1 \leq p \leq \infty$ ,*

$$(2.1.13) \quad \|\chi^{(k)}\|_{L_p(I)} \leq C(s) \left( \frac{h}{C_{QU}} \right)^{-(k-\ell)} \|\chi^{(\ell)}\|_{L_p(I)}, \text{ for } \chi \in S_h^{\mu,s+1}(I).$$

As a final comment to this section, the reader will have noticed that care has been taken to avoid constants depending on  $p$  and  $q$  in Theorems 2.1.1 and 2.1.2. It is important to, at least, keep uniform control of such constants. As we shall see in the multi-dimensional situation, one may want to apply the results with  $p$  depending on  $h$ .

## 2.2. On approximation theory, and duality.

We shall merely give some scattered remarks. Of course, of central importance (for us) is the approximation of smooth functions. Thus, with  $\$_h = \$_h^{\mu,r}(I)$  we want to ascertain that, e.g.,

$$(2.2.1) \quad \min_{\chi \in \$_h} \|v - \chi\|_{L_\infty(I)} \leq Ch^r \|v\|_{W_\infty^r(I)}$$

where  $C$  is a constant which does not depend on  $h$  or  $v$ . The case of  $\mu = 0$  (or,  $\mu = -1$ , of course) is simple: On each  $I_i$ , we choose the two endpoints  $x_{i,1} = x_i$  and  $x_{i,r} = x_{i+1}$  and then  $r-2$  additional points  $x_{i,j}$ ,  $j = 2, \dots, r-1$  forming an equispaced partition of  $\bar{I}_i$ . Letting  $Int(v)$  be the Lagrange interpolant of polynomial degree  $r-1$  to  $v$  at these points, almost any elementary text in Numerical Analysis tells us that there is a constant  $C = C(r)$  such that

$$(2.2.2) \quad \|v - Int(v)\|_{L_\infty(I_i)} \leq Ch_i^r \|v^{(r)}\|_{L_\infty(I_i)}.$$

Since the mesh points  $x_i$  are interpolation points,  $Int(v)$  will belong to  $\$_h^{\mu,r}(i)$  when  $\mu = 0$  (or  $\mu = -1$ ). Also, if  $\$_h$  is replaced by  $\overset{\circ}{\$}_h$  there is no additional problem if  $v(0) = v(1) = 0$ ; the piecewise Lagrange interpolant will be in  $\overset{\circ}{\$}_h$ .

A similar classical procedure which takes into account only what happens on  $I_i$  for the approximation occurs in the Hermite case, (or, rather “a” Hermite case), when  $r = 2(\mu+1)$ . Then we find  $Int(v) \in \Pi_{r-1}(I_i)$  to match  $v^{(j)}(x_i)$  and  $v^{(j)}(x_{i+1})$ , for  $j = 0, \dots, \mu$ . The combined result will lie in  $\$_h^{\mu,r}$ .

If  $2(\mu+1) < r$ , throw in a few equispaced interpolation points in between . . .

However, if  $2(\mu+1) > r$  we run out of free parameters in this game. The construction now becomes more elaborate, but the result (2.2.1) is still valid, also if  $\$_h$  is replaced by  $\overset{\circ}{\$}_h$ , i.e., boundary conditions are taken into account. We refer the reader to, say, [Powell 1981, Chapters 19–20]. The interpolant used can be “global” in the sense that the value used at one point influences the interpolant all over  $I$  (but with exponentially diminishing influence away from the point) or it can be “local”, i.e., the value of the interpolant at a point depends only on the values of the function over a few mesh-intervals around the point. For such local interpolants, “quasi-interpolants”, see in particular [deBoor and Fix 1973] and also [Powell 1981, Section 20.3].

However, there is a need not only for high-order approximation as in (2.2.1) but also for low-order approximation. Let us motivate this need via a duality argument. Let us then say we know, for  $e = u - u_h$  where  $u_h$  is the solution to (1.1.6), that

$$(2.2.3) \quad \|e\|_{W_\infty^1(I)} \leq Ch^{r-1}.$$

Now we wish to show that, for  $r \geq 3$ ,

$$(2.2.4) \quad \|e\|_{W_\infty^{-1}(I)} \leq Ch^{r+1}$$

where

$$(2.2.5) \quad \|v\|_{W_\infty^{-1}(I)} = \sup_{\|w\|_{W_1^1(I)}=1} (v, w)$$

(cf. Remark 1.3.2 and Section 1.12 where (2.2.4) = (1.1.12) was used.)

A typical duality argument now proceeds as follows:

$$(2.2.6) \quad \|e\|_{W_\infty^{-1}(I)} = \sup_{\|w\|_{W_1^1(I)}=1} (e, w).$$

For each such  $w$ , let  $\psi \in \overset{\circ}{H}{}^1(I)$  be the solution of the adjoint problem to (1.1.1), in weak form,

$$(2.2.7) \quad A(\chi, \psi) = (\chi, w), \text{ for any } \chi \in \overset{\circ}{H}{}^1(I).$$

By Fredholm's alternative, if (1.1.1) has unique solutions, so does (2.2.7). Furthermore, in this one-dimensional case, it is not hard to show the a priori estimate that

$$(2.2.8) \quad \|\psi\|_{W_1^3(I)} \leq C\|w\|_{W_1^1(I)} = C.$$

Then

$$(2.2.9) \quad (e, w) = A(e, \psi) = A(e, \psi - \chi), \text{ for any } \chi \in \overset{\circ}{S}_h,$$

and so by Hölder's inequality, and (2.2.3),

$$(2.2.10) \quad |(e, w)| \leq Ch^{r-1} \min_{\chi \in \overset{\circ}{S}_h} \|\psi - \chi\|_{W_1^1(I)}.$$

Thus, to complete the program of deriving (2.2.4) it would suffice that, cf. (2.2.8),

$$(2.2.11) \quad \min_{\chi \in \overset{\circ}{S}_h} \|\psi - \chi\|_{W_1^1(I)} \leq Ch^2 \|\psi\|_{W_1^3(I)},$$

where  $\psi(0) = \psi(1) = 0$ . We note that the Hermite interpolant given above would not work in general: it requires point-values at mesh-points for certain derivatives of  $\psi$  and such would in general not be available if  $\psi$  is merely in  $W_1^3(I)$ .

A solution lies in first smoothing  $\psi$  before applying the "interpolant". This is easy if one disregards the boundary conditions  $\chi(0) = \chi(1) = 0$ . Let  $K$  be a "smoothing kernel", a  $C_0^\infty$  function with certain properties to be given below. Assume that  $\psi$  is extended over the boundary  $\partial I$  in some suitable way, stable in  $W_p^m$  for  $m$  as large as needed, cf. [Nečas 1967, p. 75] or [Friedman 1969, p. 10] where the rather well-known and simple Lions–Nikolskii–Peetre extension operator is given. Setting then

$$(2.2.12) \quad Sm_h(\psi)(x) = \frac{1}{h} \int \psi(y) K\left(\frac{x-y}{h}\right) dy$$

it is not hard to see that a “high–order” interpolant applied to  $Sm_h(\psi)(x)$  does the trick, provided “moment conditions” such as

$$(2.2.13) \quad \int K(x-y)y^\ell dy = x^\ell, \quad \ell = 0, 1, \dots, L$$

for  $L$  sufficiently high are fulfilled. (I.e., the smoothing operator reproduces polynomials up to a certain order.) We refer to [Hilbert 1973] or [Strang 1973] for details.

Finally, if one must respect boundary conditions (as in (2.2.11) as stated!), one uses instead of  $Int(Sm_h(\psi))$  as an approximation a perturbed one,  $\widetilde{Int}(Sm_h(\psi))$ , which equals  $\psi(0)$  at  $x = 0$  (or  $\psi(1)$  at  $x = 1$ ). In the case of (2.2.11), e.g., one checks that (for  $Int$  being the local quasi–interpolant, cf. [deBoor and Fix 1973], e.g.)

$$(2.2.14) \quad |\widetilde{Int}(v)(x) - Int(v)(x)| \leq \begin{cases} C|Int(v)(0)|, & \text{for } 0 \leq x \leq Ch, \\ 0, & \text{for } x > Ch. \end{cases}$$

Since, if  $\psi(0) = 0$ ,

$$(2.2.15) \quad |Int(Sm_h(\psi))(0)| \leq Ch^2\|\psi\|_{W_\infty^2(I)} \leq Ch^2\|\psi\|_{W_1^3(I)},$$

we obtain upon using inverse estimates (there is no problem if  $\mu = 0$  or  $-1$ ), that

$$\begin{aligned} (2.2.16) \quad & \|\psi - \widetilde{Int}(Sm_h(\psi))\|_{W_1^1(I)} \\ & \leq \|\psi - Int(Sm_h(\psi))\|_{W_1^1(I)} \\ & \quad + \|(Int - \widetilde{Int})(Sm_h(\psi))\|_{W_1^1(0, Ch)} \\ & \leq Ch^2\|\psi\|_{W_1^3(I)} + \frac{C}{h}\|(Int - \widetilde{Int})(Sm_h(\psi))\|_{L_1(0, Ch)} \\ & \leq Ch^2\|\psi\|_{W_1^3(I)} + C\|(Int - \widetilde{Int})(Sm_h(\psi))\|_{L_\infty(0, Ch)} \\ & \leq Ch^2\|\psi\|_{W_1^3(I)} + C|Int(Sm_h(\psi))(0)| \\ & \leq Ch^2\|\psi\|_{W_1^3(I)} + Ch^2\|\psi\|_{W_\infty^2(I)} \quad (\text{since } \psi(0) = 0) \\ & \leq Ch^2\|\psi\|_{W_1^3(I)} \quad (\text{Sobolev in one dimension}). \end{aligned}$$

Applying a cut–off argument at  $x = 1/2$  and similar techniques at  $x = 1$ , this then shows (2.2.11) in general even when respecting the boundary conditions  $\chi(0) = \chi(1) = 0$ .

### 2.3. Superapproximation.

Recall that this played a major role in Section 1.3. We shall assume that we have at our disposal a high order local approximation operator  $I_h$  into  $\$h = S_h^{\mu, s}(I)$  (or, into  $\overset{\circ}{S}_h$  if boundary conditions are to be respected). This operator is a linear operator such that

$$(2.3.1) \quad I_h\chi \equiv \chi \text{ for } \chi \in \$h.$$

It is a high-order local approximation in the sense that there exist two constants  $C_1$  and  $C_2$  such that

$$(2.3.2) \quad \|v - I_h(v)\|_{W_\infty^k(I_i)} \leq C_1 h^{s-k} \|v^{(s)}\|_{L_\infty(I'_i)},$$

for  $k = 0, 1, \dots, s-1$ , any  $v \in W_\infty^s(I'_i)$ ,

where

$$(2.3.3) \quad I'_i = \{x \in I : \text{dist}(x, I_i) \leq C_2 h\} \cap I.$$

Furthermore,

$$(2.3.4) \quad I_h(v) \text{ has support in } (\text{Supp}(v) + C_2 h) \cap I,$$

where for any set  $B$  and real  $d > 0$ ,  $B + d = \{x \in R : \text{dist}(x, B) \leq d\}$ .

We shall assume that, cf. (1.1.7),

the meshes are quasi-uniform.

The above approximation operator  $I_h$  is in general the quasi-interpolant of [de Boor and Fix 1973].

Remark 2.3.1. The case of  $\mu = 0$  (or,  $\mu = -1$ ) is much simpler and is left to the reader. The operator  $I_h$  can then be taken completely local on each  $I_i$ , namely, as the Lagrange interpolation operator on equidistanted points including  $x_i$  and  $x_{i+1}$ . The quasi-uniform condition is then not necessary, as the reader should check. (Actually, the quasi-uniformity is not necessary in any case in which the approximation operator is completely local on each  $I_i$ ; this happens e.g. in Hermite cases.)  $\square$

We shall use piecewise norms with respect to the partitions  $\tau$ ,

$$(2.3.5) \quad \|v\|_{W_p^{k,h}(I)} = \left( \sum_{i=0}^{N-1} \|v\|_{W_p^k(I_i)}^p \right)^{1/p},$$

with the usual modification if  $p = \infty$ . We have the following as the major (very complete!) result on superapproximation in one space dimension.

**Theorem 2.3.1.** *There exist constants  $C_3$  and  $C_4$  depending on  $C_1$ ,  $C_2$  above and on  $s$  and  $C_{QU}$  (in (1.1.7)) such that the following holds.*

*Let  $\omega$  be a smooth function such that*

$$(2.3.6) \quad \|\omega\|_{W_\infty^k(I)} \leq \Lambda d^{-k}, \text{ for } k = 0, 1, \dots, s,$$

*where  $d \geq h$ . Then for any  $\chi \in S_h^{\mu,s}(I)$  there exists  $\psi \in \$_h^{\mu,s}(I)$  such that for any  $k \leq s-1$  and any  $1 \leq p \leq \infty$ ,*

$$(2.3.7) \quad \|\omega\chi - \psi\|_{W_p^{k,h}(I)} \leq C_3 \Lambda \left( \frac{h}{d} \right) \|\chi\|_{W_p^{k,h}((\text{Supp}(\omega) + C_4 h) \cap I)}.$$

Furthermore,

$$(2.3.8) \quad \text{Supp}(\psi) \subseteq (\text{Supp}(\omega) + C_4 h) \cap I.$$

Proof: The natural choice is  $\psi = I_h(\omega\chi)$ . We are now faced with the problem that, in general,  $\omega\chi \notin W_\infty^s(I'_i)$  so that the error estimate (2.3.2) is not immediately

applicable. We employ the following trick taken from [Douglas, Dupont and Wahlbin 1975]: With each interior mesh point  $x_i$  we associate the functions

$$(2.3.9) \quad v_{ik}(x) = (x - x_i)_+^k, \quad k = \mu + 1, \dots, s - 1.$$

Starting from the left we then determine a linear combination  $w$  of the  $v_{ik}$  so that  $\omega\chi + w \in C^{s-1}(I)$ ; i.e., we have soaked up the jumps in certain derivatives. Then  $\omega\chi + w \in W_\infty^s(I)$ . Now  $w \in \$_h(I)$  so that  $I_h w = w$  and so,

$$(2.3.10) \quad I_h(\omega\chi) = I_h(\omega\chi + w) - w.$$

We now use the approximation hypothesis (2.3.2) so that

$$\begin{aligned} (2.3.11) \quad \|\omega\chi - I_h(\omega\chi)\|_{W_\infty^k(I_i)} &= \|\omega\chi + w - I_h(\omega\chi + w)\|_{W_\infty^k(I_i)} \\ &\leq C_1 h^{s-k} \|(\omega\chi + w)^{(s)}\|_{L_\infty(I'_i)} \\ &= C_1 h^{s-k} \|(\omega\chi)^{(s)}\|_{L_\infty^h(I'_i)} \\ &= C_1 h^{s-k} \|(\omega\chi)^{(s)}\|_{L_\infty^h(I'_i \cap \text{Supp}(\omega))}, \end{aligned}$$

where the notation  $L_\infty^h$  means that the  $s^{\text{th}}$  derivative is taken piecewise over the partition. By Leibniz' formula, and (2.3.6), since  $\chi^{(s)} \equiv 0$  (in the piecewise sense),

$$\begin{aligned} (2.3.12) \quad \|(\omega\chi)^{(s)}\|_{L_\infty^h(I'_i \cap \text{Supp}(\omega))} \\ &\leq C(s) \sum_{\ell=0}^{s-1} \Lambda d^{-(s-\ell)} \|\chi\|_{W_\infty^{\ell,h}(I'_i \cap \text{Supp}(\omega))}. \end{aligned}$$

Hence, using the inverse estimates of Theorem 2.1.3,

$$\begin{aligned} (2.3.13) \quad \|(\omega\chi)^{(s)}\|_{L_\infty^h(I'_i \cap \text{Supp}(\omega))} \\ &\leq C \left( \sum_{\ell=0}^{s-1} \Lambda d^{-(s-\ell)} h^{-(\ell-k)} \right) \|\chi\|_{W_\infty^{k,h}(I'_i \cap \text{Supp}(\omega) + h)} \\ &\leq C \Lambda d^{-s} h^k \left( \frac{d}{h} \right)^{s-1} \|\chi\|_{W_\infty^{k,h}(I'_i \cap \text{Supp}(\omega) + h)}. \end{aligned}$$

Hence, from (2.3.11),

$$\begin{aligned} (2.3.14) \quad \|\omega\chi - I_h(\omega\chi)\|_{W_\infty^k(I_i)} \\ &\leq C \Lambda d^{-s} h^s \left( \frac{d}{h} \right)^{s-1} \|\chi\|_{W_\infty^{k,h}(I'_i \cap \text{Supp}(\omega) + k)} \\ &\leq C \Lambda \left( \frac{h}{d} \right) \|\chi\|_{W_\infty^{k,h}(I'_i \cap \text{Supp}(\Omega) + h)}. \end{aligned}$$

By use of Hölder's inequality and the inverse estimates of Theorem 2.1.1, we then obtain

$$\begin{aligned}
 (2.3.15) \quad & \|\omega\chi - I_h(\omega\chi)\|_{W_p^k(I_i)} \leq h^{1/p} \|\omega\chi - I_h(\omega\chi)\|_{W_\infty^k(I_i)} \\
 & \leq Ch^{1/p}\Lambda\left(\frac{h}{d}\right) \|\chi\|_{W_\infty^{k,h}(I'_i \cap \text{Supp}(\omega) + h)} \\
 & \leq C\Lambda\left(\frac{h}{d}\right) \|\chi\|_{W_p^{k,h}(I'_i \cap \text{Supp}(\omega) + 2h)}.
 \end{aligned}$$

Raising to the  $p^{\text{th}}$  power and summing, the estimate (2.3.7) obtains. The statement about the support of  $\psi$  is easily verified.  $\square$

#### 2.4. A typical combination of inverse estimates and approximation theory used in Chapter 1.

Consider the operator  $P$ , the  $L_2$ -projection into  $\$_h$ . At various points in Chapter 1 we used the following: Assume that the meshes are quasi-uniform, and that the operator  $P$  is stable in  $L_p(I)$ ,  $1 \leq p \leq \infty$ , i.e.,

$$(2.4.1) \quad \|Pv\|_{L_p(I)} \leq C_0 \|v\|_{L_p(I)}$$

where  $C_0$  does not depend on  $h$  or  $v$ . (This will be shown for  $\mu \geq 0$  and quasi-uniform meshes in Chapter 3; of course, the case  $\mu = -1$  is trivial.)

We then asserted, and used, that

$$(2.4.2) \quad \|Pv\|_{W_p^1(I)} \leq C \|v\|_{W_p^1(I)}.$$

(Again, this is trivial if  $\mu = -1$ .) In general for quasi-uniform meshes, one would proceed as follows to deduce (2.4.2) from (2.4.1).

Now,

$$(2.4.3) \quad \|v - Pv\|_{W_p^1(I)} \leq \|v - \chi\|_{W_p^1(I)} + \|\chi - Pv\|_{W_p^1(I)}.$$

Now since the mesh is quasi-uniform, by inverse estimates we have, since  $\chi - Pv \in \$_h$ ,

$$\begin{aligned}
 (2.4.4) \quad & \|\chi - Pv\|_{W_p^1(I)} \leq C_{inv}h^{-1} \|\chi - Pv\|_{L_p(I)} \\
 & \leq C_{inv}h^{-1} \|\chi - v\|_{L_p(I)} + C_{inv}h^{-1} \|v - Pv\|_{L_p(I)}.
 \end{aligned}$$

But since  $\chi \in \$_h$ ,  $P\chi = \chi$ , and we have  $v - Pv = (v - \chi) - P(v - \chi)$  so that

$$(2.4.5) \quad \|v - Pv\|_{L_p(I)} \leq \|(v - \chi) - P(v - \chi)\|_{L_p(I)} \leq (1 + C_0) \|v - \chi\|_{L_p(I)}.$$

From the above, for any  $\chi \in \$_h$ ,

$$(2.4.6) \quad \|v - Pv\|_{W_p^1(I)} \leq \|v - \chi\|_{W_p^1(I)} + C_{inv}(1 + C_0)h^{-1} \|v - \chi\|_{L_p(I)}.$$

The argument to show (2.4.2) is now capped off by “low order” approximation theory: we may find  $\chi \in \$_h$  such that

$$(2.4.7) \quad \|v - \chi\|_{W_p^1(I)} + h^{-1} \|v - \chi\|_{L_p(I)} \leq C_{app} \|v\|_{W_p^1(I)}.$$

### Chapter 3. Local and global properties of $L_2$ -projections.

We shall prove the relevant estimates used in Chapter 1 in any number of space dimensions. We then need appropriate inverse estimates and superapproximation results which we proceed to state and discuss in the first section.

Let us remark that the  $L_2$ -projection occurs in its own right in Numerical Analysis and that hence the results of this chapter, and those of Chapter 4, are of independent interest.

#### 3.1. Assumptions.

We shall treat in detail only the case when no imposed boundary conditions are present (this is the case used in Chapter 1 in the one-dimensional situation). Let thus  $T_h$ ,  $0 < h < 1/2$ , be a sequence of partitions of domains  $\mathcal{D}_h \subset\subset R^n$  into disjoint open “finite elements”  $\tau_i^h$ ,  $\mathcal{D}_h = \bigcup_{i=0}^{N(h)} \bar{\tau}_i^h$ .

The parameter  $h$  may be thought of as  $\max_i(\text{diam}(\tau_i^h))$ , and, to fix thoughts,  $\mathcal{D}_h$  as approximations to a fixed domain  $\mathcal{D}$ .

Let now  $S_h = S_h(\mathcal{D}_h)$  be a family of finite dimensional function spaces on  $\mathcal{D}_h$ . A very typical case is that of  $\tau_i^h$  being  $n$ -simplices in  $R^n$  and  $S_h$  a space of continuous functions on  $\mathcal{D}_h$  such that for  $\chi \in S_h$ ,  $\chi$  restricted to each  $\tau_i^h$  is a polynomial of total degree  $\leq r - 1$ . These are the Lagrange finite element spaces. We refer to [Ciarlet 1991] for generalities about finite element spaces.

We shall use some special notation relative to the domains  $\mathcal{D}_h$ . If  $\Omega_0 \subseteq \Omega_1 \subseteq \mathcal{D}_h$ , then

$$(3.1.1) \quad \partial_<(\Omega_0, \Omega_1) = \text{dist}(\partial\Omega_0 \setminus \partial\mathcal{D}_h, \partial\Omega_1 \setminus \partial\mathcal{D}_h).$$




Figure 3.1.1.

We let  $\mathcal{C}_\infty^<(\bar{\Omega}) = \{v \in \mathcal{C}_\infty(\bar{\Omega}) : \partial_<(\text{supp}(v), \Omega) > 0\}$  and  $S_h^<(\Omega) = \{v \in S_h : \partial_<(\text{supp}(v), \Omega) > 0\}$ .

We first state our superapproximation assumption.

Assumption 3.1.1. (Superapproximation) There exist constants  $c$  and  $C$  and a number  $L$ , independent of  $h$ , such that the following holds: Let  $\Omega_0 \subseteq \Omega_1 \subseteq \mathcal{D}_h$  with  $d = \partial_<(\Omega_0, \Omega_1) > ch$ . Let further  $\omega \in \mathcal{C}_\infty^<(\bar{\Omega}_0)$  with

$$(3.1.2) \quad \|\omega\|_{W_\infty^\ell(\Omega_0)} \leq \Lambda d^{-\ell}, \quad \ell = 0, \dots, L.$$

Then for any  $\chi \in S_h$  there exists  $\psi \in S_h^<(\Omega_1)$  such that

$$(3.1.3) \quad \|\omega\chi - \psi\|_{L_2(\Omega_1)} \leq C\Lambda\left(\frac{h}{d}\right)\|\chi\|_{L_2(\Omega_1)}. \quad \square$$

We shall also need an inverse estimate:

**Assumption 3.1.2.** (Inverse estimate) There exists a constant  $C$  independent of  $h$  such that for any element  $\tau_i^h$ , for any  $1 \leq p \leq \infty$ ,  $1 \leq q \leq \infty$ ,

$$(3.1.4) \quad \|\chi\|_{L_p(\tau_i^h)} \leq Ch^{-n(\frac{1}{q} - \frac{1}{p})}\|\chi\|_{L_q(\tau_i^h)}, \text{ for } \chi \in S_h. \quad \square$$

We note that for  $q < p$ , the estimate (3.1.4) is valid with  $L_p(\tau_i^h)$  and  $L_q(\tau_i^h)$  replaced by  $L_p(\mathcal{D}_h)$ , respectively  $L_q(\mathcal{D}_h)$ , cf. the proof of Theorem 2.1.2.

An inverse estimate in general is proven essentially as in Section 2.1 by mapping to a standard (reference) element of unit size and appealing to finite-dimensionality. In the form we have stated (3.1.4), one has to avoid degeneracies in meshes such as triangles getting thin. We refer to [Ciarlet 1991, Theorem 17.2, p. 135] for more details.

Our superapproximation Assumption 3.1.1 can often be proven by following the lines of Section 2.3. E.g., in the Lagrange case described above, there is an interpolation operator  $I_h$  locally determined on each  $\tau_i^h$  which preserves polynomials of total degree  $\leq r - 1$  such that

$$(3.1.5) \quad \|v - I_h v\|_{L_\infty(\tau_i^h)} \leq Ch^r |v|_{W_\infty^r(\tau_i^h)},$$

in case certain geometry constraints are not violated, see [Ciarlet 1991, Theorem 16.2, p. 128]. Here  $|\cdot|_{W_\infty^r}$  denotes the  $r^{\text{th}}$  semi-norm involving only  $r^{\text{th}}$  derivatives. Applying this to  $v = \omega\chi$  and noting that  $r^{\text{th}}$  derivatives of  $\chi$  vanish, and using inverse estimates, one easily obtains our Assumption 3.1.1.

The systematic derivation of error estimates such as (3.1.5) using polynomial invariance of  $I_h$  goes back to [Bramble and Hilbert 1970], the “Bramble–Hilbert Lemma”. (In this respect, cf. also (2.2.13).) For the rather laborious derivation of error estimates before the Bramble–Hilbert Lemma, the reader may ponder the following quote from [Prenter 1975, p. 127] (who did not employ the Bramble–Hilbert Lemma): “Waning sadism forbids us to go further.”

A key idea in the Bramble–Hilbert Lemma, namely that  $\inf_{\pi \in \Pi_{s-1}} \|v - \pi\|_{W_p^s(\Omega)}$  is equivalent to  $|v|_{W_p^s(\Omega)}$ , can be traced at least as far back as [Deny and Lions 1953–4, Théorème 5.1]. Of course, there it had nothing to do with finite element approximation theory!

In other cases, Assumption 3.1.1 is less trivial. E.g., if (in some piece over  $\mathcal{D}_h \subset \subset R^2$ ) the mesh is rectangular and  $S_h$  then consists of continuous bilinear functions  $a_0 + a_1x + a_2y + a_3xy$  on the rectangle  $\tau_i^h$ , then the counterpart of (3.1.5) would not be enough since  $\frac{\partial^2}{\partial x \partial y}\chi \neq 0$  in general, for  $\chi \in S_h$ . What is needed then is the so-called “sharp” form of the Bramble–Hilbert Lemma [Bramble and Hilbert 1971], cf. also [Nitsche and Schatz 1974, Appendix] and [Bramble, Nitsche and Schatz 1975, Appendix]. In our present case of bilinears this means that

$$(3.1.6) \quad \|v - I_h(v)\|_{L_\infty(\tau_i^h)} \leq Ch^2 |v|'_{W_\infty^2(\tau_i^h)}$$

where  $|v|'_{W_\infty^2(\tau_i^h)} = |\frac{\partial^2 v}{\partial x^2}|_{L_\infty(\tau_i^h)} + |\frac{\partial^2 v}{\partial y^2}|_{L_\infty(\tau_i^h)}$ , i.e., *only those derivatives which annihilate  $\chi \in S_h$  are involved.* Then superapproximation is easily proved assuming suitable inverse properties.

### 3.2. Estimates for $L_2$ -projections.

Assuming that the superapproximation Assumption 3.1.1 holds, we have first the following:

**Lemma 3.2.1.** *There exist positive constants  $c_0$ ,  $c_1$  and  $C$  such that the following holds. Let  $\Omega_0 \subseteq \Omega_1 \subseteq \mathcal{D}_h$  with  $d = \partial_<(\Omega_0, \Omega_1) \geq c_0 h$ . Assume further that  $v_h \in S_h$  is such that*

$$(3.2.1) \quad (v_h, \chi) = 0, \text{ for all } \chi \in S_h^<(\Omega_1).$$

Then

$$(3.2.2) \quad \|v_h\|_{L_2(\Omega_0)} \leq C e^{-c_1 d/h} \|v_h\|_{L_2(\Omega_1)}.$$

Of course, if  $S_h$  consists of discontinuous elements this is trivial!

Proof: Let  $\delta = ch$  with  $c$  as in Assumption 3.1.1, i.e., the “separation” between domains required there. Let further  $\Omega_0 \subseteq \Omega_\delta \subseteq \Omega_{2\delta}$  with  $\delta = \partial_<(\Omega_0, \Omega_\delta) = \partial_<(\Omega_\delta, \Omega_\delta)$ . Let  $\omega \in C_\infty^<(\Omega_\delta)$  be nonnegative with

$$(3.2.3) \quad \omega \equiv 1 \text{ on } \Omega_0,$$

and

$$(3.2.4) \quad \|\omega\|_{W_\infty^\ell} \leq \Lambda \delta^{-\ell}, \quad \ell = 0, \dots, L.$$

(It is well known how to construct such a function.) Then

$$(3.2.5) \quad \|v_h\|_{L_2(\Omega_0)}^2 \leq (v_h, \omega v_h).$$

For any  $\chi \in S_h^<(\Omega_1)$ , we have

$$(3.2.6) \quad (v_h, \omega v_h) = (v_h, \omega v_h - \chi).$$

Assuming that  $\Omega_{2\delta} \subseteq \Omega_1$ , we choose  $\chi$  according to superapproximation, in particular with  $\chi \in S_h^<(\Omega_{2\delta})$  and obtain from (3.1.3),

$$(3.2.7) \quad \begin{aligned} \|v_h\|_{L_2(\Omega_0)}^2 &\leq C \|v_h\|_{L_2(\Omega_{2\delta})} \Lambda \left( \frac{h}{\delta} \right) \|v_h\|_{L_2(\Omega_{2\delta})} \\ &\leq \left( C \Lambda \frac{h}{\delta} \right) \|v_h\|_{L_2(\Omega_{2\delta})}^2. \end{aligned}$$

Repeat the argument with domains  $\Omega_{2\delta} \subseteq \dots \subseteq \Omega_{2N\delta}$  as long as they remain inside  $\Omega_1$ , i.e., since  $\delta = ch$ , for  $N \geq \text{const. } d/h$ . Increasing  $\delta$  in Assumption 3.1.1 if necessary, we may assume  $C\Lambda h/\delta < 1$  and set  $C\Lambda h/\delta = e^{-c'_1}$ ; then

$$(3.2.8) \quad \|v_h\|_{L_2(\Omega_0)}^2 \leq e^{-\text{const. } c'_1 d/h} \|v_h\|_{L_2(\Omega_1)}^2.$$

The desired result follows.  $\square$

We next investigate the discrete  $\delta$ -function centered at  $x_0 \in \mathcal{D}_h$ , i.e., the function  $\delta = \delta_h^{x_0} \in S_h$  such that

$$(3.2.9) \quad (\chi, \delta) = \chi(x_0), \text{ for all } \chi \in S_h.$$

(Here we assume that  $S_h \subseteq C^0(\mathcal{D}_h)$ ; the case of totally discontinuous elements is much simpler and left to the reader.) From the Fundamental Theorem of Linear Algebra, it is easily seen that there exists a unique such discrete  $\delta$ -function centered at  $x_0$ .

Assuming now both Assumption 3.1.1 and 3.1.2 from Section 3.1, we have the following:

**Theorem 3.2.2.** *There exist positive constants  $C$  and  $c_2$  such that*

$$(3.2.10) \quad |\delta_h^{x_0}(x)| \leq Ch^{-n}e^{-c_2|x-x_0|/h}.$$

Let us remark that this estimate was heavily used in Section 1.5.

Proof: From Lemma 3.2.1 we have for  $\Omega_0$  not containing  $x_0$ , and  $d = \text{dist}(\Omega_0, x_0) \geq ch$  with  $c$  sufficiently large,

$$(3.2.11) \quad \|\delta_h\|_{L_2(\Omega_0)} \leq Ce^{-c_1 d/h} \|\delta_h\|_{L_2(\mathcal{D}_h)}.$$

This is true by default if  $\Omega_0$  contains  $x_0$  (then  $d = 0$ ). Thus, by use of the inverse Assumption 3.1.2,

$$(3.2.12) \quad |\delta_h(x_0)| \leq Ch^{-n/2}e^{-c_1 d/h} \|\delta_h\|_{L_2(\mathcal{D}_h)}.$$

From (3.2.9) and again using inverse assumptions,

$$(3.2.13) \quad \|\delta_h\|_{L_2(\mathcal{D}_h)}^2 = \delta_h(x_0) \leq Ch^{-n/2} \|\delta_h\|_{L_2(\mathcal{D}_h)}.$$

The desired result follows.  $\square$

From Theorem 3.2.2 we obtain the  $L_p$ -stability of the  $L_2$ -projection  $P$  into  $S_h$ .

**Theorem 3.2.3.** *There exists a constant  $C$  such that for any  $1 \leq p \leq \infty$ ,*

$$(3.2.14) \quad \|Pv\|_{L_p(\mathcal{D}_h)} \leq C\|v\|_{L_p(\mathcal{D}_h)}, \text{ for all } v \in L_p(\mathcal{D}_h).$$

Proof: We have

$$(3.2.15) \quad Pv(x_0) = (Pv, \delta_h^{x_0}) = (v, \delta_h^{x_0}).$$

Thus, from (3.2.10),

$$(3.2.16) \quad |Pv(x_0)| \leq C\|v\|_{L_\infty(\mathcal{D}_h)} \int_{\mathcal{D}_h} h^{-n}e^{-c_1|x-x_0|/h} dx \leq C\|v\|_{L_\infty(\mathcal{D}_h)},$$

and so

$$(3.2.17) \quad \|Pv\|_{L_\infty(\mathcal{D}_h)} \leq C\|v\|_{L_\infty(\mathcal{D}_h)}.$$

By duality (note that  $Pv$  is perfectly well defined for  $v \in L_1(\mathcal{D}_h)$ , for “typical” finite element spaces),

$$\begin{aligned} (3.2.18) \quad \|Pv\|_{L_1(\mathcal{D}_h)} &= \sup_{\|w\|_{L_\infty(\mathcal{D}_h)}=1} (Pv, w) \\ &= \sup(v, Pw) \leq C\|v\|_{L_1(\mathcal{D}_h)}\|w\|_{L_\infty(\mathcal{D}_h)} \\ &\leq C\|v\|_{L_1(\mathcal{D}_h)}. \end{aligned}$$

The Riesz–Thorin interpolation theorem (cf. [Bennett and Sharpley 1988, p. 192]) now gives the stability of the  $L_2$ -projection with respect to any  $L_p$ -norm.  $\square$

Finally we derive a general result used in Section 1.9 (cf. Lemma 1.9.1) and in Section 1.11 (Lemma 1.11.1).

**Theorem 3.2.4.** Under the Assumptions 3.1.1 and 3.1.2, there exist positive constants  $c_3$ ,  $c_4$  and  $C$  such that the following holds. Assume that  $v$  is a continuous function and  $v_h \in S_h$  is such that

$$(3.2.19) \quad (v - v_h, \chi) = 0, \text{ for all } \chi \in S_h^<(\Omega_d(x_0))$$

where  $\Omega_d(x_0) = \{x : |x - x_0| \leq d\} \cap I$  with  $d \geq c_3 h$ . Then

$$(3.2.20) \quad \begin{aligned} |(v - v_h)(x_0)| &\leq C \min_{\chi \in S_h} \|v - \chi\|_{L_\infty(\Omega_d(x_0))} \\ &\quad + Cd^{-n/2} e^{-c_4 d/h} \|v - v_h\|_{L_2(\Omega_d(x_0))}. \end{aligned}$$

Proof: Let  $\tilde{P}$  denote the  $L_2$ -projection into  $S_h^<(\Omega_d(x_0))$ . We may assume that  $\Omega_d(x_0)$  is a union of elements. By Theorem 3.2.3, this is stable in any  $L_p$ -space. (It is easy to see that the size of  $D_h$  doesn't enter in any essential way in the proof of Theorem 3.2.3, cf. in particular (3.2.16).) Then

$$(3.2.21) \quad (v - v_h)(x_0) = (v - \tilde{P}v)(x_0) + (\tilde{P}v - v_h)(x_0).$$

Here, by the stability of  $\tilde{P}$ ,

$$(3.2.22) \quad |(v - \tilde{P}v)(x_0)| \leq C \|v\|_{L_\infty(\Omega_d(x_0))}.$$

Since

$$(3.2.23) \quad (\tilde{P}v - v_h, \chi) = (v - v_h, \chi) = 0, \text{ for } \chi \in S_h^<(\Omega_d(x_0)),$$

and  $\tilde{P}v - v_h \in S_h(\Omega_d(x_0))$ , our inverse estimate and Lemma 3.2.1 lead to, for  $x_0 \in \tau_{i_0}^h$ ,

$$(3.2.24) \quad \begin{aligned} |(\tilde{P}v - v_h)(x_0)| &\leq Ch^{-n/2} \|\tilde{P}v - v_h\|_{L_2(\tau_{i_0}^h)} \\ &\leq Ch^{-n/2} e^{-c_1 d/h} \|\tilde{P}v - v_h\|_{L_2(\Omega_d(x_0))}. \end{aligned}$$

Now,

$$(3.2.25) \quad \begin{aligned} \|\tilde{P}v - v_h\|_{L_2(\Omega_d(x_0))} &\leq \|\tilde{P}v - v\|_{L_2(\Omega_d(x_0))} + \|v - v_h\|_{L_2(\Omega_d(x_0))} \\ &\leq \|v\|_{L_2(\Omega_d(x_0))} + \|v - v_h\|_{L_2(\Omega_d(x_0))} \\ &\leq Cd^{n/2} \|v\|_{L_\infty(\Omega_d(x_0))} + \|v - v_h\|_{L_2(\Omega_d(x_0))}. \end{aligned}$$

Thus, using also elementary calculus (with a change of constants),

$$(3.2.26) \quad \begin{aligned} |(\tilde{P}v - v_h)(x_0)| &\leq Ch^{-n/2} d^{n/2} e^{-c_1 d/h} \|v\|_{L_\infty(\Omega_d(x_0))} \\ &\quad + Ch^{-n/2} e^{-c_1 d/h} \|v - v_h\|_{L_2(\Omega_d(x_0))} \\ &\leq C \|v\|_{L_\infty(\Omega_d(x_0))} + Cd^{-n/2} e^{-c_4 d/h} \|v - v_h\|_{L_2(\Omega_d(x_0))}. \end{aligned}$$

Combining (3.2.26) and (3.2.22) into (3.2.21),

$$(3.2.27) \quad |(v - v_h)(x_0)| \leq C \|v\|_{L_\infty(\Omega_d(x_0))} + Cd^{-n/2} e^{-c_4 d/h} \|v - v_h\|_{L_2(\Omega_d(x_0))}.$$

Writing now  $v - v_h = (v - \chi) + (\chi - v_h)$  for any  $\chi \in S_h$ , we obtain the theorem.  $\square$

With respect to the local results for the  $L_2$ -projection in several space dimensions given here, original work can be found in [Descloux 1972], [Nitsche and Schatz 1972]

and [Douglas, Dupont and Wahlbin 1975b]. Our treatment above has followed [Nitsche and Schatz 1972] with some ideas from [Schatz and Wahlbin 1983] thrown in to show exponential decay.

In our development above we have kept the assumption of global quasi-uniformity of meshes, as implicit in Section 3.1, for simplicity. It is apparent from the proofs that there is room for extensions of the results to less regular meshes. For an investigation of the technique of [Descloux 1972] in this regard, see [Crouzeix and Thomée 1985].

(There was at one time an on-going battle to show general  $L_p$ -stability of the  $L_2$ -projection in one space dimension under no (or “minimal”) mesh-conditions. The interested reader will find references to some work in [deBoor 1981].)

## Chapter 4. Introduction to several space dimensions: some results about superconvergence in $L_2$ -projections.

From Chapter 3 we now have recourse to rather sharp results concerning the  $L_2$ -projection. In this brief chapter we shall give examples of superconvergence results for the  $L_2$ -projection. The chapter is intended as an introductory one to multi-dimensional superconvergence results for solutions to elliptic boundary value problems: the technicalities there become more formidable. However, as already remarked, the  $L_2$ -projection is of independent interest in Numerical Analysis.

In our first section we shall investigate the existence of local superconvergence points for function values in general.

### 4.1. Negative norm estimates and existence of general superconvergence points for function values.

This section is somewhat analogous to Section 1.4. If the finite element space  $S_h$  allows us to construct a nonnegative function  $B \in S_h$ , then

$$(4.1.1) \quad (v - Pv, B) = 0$$

so that  $v - Pv$  vanishes at some point in the support of  $B$ . (If  $S_h$  consists of discontinuous functions, then  $B \equiv 1$  on an element  $\tau_i^h$ ,  $B \equiv 0$  outside, may be taken.) Very often, this may be accomplished with  $\text{Supp}(B)$ , and thus a zero of  $v - Pv$ , in a ball of radius  $\leq Ch$ .

We shall next look at negative norm estimates for the  $L_2$ -projection.

$$(4.1.2) \quad \|v - Pv\|_{W_\infty^{-r}(\mathcal{D}_h)} = \sup_{\|w\|_{W_1^r(\mathcal{D}_h)}=1} (v - Pv, w).$$

For any  $\chi \in S_h$ , we have upon using various approximation assumptions (which we leave to the reader to make explicit)

$$\begin{aligned} (4.1.3) \quad (v - Pv, w) &= (v - Pv, w - \chi) \\ &\leq \|v - Pv\|_{L_\infty(\mathcal{D}_h)} \|w - \chi\|_{L_1(\mathcal{D}_h)} \\ &\leq Ch^{2r} \|v\|_{W_\infty^r(\mathcal{D}_h)} \|w\|_{W_1^r(\mathcal{D}_h)} \end{aligned}$$

Thus, for  $v$  smooth,

$$(4.1.4) \quad \|v - Pv\|_{W_\infty^{-r}(\mathcal{D}_h)} \leq Ch^{2r}.$$

We shall now show that this negative norm estimate leads to a rather “dense” occurrence of superconvergence points.

Now, if say  $B_d$  is an open ball of radius  $d$  in  $R^n$ , then we may construct  $w$  such that  $w > 0$  in  $B_d$ ,  $\text{supp } w = B_d$ ,

$$(4.1.5) \quad |w|_{W_\infty^\ell(B_d)} \leq Cd^{-\ell}, \quad \ell = 0, \dots, r$$

and, with  $c > 0$ ,

$$(4.1.6) \quad \|w\|_{L_1(B_d)} \geq cd^n.$$

Thus, since by (4.1.5)  $\|w\|_{W_1^r(B_d)} \leq \|w\|_{W_1^r(B_d)} \leq Cd^{n-r}$ ,

$$(4.1.7) \quad |(v - Pv, w)| \leq Ch^{2r} d^{n-r}.$$

Assuming now that  $v - Pv$  does not vanish on  $B_d$  we have since  $w > 0$  (taking  $v - Pv > 0$ , say),

$$(4.1.8) \quad \begin{aligned} \min_{x \in B_d} (v - Pv)(x) &\leq \int_{B_d} (v - Pv)(y)w(y)dy/\|w\|_{L_1(B_d)} \\ &\leq Ch^{2r}d^{-r} = Ch^r \left(\frac{h}{d}\right)^r. \end{aligned}$$

Thus, in any ball of radius  $d = h^{1-\sigma/r}$ , there is a superconvergent point of order  $\sigma$  (at least). (Here we have taken  $h^r$  to be the general best order of approximation in  $L_p$ , the so-called “optimal order”.) Of course, the above analysis gives no clue as to how to locate such superconvergence points a priori. They do, again, give us a hunting license.

#### 4.2. Superconvergence in $L_2$ -projections on $n$ -dimensional tensor product spaces.

Our second theme will be to show that any superconvergence result we know for one-dimensional  $L_2$ -projections will “automatically” carry over to  $n$ -dimensional  $L_2$ -projections on finite element spaces which are locally tensor-products.

We shall give the argument in detail for  $n = 2$ .

Let  $\mathcal{D}_h$  be plane domains and  $\Omega_0 = \Omega_0(h) \subseteq \Omega_1 = \Omega_1(h)$  two rectangles in  $\mathcal{D}_h$  such that in  $\Omega_1$ , the finite element space  $S_h(\Omega_1)$  is given as linear combinations of  $\chi_1(x)\chi_2(y)$  where  $\chi_i$ ,  $i = 1, 2$ , belong to one-dimensional finite element spaces (with partitions matching the boundaries of  $\Omega_1$ ). In “obvious” notation we may then write

$$(4.2.1) \quad S_h(\Omega_1) = S_h^x(I_x) \otimes S_h^y(I_y).$$

Let now  $v$  be a given function and  $v_h$  a function in  $S_h(\Omega_1)$  such that

$$(4.2.2) \quad (v - v_h, \chi) = 0, \text{ for } \chi \in S_h^<(\Omega_1).$$

Let  $P_x$  denote the one-dimensional  $L_2$ -projection into  $S_h^x(I_x)$  and similarly  $P_y$  for  $S_h^y(I_y)$ . Writing  $P_x \otimes P_y = (P_x \otimes I)(I \otimes P_y)$  it is easy to see that, in fact,  $P_x \otimes P_y = P$ , the  $L_2$ -projection into  $S_h(\Omega_1)$ . Thus,

$$(4.2.3) \quad (v_h - Pv, \chi) = 0, \text{ for } \chi \in S_h^<(\Omega_1)$$

so that assuming that

$$(4.2.4) \quad \partial_<(\Omega_0, \Omega_1) \geq C_1 h \ln 1/h,$$

and also that

$$(4.2.5) \quad \|v - v_h\|_{L_1(\mathcal{D}_h)} \leq C,$$

we have from Theorem 3.2.4 (making the implicit assumptions that it is applicable!), that

$$(4.2.6) \quad \|v_h - Pv\|_{L_\infty(\Omega_0)} \leq Ch^k$$

for any  $k$  provided  $C_1 = C_1(k)$  in (4.2.4) is large enough. To estimate  $v - v_h$  it now remains to look at  $v - P_x \otimes P_y v = (v - (P_x \otimes I)v) + (P_x \otimes I)(v - I \otimes P_y v)$ . We obtain:

**Theorem 4.2.1.** *With assumptions as given above, if  $\bar{x} = \bar{x}(h)$  is a superconvergence point for function values for  $P_x$  and  $\bar{y} = \bar{y}(h)$  a superconvergence point for function values for  $P_y$ , then, if  $(\bar{x}, \bar{y}) \in \Omega_0$ , it is a superconvergent point for  $v - v_h$ .*

The reader will have no trouble in extending this result to  $n$ -dimensional tensor products.

Also, assuming inverse estimates for derivatives, the result is true for superconvergence points in derivatives. (It is easy to “descend” in the results of Chapter 1 to give superconvergence results for derivatives in one-dimensional  $L_2$ -projections. We leave this to the interested reader.) Likewise, it is not hard to consider superconvergence for derivatives of suitable difference quotients on translation invariant meshes, cf. Sections 1.11 and Chapter 8.

### 4.3. Superconvergence by symmetry in $L_2$ -projections.

The results in this section are due to [Schatz, Sloan and Wahlbin 1994].

We continue to assume the results of Chapter 3. Assume now further that the finite element spaces are symmetric about points  $x_0 = x_0(h)$  in a  $C_1 h \ln 1/h$  neighborhood with respect to the anti-podal map (cf. (1.6.9)): If  $N_1 = \{x : |x - x_0| \leq C_1 h \ln 1/h\}$  we assume that  $N_1 \subseteq \mathcal{D}_h$  and that with  $\bar{\chi}(x) := \chi(x_0 - (x - x_0))$ ,

$$(4.3.1) \quad \bar{\chi}(x) \in S_h(N_1) \text{ for any } \chi \in S_h(N_1).$$

We shall give some examples taken from [Schatz, Sloan and Wahlbin 1994] at the end of this section. Note that  $x_0$  may vary with  $h$ . We also assume the approximation property that

$$(4.3.2) \quad \min_{\chi \in S_h} \|v - \chi\|_{L_\infty(N_1)} \leq Ch^r |v|_{W_\infty^r(N_2)}$$

where  $N_2 = N_1 + Ch$  and  $|\cdot|_{W_\infty^r(N_2)}$  denotes the seminorm involving only  $r^{\text{th}}$  order derivatives ( $r$  is again the “optimal” order in  $L_p$ ). This dependence on the seminorm only is crucial for our arguments. We further assume that

$$(4.3.3) \quad \|v - v_h\|_{L_2(\mathcal{D}_h)} \leq C.$$

With  $C_1$  sufficiently large we then have:

**Theorem 4.3.1.** *Assume that  $S_h$  consists of continuous functions. Assume further that  $r$  is odd and that the assumptions described above hold. Then for  $v$  and  $v_h \in S_h$  such that*

$$(4.3.4) \quad (v - v_h, \chi) = 0, \text{ for } \chi \in S_h^<(N_1),$$

*we have for  $v$  sufficiently smooth and  $x_0$  a symmetry point,*

$$(4.3.5) \quad |(v - v_h)(x_0)| \leq Ch^{r+1} \ln 1/h.$$

Proof: The key is to consider the even part of  $v$  and  $v_h$ ,

$$(4.3.6) \quad v_{\text{even}}(x) := (v(x) + \bar{v}(x))/2$$

and similarly for  $v_h$ . Then  $v_{h,\text{even}} \in S_h(N_1)$  by (4.3.1). It also follows from (4.3.1) by a simple transformation of variables argument that

$$(4.3.7) \quad (v_{\text{even}} - v_{h,\text{even}}, \chi) = 0, \text{ for } \chi \in S_h^<(N_1).$$

Since  $v(x_0) - v_h(x_0) = (v_{even} - v_{h,even})(x_0)$ , Theorem 3.2.4 with (4.3.3) tells us that

$$(4.3.8) \quad |(v - v_h)(x_0)| \leq \min_{\chi \in S_h} \|v_{even} - \chi\|_{L_\infty(N_1)} + Ch^{r+1}$$

provided  $C_1$  is large enough.

Using now (4.3.2) and the fact that  $r$  is odd,

$$(4.3.9) \quad \begin{aligned} \min_{\chi \in S_h} \|v_{even} - \chi\|_{L_\infty(N_1)} &\leq Ch^r |v_{even}|_{W_\infty^r(N_2)} \\ &\leq Ch^{r+1} \ln 1/h |v|_{W_\infty^{r+1}(N_2)}. \end{aligned}$$

This proves the theorem.  $\square$

For  $r$  even, one similarly obtains  $O(h^r \ln 1/h)$  superconvergent estimates for derivatives at symmetry points  $x_0$ . The key is now to consider the odd part,  $v_{odd} - v_{h,odd}$ . The details are easy for quasi-uniform meshes. (In Chapter 7 we shall show that for  $C^0$  elements, if a symmetry point is such that the derivative is not continuous there, then averaging works.)

Finally, we shall elucidate locally symmetric meshes, following [Schatz, Sloan and Wahlbin 1994]. Here, of course, one assumes that the piecewise polynomial spaces or, more general isoparametric ones, cf. [Ciarlet 1991, Chapter VI], are “the same” on each element. If one takes a mesh locally symmetric about a point  $x_0$  on the line, then extends that mesh to the upper half-plane in any manner whatsoever, and then reflects that mesh via the antipodal map about  $x_0$  to create a mesh in the plane, one obtains a mesh locally symmetric about  $x_0$ . ( $\bullet$  denotes the symmetry point(s).)




Figure 4.3.1.

Another example, not constructed by the above method, is as follows:




Figure 4.3.2.

Perhaps more traditional examples are as follows:  
Rectangular grids:




Figure 4.3.3.

A triangular grid of the criss-cross type:




Figure 4.3.4.

If the mesh is locally composed of congruent axes-parallel parallelepipeds in  $R^3$ , each parallelepiped has 27 symmetry points:




Figure 4.3.5.

## Chapter 5. Second order elliptic boundary value problems in any number of space dimensions: preliminary considerations on local and global estimates and presentation of the main technical tools for showing superconvergence.

### 5.1. Introduction.

Let  $\mathcal{D}$  be a bounded domain in  $R^n$ . The basic boundary value problem to be approximated is the following: Find  $u$  such that

$$(5.1.1) \quad -\sum_{i,j=1}^n \frac{\partial}{\partial x_j} \left( a_{ij}(x) \frac{\partial u}{\partial x_i} \right) - \sum_{i=1}^n \frac{\partial}{\partial x_i} (a_i(x)u) + a(x)u = f \text{ in } \mathcal{D},$$

some boundary condition on  $\partial\mathcal{D}$ .

We shall assume that the coefficients  $a_{ij}$ ,  $a_i$  and  $a$  are sufficiently smooth (on  $\mathcal{D}$ , or the relevant domain under consideration) and that the  $a_{ij}$  satisfy the uniform ellipticity condition

$$(5.1.2) \quad \sum_{i,j=1}^n a_{ij}(x) \xi_i \xi_j \geq c_{ell} |\xi|^2, \text{ for } \xi \in R^n,$$

with  $c_{ell} > 0$  independent of  $x$  (for  $x \in \mathcal{D}$ , or  $x$  in the domain under consideration). The formal bilinear form corresponding to the elliptic operator in (5.1.1) is

$$(5.1.3) \quad A(u, w) = \int \left( \sum_{i,j=1}^n a_{ij} \frac{\partial v}{\partial x_i} \frac{\partial w}{\partial x_j} + \sum_{i=1}^n a_i v \frac{\partial w}{\partial x_i} + avw \right) dx.$$

In examples later on, we shall be specific about the boundary condition on  $\partial\mathcal{D}$ .

Let  $S_h = S_h(\mathcal{D}) \subseteq W_\infty^1(\mathcal{D})$ ,  $0 < h < 1/2$ , be a one-parameter family of “finite-element” spaces (the “ $h$ -method”). For  $\Omega \subseteq \mathcal{D}$  we let  $S_h$  denote the restrictions of functions in  $S_h$  to  $\Omega$  and we let  $S_h^{comp}(\Omega)$  denote the set of those functions in  $S^h(\mathcal{D})$  with compact support in the interior of  $\Omega$ . Various assumptions concerning  $S_h$  will be introduced as we go along.

In our general treatment we shall only consider domains which are compactly contained in  $\mathcal{D}$ . In the main technical development we then do not have to consider the actual boundary conditions on  $\mathcal{D}$ ; of course, in various examples, we shall be specific.

Our *general starting point* is then that we have a function  $u$  and a function  $u_h \in S_h$  such that for domains  $\Omega \subset\subset \mathcal{D}$  (*which may depend on  $h$* ),

$$(5.1.4) \quad A(u - u_h, \chi) = F(\chi), \text{ for } \chi \in S_h^{comp}(\Omega).$$

Here  $F(\chi)$  is a bounded linear functional on  $\overset{\circ}{W}_1^1(\Omega)$ .

If  $u_h$  is a “usual” finite element approximation to the solution of (5.1.1), then most often  $F \equiv 0$  (if we disregard numerical integration, as we shall unless explicitly stated to the contrary). As we shall see, however, it will be important to allow a general right hand side in (5.1.4).

In general, we shall assume that there is an integer  $r \geq 2$  such that for smooth functions  $v$  on  $\mathcal{D}$ ,

$$(5.1.5) \quad \min_{\chi \in S_h} \|v - \chi\|_{L_\infty(\Omega)} \leq C(u)h^r.$$

If  $\Omega \subset\subset \mathcal{D}$  with enough separation between  $\Omega$  and  $\mathcal{D}$ , this does not take into consideration essential boundary conditions (such as Dirichlet boundary conditions). We shall call this integer  $r$  the *optimal order*, assuming that no higher order of approximation is possible for “general”  $u$ , cf. (1.1.14). Similarly, we shall assume that

$$(5.1.6) \quad \min_{\chi \in S_h} \|v - \chi\|_{W_\infty^1(\Omega)} \leq C(u)h^{r-1}$$

and no better in general. We refer to [Ciarlet 1991] for basics.

It is now rather obvious what we shall call *superconvergence of order  $\sigma > 0$*  cf. (1.1.15) and (1.1.16). Again, we shall also consider postprocessing.

The main technical tools for proving superconvergence will be presented in Section 5.5. The intervening sections are devoted to elucidating these tools.

## 5.2. Existence of superconvergence points in general: an example (also an example of a multi-dimensional duality argument).

Let  $\mathcal{D}$  be a bounded domain in  $R^n$  with sufficiently smooth boundary. Consider the Neumann problem of finding  $u$  such that

$$(5.2.1) \quad \begin{cases} -\Delta u + u = f & \text{in } \mathcal{D}, \\ \frac{\partial u}{\partial n} = 0 & \text{on } \partial\mathcal{D}. \end{cases}$$

The weak formulation of this problem is to find  $u \in H^1(\mathcal{D})$  such that

$$(5.2.2) \quad A(u, \chi) \equiv \int_{\mathcal{D}} (\nabla u \cdot \nabla \chi + u \chi) dx = (f, \chi) \equiv \int_{\mathcal{D}} f \chi, \text{ for all } \chi \in H^1(\mathcal{D}).$$

(Note that the “natural” boundary conditions  $\frac{\partial u}{\partial n} = 0$  on  $\partial\mathcal{D}$  are not seen in the weak formulation, cf. e.g., [Nečas 1967, Example 2.8, p. 33].)

In our case the form  $A$  is coercive over  $H^1(\mathcal{D})$ , indeed,

$$(5.2.3) \quad A(v, v) = \|v\|_{H^1(\mathcal{D})}^2, \text{ for } v \in H^1(\mathcal{D})$$

so that the Riesz representation theorem guarantees a unique solution for any  $f$  in  $L_2(\mathcal{D})$  (say).

One may now introduce finite elements partitions of  $\mathcal{D}$ ; if numerical integration is not taken into account we may assume that the elements that meet  $\partial\mathcal{D}$  are curved to fit  $\partial\mathcal{D}$  exactly. Under conditions on quasi-uniformity (and other general standard conditions that we shall not give here) it may be shown (see [Schatz and Wahlbin 1994, Theorem 4.1], cf. also [Scott 1976] for an original, somewhat weaker, result when  $n = 2$ ), that with  $u_h \in S_h$  the finite element approximation given by

$$(5.2.4) \quad A(u_h, \chi) = (f, \chi), \text{ for all } \chi \in S_h,$$

we have for  $e = u - u_h$ ,

$$(5.2.5) \quad \|e\|_{W_\infty^1(\mathcal{D})} \leq Ch^{r-1},$$

where  $C$  depends on the smoothness of  $u$  (i.e.,  $f$ ), the coefficients and various parameters describing  $S_h$ ; however, it can be taken independent of  $h$ .

We now make explicit the approximation hypothesis that

$$(5.2.6) \quad \min_{\chi \in S_h} \|w - \chi\|_{W_p^1(\mathcal{D})} \leq Ch^{r-1}\|w\|_{W_p^r(\mathcal{D})}, \text{ for } 1 \leq p \leq \infty,$$

where  $C$  is independent of  $w$ ,  $h$  and  $p$  (see [Ciarlet 1991]; in certain cases smoothing à la [Hilbert 1973] or [Strang 1973], cf. also Section 2.2, may be required).

We shall estimate, via a typical duality argument, the following negative norm,

$$(5.2.7) \quad \|e\|_{W_p^{-(r-2)}(\mathcal{D})} \equiv \sup_{\|v\|_{W_{p'}^{r-2}(\mathcal{D})}=1} (e, v)$$

where  $p$  is close to  $\infty$  and  $\frac{1}{p} + \frac{1}{p'} = 1$ , and thus  $p'$  is close to 1. For any  $v$  as above, let

$$(5.2.8) \quad \begin{cases} -\Delta w + w = v & \text{in } \mathcal{D}, \\ \frac{\partial w}{\partial n} = 0 & \text{on } \partial\mathcal{D}. \end{cases}$$

We then have the following a priori estimate, for  $\partial\mathcal{D}$  smooth enough,

$$(5.2.9) \quad \|w\|_{W_{p'}^r(\mathcal{D})} \leq C \frac{1}{p' - 1} \|v\|_{W_{p'}^{r-2}(\mathcal{D})}.$$

We refer the reader to [Gilbarg and Trudinger 1983, Chapter 9] for tracing constants in a priori estimates to see that (5.2.9) is true. (The argument there is for a Dirichlet problem but easily adapted to our present Neumann problem. The reader who pursues this will connect with the constant in the Calderon-Zygmund Theorem and, ultimately, with that in  $L_p$ -estimates for the Hilbert transform.) The point is that, in contrast to the one-dimensional situation, we cannot take  $p' = 1$ . We now have

$$(5.2.10) \quad (e, v) = A(e, w) = A(e, w - \chi), \text{ for any } \chi \in S_h(\mathcal{D}).$$

Thus, by Hölder's inequality and (5.2.5), (5.2.6), (5.2.9), for a suitable choice of  $\chi \in S_h$ ,

$$\begin{aligned} (5.2.11) \quad |(e, v)| &\leq Ch^{r-1} \|w - \chi\|_{W_{p'}^1(\mathcal{D})} \\ &\leq Ch^{2r-2} \|w\|_{W_{p'}^r(\mathcal{D})} \leq \frac{Ch^{2r-2}}{(p' - 1)} \|v\|_{W_{p'}^{r-2}(\mathcal{D})} \\ &\equiv \frac{Ch^{2r-2}}{(p' - 1)}. \end{aligned}$$

Thus

$$(5.2.12) \quad \|e\|_{W_p^{-(r-2)}(\mathcal{D})} \leq Ch^{2r-2} p,$$

for  $p < \infty$ .

Analogously to our investigation in Section 4.1, cf. (4.1.4) we shall find that, now for  $r \geq 3$ , the negative norm estimate (5.2.12) leads to an abundance of (nonidentifiable) superconvergence points.

Let thus  $B_d$  be a ball of radius  $d$ , inside  $\mathcal{D}$ , and  $v$  a smooth nonnegative function with support in  $B_d$  such that

$$(5.2.13) \quad v = 1 \quad \text{on } B_{d/2}$$

and

$$(5.2.14) \quad \|v\|_{W_\infty^\ell(B_d)} \leq \Lambda d^{-\ell}, \quad \ell = 0, 1, \dots, r-2.$$

Then assuming that  $e$  does not vanish on  $B_d$ , say  $e > 0$  there, with  $\frac{1}{p} + \frac{1}{p'} = 1$ ,

$$(5.2.15) \quad \min_{x \in B_d} e(x) \leq \int_{B_d} e(y)v(y)/\|v\|_{L_1(B_d)}.$$

By (5.2.13),

$$(5.2.16) \quad \|v\|_{L_1(B_d)} \geq cd^n.$$

From (5.2.12) we then obtain

$$(5.2.17) \quad \min_{x \in B_d} e(x) \leq Cd^{-n}h^{2r-2}p\|v\|_{W_{p'}^{r-2}(B_d)}$$

and using (5.2.14) and Hölder's inequality, this is bounded as

$$\begin{aligned} (5.2.18) \quad \min_{x \in B_d} e(x) &\leq Cd^{-n}h^{2r-2}pd^{-(r-2)}d^{n/p'} \\ &= Ch^r \left(\frac{h}{d}\right)^{r-2} pd^{-n/p} \\ &\leq Ch^r \left(\frac{h}{d}\right)^{r-2} \ln(1/d) \end{aligned}$$

where we have chosen  $p = \ln(1/d)$ .

We conclude the following upon choosing  $d = h^{1-\sigma/(r-2)}$ :

**Proposition 5.2.1.** *For  $r \geq 3$ , under the various assumptions above, any ball of radius  $d = h^{1-\sigma/(r-2)}$  contains a point  $\xi$  where*

$$(5.2.19) \quad |e(\xi)| \leq Ch^{r+\sigma} \ln(1/h),$$

i.e., a point of superconvergence of (almost) order  $\sigma$ .

A similar investigation considering any first derivative  $\frac{\partial e}{\partial x_i}$  and  $\|\frac{\partial e}{\partial x_i}\|_{W_p^{-(r-1)}}$ , for  $p$  close to  $\infty$  (using now a priori estimates for the problem  $-\Delta w + w = \partial v/\partial x$  in  $\mathcal{D}$ , with homogeneous Neumann conditions on  $\partial\mathcal{D}$ ), gives that in any ball of size  $d = h^{1-\sigma/(r-1)}$ , there is a point where  $\partial e/\partial x_i$  changes sign or, a point superconvergent (almost) to order  $\sigma$  (i.e.,  $\partial e/\partial x_i = 0(h^{r-1+\sigma} \ln(1/h))$ ). Note that  $\partial e/\partial x_i$  may not be continuous in general.

The arguments given above in a simple case of “natural” boundary conditions may rather easily be extended to variable coefficient operators. They may also be extended to some essential boundary conditions such as homogeneous Dirichlet conditions on smooth domains, provided boundary data approximations are performed to sufficiently high accuracy, cf. [Schatz, Sloan and Wahlbin, Example 3.2]. (One notes that it is *not* in general enough to employ isoparametric elements at the boundary; these only approximate the boundary to order  $O(h^r)$ . Super-parametric approximations are required to obtain (5.2.12).) Due to the use of duality arguments, the arguments need to be carefully reconsidered when considering nonsmooth boundaries (such as polygonal domains) and nonsmooth coefficients; the a priori estimates used above in situations with sufficient smoothness do not necessarily hold in such cases.

### 5.3. General comments on local a priori error estimates.

The main technical tool for investigating superconvergence in the situation introduced in Section 5.1 will be given in Section 5.5. This section will be devoted to generalities and loose discussion, and the aim is to “appreciate” the rather formidable general tool to come . . .

We note that our main tool in the one-dimensional situation, viz. that  $u'_h \simeq P(u')$  where  $P$  is the  $L_2$ -projection into  $\$_h = \$_h^{\mu-1, r-1}$ , is now lost. To our knowledge, no one has been able to recover a suitable general analogue of it in several dimensions. (Only the case of tensor-product elements will build on one-dimensional results.)

Let us thus consider first the following simple situation, corresponding to the one in Lemma 3.2.1. Let  $u_h \in S_h$  be a function such that

$$(5.3.1) \quad A(u_h, \chi) = 0, \text{ for } \chi \in S_h^{\text{comp}}(\Omega),$$

where  $\Omega \subset\subset \mathcal{D}$ . For further simplicity here, we shall take the “simplest” case,

$$(5.3.2) \quad A(v, w) \equiv D(v, w) \equiv \int \nabla v \cdot \nabla w dx,$$

corresponding to Poisson’s equation in (5.1.1).

Now, a rather well-known result about harmonic functions is the following: Let  $B_0$  be any ball of radius  $r$  and  $B_d$  a concentric ball of radius  $r + d$ . (The base radius  $r$  is rather immaterial here; what matters is the difference  $d$ .) There is then a universal constant  $C$  such that if  $u$  is harmonic in  $B_d$ , then

$$(5.2.3) \quad \|\nabla u\|_{L_2(B_0)} \leq Cd^{-1}\|u\|_{L_2(B_d)}.$$

We would now wish to prove a corresponding result for  $u_h \in S_h$  a “discrete harmonic function”, i.e., satisfying

$$(5.3.4) \quad D(u_h, \chi) = 0, \text{ for } \chi \in S_h^{\text{comp}}(B_d).$$

A first step is to try to prove the continuous case (5.3.3) by use of the similar property,

$$(5.3.5) \quad D(u, \chi) = 0, \text{ for } \chi \in \overset{\circ}{H}_1(B_d).$$

(I.e., we eschew, at least for the moment, any use of estimates for fundamental solutions or of “high-powered” Harmonic Analysis.)

Here is one “elementary” argument: Let  $\omega$  be a “cut-off” function,

$$(5.3.6) \quad \omega \equiv 1 \text{ on } B_0,$$

$$(5.3.7) \quad \text{Supp}(\omega) \subset\subset B_d,$$

and

$$(5.3.8) \quad \|\omega\|_{W_\infty^\ell(B_d)} \leq Cd^{-\ell}, \ell = 0, 1.$$

It is well-known how to construct such a function (by use of  $\exp(-1/x^2)$ , e.g.). Then

$$(5.3.9) \quad \|\nabla u\|_{L_2(B_0)}^2 = \int_{B_0} \nabla u \cdot \nabla u \leq \int_{B_d} \omega^2 \nabla u \cdot \nabla u = \|\omega \nabla u\|_{L_2(B_d)}^2.$$

Now,

$$(5.3.10) \quad \|\omega \nabla u\|_{L_2(B_d)}^2 = \int_{B_d} \nabla u \nabla (\omega^2 u) - \int_{B_d} \nabla u \cdot 2\omega (\nabla \omega) u$$

and from (5.3.5), the first term on the right vanishes!

Thus, by Cauchy-Schwarz' inequality, and (5.3.8),

$$(5.3.11) \quad \begin{aligned} \|\omega \nabla u\|_{L_2(B_d)}^2 &\leq \left| \int_{B_d} (\omega \nabla u) \cdot (\nabla \omega) u \right| \\ &\leq \|\omega \nabla u\|_{L_2(B_d)} C d^{-1} \|u\|_{L_2(B_d)} \end{aligned}$$

and we have proven that

$$(5.3.12) \quad \|\omega \nabla u\|_{L_2(B_d)} \leq C d^{-1} \|u\|_{L_2(B_d)}$$

which of course implies (5.3.3) by use of (5.3.9).

Well, let us now see how this travels to the discrete situation, (5.3.4)! Equations (5.3.9) and (5.3.10) are fine, with  $u$  replaced by  $u_h$ . However, now we can't scrap the first term on the right of (5.3.10) since  $\omega^2 u_h$  does not in general belong to  $S_h^{comp}$ . Instead of (5.3.10) we have for any  $\chi \in S_h^{comp}(B_d)$ ,

$$(5.3.13) \quad \|\omega \nabla u_h\|_{L_2(B_d)}^2 = \int_{B_d} \nabla u_h \nabla (\omega^2 u_h - \chi) - 2 \int_{B_d} \omega \nabla u_h (\nabla \omega) u_h.$$

This is where superapproximation enters; this point is one of the essential discoveries in [Nitsche and Schatz 1974]. So, we now take a little bit more care and, say, restrict the cut-off function  $\omega$  to have support in  $B_{d/2}$ . If we then let  $d \geq c_1 h$  for  $c_1$  large enough ( $d$  sufficiently large compared to element diameters), we make the appropriate superapproximation assumption: There exists  $\chi \in S_h^{comp}(B_d)$  such that

$$(5.3.14) \quad \|\omega^2 u_h - \chi\|_{H^1(B_d)} \leq C(h/d) \|u_h\|_{H^1(B_d)},$$

cf. Theorem 2.3.1. From (5.3.13) we then have

$$(5.3.15) \quad \begin{aligned} \|\omega \nabla u_h\|_{L_2(B_d)}^2 &\leq \|\nabla u_h\|_{L_2(B_d)} C(h/d) \|u_h\|_{H^1(B_d)} \\ &\quad + C d^{-1} \|\omega \nabla u_h\|_{L_2(B_d)} \|u_h\|_{L_2(B_d)}. \end{aligned}$$

It follows from this and (5.3.9) (with  $u$  replaced by  $u_h$ ), that

$$(5.3.16) \quad \|u_h\|_{H^1(B_0)} \leq C_1 \left( \frac{h}{d} \right)^{1/2} \|u_h\|_{H^1(B_d)} + C_1 d^{-1} \|u_h\|_{L_2(B_d)}.$$

Now, taking a little bit more care so that the original cut-off has support in  $B_{d/4}$ , say, we have upon repeating the argument with the obvious changes ( $B_0$  replaced by  $B_{d/2}$ ),

$$(5.3.17) \quad \|u_h\|_{H^1(B_0)} \leq C_2 \left( \frac{h}{d} \right) \|u_h\|_{H^1(B_d)} + C_2 d^{-1} \|u_h\|_{L_2(B_d)}.$$

To cap off the argument, we now assume an inverse estimate for derivatives: Shrinking  $B_d$  a little bit if necessary to coincide with a mesh-domain,

$$(5.3.18) \quad \|u_h\|_{H^1(B_d)} \leq C_3 h^{-1} \|u_h\|_{L_2(B_d)},$$

cf. Theorem 1.2.4. We then obtain the desired estimate

$$(5.3.19) \quad \|u_h\|_{H^1(B_0)} \leq C_4 d^{-1} \|u_h\|_{L_2(B_d)},$$

for  $d \geq c_1 h$ .

[Nitsche and Schatz 1974] now go on to show, e.g., that if

$$(5.3.20) \quad A(u - u_h, \chi) = 0, \text{ for } \chi \in S_h^{comp}(B_d),$$

then

$$(5.3.21) \quad \begin{aligned} & \|u - u_h\|_{H^1(B_0)} \\ & \leq C \min_{\chi \in S_h} \left( \|u - \chi\|_{H^1(B_d)} + d^{-1} \|u - \chi\|_{L_2(B_d)} \right) \\ & \quad + C d^{-1} \|u - u_h\|_{L_2(B_d)}. \end{aligned}$$

The argument for showing (5.3.21) goes briefly as follows: We first assume that  $B_0$  is a ball of radius  $d$  and  $B_d$  one of radius  $2d$ . Let  $\omega \in C_0^\infty(B_d)$  be a cutoff function roughly as before, but now with

$$(5.3.22) \quad \omega \equiv 1 \quad \text{on } B_{d/2}.$$

Set  $\tilde{u} := \omega u$  and let  $\tilde{u}_h \in S_h^{comp}(B_d)$  be the Ritz–Galerkin projection into  $S_h^{comp}(B_d)$ , i.e., given by

$$(5.3.23) \quad D(\tilde{u} - \tilde{u}_h, \chi) = 0, \text{ for } \chi \in S_h^{comp}(B_d).$$

As is well known and trivial to prove, with the semi-norm  $|u|_{\overset{\circ}{H}^1} = \|\nabla u\|_{L_2}$ ,

$$(5.3.24) \quad |\tilde{u}_h|_{\overset{\circ}{H}^1(B_d)} \leq |\tilde{u}|_{\overset{\circ}{H}^1(B_d)} \leq C \left( |u|_{\overset{\circ}{H}^1(B_d)} + d^{-1} \|u\|_{L_2(B_d)} \right).$$

We now write

$$(5.3.25) \quad u - u_h = (\tilde{u} - \tilde{u}_h) + (\tilde{u}_h - u_h) \text{ on } B_0.$$

Since  $\tilde{u} - \tilde{u}_h \in \overset{\circ}{H}^1(B_d)$  we have by Poincaré–Friedrichs' inequality and then using (5.3.24),

$$(5.3.26) \quad \begin{aligned} \|\tilde{u} - \tilde{u}_h\|_{H^1(B_0)} & \leq C(1+d)|\tilde{u} - \tilde{u}_h|_{\overset{\circ}{H}^1(B_d)} \\ & \leq C(1+d)|\tilde{u}|_{\overset{\circ}{H}^1(B_d)} \\ & \leq C \left( |u|_{\overset{\circ}{H}^1(B_d)} + d^{-1} \|u\|_{L_2(B_d)} \right). \end{aligned}$$

It remains to estimate the second term on the right of (5.3.25).

Now, by (5.3.22), for  $\chi \in S_h^{comp}(B_{d/2})$ ,

$$(5.3.27) \quad D(\tilde{u}_h - u_h, \chi) = D(\tilde{u} - u, \chi) = 0,$$

i.e.,  $\tilde{u}_h - u_h \in S_h$  is discrete harmonic in  $B_{d/2}$ . By (5.3.19) then, and again using that  $\omega \equiv 1$  on  $B_{d/2}$ ,

$$(5.3.28) \quad \begin{aligned} \|\tilde{u}_h - u_h\|_{H^1(B_0)} & \leq \frac{C}{d} \|\tilde{u}_h - u_h\|_{L_2(B_{d/2})} \\ & \leq \frac{C}{d} \left( \|\tilde{u}_h - \tilde{u}\|_{L_2(B_{d/2})} + \|u - u_h\|_{L_2(B_{d/2})} \right). \end{aligned}$$

By the same argument as in (5.3.26) we have

$$(5.3.29) \quad \frac{C}{d} \|\tilde{u}_h - \tilde{u}\|_{L_2(B_{d/2})} \leq C |\tilde{u}_h - \tilde{u}|_{\dot{H}^1(B_d)} \leq C \left( |u|_{\dot{H}^1(B_d)} + d^{-1} \|u\|_{L_2(B_d)} \right).$$

From (5.3.25), (5.3.26) and (5.3.28), (5.3.29) we have

$$(5.3.30) \quad \|u - u_h\|_{H^1(B_1)} \leq C \left( |u|_{\dot{H}^1(B_d)} + d^{-1} \|u\|_{L_2(B_d)} \right) + Cd^{-1} \|u - u_h\|_{L_2(B_d)}.$$

Recall that this was proven under the assumption that  $B_0$ ,  $B_d$  were concentric balls of radii  $d$ ,  $2d$ , respectively. (We did use Poincaré–Friedrichs' inequality.) By squaring and summing, (5.3.30) holds with the original setup of  $B_0$  any ball and  $B_d$  the concentric one separated by a distance  $d$ . The desired estimate (5.3.21) now follows by writing  $u - u_h = (u - \chi) + (\chi - u_h)$ , for any  $\chi \in S_h$ .

We shall informally refer to the first term on the right of (an estimate such as) (5.3.21) as “the best local approximation”, and the weaker norm last term as the “slush term”. Note that if we try to take local approximability into account in a very sharp fashion, i.e., taking  $d$  smaller (while still  $\geq c_1 h$ ), we pay for this in the slush term. Also note that the slush term in the similar result for the  $L_2$ -projection, (3.2.14), enjoys exponential decay  $\exp(-d/h)$ . The “decay” in (5.3.21) is much weaker, and [Nitsche and Schatz 1974] thus felt obliged to further weaken the slush-term into one involving negative norms:

$$(5.3.31) \quad \begin{aligned} & \|u - u_h\|_{H^1(B_0)} \\ & \leq C \min_{\chi \in S_h} (\|u - \chi\|_{H^1(B_d)} + d^{-1} \|u - \chi\|_{L_2(B_d)}) \\ & \quad + Cd^{-1-s} \|u - u_h\|_{H^{-s}(B_d)}, \end{aligned}$$

cf. (1.1.10).

Correspondingly in the  $L_2$ -norm, again of course under suitable assumptions,

$$(5.3.32) \quad \begin{aligned} \|u - u_h\|_{L_2(B_0)} & \leq C \min_{\chi \in S_h} (\|u - \chi\|_{L_2(B_d)} + h \|u - \chi\|_{H^1(B_d)}) \\ & \quad + Cd^{-s} \|u - u_h\|_{H^{-s}(B_d)}. \end{aligned}$$

The inclusion of the term  $h \|u - \chi\|_{H^1(B_d)}$  on the right reflects the fact that in general the  $H^1$ -projection is not a stable operator in  $L_2$ . E.g., in the one-dimensional situation for the problem  $-u'' = f$  on  $I$ ,  $u(0) = u(1) = 0$ , with piecewise linear  $u_h$  is the interpolant of  $u$ . Pondering then the following figure




Figure 5.3.1

for  $h = 1/2$ , it is obvious that the  $\overset{\circ}{H}{}^1$ -projection cannot be a stable operator in *any*  $L_p$ -norm except for  $p = \infty$ . (This counterexample was noted in [Babuška and Osborn 1980, page 58]; if interelement continuity is  $C^1$  or better,  $L_2$ -stability holds, cf. [Schatz and Wahlbin 1977, Appendix 2] for the local estimate, and [Babuška and Aziz 1972, Theorem 6.3.8] for the easier global case.)

[Nitsche and Schatz 1974] also gave corresponding  $L_2$ -based estimates in the case of  $u - u_h$  satisfying (5.1.4), i.e.,

$$(5.3.33) \quad A(u - u_h, \chi) = F(\chi), \text{ for } \chi \in S_h^{\text{comp}}(\Omega).$$

The argument is, in brief, as follows. Again, first let  $B_0$  be a ball of radius  $d$  and  $B_d$  a ball of radius  $2d$ . Now  $F(\cdot)$  is a functional on  $\overset{\circ}{H}{}^1(B_d)$ , and we set

$$(5.3.34) \quad |||F|||_{H^{-1}(B_d)} = \sup_{\substack{v \in \overset{\circ}{H}{}^1(B_d) \\ |v|_{\overset{\circ}{H}{}^1(B_d)} = 1}} F(v).$$

Again, let us assume that  $A(v, w) = D(v, w) \equiv \int \nabla v \cdot \nabla w$ , for simplicity. Solve now the problem  $-\Delta w = f$  in  $B_d$ ,  $w = 0$  on  $\partial B_d$ , i.e., find  $w \in \overset{\circ}{H}{}^1(B_d)$  such that

$$(5.3.35) \quad D(w, v) = F(v), \text{ for all } v \in \overset{\circ}{H}{}^1(B_d).$$

This problem has a unique solution by Riesz representation theorem.

Then

$$(5.3.36) \quad D(u + w - u_h, \chi) = 0, \text{ for } \chi \in S_h^{\text{comp}}(B_d).$$

Thus by the triangle inequality and (5.3.31), taking  $\chi = 0$  there for the moment,

$$\begin{aligned}
 (5.3.37) \quad \|u - u_h\|_{H^1(B_0)} &\leq \|(u + w) - u_h\|_{H^1(B_0)} + \|w\|_{H^1(B_0)} \\
 &\leq C \left( |u + w|_{\overset{\circ}{H}^1(B_d)} + d^{-1} \|u + w\|_{L_2(B_d)} \right) \\
 &\quad + Cd^{-1} \|u + w - u_h\|_{L_2(B_d)} + \|w\|_{H^1(B_d)} \\
 &\leq C \left( |u|_{\overset{\circ}{H}^1(B_d)} + d^{-1} \|u\|_{L_2(B_d)} \right) \\
 &\quad + Cd^{-1} \|u - u_h\|_{L_2(B_d)} \\
 &\quad + C \left( |w|_{\overset{\circ}{H}^1(B_d)} + d^{-1} \|w\|_{L_2(B_d)} \right).
 \end{aligned}$$

Setting  $v = w$  in (5.3.35) we find that

$$(5.3.38) \quad |w|_{\overset{\circ}{H}^1(B_d)} \leq |||F|||_{H^{-1}(B_d)}$$

and by Poincaré–Friedrichs' inequality, thus

$$(5.3.39) \quad d^{-1} \|w\|_{L_2(B_d)} \leq C |w|_{\overset{\circ}{H}^1(B_d)} \leq C |||F|||_{H^{-1}(B_d)}.$$

So, from (5.3.37),

$$\begin{aligned}
 (5.3.40) \quad \|u - u_h\|_{H^1(B_0)} &\leq C \left( |u|_{\overset{\circ}{H}^1(B_d)} + d^{-1} \|u\|_{L_2(B_d)} \right) \\
 &\quad + Cd^{-1} \|u - u_h\|_{L_2(B_d)} + C |||F|||_{H^{-1}(B_d)}.
 \end{aligned}$$

This was proven for  $B_0$  and  $B_d$  balls of radii  $d$  and  $2d$ . To get it for a general  $B_0$  of radius  $r$ , and  $B_d$  of radius  $r + d$ , one again squares and sums. One then needs the following lemma, which is not completely obvious.

**Lemma 5.3.1.** *If  $\Omega_j$ ,  $j = 1, \dots, J$  are such that  $\bar{\Omega}_j$  are disjoint sets, we have*

$$(5.3.41) \quad |||F|||_{H^{-1}(\bigcup_1^J \Omega_j)}^2 \geq \sum_{j=1}^J |||F|||_{H^{-1}(\Omega_j)}^2.$$

Assuming this lemma for the moment, we cover  $B_0$  by (finite) families of disjoint balls of radius  $d$  (how many families? — solve the sphere packing problem!). The result (5.3.40) is then extended to general  $B_0$  and  $B_d$  as before. (Indeed, to any domain  $\Omega_0$  and  $\Omega_d = \{x : \text{dist}(x, \Omega_0) \leq d\}$ .)

Writing  $u - u_h = (u - \chi) - (u_h - \chi)$  for  $\chi \in S_h$  then gives for  $u$  and  $u_h$  satisfying (5.3.33) (i.e., (5.1.4)),

$$\begin{aligned}
 (5.3.42) \quad \|u - u_h\|_{H^1(B_0)} &\\
 &\leq C \min_{\chi \in S_h} \left( |u - \chi|_{\overset{\circ}{H}^1(B_d)} + d^{-1} \|u - \chi\|_{L_2(B_d)} \right) \\
 &\quad + Cd^{-1} \|u - u_h\|_{L_2(B_d)} + C |||F|||_{H^{-1}(B_d)}.
 \end{aligned}$$

(Again, the term  $\|u - u_h\|_{L_2(B_d)}$  can be replaced by a negative norm at the expense of additional inverse powers of  $d$ .)

Remark 5.3.1. In (5.3.33),  $F(\cdot)$  is seen only by how it acts on  $\chi \in S_h^{comp}(\Omega)$ . Since  $S_h^{comp}(\Omega) \subseteq \overset{\circ}{H}{}^1(\Omega)$ , an argument involving the Hahn–Banach extension theorem then shows that  $|||F|||_{H^{-1}(B_d)}$  in (5.3.43) may be replaced by

$$(5.3.43) \quad \max_{\substack{\chi \in S_h^{comp}(B_d) \\ |\chi|_{\overset{\circ}{H}{}^1(B_d)} = 1}} F(\chi). \quad \square$$

It remains to prove Lemma 5.3.1.

Proof of Lemma 5.3.1. We shall prove it for  $J = 2$ ; it will be obvious that the proof generalizes to any  $J$ .

If  $\Omega_1$  and  $\Omega_2$  are two balls such that  $\overline{\Omega}_1$  and  $\overline{\Omega}_2$  are disjoint, we let  $v_j \in \overset{\circ}{H}{}^1(\Omega_j)$ , with  $|v_j|_{\overset{\circ}{H}{}^1(\Omega_j)} = 1$ , for  $j = 1, 2$ . Setting (for simplicity in notation),

$$\theta_j = |||F|||_{H^{-1}(\Omega_j)}, \quad j = 1, 2$$

we have with  $\Omega = \Omega_1 \cup \Omega_2$  and

$$v := \theta_1 v_1 + \theta_2 v_2 \in \overset{\circ}{H}{}^1(\Omega).$$

Then

$$|||F|||_{H^{-1}(\Omega)}^2 \geq \left( \frac{F(v)}{|v|_{\overset{\circ}{H}{}^1(\Omega)}} \right)^2 = \frac{(\theta_1 F(v_1) + \theta_2 F(v_2))^2}{\theta_1^2 + \theta_2^2}.$$

We take the supremum over such  $v_1$  and  $v_2$  as above and obtain

$$|||F|||_{H^{-1}(\Omega)}^2 \geq \frac{(\theta_1^2 + \theta_2^2)^2}{\theta_1^2 + \theta_2^2} = |||F|||_{H^{-1}(\Omega_1)}^2 + |||F|||_{H^{-1}(\Omega_2)}^2. \quad \square$$

#### 5.4. General comments on $L_\infty$ estimates.

In the years 1975–1976 the following basic papers on  $L_\infty$ -estimates in finite element methods in multi-dimensions on general meshes were published: [Natterer 1975], [Nitsche 1975], [Frehse and Rannacher 1976] and [Scott 1976].

To this was quickly added an investigation into the harder problem of *localized* maximum norm estimates, [Schatz and Wahlbin 1977]. (There had been previous papers on local maximum norm estimates in several space dimensions, on uniform or translation invariant meshes, cf. [Bramble and Thomée 1974], [Bramble, Nitsche and Schatz 1975] and [Bramble and Schatz 1976].)

In the next section we shall give — finally! — our basic tool to be used in our superconvergence investigations. We therefore feel no need to review the papers mentioned above. However, some readers may appreciate seeing how the local  $H^1$  estimates of [Nitsche–Schatz 1974] actually lead to global maximum norm error estimates. So, as a longish aside, we shall give the argument in the plane case of two space dimensions and the particular case of piecewise linear functions on triangular finite elements. (In this case much technical detail is simplified; cf. [Schatz and Wahlbin 1982, Remark 5.3].)

Let, for simplicity,  $\mathcal{D}$  be a convex plane domain with a smooth boundary  $\partial\mathcal{D}$ . Let  $\tau_h = \{\tau_i^h\}_{i=1}^{N(h)}$  be disjoint open triangles such that

$$(5.4.1) \quad \overline{\mathcal{D}}_h = \bigcup_{i=1}^{N(h)} \overline{\tau_i^h} \subseteq \overline{\mathcal{D}}$$

and such that

$$(5.4.2) \quad \text{dist}(\mathcal{D}_h, \partial\mathcal{D}) \leq Ch^2.$$

We assume that the triangles are edge-to-edge, i.e., no vertex of one triangle falls in the interior of the edge of another. We also assume that with  $h = \max_i \text{diam}(\tau_i^h)$ , the family of triangulations is quasi-uniform in the sense that with  $\rho_i^h$  the radius of the maximal inscribed disc in  $\tau_i^h$ ,

$$(5.4.3) \quad \rho_i^h \geq c_0 h$$

where  $c_0 > 0$  is independent of  $i$  and  $h$ .

The problem we shall consider is, again for simplicity, that of finding  $u$  such that

$$(5.4.4) \quad \begin{aligned} -\Delta u &= f && \text{in } \mathcal{D}, \\ u &= 0 && \text{on } \partial\mathcal{D}. \end{aligned}$$

The weak form of (5.4.3) namely, to find  $u \in \overset{\circ}{H}^1(\mathcal{D})$  such that

$$(5.4.5) \quad D(u, \chi) \equiv \int_{\mathcal{D}} \nabla u \cdot \nabla \chi = (f, \chi), \text{ for } \chi \in \overset{\circ}{H}^1(\mathcal{D}),$$

incorporates the essential Dirichlet boundary conditions. Our piece-wise linear finite element family is now

$$(5.4.6) \quad \overset{\circ}{S}_h(\mathcal{D}) = \{\chi \in C^0(\mathcal{D}_h) : \chi = 0 \text{ on } \partial\mathcal{D}_h, \chi|_{\tau_i^h} = c_0 + c_1 x + c_2 y\}.$$

We regard the functions in  $\overset{\circ}{S}_h(\mathcal{D})$  as  $\equiv 0$  in the skin-layer  $\mathcal{D} \setminus \mathcal{D}_h$ . The approximation  $u_h \in \overset{\circ}{S}_h$  is sought as

$$D(u_h, \chi) = (f, \chi), \text{ for } \chi \in \overset{\circ}{S}_h,$$

i.e.,

$$(5.4.7) \quad D(u - u_h, \chi) = 0, \text{ for } \chi \in \overset{\circ}{S}_h.$$

Let us now fix  $x_0 \in \mathcal{D}$ . If  $x_0 \in \mathcal{D} \setminus \mathcal{D}_h$ , obviously by (5.4.2),  $|(u - u_h)(x_0)| \leq Ch^2$  if  $u$  is smooth. We assume now that  $x_0 \in \mathcal{D}_h$ . Using a suitable  $\chi \in \overset{\circ}{S}_h$ , we have if  $x_0 \in \tau_{i_0}^h \equiv \tau_0$ ,

$$(5.4.8) \quad \begin{aligned} |(u - u_h)(x_0)| &\leq |(u - \chi)(x_0)| + |(\chi - u_h)(x_0)| \\ &\leq |(u - \chi)(x_0)| + Ch^{-1} \|\chi - u_h\|_{L_2(\tau_0)} \end{aligned}$$

where we used an easily verifiable inverse estimate, cf. [Ciarlet 1991]. Thus, again by the triangle inequality and also Hölder's inequality,

$$(5.4.9) \quad |(u - u_h)(x_0)| \leq C \|u - \chi\|_{L_\infty(\tau_0)} + Ch^{-1} \|u - u_h\|_{L_2(\tau_0)}.$$

Hence,

$$(5.4.10) \quad |(u - u_h)(x_0)| \leq C \min_{\chi \in \overset{\circ}{S}_h} \|u - \chi\|_{L_\infty(\mathcal{D})} + Ch^{-1} \|u - u_h\|_{L_2(\tau_0)}.$$

Now for a duality argument:

$$(5.4.11) \quad \|u - u_h\|_{L_2(\tau_0)} = \sup_{\|v\|_{L_2(\tau_0)}=1} (u - u_h, v).$$

For each such  $v$ , let  $-\Delta w = v$  in  $\mathcal{D}$ ,  $w = 0$  on  $\partial\mathcal{D}$ . Then

$$(5.4.12) \quad (u - u_h, v) = D(u - u_h, w).$$

With  $w_h \in \overset{\circ}{S}_h$  defined by

$$(5.4.13) \quad D(\chi, w - w_h) = 0, \text{ for } \chi \in \overset{\circ}{S}_h$$

we then have, for any  $\chi \in \overset{\circ}{S}_h$ ,

$$(5.4.14) \quad (u - u_h, v) = D(u - \chi, w - w_h).$$

So,

$$(5.4.15) \quad |(u - u_h, v)| \leq C \min_{\chi \in \overset{\circ}{S}_h} \|u - \chi\|_{W_\infty^1(\mathcal{D})} \|w - w_h\|_{W_1^1(\mathcal{D})}.$$

Next, let

$$(5.4.16) \quad \Omega_j = \{x : 2^{-j} \leq |x - x_0| \leq 2^{-j+1}\} \cap \mathcal{D},$$

and let  $\mathcal{J}_0$  be such that  $2^{-\mathcal{J}_0} = c_0 h$  with  $c_0$  large enough (or, the nearest integer to accomplish this). Also let

$$(5.4.17) \quad d_j = 2^{-j},$$

and

$$(5.4.18) \quad \Omega_{\mathcal{J}_0} = \{x : |x - x_0| \leq 2^{-\mathcal{J}_0}\} \cap \mathcal{D}.$$

Then

$$(5.4.19) \quad \|w - w_h\|_{W_1^1(\mathcal{D})} = \|w - w_h\|_{W_1^1(\Omega_{\mathcal{J}_0})} + \sum_{j=I}^{\mathcal{J}_0-1} \|w - w_h\|_{W_1^1(\Omega_j)}$$

where  $I$  is small enough so that  $\cup \Omega_j$  covers  $\mathcal{D}$ . Here,

$$(5.4.20) \quad \begin{aligned} \|w - w_h\|_{W_1^1(\Omega_{\mathcal{J}_0})} &\leq Ch \|w - w_h\|_{H^1(\mathcal{J}_0)} \\ &\leq Ch \|w - w_h\|_{H^1(\mathcal{D})} \leq Ch^2 \|w\|_{H^2(\mathcal{D})} \leq Ch^2, \end{aligned}$$

where we used that  $w_h$  is the  $\overset{\circ}{H}^1$  projection of  $w$  so that  $\|w - w_h\|_{H^1(\mathcal{D})} = \min_{\chi \in \overset{\circ}{S}_h} \|w - \chi\|_{H^1(\mathcal{D})}$  and then approximation theory: take  $\chi$  as the interpolant at vertices of  $w$ ; note that  $w \in H^2(\mathcal{D})$  since  $\partial\mathcal{D}$  is smooth. (See any standard textbook on Partial Differential Equations.) Then  $w \in \mathcal{C}^0(\mathcal{D})$  by Sobolev's theorem and the piecewise linear interpolant  $\chi$  of  $w$  makes sense and is well-known to satisfy

$\|w - \chi\|_{H^1(\mathcal{D})} \leq Ch\|w\|_{H^2(\mathcal{D})}$ . The reader is encouraged to check this with respect to the “skin–layer”  $\mathcal{D} \setminus \mathcal{D}_h$ , where  $\chi \equiv 0$ .

For  $j < J_0$  we have by local  $H^1$ –error estimates, see (5.3.20) (cf. also [Schatz and Wahlbin 1982, Theorem 4.1] for the extension needed in case the domains involved abut on the boundary  $\partial\mathcal{D}$ ),

$$\begin{aligned} (5.4.21) \quad & \|w - w_h\|_{W_1^1(\Omega_j)} \leq d_j \|w - w_h\|_{H^1(\Omega_j)} \\ & \leq Cd_j \left\{ \min_{\chi \in \overset{\circ}{S}_h} (\|w - \chi\|_{H^1(\Omega'_j)} + d_j^{-1} \|w - \chi\|_{L_2(\Omega'_j)}) \right. \\ & \quad \left. + d_j^{-1} \|w - w_h\|_{L_2(\Omega'_j)} \right\} \\ & \leq Cd_j \min_{\chi \in \overset{\circ}{S}_h} (\|w - \chi\|_{H^1(\Omega'_j)} + d_j^{-1} \|w - \chi\|_{L_2(\Omega'_j)}) \\ & \quad + C\|w - w_h\|_{L_2(\mathcal{D})}. \end{aligned}$$

Here,  $\Omega'_j = \Omega_{j-1} \cup \bar{\Omega}_j \cup \Omega_{j+1}$  (recall that  $d_j \geq c_0 h$  with  $c_0$  large enough). Again by standard duality argument over  $L_2$ , and standard approximation theory (and checking the skin layer  $\mathcal{D} \setminus \mathcal{D}_h$ ),

$$(5.4.22) \quad \|w - w_h\|_{L_2(\mathcal{D})} \leq Ch^2 \|w\|_{H^2(\mathcal{D})} \leq Ch^2$$

where we again used a standard a priori estimate in the last step. Furthermore, again by approximation theory, taking  $\chi$  to be the piece–wise linear interpolant,

$$(5.4.23) \quad \min_{\chi \in \overset{\circ}{S}_h} (\|w - \chi\|_{H^1(\Omega'_j)} + d_j^{-1} \|w - \chi\|_{L_2(\Omega'_j)}) \leq Ch\|w\|_{H^2(\Omega'_j)},$$

with  $\Omega''_j = \Omega_{j-2} \cup \bar{\Omega}_{j-1} \cup \bar{\Omega}_j \cup \bar{\Omega}_{j+1} \cup \Omega_{j+2}$ . Since  $c_0$  is large enough is assumed, the support of  $v$  ( $= \tau_0$ ) may be assumed  $\simeq d_j$  away from  $\Omega''_j$ . Now,

$$(5.4.24) \quad w(x) = \int_{\tau_0} v(y) G^{(x)}(y) dy$$

where  $G^{(x)}(y)$  is the Green’s function for the problem  $-\Delta w = v$  in  $\mathcal{D}$ ,  $w = 0$  on  $\partial\mathcal{D}$ . It is well known that for any  $k^{\text{th}}$  derivative with respect to  $x$ ,  $k \geq 1$ ,  $D_x^k$ , we have (remember,  $n = 2$ )

$$(5.4.25) \quad |D_x^k G^{(x)}(y)| \leq C|x - y|^{-k}$$

since  $\partial\mathcal{D}$  is smooth (cf. [Krasovskii 1967]). Hence, using also Hölder’s inequality,

$$\begin{aligned} (5.4.26) \quad & \|w\|_{H^2(\Omega''_j)} \leq Cd_j \|w\|_{W_\infty^2(\Omega''_j)} \leq Cd_j \int_{\tau_0} |v(y)| d_j^{-2} dy \\ & \leq Cd_j h \|v\|_{L_2(\tau_0)} d_j^{-2} = Ch d_j^{-1}. \end{aligned}$$

Then, from (5.4.23),

$$(5.4.27) \quad \min_{\chi \in \overset{\circ}{S}_h} (\|w - \chi\|_{H^1(\Omega'_j)} + d_j^{-1} \|w - \chi\|_{L_2(\Omega'_j)}) \leq Ch^2 d_j^{-1},$$

and so from (5.4.21) and (5.4.22),

$$(5.4.28) \quad \|w - w_h\|_{W_1^1(\Omega_j)} \leq Ch^2.$$

Combining this with (5.4.20), we have from (5.4.19), and since  $2^{-J_0} = c_0 h$ , i.e.,  $J_0 \simeq \ln(1/h)$ ,

$$(5.4.29) \quad \|w - w_h\|_{W_1^1(\mathcal{D})} \leq Ch^2 + C \sum_{j=I}^{J_0-1} h^2 \leq Ch^2 \ln(1/h).$$

Hence from (5.4.15),

$$(5.4.30) \quad |(u - u_h, v)| \leq C \min_{\chi \in \overset{\circ}{S}_h} \|u - \chi\|_{W_\infty^1(\mathcal{D})} h^2 \ln(1/h)$$

so that, finally, from (5.4.11) and (5.4.10),

$$(5.4.31) \quad |(u - u_h)(x_0)| \leq C \ln(1/h) \min_{\chi \in \overset{\circ}{S}_h} \{\|u - \chi\|_{L_\infty(\mathcal{D})} + h \|u - \chi\|_{W_\infty^1(\mathcal{D})}\},$$

or, if  $u$  is smooth enough, with  $C = C(u)$ ,

$$(5.4.32) \quad \|u - u_h\|_{L_\infty(\mathcal{D})} \leq Ch^2 \ln 1/h.$$

Actually, by an argument involving integration by parts over each element in (5.4.14), and for general  $r$ , cf. [Schatz and Wahlbin 1982], we have

$$(5.4.33) \quad \|u - u_h\|_{L_\infty(\mathcal{D}_h)} \leq C (\ln(1/h))^{\bar{r}} \min_{\chi \in \overset{\circ}{S}_h} \|u - \chi\|_{L_\infty(\mathcal{D}_h)}$$

where

$$\bar{r} = \begin{cases} 1, & \text{if } r = 2, \\ 0, & \text{if } r \geq 3. \end{cases}$$

We are now ready to describe:

## 5.5. The main technical tools for proving superconvergence in second order elliptic problems in several space dimensions.

As an introduction here, let us recall (1.3.7),

$$|(\theta', \chi')| \leq Ch^r \|\chi\|_{W_1^1(I)}, \text{ for } \chi \in \overset{\circ}{S}_h.$$

From this, in the one-dimensional situation, we were able to deduce that, (1.3.13),

$$\|\theta'\|_{L_\infty(I)} \leq Ch^r.$$

What we shall describe are localized generalizations of this to any number of space dimensions. Or, we can view them as generalizations of the local  $H^1$ -estimate (5.3.43) to  $W_\infty^1$  and  $L_\infty$ . They are proven “essentially” following the scheme in Section 5.3: First, consider “discrete harmonic” functions (or, more generally functions satisfying the discrete analogue of  $Lu = 0$ ,  $L$  a second order elliptic operator). Show that one may bound the  $W_\infty^1(B_0)$ -norm in terms of  $L_\infty(B_d)$ , or  $L_2(B_d)$ , or negative norms, at the expense of inverse powers of  $d$ . (Decidedly nontrivial, now!) Then let  $A(u - u_h, \chi) = 0$ , for  $\chi \in S_h^{comp}(B_d)$ . Use a cut-off function  $\omega$  and consider  $\tilde{u} = \omega u$  and  $\tilde{u}_h$ , a local Ritz-Galerkin projection, this time actually w.r.t. a local Neumann problem (cf. (5.3.22) of seq.). To treat the difference  $\tilde{u} - \tilde{u}_h$  requires so to speak *global*  $L_\infty$ -based estimates for this local problem, cf. (5.4.32) and (5.4.33). Then  $u_h - \tilde{u}_h \in S_h$  is locally “discrete harmonic” and already taken care of. The right hand side functional  $F(\cdot)$  has to be worked in, of course. The details are longish and given in [Schatz and Wahlbin 1994].

Let thus  $\Omega_0$  be a basic domain compactly contained in  $\mathcal{D}$ , and  $\Omega_d$  with  $\Omega_0 \subset\subset \Omega_d \subseteq \mathcal{D}$  where  $d = \text{dist}(\partial\Omega_0, \partial\Omega_d)$ , and  $u$  and  $u_h$  given functions with  $u_h \in S_h$  such that

$$(5.5.1) \quad A(u - u_h, \chi) = F(\chi), \text{ for } \chi \in S_h^{\text{comp}}(\Omega_d).$$

Let with  $\frac{1}{q} + \frac{1}{q'} = 1$ ,

$$(5.5.2) \quad \|v\|_{W_q^{-s}(\Omega)} = \sup_{\substack{\varphi \in C_0^\infty(\Omega) \\ |\varphi|_{W_{q'}^s(\Omega)} = 1}} (v, \varphi)$$

and

$$(5.5.3) \quad |||F|||_{-1, \infty, \Omega_0} = \sup_{\substack{\varphi \in C_0^\infty(\Omega) \\ |\varphi|_{W_1^1(\Omega)} = 1}} F(\varphi).$$

Under various conditions given in [Schatz and Wahlbin 1994] (“typical finite element spaces”, quasi-uniform meshes on  $\Omega_d$ ) we then have for  $d \geq c_0 h$ , with  $c_0$  sufficiently large.

**Theorem 5.5.1.** *There exist constants  $c_0$  and  $C$  such that for  $d \geq c_0 h$ ,*

$$(5.5.4) \quad \begin{aligned} & |u - u_h|_{W_\infty^1(\Omega_0)} + d^{-1} \|u - u_h\|_{L_\infty(\Omega_0)} \\ & \leq C \min_{\chi \in S_h} (|u - \chi|_{W_\infty^1(\Omega_d)} + d^{-1} \|u - \chi\|_{L_\infty(\Omega_d)}) \\ & \quad + C d^{-1-s-n/q} \|u - u_h\|_{W_q^{-s}(\Omega_d)} \\ & \quad + C \ln(d/h) |||F|||_{-1, \infty, \Omega_d}. \end{aligned}$$

**Remark 5.5.1.** Corresponding to Remark 5.3.1, we may replace  $|||F|||_{-1, \infty, \Omega_d}$  in (5.5.4) by

$$\max_{\substack{\chi \in S_h^{\text{comp}}(\Omega_d) \\ |\chi|_{W_1^1(\Omega_d)} = 1}} F(\chi). \quad \square$$

The constants  $C$  and  $c_0$  depend on the finite element spaces and its local properties, such as constants occurring in quasi-uniformity conditions, approximation hypotheses and superapproximation conditions. They also depend on the smoothness properties of the coefficients  $a_{ij}$ ,  $a_i$  and  $a$  (cf. (5.1.1)) and on the ellipticity constant in (5.1.2). Of course, they also depend on  $n$ ,  $s$  and  $q$ . However, they are independent of  $h$ ,  $u$ ,  $u_h$  and  $F$ ; and, no conditions on coercivity, local or global, of the form  $A(v, w)$  are involved (beyond what has already been said about ellipticity for the highest order coefficients  $a_{ij}$ .)

The corresponding result for the error in function values for  $u - u_h$  is as follows. Here we have, if  $r$  is the optimal order of approximation in  $L_p$ -spaces,

$$(5.5.5) \quad \bar{r} = \begin{cases} 1, & \text{if } r = 2, \\ 0, & \text{if } r \geq 3. \end{cases}$$

**Theorem 5.5.2.** *There exist constants  $c_0$  and  $C$  such that for  $d \geq c_0 h$ ,*

$$(5.5.6) \quad \begin{aligned} \|u - u_h\|_{L_\infty(\Omega_0)} &\leq C(\ln d/h)^{\bar{r}} \min_{\chi \in S_h} \|u - \chi\|_{L_\infty(\Omega_d)} \\ &+ Cd^{-s-n/q} \|u - u_h\|_{W_q^{-s}(\Omega_d)} \\ &+ Ch(\ln d/h)^{\bar{r}} |||F|||_{-1,\infty,\Omega_d} \\ &+ C(\ln d/h) |||F|||_{-2,\infty,\Omega_d}. \end{aligned}$$

Here, in addition to the definitions in (5.5.2) and (5.5.3),

$$(5.5.7) \quad |||F|||_{-2,\infty,\Omega} = \sup_{\substack{\varphi \in C_0^\infty(\Omega) \\ |\varphi|_{W_1^2(\Omega)}=1}} F(\varphi).$$

For the necessity of the logarithmic factor  $(\ln d/h)^{\bar{r}}$  in the case of  $r = 2$  in (5.5.6), we refer to [Haverkamp 1984]. And, for the history of Theorems 5.5.1 and 5.5.2, see the discussion in [Schatz and Wahlbin 1994].

## Chapter 6. Superconvergence in tensor–product elements.

### 6.1. Introduction.

We give some results of the general nature that superconvergence points which are present in the case of one-dimensional two-point boundary value problems automatically give rise to superconvergence points in a multi-dimensional tensor product situation. We shall give details only in the two-dimensional case; generalizations to arbitrary  $n$  will be fairly obvious. The general principle was applied to  $L_2$ -projections in Section 4.2.

Our investigation is limited in that we shall not treat the general case of operators of the form (5.1.1). For superconvergence in derivatives we restrict ourselves to the case of

$$(6.1.1) \quad -\operatorname{div}(\alpha(x, y)\nabla u) - \sum_{i=1}^n \frac{\partial}{\partial x_i}(a_i(x, y)u) + a(x, y)u = f,$$

with  $\alpha$  a scalar smooth uniformly positive function,  $\alpha \geq c_{ell} > 0$ , and with corresponding bilinear form

$$(6.1.2) \quad A(v, w) = \int \left( \alpha \nabla v \cdot \nabla w + \sum_{i=1}^n a_i v \frac{\partial w}{\partial x_i} + avw \right) dx dy.$$

For superconvergence in function values we shall only treat the case of the Laplacian,  $\alpha \equiv 1$ ,  $a_i \equiv a \equiv 0$ .

Our general starting point is that we have  $u$  and  $u_h \in S_h(\Omega_1)$  such that

$$(6.1.3) \quad A(u - u_h, \chi) = 0, \text{ for } \chi \in S_h^{\text{comp}}(\Omega_1).$$

We shall then investigate superconvergence points for  $u - u_h$  in  $\Omega_0 \subset \subset \Omega_1$ . For simplicity, we shall only consider the case when  $\operatorname{dist}(\partial\Omega_0, \partial\Omega_1)$  is of unit magnitude. The essential ideas in this chapter are those of [Douglas, Dupont and Wheeler 1974b], combined with those of Theorems 5.5.1 and 5.5.2.

### 6.2. Superconvergence in derivatives for the case of the Laplacian.

Let  $\Omega_0 = I_0^x \times I_0^y$  be a rectangle and  $\Omega_1 = I_1^x \times I_1^y$  another rectangle containing  $\Omega_0$  with  $\operatorname{dist}(\partial\Omega_0, \partial\Omega_1) = 1$ . We assume that on  $\Omega_1$ ,  $S_h(\Omega_1)$  consists of linear combinations of  $\chi^x(x)\chi^y(y)$  where  $\chi^x$  and  $\chi^y$  belong to one-dimensional finite element spaces  $S_h^x(I_1^x)$  and  $S_h^y(I_1^y)$ . We also assume that  $\Omega_1$  is adjusted so that the endpoints of  $I_1^x$  and  $I_1^y$  coincide with meshpoints. We write in common notation in the field (it differs from the “tensor–product” ideas of Algebra),

$$(6.2.1) \quad S_h(\Omega_1) = S_h^x(I_1^x) \otimes S_h^y(I_1^y).$$

We shall follow here the ideas of [Douglas, Dupont and Wheeler 1974b]. Our starting point is a function  $u$  and  $u_h \in S_h$  such that

$$(6.2.2) \quad D(u - u_h, \chi) \equiv \iint \left( \frac{\partial(u - u_h)}{\partial x} \frac{\partial \chi}{\partial x} + \frac{\partial(u - u_h)}{\partial y} \frac{\partial \chi}{\partial y} \right) dx dy = 0,$$

for  $\chi \in S_h^{\text{comp}}(\Omega_1)$ .

Let now  $\omega$  be a smooth cut-off function with  $\text{supp}(\omega) \subset \Omega_1$  and  $\omega \equiv 1$  on  $\Omega_{1/2}$  where  $\Omega_0 \subset\subset \Omega_{1/2} \subset\subset \Omega_1$  with fixed separation. Let

$$(6.2.3) \quad \tilde{u} := \omega u.$$

We let  $\overset{\circ}{S}_h^x(I_1^x)$  and  $\overset{\circ}{S}_h^y(I_1^y)$  be the subspaces vanishing at the endpoints and let  $R^x$  and  $R^y$  denote the one dimensional projections into them: E.g.,  $R^x v$  is given by

$$(6.2.4) \quad \int_{I_1^x} (R^x v)' \chi' = \int_{I_1^x} v' \chi', \quad \text{for } \chi \in \overset{\circ}{S}_h^x(I_1^x).$$

We then define

$$(6.2.5) \quad W_h := R^x \otimes R^y(\tilde{u}),$$

i.e., we first take the  $y$ -projection to obtain  $R^y(\tilde{u}(x, \cdot))(y)$  for each fixed  $x$ , and then the  $x$ -projection of that for each fixed  $y$  (or, vice versa). Clearly, due to the tensor-product structure,  $W_h \in \overset{\circ}{S}_h(\Omega_1)$ . We shall now consider  $u_h - W_h$ . For  $\chi \in S_h^{\text{comp}}(\Omega_{1/2})$ , since  $\omega \equiv 1$  on  $\Omega_{1/2}$ , we have using (6.2.4) and its counterpart for  $y$ ,

$$\begin{aligned} (6.2.6) \quad D(u_h - W_h, \chi) &= D(\tilde{u} - W_h, \chi) \\ &= \iint \frac{\partial}{\partial x}(\tilde{u} - W_h) \frac{\partial}{\partial x} \chi + \frac{\partial}{\partial y}(\tilde{u} - W_h) \frac{\partial}{\partial y} \chi \\ &= \iint \frac{\partial}{\partial x}(I - R^x \otimes R^y)(\tilde{u}) \frac{\partial \chi}{\partial x} + \frac{\partial}{\partial y}(I - R^x \otimes R^y)(\tilde{u}) \frac{\partial \chi}{\partial y} \\ &= \int_{I_1^y} dy \left( \int_{I_1^x} \frac{\partial}{\partial x}(I - R^x \otimes R^y)(\tilde{u}) \frac{\partial \chi}{\partial x} dx \right) \\ &\quad + \int_{I_1^x} dx \left( \int_{I_1^y} \frac{\partial}{\partial y}(I - R^x \otimes R^y)(\tilde{u}) \frac{\partial \chi}{\partial y} dy \right) \\ &= \int_{I_1^y} dy \left( \int_{I_1^x} \frac{\partial}{\partial x}(I - I^x \otimes R^y)(\tilde{u}) \frac{\partial \chi}{\partial x} dx \right) \\ &\quad + \int_{I_1^x} dx \left( \int_{I_1^y} \frac{\partial}{\partial y}(I - R^x \otimes I^y)(\tilde{u}) \frac{\partial \chi}{\partial y} dy \right) \\ &= \iint (I - I^x \otimes R^y) \left( \frac{\partial \tilde{u}}{\partial x} \right) \frac{\partial \chi}{\partial x} + \iint (I - R^x \otimes I^y) \left( \frac{\partial \tilde{u}}{\partial y} \right) \frac{\partial \chi}{\partial y} \\ &\equiv F(\chi), \quad \text{for } \chi \in S_h^{\text{comp}}(\Omega_{1/2}). \end{aligned}$$

Now, for  $\varphi \in C_0^\infty(\Omega_{1/2})$ ,

$$\begin{aligned} (6.2.7) \quad & \left| \iint (I - I^x \otimes R^y) \left( \frac{\partial \tilde{u}}{\partial x} \right) \frac{\partial \varphi}{\partial x} dxdy \right| \\ &= \left| \int_{I_1^x} dx \left( \int_{I_1^y} I^x \otimes (I^y - R^y) \left( \frac{\partial \tilde{u}}{\partial x} \right)(x, y) \frac{\partial \varphi}{\partial x}(x, y) dy \right) \right| \\ &\leq \max_x \left\| (I^y - R^y) \left( \frac{\partial \tilde{u}}{\partial x} \right)(x, \cdot) \right\|_{L_\infty(I^y)} \int_{I_1^x} \int_{I_1^y} \left| \frac{\partial \varphi}{\partial x}(x, y) \right| dxdy. \end{aligned}$$

We next make the assumption, if you like, that

$$(6.2.8) \quad \|(I^y - R^y)v\|_{L_\infty(I_1^y)} \leq Ch^r \|v\|_{W_\infty^r(I_1^y)},$$

so that from (6.1.9) and the symmetric argument for the second term in  $F(\cdot)$ ,

$$(6.2.9) \quad |||F|||_{-1,\infty,\Omega_{1/2}} \leq Ch^r \|u\|_{W_\infty^{r+1}(\Omega_{1/2})}.$$

Thus,  $D(u_h - W_h, \chi) = F(\chi)$ , for  $\chi \in S_h^{comp}(\Omega_{1/2})$  with  $F(\cdot)$  satisfying (6.2.9). We may then use Theorem 5.5.1 (with “ $u = 0$ ”) and find that

$$(6.2.10) \quad \|u_h - W_h\|_{W_\infty^1(\Omega_0)} \leq C\|u_h - W_h\|_{W_q^{-s}(\Omega_{1/2})} + C(\ln 1/h)h^r.$$

Hence, by use of the triangle inequality and that  $\omega \equiv 1$  on  $\Omega_{1/2}$ , we have for any point  $\bar{x}, \bar{y} \in \Omega_0$ , with  $\partial = \partial/\partial x$  or  $\partial/\partial y$ ,

$$\begin{aligned} (6.2.11) \quad & |\partial(u - u_h)(\bar{x}, \bar{y})| \leq |\partial(\tilde{u} - W_h)(\bar{x}, \bar{y})| + \|W_h - u_h\|_{W_\infty^1(\Omega_0)} \\ & \leq |\partial(\tilde{u} - W_h)(\bar{x}, \bar{y})| + C\|W_h - u_h\|_{W_q^{-s}(\Omega_{1/2})} + C(\ln 1/h)h^r \\ & \leq |\partial(\tilde{u} - W_h)(\bar{x}, \bar{y})| + C\|u - u_h\|_{W_q^{-s}(\Omega_{1/2})} \\ & \quad + C\|\tilde{u} - W_h\|_{W_q^{-s}(\Omega_{1/2})} + C(\ln 1/h)h^r \\ & \leq |\partial(\tilde{u} - W_h)(\bar{x}, \bar{y})| + C\|\tilde{u} - W_h\|_{L_\infty(\Omega_{1/2})} + C(\ln 1/h)h^r \\ & \quad + C\|u - u_h\|_{W_q^{-s}(\Omega_{1/2})}. \end{aligned}$$

We now observe that

$$\begin{aligned} (6.2.12) \quad & \tilde{u} - W_h = (I^x \otimes I^y - R^x \otimes R^y)(\tilde{u}) \\ & = (I^x - R^x) \otimes I^y(\tilde{u}) + I^x \otimes (I^y - R^y)(\tilde{u}) \\ & \quad - (I^x - R^x) \otimes (I^y - R^y)(\tilde{u}). \end{aligned}$$

Under appropriate conditions on the one-dimensional projections (cf. Chapter 1) we have for  $u$  smooth,

$$(6.2.13) \quad \|\tilde{u} - W_h\|_{L_\infty(\Omega_{1/2})} \leq Ch^r \|u\|_{W_\infty^r(\Omega_1)}.$$

Thus from (6.2.11),

$$\begin{aligned} (6.2.14) \quad & |\partial(u - u_h)(\bar{x}, \bar{y})| \\ & \leq |\partial(\tilde{u} - W_h)(\bar{x}, \bar{y})| + C(\ln 1/h)h^r + C\|u - u_h\|_{W_q^{-s}(\Omega_{1/2})}. \end{aligned}$$

We need two more ingredients to deduce that  $(\bar{x}, \bar{y})$  is a superconvergence point for  $\partial$ . First, we need an estimate for  $u - u_h$  in any negative norm. We will discuss this in the next section and for the moment we simply assume that for some  $s, q$ ,

$$(6.2.15) \quad \|u - u_h\|_{W_q^{-s}(\Omega_{1/2})} \leq C(\ln 1/h)h^r.$$

Then

$$(6.2.16) \quad |\partial(u - u_h)(\bar{x}, \bar{y})| \leq |\partial(\tilde{u} - W_h)(\bar{x}, \bar{y})| + C(\ln 1/h)h^r.$$

We now come to the heart of the matter! Going back to (6.1.14) and taking for definiteness  $\partial = \frac{\partial}{\partial x}$ , we have

$$(6.2.17) \quad \begin{aligned} \frac{\partial}{\partial x}(\tilde{u} - W_h) &= \frac{\partial}{\partial x}(I^x - R^x) \otimes I^y(\tilde{u}) + I^x \otimes (I^y - R^y)\left(\frac{\partial \tilde{u}}{\partial x}\right) \\ &\quad - \left(\frac{\partial}{\partial x}(I^x - R^x)\right) \otimes (I^y - R^y)(\tilde{u}) \\ &\equiv I_1 + I_2 + I_3. \end{aligned}$$

Here, again under assumptions on appropriate behavior of the one-dimensional projections,

$$(6.2.18) \quad |I_2| + |I_3| \leq Ch^r \|u\|_{W_\infty^{r+1}(\Omega_1)}.$$

We conclude that if  $\bar{x}$  is a superconvergence point for the first derivative in the one-dimensional projection  $R^x$  of order  $\sigma$ , then  $(\bar{x}, y)$ , for any  $y$ , if it belongs to  $\Omega_0$ , is a superconvergence point for  $\frac{\partial}{\partial x}(u - u_h)$  of order  $\bar{\sigma} = \min(\sigma, 1)$  (modulo a  $\ln 1/h$ -factor).

We collect the above and its obvious consequence in a somewhat informally stated theorem:

**Theorem 6.2.1.** *Assume that the one-dimensional projections  $R^x$  and  $R^y$  are stable in  $L_\infty$  and  $W_\infty^1$  and that  $\dot{S}_h^x$  and  $\dot{S}_h^y$  have approximation to orders  $r$  and  $r-1$ , for function values and derivatives, respectively. Assume also that  $S_h(\Omega_1)$  is the tensor product  $S_h(I^x) \otimes S_h(I^y)$ . Assume further that for  $u$  and  $u_h$  satisfying*

$$(6.2.19) \quad D(u - u_h, \chi) = 0, \text{ for } \chi \in S_h^{\text{comp}}(\Omega_1),$$

*we have for some  $s$  and  $q$ ,*

$$(6.2.20) \quad \|u - u_h\|_{W_q^{-s}(\Omega_1)} \leq C(\ln 1/h)h^r.$$

*Then:*

a) if  $\bar{x} \in I_0^x$  is a superconvergence point of order  $\sigma$  for the first derivative for  $R^x$ , then for any  $(\bar{x}, y) \in \Omega_0$ ,

$$(6.2.21) \quad \left| \frac{\partial}{\partial x}(u - u_h)(\bar{x}, y) \right| \leq C(\ln 1/h)h^{r-1+\bar{\sigma}}, \quad \bar{\sigma} = \min(1, \sigma).$$

b) if  $\bar{y} \in I_0^y$  is a superconvergence point of order  $\sigma$  for the first derivative of  $R^y$ , then for any  $(x, \bar{y}) \in \Omega_0$ ,

$$(6.2.22) \quad \left| \frac{\partial}{\partial y}(u - u_h)(x, \bar{y}) \right| \leq C(\ln 1/h)h^{r-1+\bar{\sigma}}, \quad \bar{\sigma} = \min(1, \sigma).$$

c) If  $\bar{x} \in I_0^x$  and  $\bar{y} \in I_0^y$  are similarly superconvergence points for first derivatives of order  $\sigma_x, \sigma_y$ , respectively, then

$$(6.2.23) \quad |\nabla(u - u_h)(\bar{x}, \bar{y})| \leq C(\ln 1/h)h^{r-1+\bar{\sigma}}, \quad \bar{\sigma} = \min(1, \sigma_x, \sigma_y).$$

Remark 6.2.1. Clearly, if

$$(6.2.24) \quad \|u - u_h\|_{W_q^{-s}(\Omega_1)} \leq Ch^{r-1+\tau}$$

with  $\tau > 0$ , corresponding superconvergence results hold but with  $\bar{\sigma}$  replaced by  $\min(\bar{\sigma}, \tau)$ .  $\square$

### 6.3. Negative norm estimates for $u - u_h$ : Examples.

The term  $\|u - u_h\|_{W_q^{-s}(\Omega_1)}$  in the previous section is the only one that takes into account what happens outside of  $\Omega_1$ : all other considerations are local to  $\Omega_1$ . We shall give four brief examples of how to estimate it. For future purposes we shall also give estimates of higher order than immediately necessary in the previous section. Note that with our definition,  $\|v\|_{W_q^{-s}(\Omega_1)} \leq \|v\|_{W_q^{-s}(\Omega_2)}$  for  $\Omega_1 \subseteq \Omega_2$ .

Example 6.3.1. Homogeneous Neumann problems in smooth domains in  $R^n$ .

In Section 5.2 we considered a model Neumann problem,  $-\Delta u + u = f$  in  $\mathcal{D}$ ,  $\partial u / \partial n = 0$  on  $\partial\mathcal{D}$ , with  $f$  and  $\partial\mathcal{D}$  smooth. Assuming a perfect match for elements meeting  $\partial\mathcal{D}$ , we showed that

$$(6.3.1) \quad \|u - u_h\|_{W_q^{-(r-2)}(\mathcal{D})} \leq Ch^{2r-2}q, \text{ for } q \geq 2 \text{ (say).}$$

The argument easily extends to general operators of the form (5.1.1) with conormal derivative conditions on  $\partial\mathcal{D}$ , provided the problem has uniqueness (and smooth coefficients).  $\square$

Example 6.3.2. Dirichlet problems in smooth domains in  $R^n$ .

Here we consider (5.1.1) with homogeneous Dirichlet boundary conditions,  $u = 0$  on  $\partial\mathcal{D}$ , and smooth data. We impose these essential boundary conditions so that  $\overset{\circ}{S}_h(\mathcal{D}_h) \subseteq \overset{\circ}{H}^1(\mathcal{D})$ , say. Using isoparametric elements at the boundary which approximate it to order  $h^r$ , it was shown in [Schatz and Wahlbin 1982] that

$$(6.3.2) \quad \|u - u_h\|_{L_\infty(\mathcal{D})} \leq C(\ln 1/h)^{\bar{r}} h^r.$$

This estimate suffices for the purposes of Section 6.2.

For the Laplacian, an easy argument involving the maximum principle will convince the reader that it is hopeless in general to try for higher order accuracy in any negative norm for isoparametric elements (if  $\mathcal{D}_h \subseteq \mathcal{D}$ , say). Note in this connection that in general for isoparametrics it is not true that

$$(6.3.3) \quad \min_{\chi \in \overset{\circ}{S}_h(\mathcal{D})} \|v - \chi\|_{H^1(\mathcal{D})} \leq Ch^{r-1} \|v\|_{H^r(\mathcal{D})},$$

except in the case of piecewise linear,  $r = 2$ .  $\square$

If one approximates the smooth boundary to order  $O(h^{2r-2})$ , e.g., by use of “superparametric” modifications at the boundary, one obtains essentially (6.3.1) also in this case. The argument is the same.  $\square$

The final two problems are more special.

Example 6.3.3. Dirichlet problems in smooth plane domains.

In [Scott 1975] a special way of treating Dirichlet problems in plane domains without use of isoparametric or superparametric modification at the boundary was given and shown to satisfy

$$(6.3.4) \quad \|u - u_h\|_{W_2^{-(r-2)}(\mathcal{D})} \leq Ch^{2r-2}. \quad \square$$

Example 6.3.4. A Dirichlet problem in a plane polygonal domain.

Consider the problem

$$(6.3.5) \quad \begin{aligned} -\Delta u &= f && \text{in } \mathcal{D}, \\ u &= 0 && \text{on } \partial\mathcal{D}, \end{aligned}$$

where  $\mathcal{D}$  is a plane *polygonal* domain. As is well known, even if  $f$  is smooth ( $f \equiv 1$  is the classical torsion engineering problem), the solution  $u$  has singularities at corners of the domain. E.g., even if  $f \equiv 0$  near a corner of interior angle  $\alpha$ , the solution behaves like

$$(6.3.6) \quad u(r, \theta) = \text{const. } r^{\pi/\alpha} \sin(\theta\pi/\alpha) + \dots,$$

expressed in polar coordinates at the corner. Thus, although we have no trouble fitting the boundary exactly here, the relevant a priori estimates for a duality argument may be lacking: In fact, if  $\alpha$  is the maximum angle, then  $O(h^{2\pi/\alpha})$  is the best we can get in *any negative norm* if we use globally quasi-uniform elements and the constant in (6.3.6) is nonzero, see [Wahlbin 1984]. However, the exact behavior indicated in (6.3.6) is known, see [Grisvard 1985, Theorem 5.1.3.5], and if one uses suitable mesh refinements towards the corners, cf. e.g. [Babuška 1970], one obtains

$$(6.3.7) \quad \min_{\chi \in \overset{\circ}{S}_h(\mathcal{D})} \|u - \chi\|_{H^1(\mathcal{D})} \leq Ch^{r-1} \|f\|_{H^{r-2}(\mathcal{D})}.$$

A standard duality argument *then* gives

$$(6.3.8) \quad \|u - u_h\|_{W_2^{-(r-2)}(\mathcal{D})} \leq Ch^{2r-2}.$$

If one thus assumes that the patch of tensor-product elements considered in Section 6.2 sits away from corners and has quasi-uniform partitions, the theory there (and that to come) will apply.  $\square$

#### 6.4. Superconvergence in derivatives for the case of (6.1.1).

Recall that in (6.1.1) we imposed that the highest order term in the elliptic operator is of the form  $-\text{div}(\alpha(x, y)\nabla u)$ , with  $\alpha$  a scalar function. Again following essentially [Douglas, Dupont and Wheeler 1974b] and Section 5.5, or, viewed alternatively, extending our one-dimensional results of Section 1.3 to several dimensions, we shall show that the results of Section 6.2 hold for a general form of the type (6.1.2). We shall introduce the necessary superapproximation hypothesis as we go along.

As in Section 6.2 we let  $\Omega_0 \subset\subset \Omega_{1/2} \subset\subset \Omega_1$  be rectangles with  $S_h(\Omega_1)$  having a tensor-product structure. Again,  $\omega \equiv 1$  on  $\Omega_{1/2}$  and has compact support in  $\Omega_1$ .

This time, we define a comparison function  $Z_h \in \overset{\circ}{S}_h(\Omega_1)$  by

$$(6.4.1) \quad D(\tilde{u} - Z_h, \chi) = 0, \text{ for } \chi \in \overset{\circ}{S}_h(\Omega_1).$$

Now, for  $\chi \in S_h^{comp}(\Omega_{1/4})$  (what  $\Omega_{1/4}$  should be is obvious) and  $\psi \in S_h^{comp}(\Omega_{1/2})$ ,

$$\begin{aligned}
(6.4.2) \quad D(Z_h - u_h, \chi) &= D(u - u_h, \chi) \\
&= \int \nabla(u - u_h) \cdot \nabla \chi \\
&= \int \alpha \nabla(u - u_h) \cdot \left( \frac{1}{\alpha} \nabla \chi \right) \\
&= \int \alpha \nabla(u - u_h) \cdot \nabla \left( \frac{1}{\alpha} \chi \right) + \int \nabla(u - u_h) \cdot \frac{1}{\alpha} (\nabla \alpha) \chi \\
&= \int \alpha \nabla(u - u_h) \nabla \left( \frac{1}{\alpha} \chi - \psi \right) - \int \sum_{i=1}^n a_i(u - u_h) \frac{\partial \psi}{\partial x_i} \\
&\quad - \int a(u - u_h) \psi - \int (u - u_h) \operatorname{div} \left( \frac{1}{\alpha} (\nabla \alpha) \chi \right),
\end{aligned}$$

since  $A(u - u_h, \psi) = 0$ .

We now make the superapproximation hypothesis that we can find  $\psi \in S_h^{comp}(\Omega_{1/2})$  such that

$$(6.4.3) \quad \left\| \frac{1}{\alpha} \chi - \psi \right\|_{W_1^1(\Omega_{1/2})} \leq Ch \|\chi\|_{W_1^1(\Omega_{1/2})}, \text{ for } \chi \in S_h^{comp}(\Omega_{1/4}).$$

We then have

$$(6.4.4) \quad D(Z_h - u_h, \chi) = F(\chi), \text{ for } \chi \in S_h^{comp}(\Omega_{1/4})$$

where

$$(6.4.5) \quad \sup_{\substack{\chi \in S_h^{comp}(\Omega_{1/4}) \\ |\chi|_{W_1^1(\Omega_{1/4})}=1}} F(\chi) \leq C \left( h \|u - u_h\|_{W_\infty^1(\Omega_{1/2})} + \|u - u_h\|_{L_\infty(\Omega_{1/2})} \right).$$

Thus, from Theorem 5.5.1, cf. Remark 5.5.1,

$$\begin{aligned}
(6.4.6) \quad &\|Z_h - u_h\|_{W_\infty^1(\Omega_0)} \\
&\leq C(\ln 1/h) \left( h \|u - u_h\|_{W_\infty^1(\Omega_{1/2})} + \|u - u_h\|_{L_\infty(\Omega_{1/2})} \right) \\
&\quad + C \|Z_h - u_h\|_{W_q^{-s}(\Omega_{1/2})} \\
&\leq C(\ln 1/h) \left( h \|u - u_h\|_{W_\infty^1(\Omega_{1/2})} + \|u - u_h\|_{L_\infty(\Omega_{1/2})} \right) \\
&\quad + C \|Z_h - \tilde{u}\|_{W_q^{-s}(\Omega_{1/2})} + C \|u - u_h\|_{W_q^{-s}(\Omega_{1/2})}.
\end{aligned}$$

We next attack the negative norm term involving  $Z_h - \tilde{u}$ . We have  $H^2$ -regularity for the Laplace operator on a rectangle, i.e., if

$$\begin{aligned}
(6.4.7) \quad &-\Delta u = f \quad \text{in } \Omega_1, \\
&u = 0 \quad \text{on } \partial\Omega_1
\end{aligned}$$

then

$$(6.4.8) \quad \|u\|_{H^2(\Omega_1)} \leq C \|f\|_{L_2(\Omega_1)}.$$

A standard duality argument then gives (under natural assumptions)

$$(6.4.9) \quad \|Z_h - \tilde{u}\|_{L_2(\Omega_1)} \leq Ch^r,$$

for  $u$  smooth enough.

Further, by Theorems 5.5.1 and 5.5.2, with  $F \equiv 0$ , we have since  $A(u - u_h, \chi) = 0$ , for  $\chi \in S_h^{comp}(\Omega_1)$ ,

$$\begin{aligned} (6.4.10) \quad & h\|u - u_h\|_{W_\infty^1(\Omega_{1/2})} + \|u - u_h\|_{L_\infty(\Omega_{1/2})} \\ & \leq C(\ln 1/h) \min_{\chi \in S_h} \left( h\|u - \chi\|_{W_\infty^1(\Omega_1)} + \|u - \chi\|_{L_\infty(\Omega_1)} \right) \\ & \quad + C\|u - u_h\|_{W_q^{-s}(\Omega_1)} \\ & \leq Ch^r(\ln 1/h) + C\|u - u_h\|_{W_q^{-s}(\Omega_1)}, \end{aligned}$$

again under natural approximation hypotheses.

From (6.4.6), (6.4.9) and (6.4.10) then,

$$(6.4.11) \quad \|Z_h - u_h\|_{W_\infty^1(\Omega_0)} \leq Ch^r(\ln 1/h)^2 + C\|u - u_h\|_{W_q^{-s}(\Omega_1)}.$$

Thus, with  $\partial = \partial/\partial x$  or  $\partial/\partial y$ ,

$$\begin{aligned} (6.4.12) \quad & |\partial(u - u_h)(\bar{x}, \bar{y})| \leq |\partial(\tilde{u} - Z_h)(\bar{x}, \bar{y})| + \|Z_h - u_h\|_{W_\infty^1(\Omega_0)} \\ & \leq |\partial(\tilde{u} - Z_h)(\bar{x}, \bar{y})| + Ch^r(\ln 1/h)^2 + C\|u - u_h\|_{W_q^{-s}(\Omega_1)}. \end{aligned}$$

Hence, if we have a suitable negative norm estimate for  $u - u_h$ , apart from an additional logarithmic factor, the conclusions of Theorem 6.2.1 given there for the Laplacian remain unchanged for the more general bilinear form (6.1.2)  $A(\cdot, \cdot)$  in (6.1.3).

## 6.5. Superconvergence in function values for the Laplacian and $r \geq 3$ .

Provided  $r \geq 3$  one similarly obtains that superconvergence for function values on tensor product meshes is inherited from the one-dimensional case, if the underlying partial differential operator is the Laplacian. We shall merely point out where modifications are necessary in Section 6.2.

For  $F(\cdot)$  in (6.2.6), one now also needs an estimate for  $\|F\|_{-2,\infty,\Omega_{1/2}}$ . This will be inherited from  $W_\infty^{-1}$ -error estimates for the one-dimensional cases, and this is why  $r \geq 3$  is required to get

$$(6.5.1) \quad \|F\|_{-2,\infty,\Omega_{1/2}} \leq Ch^{r+1}.$$

Of course one then applies Theorem 5.5.2 to obtain, corresponding to (6.2.10),

$$(6.5.2) \quad \|u_h - W_h\|_{L_\infty(\Omega_0)} \leq C\|u_h - W_h\|_{W_q^{-s}(\Omega_{1/2})} + C(\ln 1/h)h^{r+1}.$$

Corresponding to (6.2.11) one then has

$$\begin{aligned} (6.5.3) \quad & |(u - u_h)(\bar{x}, \bar{y})| \leq C|(\tilde{u} - W_h)(\bar{x}, \bar{y})| + C\|\tilde{u} - W_h\|_{W_q^{-s}(\Omega_{1/2})} \\ & \quad + C(\ln 1/h)h^{r+1} + C\|u - u_h\|_{W_q^{-s}(\Omega_{1/2})}. \end{aligned}$$

Again, from (6.2.12), one easily shows for  $r \geq 3$ , that  $\tilde{u} - W_h$  is bounded by  $Ch^{r+1}$  in negative norms. Thus, if also  $u - u_h$  is so bounded in some negative norm,

$$(6.5.4) \quad |(u - u_h)(\bar{x}, \bar{y})| \leq C|(\tilde{u} - W_h)(\bar{x}, \bar{y})| + C(\ln 1/h)h^{r+1}.$$

Finally, as in (6.2.17) et seq. we now see that  $u - u_h$  inherits the one-dimensional superconvergence points from  $W_h = R^x \otimes R^y(\tilde{u})$ .

## Chapter 7. Superconvergence by local symmetry.

The present chapter basically reviews [Schatz, Sloan and Wahlbin 1994].

### 7.1. Introduction.

As with tensor-product elements, we have already met the main idea in the simpler case of  $L_2$ -projections in Chapter 4 (Section 4.3, in which the reader may find illuminating pictures). We have also met the idea that symmetry implies superconvergence in the one-dimensional case of two-point boundary value problems in Sections 1.6 and 1.9. We now place ourselves in the situation of the general bilinear form in (5.1.3) and assume that

$$(7.1.1) \quad A(u - u_h, \chi) = 0, \text{ for } \chi \in S_h^{\text{comp}}(B_d),$$

where  $B_d = B_d(x_0) \subseteq \mathcal{D}$  is a ball of radius  $d = d(h)$  around the point(s)  $x_0 = x_0(h)$ , which in general will vary with  $h$ .

Our symmetry assumption is that if  $\chi \in S_h(B_d)$ , then

$$(7.1.2) \quad \bar{\chi}(x) \equiv \chi(x_0 - (x - x_0)) \in S_h(B_d).$$

We shall also require that

$$(7.1.3) \quad \min_{\chi \in S_h} \left( \|v - \chi\|_{L_\infty(B_{d/2})} + h|v - \chi|_{W_\infty^1(B_{d/2})} \right) \leq Ch^r |v|_{W_\infty^r(B_d)},$$

in which now a crucial point is that the  $W_\infty^r$ -seminorm on the right involves only the pure  $r^{\text{th}}$  derivatives of  $v$ .

Let us point out that the symmetry assumption (7.1.2) refers only to the finite element spaces. Symmetry is not required of the coefficients of the underlying second order partial differential operator in (5.1.3).

Remark 7.1.1. Anticipating the results below, we shall show that superconvergence at symmetry points occurs as follows:

for  $r$  odd, in function values,

for  $r$  even, in gradients.

Going back to Chapter 1 and, say, Hermite cubics in the uniform mesh case ( $r = 4$ ), we there have superconvergence for first derivatives (on interior subintervals) at the mesh and midpoints, i.e., the local symmetry points. However, for function values, the superconvergence points are not symmetry points (see Example 1.8.1, eq. (1.8.16)). Thus, the dichotomy between odd and even  $r$  is real.  $\square$

### 7.2. The case of a symmetric form with constant coefficients.

To bring out the central *idea* in this chapter, we first consider the case of the Laplacian, i.e.,

$$(7.2.1) \quad D(u - u_h, \chi) \equiv \int \sum_{i=1}^n \frac{\partial(u - u_h)}{\partial x_i} \frac{\partial \chi}{\partial x_i} = 0, \text{ for } \chi \in S_h^{\text{comp}}(B_d),$$

where the underlying partial differential operator is also “symmetric” with respect to the antipodal map about  $x_0$ . A change of variables via the antipodal map then establishes that

$$(7.2.2) \quad D(\bar{u} - \bar{u}_h, \chi) = D(u - u_h, \bar{\chi}) = 0, \text{ for } \chi \in S_h^{\text{comp}}(B_d),$$

since  $\bar{x} \in S_h^{comp}(B_d)$ . (The same conclusion clearly applies if the  $a_{ij}$  are constant,  $a_i \equiv 0$  and  $a$  is constant.)

We now separate the argument according to whether  $r$  is odd or even.

### 7.2.a. Odd $r$ .

Let  $e = u - u_h$  and

$$(7.2.3) \quad e_{even}(x) = (e(x) + \bar{e}(x))/2;$$

note that  $e_{even} = u_{even} + u_{h,even}$  where  $u_{h,even} \in S_h(B_d)$  by the symmetry assumption (7.1.2). Thus,

$$(7.2.4) \quad D(e_{even}, \chi) = 0, \text{ for } \chi \in S_h^{comp}(B_d).$$

We then apply Theorem 5.5.2 and the approximation assumption (7.1.3) to obtain that, provided  $d \geq c_0 h$ ,

$$\begin{aligned} (7.2.5) \quad |e_{even}(x_0)| &\leq C \min_{\chi \in S_h} \|u_{even} - \chi\|_{L_\infty(B_{d/2})} \\ &\quad + Cd^{-s-n/q} \|e_{even}\|_{W_q^{-s}(B_{d/2})} \\ &\leq Ch^r |u_{even}|_{W_\infty^r(B_d)} + Cd^{-s-n/q} \|u - u_h\|_{W_q^{-s}(\mathcal{D})}. \end{aligned}$$

We now assume that we have a suitable negative norm estimate,

$$(7.2.6) \quad \|u - u_h\|_{W_q^{-s}(\mathcal{D})} \leq Ch^{r+\tau},$$

for some  $\tau > 0$ . Such estimates were discussed in Section 6.3. Using then that  $e_{even}(x_0) = (u - u_h)(x_0)$  and Taylor's theorem, we have if  $u \in W_\infty^{r+1}(B_d)$ ,

$$(7.2.7) \quad |(u - u_h)(x_0)| \leq Ch^r d + Cd^{-s-n/q} h^{r+\tau}.$$

So far we haven't nailed down how  $d$  may depend on  $h$ . The most favorable case (from the above analysis!) is clearly to take (asymptotically),

$$(7.2.8) \quad h^r d = d^{-s-n/q} h^{r+\tau},$$

i.e.,

$$(7.2.9) \quad d = h^\sigma, \text{ with } \sigma = \tau/(s + n/q + 1).$$

We then have:

**Theorem 7.2.1.** *Assuming that  $r$  is odd, (7.2.1), the negative norm estimate (7.2.6), the symmetry assumption (7.1.2) with (7.2.9) where  $\sigma \leq 1$ , that Theorem 5.5.2 applies, the approximation assumption (7.1.3), and that  $u \in W_\infty^{r+1}(B_d)$ , we have*

$$(7.2.10) \quad |(u - u_h)(x_0)| \leq Ch^{r+\sigma}.$$

Let us remark that superconvergence of lower orders than  $\sigma$  above results if the symmetry assumption (7.1.2) is satisfied on balls  $B_d$  of radii smaller than those required in (7.2.9).

Example 7.2.1. The most “favorable” negative norm estimates we can reasonably expect occur in Example 6.3.1, or, Example 6.3.2 with superparametric modifications near the smooth boundary. Then

$$\|u - u_h\|_{W_q^{-(r-2)}(\mathcal{D})} \leq Ch^{2r-2}q.$$

We take  $q = \ln 1/h$ ,  $s = r - 2$ , ( $\tau \simeq r - 2$ ), or, more precisely,

$$d = h^\sigma, \quad \sigma = (r-2)/(r-1) = 1 - 1/(r-1),$$

and obtain from (7.2.5) and (7.2.7) that

$$|(u - u_h)(x_0)| \leq Ch^{r+1-1/(r-1)} \ln 1/h.$$

For  $r$  large we thus have superconvergence of close to a full extra order, while requiring symmetry only in a ball of very small radius,  $h^{1-1/(r-1)}$ , around  $x_0$ .  $\square$

Remark 7.2.1. In the case that the mesh is symmetric around  $x_0$  in an  $O(1)$  neighborhood, a recent argument, [Schatz 1995], shows that we have (7.2.10) with  $\sigma = 1$ . This remark also applies to Sections 7.2.b and 7.3 below.  $\square$

### 7.2.b. Even $r$ .

We shall assert under conditions similar to the ones above for odd  $r$ , that, now for even  $r$ ,  $\nabla u_h$  is a superconvergent approximation to  $\nabla u$  at the symmetry point  $x_0$ . In general, if the finite element spaces are merely continuous (which is our minimal assumption, but also very frequent in practice), if  $x_0$  is a point on the boundary of two elements,  $\nabla u_h$  is not continuous here. To include this case in our analysis we proceed as follows:

Let  $\ell$  denote a direction, i.e., unit vector, through  $x_0$  such that  $\nabla u_h(x_0 + \varepsilon\ell)$  is unique for  $\varepsilon \neq 0$  and small in magnitude and the right and left hand limits

$$(7.2.11) \quad \lim_{\varepsilon \rightarrow 0^\pm} \nabla u_h(x_0 + \varepsilon\ell) \quad \text{exist.}$$

We then set

$$(7.2.12) \quad \widehat{\nabla} u_h(x_0) = \frac{1}{2} \left( \lim_{\varepsilon \rightarrow 0^+} \nabla u_h + \lim_{\varepsilon \rightarrow 0^-} \nabla u_h \right).$$

We have as before,

$$(7.2.13) \quad D(u_{odd} - u_{h,odd}, \chi) = 0, \quad \text{for } \chi \in S_h^{comp}(B_d).$$

By Theorem 5.5.1 we then have ( $e = u - u_h$ ),

$$\begin{aligned} (7.2.14) \quad & \| \nabla(u_{odd} - u_{h,odd}) \|_{L_\infty(B_{d/2})} \\ & \leq C \min_{\chi \in S_h} \left( \|u_{odd} - \chi\|_{W_\infty^1(B_{d/2})} + d^{-1} \|u_{odd} - \chi\|_{L_\infty(B_{d/2})} \right) \\ & \quad + Cd^{-1-s-n/q} \|e_{odd}\|_{W_q^{-s}(B_{d/2})} \\ & \leq Ch^{r-1}d + Cd^{-1-s-n/q}h^{r-1+\tau'}, \end{aligned}$$

where we have assumed that the approximation hypothesis (7.1.3) holds and that

$$(7.2.15) \quad \|e\|_{W_q^{-s}(\mathcal{D})} \leq Ch^{r-1+\tau'},$$

cf. again Section 6.2 for such negative norm estimates.

Again performing a balancing act à la (7.2.8), we have

$$(7.2.16) \quad \|\nabla(u_{odd} - u_{h,odd})\|_{L_\infty(B_{d/4})} \leq Ch^{r-1-\sigma'},$$

if

$$(7.2.17) \quad d = h^{\sigma'}, \quad \sigma' = \tau'/(s + n/q + 2) \leq 1.$$

If the symmetry point  $x_0$  is a point of continuity for  $\nabla u_h$ , (7.2.16) gives immediately that

$$(7.2.18) \quad |\nabla u(x_0) - \nabla u_h(x_0)| \leq Ch^{r-1+\sigma'}.$$

If  $\nabla u_h$  is not continuous at  $x_0$ , we have from (7.2.16) for  $|\varepsilon| \neq 0$  small (with  $x_0 \pm \varepsilon\ell$  continuity points for  $\nabla u_h$ ),

$$(7.2.19) \quad \begin{aligned} & \left| \frac{1}{2} \{(\nabla u)(x_0 + \varepsilon\ell) + (\nabla u)(x_0 - \varepsilon\ell)\} \right. \\ & \quad \left. - \frac{1}{2} \{(\nabla u_h)(x_0 + \varepsilon\ell) + (\nabla u_h)(x_0 - \varepsilon\ell)\} \right| \\ & \leq Ch^{r-1-\sigma'}. \end{aligned}$$

Letting  $\varepsilon \rightarrow 0+$  we obtain

$$(7.2.20) \quad |\nabla u(x_0) - \hat{\nabla} u_h(x_0)| \leq Ch^{r-1+\sigma'}.$$

Hence,

**Theorem 7.2.2.** *Assuming that  $r$  is even, the negative norm estimate (7.2.15), the symmetry assumption (7.1.2) with (7.2.17) where  $\sigma' \leq 1$ , that Theorem 5.5.1 applies, the approximation assumption (7.1.3) and that  $u \in W_\infty^{r+1}(B_d)$ , we have*

$$(7.2.21) \quad |\nabla u(x_0) - \hat{\nabla} u_h(x_0)| \leq Ch^{r-1+\sigma'}.$$

Here  $\hat{\nabla} u_h$  is given in (7.2.12); it coincides with  $\nabla u_h$  in case  $x_0$  is a continuity point for  $\nabla u_h$ .

Again, if symmetry holds only in a smaller domain than dictated now by (7.2.17), superconvergence of lower order obtains.

Example 7.2.2. In the most favorable cases considered in Example 7.2.1, we now have

$$|(\nabla u - \hat{\nabla} u_h)(x_0)| \leq Ch^{r-1+(1-1/r)} \ln 1/h,$$

while imposing symmetry only in a ball of radius  $h^{1-1/r}$  around  $x_0$ .  $\square$

Example 7.2.3. If we consider a Dirichlet problem over a smooth domain but use only isoparametric modifications at the boundary, then we have (Example 6.3.2),

$$\|u - u_h\|_{L_\infty(\mathcal{D})} \leq C(\ln 1/h)^{\bar{r}} h^r.$$

Taking  $\tau' \simeq 1$ ,  $s = 0$ ,  $\sigma' = 1/2$  we may then assert that

$$|(\nabla u - \hat{\nabla} u_h)(x_0)| \leq Ch^{(r-1)+1/2} (\ln 1/h)^{\bar{r}},$$

for meshes symmetric in an  $O(h^{1/2})$  neighborhood of  $x_0$ .  $\square$

### 7.3. The general case (5.1.3) with variable smooth coefficients.

#### 7.3.a. Odd $r$ .

Now

$$(7.3.1) \quad A(e, \chi) = 0, \text{ for } \chi \in S_h^{\text{comp}}(B_d).$$

Thus, by the antipodal change of variables about  $x_0$ ,  $y = x_0 - (x - x_0)$ ,

$$(7.3.2) \quad \begin{aligned} A(\bar{e}, \chi) &= \int \left( \sum_{i,j=1}^n \bar{a}_{ij} \frac{\partial e}{\partial y_i} \frac{\partial \bar{\chi}}{\partial y_j} - \sum_{i=1}^n \bar{a}_i e \frac{\partial \bar{\chi}}{\partial y_i} + \bar{a} e \bar{\chi} \right) dy \\ &\equiv B(e, \bar{\chi}), \quad \text{for } \chi \in S_h^{\text{comp}}(B_d). \end{aligned}$$

Thus, using now that  $A(e, \chi) = 0$ , and by symmetry,  $A(e, \bar{\chi}) = 0$ , for  $\chi \in S_h^{\text{comp}}(B_d)$ ,

$$(7.3.4) \quad \begin{aligned} A(e_{\text{even}}, \chi) &= \frac{1}{2} A(e, \chi) + \frac{1}{2} B(e, \bar{\chi}) \\ &= \frac{1}{2} B(e, \bar{\chi}) - \frac{1}{2} A(e, \bar{\chi}) \\ &= \frac{1}{2} \int \sum_{i,j=1}^n (\bar{a}_{ij} - a_{ij}) \frac{\partial e}{\partial y_i} \frac{\partial \bar{\chi}}{\partial y_j} dy \\ &\quad - \frac{1}{2} \int \sum_{i=1}^n (\bar{a}_i + a_i) e \frac{\partial \bar{\chi}}{\partial y_i} dy + \frac{1}{2} \int (\bar{a} - a) e \bar{\chi} dy \\ &\equiv F_1(\bar{\chi}) + F_2(\bar{\chi}) + F_3(\bar{\chi}) \equiv F(\bar{\chi}), \quad \text{for } \chi \in S_h^{\text{comp}}(B_d). \end{aligned}$$

Here, for  $\varphi \in C_0^\infty(B_{d/4})$ , by the mean-value theorem applied to  $\bar{a}_{ij} - a_{ij}$ ,

$$(7.3.5) \quad |F_1(\varphi)| \leq Cd \|e\|_{W_\infty^1(B_{d/4})} |\varphi|_{W_1^1(B_{d/4})}.$$

By integration by parts and Poincaré–Friedrichs' inequality, again for  $\varphi \in C_0^\infty(B_{d/4})$ ,

$$(7.3.6) \quad \begin{aligned} |F_2(\varphi)| &= \left| \sum_{i=1}^n \left( \frac{\partial}{\partial y_i} (\bar{a}_i + a_i) e \right) \varphi dy \right| \\ &\leq C \|e\|_{W_\infty^1(B_{d/4})} \|\varphi\|_{L_1(B_{d/4})} \\ &\leq Cd \|e\|_{W_\infty^1(B_{d/4})} |\varphi|_{W_1^1(B_{d/4})} \end{aligned}$$

and the same trivially goes for  $F_3(\varphi)$ . So,

$$(7.3.7) \quad |||F|||_{-1,\infty,d/4} \leq Cd \|e\|_{W_\infty^1(B_{d/4})}.$$

Similarly, for  $\varphi \in C_0^\infty(B_{d/4})$ , after integrating by parts and then once more using Poincaré–Friedrichs' inequality and the mean-value theorem,

$$(7.3.8) \quad \begin{aligned} |F_1(\varphi)| &= \frac{1}{2} \left| \sum_{i,j=1}^n e \left( \frac{\partial}{\partial y_i} (\bar{a}_{ij} - a_{ij}) \frac{\partial \varphi}{\partial y_j} + (\bar{a}_{ij} - a_{ij}) \frac{\partial^2 \varphi}{\partial y_i \partial y_j} \right) dy \right| \\ &\leq C \|e\|_{L_\infty(B_{d/4})} d |\varphi|_{W_1^2(B_{d/4})}. \end{aligned}$$

Treating  $F_2$  and  $F_3$  in obvious analogous ways, we find that

$$(7.3.9) \quad |||F|||_{-2,\infty,d/4} \leq Cd \|e\|_{L_\infty(B_{d/4})}.$$

So, from Theorem 5.5.2, by (7.3.7) and (7.3.9),

$$(7.3.10) \quad \begin{aligned} |e(x_0)| &\leq C \min_{\chi \in S_h} \|u_{even} - \chi\|_{L_\infty(E_{d/2})} \\ &+ Cd^{-s-n/q} \|e\|_{W_q^{-s}(\mathcal{D})} \\ &+ C(\ln 1/h) \left( hd \|e\|_{W_\infty^1(B_{d/4})} + d \|e\|_{L_\infty(B_{d/4})} \right). \end{aligned}$$

By use of an inverse estimate and approximation theory, using a suitable  $\chi \in S_h$ ,

$$(7.3.11) \quad \begin{aligned} h \|e\|_{W_\infty^1(B_{d/4})} &\leq h \|u - \chi\|_{W_\infty^1(B_{d/4})} + h \|\chi - u_h\|_{W_\infty^1(B_{d/4})} \\ &\leq Ch \|u - \chi\|_{W_\infty^1(B_{d/4})} + C \|\chi - u_h\|_{L_\infty(B_{d/2})} \\ &\leq C \left( h \|u - \chi\|_{W_\infty^1(B_{d/4})} + \|u - \chi\|_{L_\infty(B_{d/2})} \right) \\ &\quad + C \|e\|_{L_\infty(B_{d/2})} \\ &\leq Ch^r + C \|e\|_{L_\infty(B_{d/2})}. \end{aligned}$$

Thus from (7.3.10), using approximation theory (7.1.3) and Taylor's theorem for the first term there (assuming  $u \in W_\infty^{r+1}(B_d)$ ),

$$(7.3.12) \quad \begin{aligned} |e(x_0)| &\leq Ch^r d(\ln 1/h) + Cd^{-s-n/q} \|e\|_{W_q^{-s}(\mathcal{D})} \\ &\quad + C(\ln 1/h) d \|e\|_{L_\infty(B_{d/2})}. \end{aligned}$$

Now again falling back to Theorem 5.5.2 (with  $F \equiv 0$ ) for the last term on the right of (7.3.12),

$$(7.3.13) \quad \begin{aligned} |e(x_0)| &\leq Ch^r d(\ln 1/h) + Cd^{-s-n/q} \|e\|_{W_q^{-s}(\mathcal{D})} \\ &\quad + C(\ln 1/h) d^{1-s-n/q} \|e\|_{W_q^{-s}(\mathcal{D})}. \end{aligned}$$

We conclude, somewhat informally stated:

**Theorem 7.3.1.** *Under the “the same” assumptions, the conclusion of Theorem 7.2.1 remains valid for the case of the general bilinear form (5.1.3), modulo a multiplicative factor  $\ln 1/h$ .*

### 7.3.b. Even $r$ .

Similarly (we leave the argument to the reader, cf. [Schatz, Sloan and Wahlbin 1994]), we assert that the conclusions of Theorem 7.2.2 remain valid for the more general bilinear form (5.1.3), again modulo a multiplicative factor  $\ln 1/h$ .

Of course, now we also used Theorem 5.5.1.

Finally we remark that Examples 7.2.1–7.2.3, which were based on the negative norm estimates of Examples 6.3.1 and 6.3.2, apply essentially as before. We leave it to the reader to consider the negative norm estimates in Examples 6.3.3 and 6.3.4 in the present context.

## 7.4. Historical remarks.

We shall rediscover a few known results through our present symmetry techniques. We shall also explain why an earlier attempt at a general principle fails. Details will be given only in the plane, although the results are relevant in any number of space dimensions.

### 7.4.i. Piecewise linear and quadratic triangles, and beyond.

Assume that the mesh is locally, in an  $O(1)$  neighborhood, say, generated by three linearly independent directions.




Figure 7.4.1

Then meshpoints and midpoints on common diagonals between two triangles are symmetry points. In particular then, if globally  $\|u - u_h\|_{W_q^{-s}(\mathcal{D})} \leq Ch^r$  for some negative norm, the average of the constant vectors  $\nabla u_h$  over two adjoining triangles is a superconvergent approximation to  $\nabla u$  at the midpoint of the shared diagonal (cf. Remark 7.2.1).

The history of this particular result can be traced back at least as far as [Oganesyan and Rukhovetz 1969, eq. (3.18)] in the  $L_2$ -based case for piecewise linear elements. Phrased in our present language,

$$(7.4.1) \quad |A(u - u^I, \chi)| \leq Ch^r \|\chi\|_{W_1^1}, \text{ for } \chi \in S_h^{comp}(\Omega_1),$$

where  $r = 2$  and  $u^I$  is the natural piecewise linear interpolant at nodes. Of course, having this we automatically have

$$(7.4.2) \quad |A(u_h - u^I, \chi)| \leq Ch^r \|\chi\|_{W_1^1}, \text{ for } \chi \in S_h^{comp}(\Omega_1)$$

and our present techniques, viz., Theorem 5.5.1, gives that  $u_h$  is “superclose” to  $u^I$  in  $W_\infty^1(\Omega_0)$ . To then find out about superconvergence for  $\nabla(u - u_h)$ , it simply remains to consider the locally defined error in  $\nabla(u - u^I)$ . This is frequently elementary.

To verify (7.4.1) is often a slightly laborious task involving taking into account local cancellation effects due to symmetry. In the piecewise linear situation, typical examples can be found in [Chen and Liu 1987] and [Lin and Xu 1985]. In other words, to state (7.4.1) as a hypothesis leaves a major part of the work to the reader.

The hypothesis (7.4.1) and its counterpart for function values were formalized e.g. in [Zhu 1983, eqs. (2.11) and (2.12)]. He then proceeded “essentially” as in our Theorems 5.5.1 and 5.5.2. Earlier, [Zlámal 1977, 1978a,b] and [Lesaint and Zlámal 1979] had used the same idea in a discrete  $\ell_1$ -setting for the error in higher order quadrilateral elements.

For our additional results about averaging from two sides about a symmetric nodal point in the piecewise linear situation, cf. also [Levine 1985].

However, an objection against basing a “general” theory of superconvergence on (7.4.1) is that it is not a general enough principle. For uniform triangular meshes we shall “see” via computational investigations in Chapter 12 that, beyond quadratics, the standard interpolant is not superclose to the finite element approximation. Furthermore, a little known but in my opinion important paper, [Li 1990], proves that on uniform triangular meshes, for  $C^0$  Lagrange cubics or Hermite cubics, the Galerkin finite element solution (for the Laplacian on a rectangle) and the standard interpolants are not superclose in  $H^1$  or  $L_2$ . Thus, basing a general theory of superconvergence on the hope that the Galerkin finite element approximation is close to a standard interpolant is doomed to fail.

#### 7.4.ii. More on $C^0$ quadratics on triangles.

Again we have a mesh consisting of triangles involving three linearly independent directions. Now  $r = 3$  and from our theory in this chapter, mesh and midpoints are superconvergent in function values.

Letting then  $u^I$  denote the piecewise quadratic interpolant of  $u$  in one variable on any side of  $\tau$ , by stability  $u_h - u^I$  is superconvergent over that side in function values, and thus, from inverse assumptions,  $\frac{\partial}{\partial t}(u_h - u^I)$  is superconvergent for the tangential derivative  $\frac{\partial}{\partial t}$  along that side. Hence superconvergence for  $\frac{\partial}{\partial t}(u - u_h)$  occurs at the same points as for  $\frac{\partial}{\partial t}(u - u^I)$ . These are trivially found to be the Gauss points: If  $-\hat{h} \leq t \leq \hat{h}$  describe a side of  $\tau$ , they are the zeroes of the normalized second degree Legendre polynomial, i.e.,

$$(7.4.3) \quad t = \pm \hat{h}/\sqrt{3}.$$

We have rediscovered a result which can be traced at least as far back as [Andreev 1984b].

We note that our “fast” derivation above was based on the fact that there were enough symmetry points present on an element to determine the “natural” local interpolant. This will be the key also in the following two examples.

#### 7.4.iii. Incomplete cubics.

We are now on rectangles, say, axis-parallel for simplicity. The incomplete cubics are given as  $A + Bx + Cy + Dx^2 + Exy + Fy^2 + Gx^2y + Hyx^2$  on each rectangle (and are, of course, globally continuous). Again,  $r = 3$ . They are determined by interpolation at the following eight points:




Figure 7.4.2

These now happen to be among the (nine) symmetry points and thus superconvergent for function values. If  $u^I$  denotes the interpolant indicated, then  $u_h - u^I$  is superconvergent there for function values, and so, by stability, “superclose” on the whole of  $\tau$ . By an inverse inequality,  $\nabla(u_h - u^I)$  is “superclose” on  $\tau$ . To detect superconvergence points for  $\nabla(u - u_h)$  it thus remains to consider such for  $\nabla(u - u^I)$ . Since  $x^2y$  and  $yx^2$  are reproduced by  $u^I$  it is enough to look at what happens for  $u = x^3$  and  $u = y^3$ . This is elementary and we find that the four interior Gauss points ( $h_x$  and  $h_y$  are the meshsizes in the  $x$  and  $y$  directions, respectively), are superconvergent for  $\nabla(u - u_h)$ .




Figure 7.4.3

We have rediscovered a result which may be traced back at least as far as to work by Zlámal and Lesaint in the late 1970's, cf. the references. Cf. also [Barlow 1976].

#### 7.4.iv. $C^0$ or $C^1$ biquadratic tensor-products.

We are again on a rectangle and we now have nine symmetry points for function values,




Figure 7.4.4

which is enough to determine a local interpolant  $u^I$  of the form  $A + Bx + Cy + Dx^2 + Exy + Fy^2 + Gx^2y + Hxy^2 + Ix^2y^2$ . (For the  $C^1$ -case, this  $u^I$  will not be globally in  $S_h$ , but that doesn't matter for our argument.)

(As for superconvergence at the four nodes, in the  $C^1$ -case we have rediscovered [Bramble and Schatz 1976, Theorem 10].) It follows as in 7.4.iii for the incomplete cubics, that the interior Gauss points are again superconvergent for gradients. (The now simple details are left to the reader.)

Again, work by Zlámal and Lesaint in the late 1970's has been rediscovered.

We may also remark that although the present situation is that of tensor products, we have *not* used the theory of Chapter 6, with its restrictions on the form of the

bilinear form  $A(\cdot, \cdot)$ . We have relied instead on superconvergence by symmetry which we have verified for a general form  $A(\cdot, \cdot)$ .

## Chapter 8. Superconvergence for difference quotients on translation invariant meshes.

### 8.1. Introduction.

We have met a simple case in Section 1.11. In this chapter we shall consider derivatives of any order, but we shall not manage to be as precise with respect to local influences as we were in the one-dimensional situation. The whole chapter is by and large based on [Nitsche and Schatz 1974, Section 6] where the theory was presented in  $L_2$ -based error estimates. Using later technology, here we work with pointwise error estimates (but cf. Remark 8.4.1 below!).

The work of Nitsche and Schatz was inspired by the finite difference theory in [Thomée and Westergren 1968]. An earlier investigation into using difference quotients to approximate derivatives in a multidimensional finite element setting is given for parabolic problems in [Fix and Nassif 1972]. Another discovery of the fact that it is better to difference than to differentiate in finite elements is in [Long and Morton 1977].

Let  $\Omega_0 \subset\subset \Omega_1 \subset\subset \Omega_2 \subset\subset \mathcal{D}$  be separated by  $d$  and let  $\ell$  be a direction, i.e., a unit vector in  $R^n$ . Let  $H > 0$  be a parameter, which will typically be a constant times  $h$ . By *translation invariance by  $H$  in the direction  $\ell$*  we shall mean the following.

Let  $T_H^\ell$  denote translation by  $H$  in the direction  $\ell$ , i.e.,

$$(8.1.1) \quad T_H^\ell v(x) = v(x + H\ell),$$

and for  $\nu$  an integer,

$$(8.1.2) \quad T_{\nu H}^\ell v(x) = v(x + \nu H\ell).$$

Then the finite element spaces are called translation invariant by  $H$  in the direction  $\ell$  if

$$(8.1.3) \quad T_{\nu H}^\ell \chi \in S_h^{comp}(\Omega_2), \text{ for } \chi \in S_h^{comp}(\Omega_1).$$

This will be required for  $|\nu| \leq M$ , some fixed  $M$ ; the dependence on  $M$  will be suppressed in our notation.

Let us give some pictorial examples, where we display what we think of as  $h$ .

Example 8.1.1.




Figure 8.1.1

Here we may take the following combinations:

$$\ell = (1, 0), H = h,$$

$$\ell = (0, 1), H = h/2,$$

$$\ell = \frac{2}{\sqrt{5}}(1, 1/2), H = h\sqrt{5}/2.$$

(Also, the direction  $\frac{2}{\sqrt{5}}(1, -1/2)$ .)

Of course, the same example works if the triangular elements are replaced by rectangular ones.  $\square$

Example 8.1.2.




Figure 8.1.2

Here

$$\ell = (1, 0), H = 2h,$$

$$\ell = (0, 1), H = h,$$

and the “diagonal” directions.  $\square$


Example 8.1.3.

Figure 8.1.3

Here,  $\ell = (0, 1)$  and  $H$  as indicated. No other direction works.  $\square$

We now consider a function  $u$  which satisfies, with  $u_h \in S_h(\Omega_1)$ ,

$$(8.1.4) \quad A(u - u_h, \chi) = 0, \text{ for } \chi \in S_h^{comp}(\Omega_2).$$

The problem is to approximate  $D_\ell^k u$ , the  $k^{\text{th}}$  derivative in the locally translation invariant direction  $\ell$ . We shall utilize a suitable difference quotient based on translations in the direction  $\ell$ ,

$$(8.1.5) \quad \partial_{\ell, H}^k v(x) = \sum_{|\nu| \leq M} C_{\nu, H} v(x + \nu H \ell).$$

The dependence of  $C_{\nu, H}$  on  $k$  and  $\ell$  will be suppressed in the notation. We shall assume that  $\partial_{\ell, H}^k$  is an  $r^{\text{th}}$  order approximation to  $D_\ell^k$  in the sense that

$$(8.1.6) \quad \|D_\ell^k \tilde{T}_h v - \partial_{\ell, H}^k v\|_{L_\infty(\Omega_1)} \leq CH^s \|v\|_{W_\infty^{k+s}(\Omega_2)}, \text{ for } 0 \leq s \leq r.$$

Since  $H \simeq h$ , we may replace  $H$  by  $h$  on the right. Here,  $\tilde{T}_h$  is a “suitable” shift operator, sometimes  $\tilde{T}_h = I$ . See Remark 8.2.1 for another, where  $\tilde{T}_h$  is translation by  $h/2$  in the  $\ell$ -direction. The meshes do not have to be translation invariant with respect to  $\tilde{T}_h$ .

## 8.2. Constant coefficient operators and unit separation, $d \simeq 1$ .

So, let  $a_{ij}$ ,  $a_i$  and  $a$  be constant in  $A$ . Then

$$(8.2.1) \quad A(T_{\nu H}^\ell v, w) = A(v, T_{-\nu H}^\ell w) \equiv A(v, (T_{\nu H}^\ell)^* w),$$

and thus, from (8.1.3) and (8.1.4),

$$(8.2.2) \quad A(\partial_{\ell, H}^k (u - u_h), \chi) = A(u - u_h, (\partial_{\ell, H}^k)^* \chi) = 0, \text{ for } \chi \in S_h^{comp}(\Omega_1).$$

From (8.1.3) again,  $\partial_{\ell, H}^k u_h \in S_h$  and Theorem 5.5.2 (with  $F \equiv 0$ ) thus implies that

$$(8.2.3) \quad \begin{aligned} \|\partial_{\ell, H}^k (u - u_h)\|_{L_\infty(\Omega_0)} &\leq C (\ln d/h)^\bar{r} \min_{\chi \in S_h} \|\partial_{\ell, H}^k u - \chi\|_{L_\infty(\Omega_1)} \\ &+ Cd^{-s-n/q} \|\partial_{\ell, H}^k (u - u_h)\|_{W_q^{-s}(\Omega_1)}. \end{aligned}$$

Here  $d \geq c_0 h$  is the separation between  $\Omega_0$  and  $\Omega_1$  (and the separation between  $\Omega_1$  and  $\Omega_2$  is assumed  $\geq$  cte  $h$ , too). By (8.1.6), with  $s = r$ , and standard approximation assumptions,

$$(8.2.4) \quad \begin{aligned} & \min_{\chi \in S_h} \|\partial_{\ell,H}^k u - \chi\|_{L_\infty(\Omega_1)} \\ & \leq \|\partial_{\ell,H}^k u - D_\ell^k \tilde{T}_h u\|_{L_\infty(\Omega_1)} + \min_{\chi \in S_h} \|D_\ell^k \tilde{T}_h u - \chi\|_{L_\infty(\Omega_1)} \\ & \leq Ch^r \|u\|_{W_\infty^{r+k}(\Omega_2)}. \end{aligned}$$

Thus, from (8.2.3) using again (8.1.6), if  $u$  is assumed smooth enough on  $\Omega_2$ ,

$$(8.2.5) \quad \begin{aligned} & \|D_\ell^k \tilde{T}_h u - \partial_{\ell,H}^k u_h\|_{L_\infty(\Omega_0)} \\ & \leq C(\ln 1/h)^{\bar{r}} h^r + Cd^{-s-n/q} \|\partial_{\ell,H}^k(u - u_h)\|_{W_q^{-s}(\Omega_1)}. \end{aligned}$$

It remains to attack the negative norm on the right of (8.2.5). We have

$$(8.2.6) \quad \|\partial_{\ell,H}^k(u - u_h)\|_{W_q^{-s}(\Omega_1)} = \sup_{\substack{\varphi \in C_0^\infty(\Omega_1) \\ \|\varphi\|_{W_{q'}^s(\Omega_1)}=1}} (\partial_{\ell,H}^k(u - u_h), \varphi).$$

Here,

$$(8.2.7) \quad \begin{aligned} (\partial_{\ell,H}^k(u - u_h), \varphi) &= (u - u_h, (\partial_{\ell,H}^k)^* \varphi) \\ &\leq C\|u - u_h\|_{L_\infty(\Omega_2)} \|(\partial_{\ell,H}^k)^* \varphi\|_{L_1(\Omega_2)}. \end{aligned}$$

From the dual version of (8.1.6) with  $s = 0$  (which we leave to the audience to make precise), if  $s \geq k$ ,

$$(8.2.8) \quad \|(\partial_{\ell,H}^k)^* \varphi\|_{L_1(\Omega_2)} \leq C.$$

From (8.2.5)–(8.2.8), since in our present case  $d \simeq 1$ ,

$$(8.2.9) \quad \|D_\ell^k \tilde{T}_h u - \partial_{\ell,H}^k u_h\|_{L_\infty(\Omega_0)} \leq C(\ln 1/h)^{\bar{r}} h^r + C\|u - u_h\|_{L_\infty(\Omega_2)}.$$

We now once again apply Theorem 5.5.1 to the last term to obtain with  $\Omega_2 \subset\subset \mathcal{D}$  separated by a unit distance, and  $u$  smooth enough,

$$(8.2.10) \quad \|D_\ell^k \tilde{T}_h u - \partial_{\ell,H}^k u_h\|_{L_\infty(\Omega_0)} \leq C(\ln 1/h)^{\bar{r}} h^r + C\|u - u_h\|_{W_q^{-s}(\mathcal{D})}.$$

We summarize informally:

**Theorem 8.2.1.** *Let the differential operator have constant coefficients; let  $\Omega_0 \subset\subset \Omega_2 \subset\subset \mathcal{D}$  be separated by unit distances; let the finite element spaces be translation invariant in the direction  $\ell$  on  $\Omega_2$  with parameter  $H \simeq h$  (cf. (8.1.3)); let  $\partial_{\ell,H}^k$  be a finite difference operator based on translations in the  $\ell$ -direction which approximates the directional derivative  $D_\ell^k$  to order  $h^r$  (cf. (8.1.6)); let the finite element space have approximation order  $h^r$  for function values; let  $A(u - u_h, \chi) = 0$  for  $\chi \in S_h^{\text{comp}}(\Omega_2)$ ; let  $u$  be locally smooth enough; and assume that for some  $q$  and  $s$ ,*

$$(8.2.11) \quad \|u - u_h\|_{W_q^{-s}(\mathcal{D})} \leq Ch^r.$$

*Furthermore assume that Theorem 5.5.2 is applicable. Then*

$$(8.2.12) \quad \|D_\ell^k \tilde{T}_h u - \partial_{\ell,H}^k u_h\|_{L_\infty(\Omega_0)} \leq C(\ln 1/h)^{\bar{r}} h^r,$$

where  $\bar{r} = 1$  if  $r = 2$ ,  $\bar{r} = 0$  for  $r \geq 3$ .

So, suitable difference quotients provide optimal order  $h^r$  (modulo a logarithmic term if  $r = 2$ ) approximation to  $D_\ell^k$ , for any  $k$ , some negative norm permitting, of course. Let us remark that above we have given the details for pure  $k^{\text{th}}$  derivatives in the direction  $\ell$ . There are no essential changes for mixed derivatives, but now one needs of course translation invariance in all directions involved.

Remark 8.2.1. Here we present again a rediscovery from Section 7.4.i, now rediscovered from the results of this section. Consider piecewise linear plane triangular elements on a locally “uniform” mesh such as depicted in Example 8.1.1. Let us consider e.g. the  $x$ -direction,  $\ell = (1, 0)$  with  $H = h$  as there, and first derivatives  $\partial u / \partial x$ . Let  $\partial$  be the difference operator

$$(8.2.13) \quad \partial v(x, y) = (v(x + h, y) - v(x, y))/h,$$

and let

$$(8.2.14) \quad \tilde{T}_h v(x) = v\left(x + \frac{h}{2}(1, 0)\right).$$

Then (8.1.6) is satisfied with  $r = 2$ . Thus, some negative norm permitting,

$$(8.2.15) \quad \left\| \frac{\partial u}{\partial x}\left(x + \frac{h}{2}, y\right) - \partial u_h(x, y) \right\|_{L_\infty(\Omega_0)} \leq Ch^2 \ln 1/h.$$

Let now  $x$  be a “mesh-point”; while  $y$  is a “meshpoint” shifted by  $\tilde{h}/2$ , where  $\tilde{h}$  is the mesh length in the  $y$ -direction (i.e.,  $y$  is a “midpoint”):




Figure 8.2.1

Denote  $u_h^-$  and  $u_h^+$  as indicated. Then

$$\begin{aligned} (8.2.16) \quad \partial u_h(x, y) &= (u_h(x + h, y) - u_h(x, y))/h \\ &= [(u_h(x + h, y) - u_h(x + h/2, y)) \\ &\quad + (u_h(x + h/2, y) - u_h(x, y))]/h \\ &= \frac{1}{2}[(u_h^+(x + h, y) - u_h^+(x + h/2, y))/(h/2) \\ &\quad + (u_h^-(x + h/2, y) - u_h^-(x, y))/(h/2)]. \end{aligned}$$

Combining this with (8.2.15) we deduce that the average of  $\partial u_h / \partial x$  (which is constant in each element) over two adjoining elements as above is an  $h^2 \ln 1/h$  order approximation to  $\frac{\partial u}{\partial x}(x, y)$  if  $x$  and  $y$  are both “midpoints”.

Doing the same in the  $y$ -direction, and after an easy generalization to any mesh generated locally by three fixed directions, we have rediscovered the result about superconvergence of averages of  $\nabla u_h$  at midpoints of shared sides.  $\square$

### 8.3. Constant coefficient operators and general separation $d$ .

Of course, we assume that  $d \geq c_0 h$  with  $c_0$  large enough. We shall sketch the argument; the reader can easily supply missing details. Repeating (8.2.5),

$$(8.3.1) \quad \begin{aligned} & \|D_\ell^k \tilde{T}_h u - \partial_{\ell,H}^k u_h\|_{L_\infty(\Omega_0)} \\ & \leq C(\ln 1/h)^{\bar{r}} h^r + Cd^{-s-n/q} \|\partial_{\ell,H}^k(u - u_h)\|_{W_q^{-s}(\Omega_1)}. \end{aligned}$$

As in (8.2.7), now using also Poincaré–Friedrichs' inequality, and, Hölder's inequality,

$$(8.3.2) \quad \begin{aligned} |(\partial_{\ell,H}^k(u - u_h), \varphi)| & \leq \|u - u_h\|_{L_\infty(\Omega_2)} \|(\partial_{\ell,H}^k)^* \varphi\|_{L_1(\Omega_2)} \\ & \leq C \|u - u_h\|_{L_\infty(\Omega_2)} \|\varphi\|_{W_1^k(\Omega_1)} \\ & \leq C \|u - u_h\|_{L_\infty(\Omega_2)} d^{s-k} \|\varphi\|_{W_1^s(\Omega_1)} \\ & \leq C \|u - u_h\|_{L_\infty(\Omega_2)} d^{s-k+n/q} \|\varphi\|_{W_q^s(\Omega_1)}, \text{ for } s \geq k. \end{aligned}$$

So,

$$(8.3.3) \quad \|D_\ell^k \tilde{T}_h u - \partial_{\ell,H}^k u_h\|_{L_\infty(\Omega_0)} \leq C(\ln 1/h)^{\bar{r}} h^r + d^{-k} \|u - u_h\|_{L_\infty(\Omega_2)}.$$

If one now assumes that

$$(8.3.4) \quad \|u - u_h\|_{L_\infty(\mathcal{D})} \leq Ch^r (\ln 1/h)^{\bar{r}},$$

cf. Example 6.2.2, one obtains

$$(8.3.5) \quad \|D_\ell^k \tilde{T}_h u - \partial_{\ell,H}^k u_h\|_{L_\infty(\Omega_0)} \leq C(\ln 1/h)^{\bar{r}} d^{-k} h^r.$$

E.g., for first derivatives,  $k = 1$ , if the mesh is translation invariant in an  $O(h^{1/2})$  neighborhood one obtains superconvergence of almost a half extra order.

If (8.3.4) does not hold one may of course further localize the term  $\|u - u_h\|_{L_\infty(\Omega_2)}$  in (8.3.3) by use of Theorem 5.5.2 (with  $F \equiv 0$ ). We leave this to the reader.

### 8.4. Variable coefficients.

We shall consider a fairly simple case which we hope will convince the reader that he or she can handle the general case. Let thus

$$(8.4.1) \quad A(v, w) = \int \sum_{i,j=1}^n a_{ij}(x) \frac{\partial v}{\partial x_i} \frac{\partial w}{\partial x_j} dx,$$

i.e., we consider only the highest order terms, but these are of a general form. Let further  $d \simeq 1$  and the mesh locally translation invariant in the  $x_1$ -direction with parameter  $H \equiv h$ . For  $r = 2$ , i.e., typically piecewise linears or bilinears, consider approximations to  $\partial^2 u / \partial x^2$  based on the traditional second order accurate operator

$$(8.4.2) \quad \partial^2 v(x) = (v(x + he_1) - 2v(x) + v(x - he_1))/h^2,$$

where  $e_1 = (1, 0, \dots, 0)$ . (Here,  $\tilde{T}_h \equiv I$ .)

Our aim is to show that if  $A(u - u_h, \chi) = 0$ , for  $\chi \in S_h^{comp}(\Omega_2)$ , then

$$(8.4.3) \quad \left\| \frac{\partial^2 u}{\partial x^2} - \partial^2 u_h \right\|_{L_\infty(\Omega_0)} \leq Ch^2(\ln 1/h)^3.$$

(Higher order elements are admitted but the final result is still only  $O(h^2)$  as in (8.4.3), for obvious reasons.) For this, we shall assume that in some suitable negative norm,

$$(8.4.4) \quad \|u - u_h\|_{W_q^{-s}(\mathcal{D})} \leq Ch^2.$$

We start by noting that

$$(8.4.5) \quad \partial^2 = \partial_- \partial_+$$

where  $\partial_+ v(x) = (v(x + he_1) - v(x))/h$  and  $\partial_- v(x) = (v(x) - v(x - he_1))/h$ .

We shall need two preliminary results. The first is that

$$(8.4.6) \quad \|\partial_\pm e\|_{W_\infty^1(\Omega_1)} \leq Ch \ln 1/h.$$

Here, and below,  $e = u - u_h$ . The second preliminary result is that

$$(8.4.7) \quad \|\partial_\pm e\|_{L_\infty(\Omega_1)} \leq Ch^2(\ln 1/h)^2.$$

We proceed to prove (8.4.6). For technical reasons we introduce a further domain  $\Omega_{1.5}$  between  $\Omega_1$  and  $\Omega_2$ . Consider then, for  $\chi \in S_h^{comp}(\Omega_{1.5})$ ,

$$\begin{aligned} (8.4.8) \quad A(\partial_+ e, \chi) &= \sum_{i,j=1}^n \int a_{ij}(x) \left( \partial_+ \frac{\partial e}{\partial x_i} \right) \frac{\partial \chi}{\partial x_j} \\ &= \frac{1}{h} \sum_{i,j=1}^n \int a_{ij}(x) \frac{\partial \chi}{\partial x_j}(x) \left( \frac{\partial e}{\partial x_i}(x + he_1) - \frac{\partial e}{\partial x_i}(x) \right) \\ &= \frac{1}{h} \sum_{i,j=1}^n \int \left( a_{ij}(x - he_1) \frac{\partial \chi}{\partial x_j}(x - he_1) - a_{ij}(x) \frac{\partial \chi}{\partial x_j}(x) \right) \frac{\partial e}{\partial x_i}(x) \\ &= -\frac{1}{h} \sum_{i,j=1}^n \int a_{ij}(x) \left( \frac{\partial \chi}{\partial x_j}(x) - \frac{\partial \chi}{\partial x_j}(x - he_1) \right) \frac{\partial e}{\partial x_i}(x) \\ &\quad + \frac{1}{h} \sum_{i,j=1}^n \int (a_{ij}(x - he_1) - a_{ij}(x)) \frac{\partial \chi}{\partial x_j}(x - he_1) \frac{\partial e}{\partial x_i}(x). \end{aligned}$$

Here, the first term on the right vanishes since  $\chi(x) - \chi(x - he_1) \in S_h^{comp}(\Omega_2)$  by translation invariance. Thus,

$$\begin{aligned} (8.4.9) \quad |A(\partial_+ e, \chi)| &= \left| - \sum_{i,j=1}^n \int (\partial_- a(x)) \frac{\partial \chi}{\partial x_j}(x - he_1) \frac{\partial e}{\partial x_i}(x) \right| \\ &\leq C \|e\|_{W_\infty^1(\Omega_2)} \|\chi\|_{W_1^1(\Omega_{1.5})}, \text{ for } \chi \in S_h^{comp}(\Omega_{1.5}). \end{aligned}$$

In other words,

$$(8.4.10) \quad A(\partial_+ e, \chi) = F(\chi), \text{ for } \chi \in S_h^{comp}(\Omega_{1.5}),$$

where

$$(8.4.11) \quad \sup_{\substack{\chi \in S_h^{comp}(\Omega_{1.5}) \\ \|\chi\|_{W_1^1(\Omega_{1.5})}=1}} |F(\chi)| \leq C \|e\|_{W_\infty^1(\Omega_2)} \leq Ch,$$

by use of Theorem 5.5.1 (with  $F \equiv 0$ ), local approximation and the negative norm estimate (8.4.4). Another application of Theorem 5.5.1 then gives (8.4.6).

As for (8.4.7), we have declared that

$$(8.4.12) \quad F(\varphi) \equiv - \sum_{i,j=1}^n \int \left( \partial_- a(x) \frac{\partial \varphi}{\partial x_j}(x - he_1) \frac{\partial e}{\partial x_i}(x) \right).$$

Integration by parts, freeing  $e$  from its derivatives, using Theorem 5.2.2 twice (once for  $e$ , with  $F \equiv 0$ , and once for  $A(\partial_+ e, \chi) = F(\chi)$ ), then gives (8.4.7).

We next consider  $A(\partial^2 e, \chi)$ . We have for  $\chi \in S_h^{comp}(\Omega_1)$ ,

$$\begin{aligned} (8.4.13) \quad A(\partial^2 e, \chi) &= \sum_{i,j=1}^n \int a_{ij}(x) \left( (\partial_- \partial_+) \frac{\partial e}{\partial x_i} \right) \frac{\partial \chi}{\partial x_j} \\ &= - \sum_{i,j=1}^n \int \partial_+ \left( a_{ij} \frac{\partial \chi}{\partial x_j} \right) \partial_+ \frac{\partial e}{\partial x_i} \\ &= \sum_{i,j=1}^n \int (\partial^2) \left( a_{ij} \frac{\partial \chi}{\partial x_j} \right) \frac{\partial e}{\partial x_i}. \end{aligned}$$

Now, using the discrete analogue of Leibniz' rule,  $\partial^2(fg) = (\partial^2 f)g + (\partial_+ f)(\partial_+ g) + (\partial_- f)(\partial_- g) + f(\partial^2 g)$ , when second differences fall on  $\frac{\partial \chi}{\partial x_j}$ , the result is zero, by translation invariance and (8.1.4). When second differences fall on  $a_{ij}$ , that difference quotient is bounded, since the  $a_{ij}$  are smooth. The typical middle term is

$$(8.4.14) \quad \sum_{i,j=1}^n \int \partial_\pm a_{ij} \left( \partial_\pm \frac{\partial \chi}{\partial x_j} \right) \frac{\partial e}{\partial x_i},$$

and we transfer the difference quotient falling on  $\frac{\partial \chi}{\partial x_j}$  to  $\partial_- a_{ij}$  or  $\frac{\partial e}{\partial x_i}$ . Using then (8.4.6) we have

$$(8.4.15) \quad A(\partial^2 e, \chi) = F(\chi), \text{ for } \chi \in S_h^{comp}(\Omega_1),$$

with

$$(8.4.16) \quad \|F\|_{-1,\infty,\Omega_1} \leq Ch \ln 1/h.$$

Similarly, using now (8.4.7) after integration by parts, cf. (8.4.12) et. seq.,

$$(8.4.17) \quad \|F\|_{-2,\infty,\Omega_1} \leq Ch^2 (\ln 1/h)^2.$$

We may thus apply Theorem 5.5.2 to deduce that

$$(8.4.18) \quad \|\partial^2 e\|_{L_\infty(\Omega_0)} \leq Ch^2 (\ln 1/h)^3.$$

We conclude that

$$(8.4.19) \quad \left\| \frac{\partial^2 u}{\partial x^2} - \partial^2 u_h \right\|_{L_\infty(\Omega_0)} \leq Ch^2 (\ln 1/h)^3.$$

The reader has no doubt found an induction argument hidden above, cf. (8.4.6) and (8.4.7). For the general case, see [Nitsche and Schatz 1974, Section 6] where this induction is brought out into the open in the case of  $L_2$ -based estimates. Details are not particularly hard to translate to pointwise estimates.

Remark 8.4.1. The reader will also have noticed a proliferation of  $\ln 1/h$  factors. Actually, one may avoid this by going back to an earlier technique for getting  $L_\infty$ -estimates, in the case of meshes which are now required to be translation invariant in all of  $n$  linearly independent directions. This technique involves estimating  $L_\infty$ -norms by Sobolev's inequality in  $L_2$ , but now with *difference quotients* rather than derivatives of order  $> n/2$  involved. Somewhat more regularity of  $u$  is required than in our arguments above in this chapter. We refer to [Bramble, Nitsche and Schatz 1975, Lemma 4.2 in particular] for details. Cf. also Section 11.4 below.

In fact, as remarked by [Liu 1994b] and [Schatz 1994] if one is interested in interior maximum-norm estimates only on translation invariant meshes but for a more complicated problem than the scalar one treated in Section 5.5 (e.g., a Stokes' problem or a linear elasticity problem), this technique may give a much quicker derivation than could be had by doing the analogues of the proofs of Theorems 5.5.1 and 5.5.2. (After all, these two theorems are valid for general quasi-uniform meshes.)  $\square$

## Chapter 9. On superconvergence in nonlinear problems.

For simplicity in notation, we shall consider only an equation involving highest order terms, and, for additional brevity, we shall only give a “global” argument. The reader will have no problem in furnishing details with respect to these omissions.

Let  $\mathbf{q}(w)$  denote the gradient vector  $\mathbf{q}(w) = \nabla w$  and let  $a_{ij}(w, \mathbf{q})$ ,  $i, j = 1, \dots, n$ , be smooth functions for  $(w, \mathbf{q}) \in R^{1+n}$ . The basic problem is now

$$(9.1) \quad \begin{cases} L(u) = -\sum_{i,j=1}^n \frac{\partial}{\partial x_i} \left( a_{ij}(u, \mathbf{q}(u)) \frac{\partial u}{\partial x_j} \right) = f \text{ on } \mathcal{D}, \\ \text{some linear, homogeneous boundary condition on } \partial\mathcal{D}. \end{cases}$$

We assume that there is a unique solution  $u$  to (9.1) which is as smooth as necessary on  $\bar{\mathcal{D}}$ . It is not assumed that the  $a_{ij}(w, \mathbf{q})$  are bounded on the whole of  $R^{1+n}$ , nor is it assumed that the operator based on  $\{a_{ij}(w, \mathbf{q})\}$  is uniformly elliptic on the whole of  $R^{1+n}$ .

We let  $A(u, \mathbf{q}(u))(v, w)$  denote the corresponding form which is bilinear in  $v$  and  $w$ . The finite element problem is now to find  $u_h \in S_h$  such that

$$(9.2) \quad A(u_h, \mathbf{q}(u_h))(u_h, \chi) = (f, \chi), \text{ for } \chi \in S_h$$

where, of course,  $S_h$  would respect essential boundary conditions (at least in some sense). Again we assume that there is a unique solution  $u_h$  to (9.2), at least for  $h$  sufficiently small.

Let us pause to remark that the essential ideas of this chapter are due to [Douglas and Dupont 1975], in an  $L_2$ -based case. Conditions which ensure that (9.2) has a unique solution may be found there, see also the treatise [Ženíšek 1990]. For  $L_\infty$ -estimates, cf. also [Nitsche 1977] and [Frehse and Rannacher 1978]. For superconvergence, cf. [Chen 1983a].

The main theme in this chapter is that if certain points (or postprocessors) are superconvergent for a certain linearized problem associated with (9.1), then they are automatically superconvergent for the error in the nonlinear problem. We proceed to introduce this linearized problem, which involves the derivative of  $A$  at  $u$  in the direction  $v$ .

Consider  $u + v$  where  $v$  is to be thought of as “small”. Then

$$(9.3) \quad \begin{aligned} & a_{ij}(u + v, \mathbf{q}(u) + \mathbf{q}(v)) \frac{\partial(u + v)}{\partial x_i} \\ &= a_{ij}(u, \mathbf{q}(u)) \frac{\partial u}{\partial x_i} + a_{ij}(u, \mathbf{q}(u)) \frac{\partial v}{\partial x_i} \\ &+ [a_{ij}(u + v, \mathbf{q}(u) + \mathbf{q}(v)) - a_{ij}(u, \mathbf{q}(u))] \left[ \frac{\partial u}{\partial x_i} + \frac{\partial v}{\partial x_i} \right]. \end{aligned}$$

By Taylor’s theorem,

$$(9.4) \quad \begin{aligned} & a_{ij}(u + v, \mathbf{q}(u) + \mathbf{q}(v)) - a_{ij}(u, \mathbf{q}(u)) \\ &= \frac{\partial a_{ij}}{\partial u}(u, \mathbf{q}(u))v + \sum_{k=1}^n \frac{\partial a_{ij}}{\partial q_k}(u, \mathbf{q}(u)) \frac{\partial v}{\partial x_k} + R, \end{aligned}$$

where  $R = R(x) = R(u(x), \mathbf{q}(u(x)), v(x), \mathbf{q}(v(x)))$  may be bounded by

$$(9.5) \quad |R| \leq C_1(u, v) \|v\|_{W_\infty^1(\mathcal{D})}^2,$$

with  $C_1(u, v)$  depending on  $\|u\|_{W_\infty^1(\mathcal{D})}$  and  $\|u + v\|_{W_\infty^1(\mathcal{D})}$  (through the  $a_{ij}$  and their derivatives).

Since also

$$(9.6) \quad |a_{ij}(u + v, \mathbf{q}(u) + \mathbf{q}(v)) - a_{ij}(u, \mathbf{q}(u))| \leq C_2(u, v) \|v\|_{W_\infty^1(\mathcal{D})},$$

we find from the above that

$$(9.7) \quad A(u + v, \mathbf{q}(u + v))(u + v, \chi) = A(u, \mathbf{q}(u))(u, \chi) + \tilde{A}(v, \chi) + F(\chi)$$

where  $\tilde{A}$ , which depends on  $u$ , is the bilinear form

$$(9.8) \quad \begin{aligned} \tilde{A}(v, \chi) &= A(u, \mathbf{q}(u))(v, \chi) \\ &+ \int_{\mathcal{D}} \sum_{i,j=1}^n \left( \frac{\partial a_{ij}}{\partial u}(u, \mathbf{q}(u))v + \sum_{k=1}^n \frac{\partial a_{ij}}{\partial q_k}(u, \mathbf{q}(u)) \frac{\partial v}{\partial x_k} \right) \frac{\partial u}{\partial x_i} \frac{\partial \chi}{\partial x_j}, \end{aligned}$$

and where

$$(9.9) \quad |||F|||_{-1, \infty, \mathcal{D}} \leq C_3(u, v) \|v\|_{W_\infty^1(\mathcal{D})}^2,$$

with  $C_3(u, v)$  depending (through the  $a_{ij}$ ) only on  $\|u\|_{W_\infty^1(\mathcal{D})}$  and  $\|u + v\|_{W_\infty^1(\mathcal{D})}$ .

We note that  $\tilde{A}$  is *not* gotten by simply freezing the coefficients of  $A$  at  $u$ .

Under our general assumptions, the form  $\tilde{A}(v, \chi)$  has smooth coefficients.

We now introduce a major assumption:

$$(9.10) \quad \text{The bilinear form } \tilde{A}(v, \chi) \text{ is such that the "global" variant of Theorems 5.5.1 holds.}$$

(This "global" variant is obtained by setting  $d = 1$ ,  $\Omega_0 = \Omega_d = \mathcal{D}$ , and scrapping the negative norm term, in (5.5.4), cf. [Schatz and Wahlbin 1994, Theorem 4.1] for a proof in a simple case.) That (9.10) holds must be checked in each application, cf. [Douglas and Dupont 1975]. Under various simplifying assumptions, the maximum-principle may apply to show that the corresponding linearized boundary value problem has a unique solution.

Example 9.1. Let  $n = 2$  and the partial differential operator in (9.1) equal

$$-div(a(q^2)\nabla u)$$

where  $q^2 = q^2(u) = |\nabla u|^2 = \sum_{i=1}^2 |\frac{\partial u}{\partial x_i}|^2$ . A short calculation establishes that the bilinear form  $\tilde{A}$  in (9.8) has only terms involving  $\frac{\partial v}{\partial x_k} \frac{\partial \chi}{\partial x_j}$  and that the corresponding matrix of coefficients  $\tilde{\mathbf{A}}$  is

$$a(q^2)I + 2a'(q^2)\{q_k q_i\} = \begin{bmatrix} a(q^2) + 2a'(q^2)q_1^2 & 2a'(q^2)q_1 q_2 \\ 2a'(q^2)q_1 q_2 & a(q^2) + 2a'(q^2)q_2^2 \end{bmatrix}.$$

In the special case of

$$a(q^2) = (1 + q^2)^t, \quad t \text{ real,}$$

we have the matrix

$$\tilde{\mathbb{A}} = (1 + q^2)^{t-1} \begin{bmatrix} 1 + q^2 + 2tq_1^2 & 2tq_1q_2 \\ 2tq_1q_2 & 1 + q^2 + 2tq_2^2 \end{bmatrix}.$$

A simple calculation shows that if  $q$  is bounded,  $\tilde{\mathbb{A}}_{11}$  and  $\det \tilde{\mathbb{A}}$  are both uniformly positive for  $t \geq -\frac{1}{2}$ . Hence, cf. e.g. [Shilov 1961, Section 46, Theorem 27] the form  $\tilde{A}$  is uniformly elliptic in these cases. If the original boundary conditions were  $u = 0$  on  $\partial\mathcal{D}$ , the maximum principle gives that the linearized problem has unique solutions. The case  $t = -1/2$  is the minimal surface equation, cf. [Johnson and Thomée 1975], and, for the reader interested in the history of finite elements, [Schellbach 1851]. For the case of the  $p$ -Laplacian, cf. [Barrett and Liu 1993].  $\square$

Remark 9.1. If instead we consider the Neumann problem

$$\begin{aligned} -\operatorname{div}(a(q^2(u))\nabla u) + u &= f \quad \text{in } \mathcal{D}, \\ \frac{\partial u}{\partial u} &= 0 \quad \text{on } \partial\mathcal{D}, \end{aligned}$$

with  $a(q^2) = (1 + q^2)^t$ ,  $t \geq -\frac{1}{2}$ , then [Schatz and Wahlbin 1994, Theorem 4.1] immediately applies to show that the assumption (9.10) holds.  $\square$

Remark 9.2. With  $L(u)$  as in (9.1), perhaps with lower order terms,  $L$  is called *monotone* if, in a formal sense,

$$(L(u + v) - L(u), v) \geq 0,$$

for  $u$  and  $v$  satisfying the boundary conditions under consideration. This can serve to show existence of solutions to (9.1) and (9.2) (for (9.1), in some exact function class). Under further conditions, including so-called strict monotonicity, one may assert that the linearized operator  $\tilde{L}$  corresponding to our form  $\tilde{A}$  in (9.8) is coercive and so has unique solutions. We shall not pursue the details but refer to e.g. [Lions 1969, Chapter 2, Section 2] and [Ženíšek 1990, Chapter 4].  $\square$

Let now  $\tilde{u}_h$  denote the projection of  $u$  into  $S_h$  with respect to the derivative form  $\tilde{A}$ ;

$$(9.11) \quad \tilde{A}(u - \tilde{u}_h, \chi) = 0, \quad \text{for } \chi \in S_h.$$

We assume that (cf. (9.10))

$$(9.12) \quad \|u - \tilde{u}_h\|_{W_\infty^1(\mathcal{D}_h)} \leq Ch^{r-1}.$$

(E.g., with essential boundary conditions,  $\mathcal{D}_h \subseteq \mathcal{D}$  denote mesh domains.)

The idea is now to use the above to compare  $\tilde{u}_h$  to  $u_h$ ; under certain assumptions, the difference  $\tilde{u}_h - u_h$  will be shown to be of higher order. If so, known “linear” superconvergence results about  $u - \tilde{u}_h$  automatically translate to the nonlinear case.

Taking  $u$  to be the solution of (9.1) and  $u + v = u_h$  the solution to (9.2), (9.7) gives

$$(9.13) \quad \tilde{A}(\tilde{u}_h - u_h, \chi) = \tilde{A}(u - u_h, \chi) = -F(\chi), \quad \text{for } \chi \in S_h.$$

Yet another assumption will now be introduced! Namely, that

$$(9.14) \quad \varepsilon \equiv \|u - u_h\|_{W_\infty^1(\mathcal{D}_h)} \equiv \|v\|_{W_\infty^1(\mathcal{D}_h)} = o\left(\frac{1}{\ln 1/h}\right), \text{ as } h \rightarrow 0.$$

E.g., cf. [Douglas and Dupont 1975], if it is known that  $\|u - u_h\|_{H^1(\mathcal{D}_h)} \leq Ch^{r-1}$ , (9.14) would easily follow from inverse assumptions if  $r > n/2 + 1$ . Of course, (9.14) controls the constant  $C_3(u, v)$  in (9.9).

Using the “global” form of Theorem 5.5.1, we now have via (9.9),

$$(9.15) \quad \|\tilde{u}_h - u_h\|_{W_\infty^1(\mathcal{D}_h)} \leq C\varepsilon^2(\ln 1/h).$$

Since

$$(9.16) \quad \begin{aligned} \varepsilon &= \|u - u_h\|_{W_\infty^1(\mathcal{D}_h)} \leq \|\tilde{u}_h - u_h\|_{W_\infty^1(\mathcal{D}_h)} + \|u - \tilde{u}_h\|_{W_\infty^1(\mathcal{D}_h)} \\ &\leq C\|\tilde{u}_h - u_h\|_{W_\infty^1(\mathcal{D}_h)} + Ch^{r-1}, \end{aligned}$$

where we used (9.12), setting  $\eta = \|\tilde{u}_h - u_h\|_{W_\infty^1(\mathcal{D}_h)}$ , we then have

$$(9.17) \quad \eta \leq C\eta^2(\ln 1/h) + Ch^{2(r-1)}(\ln 1/h).$$

From (9.12) and (9.14) we know that  $\eta = o(1/(\ln 1/h))$ , as  $h \rightarrow 0$ . Then (9.17) implies that

$$(9.18) \quad \|\tilde{u}_h - u_h\|_{W_\infty^1(\mathcal{D}_h)} \leq Ch^{2(r-1)}(\ln 1/h),$$

which is our final estimate in this brief chapter.

*In conclusion, if superconvergence points for derivatives or function values (or, postprocessors) are known for the linearized problem  $\tilde{A}(\tilde{u}_h - u, \chi) = 0$ , for  $\chi \in S_h$ , then under the various assumptions introduced in this chapter, the same points are superconvergent if (9.18) permits.*

Remark 9.3. (On tensor products) If  $L(u)$  in (9.1) is of the form

$$Lu = - \sum_{i=1}^n \frac{\partial}{\partial x_i} \left( \alpha(u) \frac{\partial u}{\partial x_i} \right)$$

then the linearized bilinear form is

$$\tilde{A}(v, \chi) = \int_{\mathcal{D}} \sum_{i=1}^n \left[ \alpha(u) \frac{\partial v}{\partial x_i} \frac{\partial \chi}{\partial x_i} + \alpha'(u) \frac{\partial u}{\partial x_i} v \frac{\partial \chi}{\partial x_i} \right].$$

I.e., it is of the type so that the tensor-product theory in Section 6.3 for superconvergence in gradients applies to it.

In a homogeneous Dirichlet problem, e.g., we may also allow a lower order term of the type  $a(u, \mathbf{q}(u))u$  with the theory of Section 6.3 still applying.  $\square$

Remark 9.4. (On Newton’s method) Let  $u^1$  be any approximation to  $u$ , not necessarily in  $S_h$ , and then perform a Newton step in  $S_h$ . I.e., with  $\tilde{A}^1(v, \chi)$  denoting the bilinear form in (9.8) but with  $u$  and  $\mathbf{q}(u)$  replaced by  $u^1$  and  $\mathbf{q}(u^1)$ , i.e., the derivative of  $A$  at  $u^1$  in the direction  $v$ , define  $u_h^2 \in S_h$  by

$$(9.20) \quad \tilde{A}^1(u_h^2 - u^1, \chi) = -A(u^1, \mathbf{q}(u^1))(u^1, \chi) + (f, \chi), \text{ for } \chi \in S_h.$$

We then have, using (9.11) and (9.20),

$$\begin{aligned}
 (9.21) \quad \tilde{A}(u_h^2 - \tilde{u}_h, \chi) &= \tilde{A}(u_h^2 - u^1, \chi) + \tilde{A}(u^1 - u, \chi) \\
 &= (\tilde{A} - \tilde{A}^1)(u_h^2 - u^1, \chi) + \tilde{A}^1(u_h^2 - u^1, \chi) + \tilde{A}(u^1 - u, \chi) \\
 &= [(\tilde{A} - \tilde{A}^1)(u_h^2 - u^1, \chi)] + [-A(u^1, \mathbf{q}(u^1))(u^1, \chi) \\
 &\quad + A(u, \mathbf{q}(u))(u, \chi) + \tilde{A}(u^1 - u, \chi)] \\
 &\equiv F_1(\chi) + F_2(\chi).
 \end{aligned}$$

Here,  $F_2(\chi)$  is as in (9.7), with  $v = u^1 - u$ . Setting

$$(9.22) \quad \delta = \|u^1 - u\|_{W_\infty^1(\mathcal{D}_h)}$$

(which we assume is  $\leq 1$ ) we then have from (9.9),

$$(9.23) \quad |||F_2|||_{-1, \infty, \mathcal{D}_h} \leq C\delta^2.$$

For  $F_1$ , we string it out as

$$(9.24) \quad F_1(\chi) = (\tilde{A} - \tilde{A}^1)(u_h^2 - \tilde{u}_h, \chi) + (\tilde{A} - \tilde{A}^1)(\tilde{u}_h - u, \chi) + (\tilde{A} - \tilde{A}^1)(u - u^1, \chi).$$

Setting

$$(9.25) \quad \eta = \|u_h^2 - \tilde{u}_h\|_{W_\infty^1(\mathcal{D}_h)}$$

we then clearly have

$$(9.26) \quad |||F_1|||_{-1, \infty, \mathcal{D}_h} \leq C\delta \left( \eta + \|\tilde{u}_h - u\|_{W_\infty^1(\mathcal{D}_h)} + \delta \right).$$

Assume for definiteness also the linear result that

$$(9.27) \quad \|\tilde{u}_h - u\|_{W_\infty^1(\mathcal{D}_h)} \leq Ch^{r-1},$$

cf. (9.10), (9.11) and (9.12). Then

$$(9.28) \quad |||F_1|||_{-1, \infty, \mathcal{D}_h} \leq C\delta\eta + C\delta h^{r-1} + C\delta^2.$$

It follows from (9.21), (9.23) and (9.28), under the assumption (9.10), that

$$(9.29) \quad \eta \leq C(\ln 1/h)[\delta\eta + \delta h^{r-1} + \delta^2].$$

If we now finally assume that  $\delta \ln 1/h \rightarrow 0$  as  $h \rightarrow 0$ , we have

$$(9.30) \quad \|u_h^2 - \tilde{u}_h\|_{W_\infty^1(\mathcal{D}_h)} \leq C(\ln 1/h) \left[ h^{2r-2} + \|u^1 - u\|_{W_\infty^1(\mathcal{D}_h)}^2 \right].$$

E.g., if  $\|u^1 - u\|_{W_\infty^1(\mathcal{D}_h)} \leq Kh^{r-1}$ , and if  $K = K(h)$  is not too badly growing with  $1/h$ , we have asserted that one Newton-step in  $S_h$  gives us an approximation  $u_h^2$  in the nonlinear problem which exhibits “the same superconvergence phenomena as in the linearized problem”.

Of course, the reader will have noted that this remark, indeed, this whole chapter, reflects the quadratic convergence of Newton’s method. One may, e.g., start with  $u^1$  a rough approximation, say on a “coarse” grid and iterate this remark, leading to a “multi-grid” approach. In this respect, cf. [Xu, J. 1994].  $\square$

## Chapter 10. Superconvergence in isoparametric mappings of translation invariant meshes: an example.

The example given in this chapter is a special case of the theory in [Cayco, Schatz and Wahlbin 1994].

### 10.1. Introduction.

Let  $\hat{R} = \{\hat{a} \leq \hat{x} \leq b, c \leq \hat{y} \leq d\}$  be a rectangle, and  $\varphi = \varphi(\hat{x}, \hat{y}) = (\varphi^1, \varphi^2)$  a smooth mapping  $\hat{R} \rightarrow R \subset \subset \mathcal{D} \subset \subset \mathbb{R}^2$ , which is one-to-one and onto, and has a smooth inverse.




Figure 10.1.1

Here,  $\mathcal{D}$  is our “physical domain” in the  $x - y$  plane and on the piece  $R$  of it, we have, for whatever reason, decided to install a mesh associated with the map  $\varphi$  of  $\hat{R}$ .

Let us say that we are interested in using quadratic isoparametric triangles. We would then first subdivide  $\hat{R}$  into triangles, say, translation invariant in the  $\hat{x}$  direction with parameter  $h$ , and quasi-uniform in the  $\hat{y}$  direction.




Figure 10.1.2

Let  $\hat{\tau}_i$ ,  $i = 1, \dots, I$  denote these triangles. For each  $\hat{\tau}_i$ , let

$$(10.1.1) \quad \hat{a}_j^i = \text{vertices, } j = 1, 2, 3; \quad \hat{a}_{jk}^i = \text{midpoints of sides, } 1 \leq j < k = 3.$$




Figure 10.1.3

By the mapping  $\varphi$ , these points (let us collectively call them  $\{\hat{a}\}$ ) map to points  $\{a\}$  in  $R$ :




Figure 10.1.4

On each  $\hat{\tau}_i$  we now construct the map

$$(10.1.2) \quad \varphi_{i,h} = (\varphi_{i,h}^1(\hat{x}, \hat{y}), \varphi_{i,h}^2(\hat{x}, \hat{y}))$$

where each  $\varphi_{i,h}^j$ ,  $j = 1, 2$ , is a quadratic in  $\hat{x}$  and  $\hat{y}$ ,

$$(10.1.3) \quad \varphi_{i,h}^j(\hat{x}, \hat{y}) = A_j + B_j \hat{x} + C_j \hat{y} + D_j \hat{x}^2 + E_j \hat{x}\hat{y} + F_j \hat{y}^2, \quad j = 1, 2,$$

such that the vertices  $\hat{a}_j^i$  go to  $a_j^i = \varphi(\hat{a}_j^i)$  and the midpoints  $\hat{a}_{jk}^i$  go to  $a_{jk}^i = \varphi(\hat{a}_{jk}^i)$ .

If  $\varphi$  is fixed smooth and  $h$  is small enough, each such  $\varphi_{i,h}$  exists and is invertible, see e.g. [Ciarlet 1991, Chapter VI, Theorem 37.2]. Also their Jacobians do not vanish. The main point in applying that theorem is the following: If  $\tilde{\tau}_i$  denotes the result of an *affine* map of  $\hat{\tau}_i$  respecting the vertices only, then with  $\tilde{a}_{jk}$  the corresponding midpoints of  $\tilde{\tau}_i$ , we have for the points  $a_{jk} = \varphi(\tilde{a}_{jk})$ ,

$$(10.1.4) \quad |\tilde{a}_{jk} - a_{jk}| = O(h^2).$$

In practice, for  $h$  not “small enough”, one has to watch out for the map being one-to-one and onto, cf. e.g. [Frey, Hall and Porsching 1978]. We denote by  $\varphi_h$  the total map of  $\widehat{R}$  to  $R_h$  (an approximation to  $\varphi$ ) thus constructed, and set  $\tau_i = \varphi_h(\widehat{\tau}_i)$ . In pictures,




Figure 10.1.5

Each  $\tau_i$  is thus a curved “triangle” and under our assumptions on  $\varphi$ , its “length dimensions” are  $O(h)$  in a quasi-uniform fashion for  $h$  small enough.

So far we have constructed a “mesh” on  $R$ , or, on  $R_h$  to be precise. As for the finite element function spaces in this part of the physical domain, they will be given as follows: On each  $\widehat{\tau}_i$ , consider quadratics  $\widehat{\chi}(\widehat{x}, \widehat{y}) = A + B\widehat{x} + C\widehat{y} + D\widehat{x}^2 + E\widehat{x}\widehat{y} + F\widehat{y}^2$ . Under the map  $\varphi_h$ , they naturally give rise to a function  $\chi(x, y)$  on  $\tau_i$ . Or, put otherwise,  $\chi(x, y)$  is such that when mapped to  $\widehat{R}$  by the inverse map  $\varphi_h^{-1}$ , it becomes a quadratic in  $\widehat{x}, \widehat{y}$ .

We shall refer to  $\widehat{R}$  as the “computational domain”.

If then  $\widehat{\chi}(\widehat{x}, \widehat{y})$  is a  $C^0$  quadratic on  $\widehat{R}$ , then  $\chi(x, y)$  is still a continuous function on  $R_h$ ; we refer to them as “isoparametric quadratic”. (The word *isoparametric* means that the same polynomial functions have been used in approximating  $\varphi$  as in giving  $\widehat{\chi}$ . With a slight abuse of terminology,  $\widehat{\tau}_i$  may be referred to as a “reference element” for  $\tau_i$ ; although,  $\widehat{\tau}_i$  would then commonly be blown up to unit size, which we shall not do. We refer to [Ciarlet 1991, Chapter VI] for details.)

We denote by  $S_h$  the isoparametric quadratic finite elements thus constructed, at least on  $R_h$ . Let then as usual  $u$  be given and  $u_h \in S_h$  such that

$$(10.1.5) \quad A(u - u_h, \chi) = 0, \text{ for } \chi \in S_h^{comp}(R_h).$$

Remark 10.1.1. As for the actual computation of a typical term like  $\int_{\tau} a_{12}(x, y) \frac{\partial u_h}{\partial x} \frac{\partial \chi}{\partial y} dx dy$ , with the correspondence  $(x, y) = \varphi_h(\widehat{x}, \widehat{y})$ , one has

$$(10.1.6) \quad \int_{\widehat{\tau}} \widehat{a}_{12} \left[ \frac{\partial}{\partial \widehat{x}} \widehat{u}_h \frac{\partial \widehat{x}}{\partial x} + \frac{\partial}{\partial \widehat{y}} \widehat{u}_h \frac{\partial \widehat{y}}{\partial x} \right] \left[ \frac{\partial \widehat{\chi}}{\partial \widehat{x}} \frac{\partial \widehat{x}}{\partial y} + \frac{\partial \widehat{\chi}}{\partial \widehat{y}} \frac{\partial \widehat{y}}{\partial y} \right] J(\varphi_h) d\widehat{x} d\widehat{y},$$

where  $\partial \widehat{x}/\partial x$  involves derivatives of  $\varphi_h^{-1}$ . Computations are, in practice, done on the “reference elements”  $\widehat{\tau}$  in the “computational” domain and clearly numerical integration is called for, even if the coefficients of the bilinear form were very simple. We shall not consider this numerical integration; if of high enough order, our final superconvergence results still hold.  $\square$

## 10.2. Superconvergence in difference quotients for first derivatives.

We shall first assume that “standard” theory applied to (10.1.5) gives that

$$(10.2.1) \quad h\|u - u_h\|_{W_\infty^1(R_h^0)} + \|u - u_h\|_{L_\infty(R_h^0)} \leq Ch^3,$$

where  $R_h^0 \subset\subset R_h$  with  $O(1)$  separation, for simplicity.

Letting

$$(10.2.2) \quad \widehat{e} = (u - u_h)(\varphi_h^{-1}) \equiv \widehat{u} - \widehat{u}_h,$$

we then have

$$(10.2.3) \quad h\|\widehat{e}\|_{W_\infty^1(\widehat{R}^0)} + \|\widehat{e}\|_{L_\infty(\widehat{R}^0)} \leq Ch^3.$$

(It is elementary to verify that under our assumptions that  $\varphi$  and  $\varphi_h$  are one to one, with the Jacobians and their inverses bounded,  $\varphi^{-1}$  and  $\varphi_h^{-1}$  behave “similarly” in  $L_\infty$  and  $W_\infty^1$ .)

Using now the map  $\varphi_h^{-1}$  we translate (10.1.5) to the computational domain  $\widehat{R}$  so that

$$(10.2.4) \quad \widehat{A}(\widehat{e}, \widehat{\chi}) = 0, \text{ for } \widehat{\chi} \in \widehat{S}_h^{comp}(\widehat{R}).$$

The finite element functions  $\widehat{u}_h$  and  $\widehat{\chi}$  are here ordinary  $C^0$  quadratics on the translation invariant triangular mesh on  $\widehat{R}$ , but the form  $\widehat{A}$  has only piecewise smooth coefficients on each triangle  $\widehat{\tau}$ ; the coefficients are in general discontinuous across element boundaries, cf. (10.1.6). Likewise, the function  $\widehat{u}(\widehat{x}, \widehat{y}) = u(\varphi_h(\widehat{x}, \widehat{y}))$  is now only piecewise smooth. For clarity we shall write

$$(10.2.5) \quad \widehat{A}(\cdot, \cdot) = \widehat{A}_{\varphi_h}(\cdot, \cdot).$$

Consider now the forward difference operator  $\partial_+$  in the translation invariant  $\widehat{x}$ -direction,

$$(10.2.6) \quad \partial_+ v(\widehat{x}, \widehat{y}) = \frac{1}{h}(v(\widehat{x} + h, \widehat{y}) - v(\widehat{x}, \widehat{y})).$$

We let  $\widehat{A}_\varphi$  denote the form obtained by the mapping  $\varphi^{-1}$  from  $R$  to  $\widehat{R}$ . Note that this form has smooth coefficients and satisfies our ellipticity conditions on the highest order terms, by standard arguments.

Let us look  $\widehat{A}_\varphi(\partial v, w)$ ! Consider e.g. one of the highest order terms, where  $\widehat{A}_{12}$

denotes the coefficient associated with  $\frac{\partial v}{\partial \hat{x}} \frac{\partial w}{\partial \hat{y}}$  in the form  $\hat{A}_\varphi$ .

$$\begin{aligned}
 (10.2.7) \quad & \int_{\hat{R}} \hat{A}_{12}(\hat{x}, \hat{y}) \frac{\partial}{\partial \hat{x}} (\partial_+ v) \frac{\partial w}{\partial \hat{y}} d\hat{x} d\hat{y} \\
 &= \frac{1}{h} \int_{\hat{R}} \hat{A}_{12}(\hat{x}, \hat{y}) \left[ \frac{\partial v}{\partial \hat{x}}(\hat{x} + h, \hat{y}) - \frac{\partial v}{\partial \hat{x}}(\hat{x}, \hat{y}) \right] \frac{\partial w}{\partial \hat{y}}(\hat{x}, \hat{y}) d\hat{x} d\hat{y} \\
 &= \frac{1}{h} \int_{\hat{R}} \frac{\partial v}{\partial \hat{x}}(\hat{x}, \hat{y}) \left[ \hat{A}_{12}(\hat{x} - h, \hat{y}) \frac{\partial w}{\partial \hat{y}}(\hat{x} - h, \hat{y}) \right. \\
 &\quad \left. - \hat{A}_{12}(\hat{x}, \hat{y}) \frac{\partial w}{\partial \hat{y}}(\hat{x}, \hat{y}) \right] d\hat{x} d\hat{y} \\
 &= - \int_{\hat{R}} \hat{A}_{12}(\hat{x}, \hat{y}) \frac{\partial v}{\partial \hat{x}}(\hat{x}, \hat{y}) \frac{\partial}{\partial \hat{y}} (\partial_- w)(\hat{x}, \hat{y}) d\hat{x} d\hat{y} \\
 &\quad - \int_{\hat{R}} (\partial_- \hat{A}_{12})(\hat{x}, \hat{y}) \frac{\partial v}{\partial \hat{x}}(\hat{x}, \hat{y}) \frac{\partial w}{\partial \hat{y}}(\hat{x} - h, \hat{y}) d\hat{x} d\hat{y},
 \end{aligned}$$

where  $\partial_-$  denotes the backward difference operator.

We conclude that for  $w$  with compact support,

$$(10.2.8) \quad \hat{A}_\varphi(\partial_+ v, w) = -\hat{A}_\varphi(v, \partial_- w) + F_1(v, w)$$

where, over the relevant domains,

$$(10.2.9) \quad |F_1(v, \psi)| \leq C \|v\|_{W_\infty^1} \|\psi\|_{W_1^1}$$

and, after integration by parts, also,

$$(10.2.10) \quad |F_1(v, \psi)| \leq C \|v\|_{L_\infty} \|\psi\|_{W_1^2}.$$

We apply (10.2.8) to  $v = \hat{e}$  and  $w = \hat{\chi} \in \hat{S}_h^{comp}(\hat{R}_1)$  where  $\hat{R}_1 \subset \subset \hat{R}$ , with  $O(1)$  separation. From (10.2.4) we have since  $\partial_- \hat{\chi} \in S_h^{comp}(\hat{R})$  by translation invariance,

$$(10.2.11) \quad -\hat{A}_\varphi(\hat{e}, \partial_- \hat{\chi}) = (\hat{A}_{\varphi_h} - \hat{A}_\varphi)(\hat{e}, \partial_- \hat{\chi})$$

and by the properties of  $\varphi - \varphi_h$ , cf. (10.1.6) which shows how the forms  $\hat{A}_{\varphi_h}$  and  $\hat{A}_\varphi$  involves derivatives of  $\varphi_h$  and  $\varphi$ , and their inverses, and by (10.2.3),

$$\begin{aligned}
 (10.2.12) \quad & |(\hat{A}_{\varphi_h} - \hat{A}_\varphi)(\hat{e}, \partial_- \hat{\chi})| \leq Ch^2 \|e\|_{W_\infty^1(\hat{R})} \frac{1}{h} \|\hat{\chi}\|_{W_1^1(\hat{R}_1)} \\
 & \leq Ch^3 \|\hat{\chi}\|_{W_1^1(\hat{R}_1)}.
 \end{aligned}$$

In total from (10.2.8) et. seq., for the smooth and uniformly elliptic form  $\hat{A}_\varphi$ ,

$$(10.2.13) \quad \hat{A}_\varphi(\partial_+ \hat{e}, \hat{\chi}) = F(\hat{\chi}), \text{ for } \hat{\chi} \in \hat{S}_h^{comp}(\hat{R}_1)$$

where, using again (10.2.3) in (10.2.9) and (10.2.10),

$$(10.2.14) \quad |||F|||_{-1, \infty, \hat{R}_1} \leq Ch^2$$

and

$$(10.2.15) \quad |||F|||_{-2, \infty, \hat{R}_1} \leq Ch^3.$$

It follows then from Theorem 5.5.2 that, for  $\widehat{R}_0 \subset\subset \widehat{R}_1$ , again with unit separation,

$$(10.2.16) \quad \begin{aligned} \|\partial_+ \widehat{e}\|_{L_\infty(\widehat{R}_0)} &\leq C \min_{\chi \in \widehat{\mathcal{S}}_h} \|\partial_+ \widehat{u} - \widehat{\chi}\|_{L_\infty(\widehat{R}_1)} + Ch^3 \ln 1/h \\ &\quad + C \|\partial_+ \widehat{e}\|_{W_q^{-s}(\widehat{R}_1)}. \end{aligned}$$

The general negative norm term on the right of (10.2.16) can easily be converted into  $\|\widehat{e}\|_{W_q^{-s+1}(\widehat{R}_{1.5})}$  and thus, from (10.2.3),

$$(10.2.17) \quad \|\partial_+ \widehat{e}\|_{L_\infty(R_0)} \leq C \min_{\widehat{\chi} \in \widehat{\mathcal{S}}_h} \|\partial_+ \widehat{u} - \widehat{\chi}\|_{L_\infty(\widehat{R}_1)} + Ch^3 \ln 1/h.$$

It now becomes necessary to introduce some further notation in order to be perfectly clear about the differences incurred by the mappings  $\varphi$  or  $\varphi_h$ .

The “fixed” points are  $\widehat{x}, \widehat{y}$  in the computational domain.

To treat the first term on the right of (10.2.17), we note that there  $\widehat{u}(\widehat{x}, \widehat{y}) \equiv \widehat{u}_{\varphi_h}(\widehat{x}, \widehat{y}) \equiv u(\varphi_h(\widehat{x}, \widehat{y}))$ . With  $\widehat{u}_\varphi(\widehat{x}, \widehat{y}) \equiv u(\varphi(\widehat{x}, \widehat{y}))$ , a smooth function, we have

$$(10.2.18) \quad \partial_+ \widehat{u}_{\varphi_h} - \widehat{\chi} = \partial_+(\widehat{u}_{\varphi_h} - \widehat{u}_\varphi) + (\partial_+ \widehat{u}_\varphi - \widehat{\chi}) \equiv \widehat{I}_1 + \widehat{I}_2.$$

We clearly have for, say,  $\widehat{\chi}$  the interpolant of  $\partial_+ \widehat{u}_\varphi$ ,

$$(10.2.19) \quad \|\widehat{I}_2\|_{L_\infty(\widehat{R}_1)} \leq Ch^3 \|\partial_+ \widehat{u}_\varphi\|_{W_\infty^3(\widehat{R}_1 + Ch)} \leq Ch^3,$$

if  $u \in \mathcal{C}^4$  locally.

For  $\widehat{I}_1$ , we have by Taylor expansion

$$(10.2.20) \quad \begin{aligned} &\partial_+(\widehat{u}_{\varphi_h} - \widehat{u}_\varphi) \\ &= \frac{1}{h} \left( u(\varphi_h(\widehat{x} + h, \widehat{y})) - u(\varphi(\widehat{x} + h, \widehat{y})) \right. \\ &\quad \left. - (u(\varphi_h(\widehat{x}, \widehat{y})) - u(\varphi(\widehat{x}, \widehat{y}))) \right) \\ &= \frac{1}{h} \left( \frac{\partial u}{\partial x}(\varphi(\widehat{x} + h, \widehat{y})) [\varphi_h^1(\widehat{x} + h, \widehat{y}) - \varphi^1(\widehat{x} + h, \widehat{y})] \right. \\ &\quad + \frac{\partial}{\partial y} u(\varphi(\widehat{x} + h, \widehat{y})) [\varphi_h^2(\widehat{x} + h, \widehat{y}) - \varphi^2(\widehat{x} + h, \widehat{y})] \\ &\quad - \left( \frac{\partial u}{\partial x}(\varphi(\widehat{x}, \widehat{y})) [\varphi_h^1(\widehat{x}, \widehat{y}) - \varphi^1(\widehat{x}, \widehat{y})] \right. \\ &\quad \left. \left. + \frac{\partial u}{\partial y}(\varphi(\widehat{x}, \widehat{y})) [\varphi_h^2(\widehat{x}, \widehat{y}) - \varphi^2(\widehat{x}, \widehat{y})] \right) \right) \\ &\quad + \text{HOTS}, \end{aligned}$$

where HOTS denotes terms of the form  $\frac{1}{h}0(Q)$ , with  $Q$  quadratic in  $\varphi_h - \varphi$ . Such HOTS are  $0(h^5)$ .

Shifting the derivative  $\frac{\partial u}{\partial x}(\varphi(\widehat{x} + h, \widehat{y}))$  to  $\frac{\partial u}{\partial x}\varphi(\widehat{x}, \widehat{y})$  (and the same for  $\frac{\partial u}{\partial y}$ ) introduces an error which is bounded by  $0(h^3)$ . So, it remains to consider the error in,

$$(10.2.21) \quad \partial_+(\varphi_h^i - \varphi^i), \quad i = 1, 2.$$

But, clearly, a difference quotient of the interpolant  $\varphi_h^i$  is the interpolant to the same difference quotient of  $\varphi^i$  in our present translation invariant situation. It follows that

$$(10.2.22) \quad \|\widehat{I}_1\|_{L_\infty(\widehat{R}_1)} = \|\partial_+(\widehat{u}_{\varphi_h} - \widehat{u}_\varphi)\|_{L_\infty(\widehat{R}_1)} \leq Ch^3.$$

From (10.2.17) and the above we then have

$$(10.2.23) \quad \|\partial_+\widehat{e}\|_{L_\infty(R_0)} \equiv \|\partial_+(\widehat{u}_{\varphi_h}(\widehat{x}, \widehat{y}) - \widehat{u}_h(\widehat{x}, \widehat{y}))\|_{L_\infty(\widehat{R}_0)} \leq Ch^3 \ln 1/h.$$

Let us now consider a difference operator which is an  $O(h^3)$  approximation to  $\partial v/\partial \widehat{x}$ , e.g.,

$$(10.2.24) \quad \begin{aligned} \partial v(\widehat{x}, \widehat{y}) &= \frac{1}{h} \left[ -\frac{1}{6}(v(\widehat{x} + h, \widehat{y}) - v(\widehat{x} - h, \widehat{y})) \right. \\ &\quad \left. + \frac{4}{3} \left( v\left(\widehat{x} + \frac{h}{2}, \widehat{y}\right) - v\left(\widehat{x} - \frac{h}{2}, \widehat{y}\right) \right) \right]. \end{aligned}$$

This operator is built up from linear combinations of  $\partial_+$  (and translations by  $h$  or  $h/2$ ) and is, in fact, an  $O(h^4)$  approximation to  $\partial v/\partial \widehat{x}$ , as is elementary to check by Taylor expansion.

Then from (10.2.23),

$$(10.2.25) \quad \|\partial(\widehat{u}_{\varphi_h}(\widehat{x}, \widehat{y}) - \widehat{u}_h(\widehat{x}, \widehat{y}))\|_{L_\infty(R_0)} \leq C \left( \ln \frac{1}{h} \right) h^3.$$

So far we haven't used that  $\partial$  is an accurate approximation to  $\partial/\partial \widehat{x}$ . We shall, shortly.

Now,  $\widehat{u}_{\varphi_h}$  is merely a piecewise smooth function. It remains to determine what, if anything, is approximated by  $\partial \widehat{u}_h$  down in the physical domain.

Let us consider

$$(10.2.26) \quad \begin{aligned} \frac{\partial}{\partial \widehat{x}} \widehat{u}_\varphi(\widehat{x}, \widehat{y}) &= \frac{\partial}{\partial \widehat{x}} u(\varphi(\widehat{x}, \widehat{y})) \\ &= \frac{\partial u}{\partial x}(\varphi(\widehat{x}, \widehat{y})) \frac{\partial \varphi^1}{\partial \widehat{x}} + \frac{\partial u}{\partial y}(\varphi(\widehat{x}, \widehat{y})) \frac{\partial \varphi^2}{\partial \widehat{x}} \\ &\equiv \frac{\partial u}{\partial \ell}(\varphi(\widehat{x}, \widehat{y})), \end{aligned}$$

i.e., the directional derivative of  $u$  in a certain direction determined by  $\varphi$  (not  $\varphi_h$ !) at the corresponding point in the physical domain.

Then

$$(10.2.27) \quad \begin{aligned} \frac{\partial u}{\partial \ell}(\varphi(\widehat{x}, \widehat{y})) - \partial \widehat{u}_h(\widehat{x}, \widehat{y}) &= \left( \frac{\partial}{\partial \widehat{x}} - \partial \right) \widehat{u}_\varphi(\widehat{x}, \widehat{y}) + \partial(\widehat{u}_\varphi - \widehat{u}_{\varphi_h})(\widehat{x}, \widehat{y}) + \partial \widehat{e} \\ &\equiv \widehat{J}_1 + \widehat{J}_2 + \partial \widehat{e}. \end{aligned}$$

From (10.2.25) and now using the approximation of properties of the difference quotient  $\partial$ ,

$$(10.2.28) \quad \|\widehat{J}_1 + \partial \widehat{e}\|_{L_\infty(\widehat{R}_0)} \leq Ch^3 \ln 1/h.$$

The result (10.2.22) shows that also

$$(10.2.29) \quad \|\widehat{J}_2\|_{L_\infty(\widehat{R}_0)} \leq Ch^3.$$

We conclude that

$$(10.2.30) \quad \left\| \frac{\partial u}{\partial \ell}(\varphi(\widehat{x}, \widehat{y})) - \partial \widehat{u}_h(\widehat{x}, \widehat{y}) \right\|_{L_\infty(\widehat{R}_0)} \leq Ch^3 \ln 1/h.$$

We may, of course, since  $\varphi - \varphi_h = O(h^3)$ , conclude also that

$$(10.2.31) \quad \left\| \frac{\partial u}{\partial \ell}(\varphi_h(\widehat{x}, \widehat{y})) - \partial \widehat{u}_h(\widehat{x}, \widehat{y}) \right\|_{L_\infty(\widehat{R}_0)} \leq Ch^3 \ln 1/h$$

so that we are now comparing things at the *same* point  $\varphi_h(\widehat{x}, \widehat{y})$  in the physical domain. However, the directional derivative  $\frac{\partial u}{\partial \ell}$  is based on the original smooth map  $\varphi$ , cf. (10.2.26), i.e.,

$$(10.2.32) \quad \frac{\partial u}{\partial \ell}(\varphi_h(\widehat{x}, \widehat{y})) = \frac{\partial u}{\partial x}(\varphi_h(\widehat{x}, \widehat{y})) \frac{\partial \varphi^1(\widehat{x}, \widehat{y})}{\partial \widehat{x}} + \frac{\partial u}{\partial y}(\varphi_h(\widehat{x}, \widehat{y})) \frac{\partial \varphi^2(\widehat{x}, \widehat{y})}{\partial \widehat{x}}.$$

Remark 10.2.1. In our present special case of quadratics we may proceed further as follows. If we assume in our situation of approximating  $\frac{\partial u}{\partial \ell}$  that  $\widehat{y}$  is a mesh point  $\widehat{y}_M$  and that  $\widehat{x}$  is a Gauss-point  $\widehat{x}_G$ , cf. (7.5.3), so that  $\partial \varphi_h^1 / \partial \widehat{x}$  and  $\partial \varphi_h^2 / \partial \widehat{x}$  are superconvergent approximations to  $\partial \varphi^1 / \partial \widehat{x}$  and  $\partial \varphi^2 / \partial \widehat{x}$  at those points (if the mesh is translation invariant in the  $\widehat{y}$  direction as well), then

$$(10.2.33) \quad \left| \frac{\partial u}{\partial \ell_h}(\varphi_h(\widehat{x}_G, \widehat{y}_M)) - \partial \widehat{u}_h(\widehat{x}_G, \widehat{y}_M) \right| \leq Ch^3 \ln 1/h$$

where

$$(10.2.34) \quad \frac{\partial u}{\partial \ell_h}(\cdot) = \frac{\partial u}{\partial x}(\cdot) \frac{\partial \varphi_h^1(\cdot)}{\partial \widehat{x}} + \frac{\partial u}{\partial y}(\cdot) \frac{\partial \varphi_h^2(\cdot)}{\partial \widehat{x}}.$$

Note that such a point  $(\widehat{x}_G, \widehat{y}_M)$ , when mapped by  $\varphi_h$ , is on the boundary of a physical element and that  $\frac{\partial u}{\partial \ell_h}$  is in the tangential direction along the mesh boundary at that point.  $\square$

There are no particular difficulties in treating higher order difference quotients, except, it becomes progressively less clear to identify what they “naturally” correspond to in the physical domain. Actually, it is often more appropriate to work “the other way”: If one desires an approximation to a certain higher order derivative in certain directions at a certain point in the physical domain, one should express that (in, perhaps, a complicated fashion) in the computational domain, and then use difference quotients for derivatives of  $\widehat{u}$  and also use difference quotients to approximate derivatives involved of the mapping function.

Remark 10.2.2. (On mesh-perturbations) In [Zlamal 1977, 1978a, 1978b] and in [Lesaint and Zlamal 1979] (and in a multitude of later papers), meshes were considered which are “slight” perturbations of translation invariant ones (or, slight perturbations of tensor product meshes). Basically, if one desires superconvergence of order  $O(h^s)$ ,  $s = r + \ell$ , for function values, (or,  $s = (r - 1) + \ell$  for derivatives), it is assumed that the mesh is on each element an  $O(h^s)$  perturbation of a regular (translation invariant, or, tensor-product) mesh.

(The perturbation argument is not hard although sometimes longish; at this point in our lectures it may safely be left to the reader.)

Our present curved mesh in the physical domain *cannot* for general  $\varphi$  be viewed as such a perturbation, except for  $s = 2$ , which would not suffice to prove superconvergence by such perturbation techniques for quadratics or higher elements. In the piecewise linear case, a mapped mesh can be viewed on each element as an  $O(h^2)$  perturbation of a regular mesh. Thus, the piecewise linear results on superconvergence of averages of derivatives at symmetry points carry over without much fuss to that situation.  $\square$

## Chapter 11. Superconvergence by averaging: mainly, the $K$ -operator.

### 11.1. Introduction.

In certain cases for second order elliptic partial differential equations we know that

$$(11.1.1) \quad \|u - u_h\|_{W_2^{-\ell}(\mathcal{D})} \leq Ch^{r+\ell},$$

for some  $\ell > 0$ ,  $\ell = r - 2$  being the best we can reasonably expect in the most favorable situation, see Section 6.3. Such higher rate of convergence in a negative norm suggests oscillatory behavior of the error, cf. Section 4.1 and 5.2. For meshes which are locally translation invariant in all of the axes directions, [Bramble and Schatz 1974] found a local “filter-device” to average out such local oscillations in a systematic fashion, the  $K$ -operator, which when applied to  $u_h$  gives  $K * u_h(x) = \int K(x - y)u_h(y)dy$  satisfying

$$(11.1.2) \quad \|u - K * u_h\|_{L_\infty(\Omega_0)} \leq Ch^{r+\ell}.$$

The function  $K$  depends on  $h$  in a simple scaling manner, and it also depends on  $r$  and  $\ell$ . The convolution  $K * u_h(x)$  involves the values of  $u_h$  only in an  $0(h)$  neighborhood of  $x$ . Since the function  $K$  is a piecewise polynomial, the convolution is easy to evaluate, involving only integrals of piecewise polynomials if  $S_h$  consists of piecewise polynomials. Aside from its dependence on  $r$ , the function  $K$  is independent of the actual form of  $S_h$ , and it is also independent of the underlying partial differential operator. It is, of course, enough to calculate  $K * \varphi$  for basis functions  $\varphi$  of  $S_h$ .

If superconvergent approximations are desired, say, at nodal points, evaluation of  $K * u_h$  can be expressed in terms of fixed values of  $K * \varphi$  and the particular coefficients of  $u_h$  in that basis. We refer to [Bramble and Schatz 1977, (5.10) and Appendix] for some examples.

In [Thomée 1977] the argument was extended to include  $O(h^{r+\ell})$  order approximation to any derivative of  $u$ .

For applications of the  $K$ -operator, cf. e.g. [Douglas 1984 and 1985].

In Section 11.3 we shall give Thomée’s argument; in fact, we shall follow his presentation very closely. It involves use of Fourier transforms and we therefore start by introducing a few concepts and results needed in this connection.

### 11.2. Preliminaries on Fourier transforms and multipliers.

For functions  $u \in L_1(\mathbb{R}^n)$  we set

$$(11.2.1) \quad \mathcal{F}u(\xi) \equiv \hat{u}(\xi) = \int e^{-i\langle x, \xi \rangle} u(x) dx,$$

where  $\langle x, \xi \rangle = \sum_{j=1}^n x_j \xi_j$ , with the formal inverse

$$(11.2.2) \quad \mathcal{F}^{-1}v(x) \equiv \overset{\vee}{v}(x) = (2\pi)^{-n} \int e^{i\langle x, \xi \rangle} v(\xi) d\xi.$$

Parseval’s relation is that

$$(11.2.3) \quad \int u \bar{v} dx = (2\pi)^{-n} \int \widehat{u} \overline{\widehat{v}} d\xi,$$

which also holds if  $u$  and  $v$  are in  $L_2(R^n)$  (Plancherel). Also recall that with  $D^\alpha = (\partial/\partial x_1)^{\alpha_1} \cdots (\partial/\partial x_n)^{\alpha_n}$  and  $\xi^\alpha = \xi_1^{\alpha_1} \cdots \xi_n^{\alpha_n}$ ,

$$(11.2.4) \quad \mathcal{F}(D^\alpha u)(\xi) = (i\xi)^\alpha \hat{u}(\xi),$$

that

$$(11.2.5) \quad \mathcal{F}(u(\cdot + y))(\xi) = e^{i(y, \xi)} \hat{u}(\xi),$$

that

$$(11.2.6) \quad \mathcal{F}(h^{-n} u(h^{-1} \cdot))(\xi) = \hat{u}(h\xi),$$

and that

$$(11.2.7) \quad \mathcal{F}(u * v)(\xi) = \mathcal{F}\left(\int u(\cdot - y)v(y)dy\right)(\xi) = \hat{u}(\xi)\hat{v}(\xi).$$

We have from Parseval's relation that

$$(11.2.8) \quad \begin{aligned} \|u\|_{W_2^s(R^n)} &= \left( \sum_{|\alpha| \leq s} \|D^\alpha u\|_{L_2(R^n)}^2 \right)^{1/2} \\ &\simeq \|(1 + |\xi|^2)^{s/2} \hat{u}\|_{L_2(R^n)}, \text{ for } s \geq 0, \end{aligned}$$

where  $\simeq$  denotes equivalence of norms, and, by duality, that

$$(11.2.9) \quad \|u\|_{W_2^{-s}(R^n)} = \sup_{\substack{\varphi \in \mathcal{C}_0^\infty(R^n) \\ \|\varphi\|_{W_2^s(R^n)} = 1}} (u, \varphi) \simeq \|(1 + |\xi|^2)^{-s/2} \hat{u}\|_{L_2(R^n)}.$$

We shall consider operators where  $\hat{u}(\xi) \rightarrow a(\xi)\hat{u}(\xi)$  for  $a(\xi)$  a smooth function on  $R^n$ ; we call  $a(\xi)$  the symbol of the operator. E.g.,  $a(\xi) = e^{i(y, \xi)}$  corresponds to translation by  $y$  for  $u$ . We shall say that  $a(\xi)$  is a Fourier multiplier on  $L_p$ , with  $1 \leq p \leq \infty$ , or,  $a \in M_p$ , if

$$(11.2.10) \quad M_p(a) = \sup\{\|\mathcal{F}^{-1}(a\hat{v})\|_{L_p} : v \in \mathcal{C}_0^\infty(R^n), \|v\|_{L_p} \leq 1\} < \infty.$$

Clearly, from Parseval's relation,

$$(11.2.11) \quad M_2(a) \leq \sup_\xi |a(\xi)|;$$

in fact equality holds. Also, it is easy to see that the product of two functions in  $M_p$  is again in  $M_p$ , and an elementary change of variables argument establishes that if  $a(\xi) \in M_p$ , then for  $h > 0$ ,  $a(h\xi) \in M_p$  with

$$(11.2.12) \quad M_p(a(\cdot)) = M_p(a(h\cdot)).$$

(A fact not unconnected with (11.2.6)!) Further, if  $a(\xi_1, \dots, \xi_m) \in M_p(R^m)$  and  $b(\xi_{m+1}, \dots, \xi_n) \in M_p(R^{n-m})$ , then

$$(11.2.13) \quad a(\xi_1, \dots, \xi_m)b(\xi_{m+1}, \dots, \xi_n) \in M_p(R^n).$$

Here is a simple and convenient criterion which gives that a symbol is in  $M_p$  for any  $1 \leq p \leq \infty$ . The criterion is taken from [Bramble and Schatz 1976, Lemma 2.3].

**Lemma 11.2.1.** Suppose that

$$(11.2.14) \quad \left( \prod_{j=1}^n (1 + \partial/\partial \xi_j) \right) a(\xi) \in L_q(R^n)$$

for some  $1 < q \leq 2$ . Then  $a \in M_p$ , for  $1 \leq p \leq \infty$ .

Proof: We shall show that (11.2.14) implies that  $\overset{\vee}{a}(x) \in L_1(R^n)$ . Then

$$(11.2.15) \quad \mathcal{F}^{-1}(a\widehat{u})(x) = (\overset{\vee}{a} * u)(x)$$

and so (Young's inequality, which is essentially Fubini's theorem),

$$(11.2.16) \quad \|\mathcal{F}^{-1}(a\widehat{u})\|_{L_p} \leq \|\overset{\vee}{a}\|_{L_1} \|u\|_{L_p},$$

i.e.,  $a \in M_p$ , with  $M_p(a) = \|\overset{\vee}{a}\|_{L_1}$ .

We have, cf. (11.2.4),

$$(11.2.17) \quad a(x) = \left( \prod_{j=1}^n \frac{1}{(1 - ix_j)} \right) \left( \prod_{j=1}^n \left( 1 + \frac{\partial}{\partial \xi_j} \right) a(\xi) \right)^{\vee}(x)$$

so that by Hölder's inequality, with  $q$  and  $q'$  conjugate indices,

$$(11.2.18) \quad \int |\overset{\vee}{a}(x)| dx \leq \left( \int \frac{1}{\prod_{j=1}^n (1 + x_j^2)^{q/2}} dx \right)^{1/q} \cdot \left( \int \left| \left( \prod_{j=1}^n \left( 1 + \frac{\partial}{\partial \xi_j} \right) a \right)^{\vee} \right|^{q'} dx \right)^{1/q'}$$

Since  $q > 1$  and since by Hausdorff–Young's inequality,  $\|f\|_{q'} \leq C\|f\|_q$ , for  $1 \leq q \leq 2$ , we obtain that  $\overset{\vee}{a} \in L_1$ .  $\square$

Remark 11.2.1. Hausdorff–Young's inequality is elementary for  $q = 1 : \|f\|_{L_\infty} \leq (2\pi)^{-n} \|f\|_{L_1}$ . For  $q = 2$ , it is Parseval's relation. It follows in between by the Riesz–Thorin interpolation theorem, cf. [Bennet and Sharpley 1988, Theorem 4.1.7.].  $\square$

The construction of the  $K$ -operator is based on the (smootherest)  $B$ -splines. Let  $\chi$  be the characteristic function of  $[-1/2, 1/2]$  and, for  $\ell \geq 1$ ,

$$(11.2.19) \quad \psi_{(\ell)} = \chi * \chi * \cdots * \chi,$$

convolution with  $\ell$  factors.  $\psi_{(\ell)}$  is thus the one-dimensional (smootherest)  $B$ -spline of “order”  $\ell$ ; e.g.,  $\psi_{(2)}$  is the piecewise linear basis function on  $[-1, 1]$ . We set, for  $h > 0$ ,

$$(11.2.20) \quad \psi_{(\ell),h}(x) = h^{-1} \psi_{(\ell)}(h^{-1}x).$$

We have by a simple calculation that  $\widehat{\chi}(\sigma) = \sin(\sigma/2)/(\sigma/2)$  and so,

$$(11.2.21) \quad \widehat{\psi}_{(\ell),h}(\xi) = \widehat{\psi}_{(\ell)}(h\xi) = \left( \frac{\sin(\frac{1}{2}h\xi)}{\frac{1}{2}h\xi} \right)^\ell,$$

where we used (11.2.6) and (11.2.7). Since  $\|\psi_\ell\|_{L_1} = 1$ ,

$$(11.2.22) \quad M_p(\widehat{\psi}_{(\ell),h}) \leq 1, \text{ independent of } p, \ell \text{ and } h,$$

cf. (11.2.16) and (11.2.12). For  $\alpha = (\alpha_1, \dots, \alpha_N)$  a multi-index, we set

$$(11.2.23) \quad \psi_{(\alpha)}(x) = \prod_{j=1}^n \psi_{(\alpha_j)}(x_j)$$

and correspondingly,  $\psi_{(\alpha),h}(x) = h^{-n}\psi_{(\alpha)}(h^{-1}x)$ . (For simplicity, we assume that “ $h$ ” is the same in each direction.) The analogue of (11.2.22) holds, cf. (11.2.13).

We shall have occasion to use the special multi-index

$$(11.2.24) \quad \mathbf{m} = (1, 1, \dots, 1) \in \mathbb{Z}^n.$$

Hence  $\psi_{(\ell\mathbf{m})}$  denotes the tensor-product  $B$ -spline of the same order  $\ell$  in each variable.

Finally, we shall use the difference operators in the axes directions,

$$(11.2.25) \quad \partial_h^\alpha = \partial_{h,1}^{\alpha_1} \cdots \partial_{h,n}^{\alpha_n}, \text{ where } \partial_{h,j} v(x) = \left( v\left(x + \frac{1}{2}he_j\right) - v\left(x - \frac{1}{2}he_j\right) \right)/h.$$

We have from an elementary calculation, using (11.2.5),

$$(11.2.26) \quad \mathcal{F}(\partial_h^\alpha v)(\xi) = (2i)^{|\alpha|} \prod_{j=1}^n \left( \frac{\sin(\frac{1}{2}h\xi_j)}{h} \right)^{\alpha_j} \widehat{v}(\xi).$$

From (11.2.4), (11.2.21) and (11.2.26) we see that

$$\begin{aligned} (11.2.27) \quad \mathcal{F}(D^\alpha \psi_{(\alpha+\ell\mathbf{m}),h}) &= (i\xi)^\alpha \prod_{j=1}^n \left( \frac{\sin(\frac{1}{2}h\xi_j)}{\frac{1}{2}h\xi_j} \right)^{\alpha_j + \ell} \\ &= (2i)^\alpha \prod_{j=1}^n \left( \frac{\sin(\frac{1}{2}h\xi_j)}{h} \right)^{\alpha_j} \prod_{j=1}^n \left( \frac{\sin(\frac{1}{2}h\xi_j)}{\frac{1}{2}h\xi_j} \right)^\ell \\ &= \mathcal{F}(\partial_h^\alpha \xi_{(\ell\mathbf{m}),h}), \end{aligned}$$

or,

$$(11.2.28) \quad D^\alpha \psi_{(\alpha+\ell\mathbf{m}),h} = \partial_h^\alpha \psi_{(\ell\mathbf{m}),h}.$$

In words, the derivative of a certain spline is the difference quotient of another spline. *This simple result is fundamental to our development.*

For more details about Fourier multipliers we refer to [Hörmander 1960] or [Brenner, Thomée and Wahlbin 1975, Chapter 1].

### 11.3. The $K$ -operator in general.

In the present section we have a smooth function  $u$  and another function  $v$  that is continuous. For the purposes of this section it will play no role that  $v$  is in fact equal to  $u_h$ , a finite element approximation to  $u$ , and it will consequently also be immaterial that the finite element space is translation invariant. Our main result is the following, where  $K_h^{(\alpha, \ell, p)}$  is a “simple” local function built up from tensor-product smoothest  $B$ -splines. (It is defined in (11.3.15) below.)

**Theorem 11.3.1.** *Let  $\alpha$  be a given multi-index, let  $\ell \geq 1$  and  $p \geq 1$ . Let  $u \in C^{2p+|\alpha|}(\Omega_1)$ ,  $v \in C(\Omega_2)$ , where  $\Omega_0 \subset\subset \Omega_1 \subset\subset \Omega_2$  with unit size separations. Further, set  $N_0 = [n/2] + 1$ . Then for  $h$  sufficiently small,*

$$(11.3.1) \quad \begin{aligned} \|D^\alpha u - K_h^{(\alpha, \ell, p)} * (\partial_h^\alpha v)\|_{L_\infty(\Omega_0)} \\ \leq C \left( h^{2p} \|u\|_{W_\infty^{2p+|\alpha|}(\Omega_1)} + \sum_{|\beta| \leq \ell+N_0} \|\partial_h^{\alpha+\beta}(u-v)\|_{W_2^{-\ell}(\Omega_1)} \right. \\ \left. + h^\ell \sum_{|\beta| \leq \ell} \|\partial_h^{\alpha+\beta}(u-v)\|_{L_\infty(\Omega_1)} \right). \end{aligned}$$

Note that  $\partial_h^\alpha$  is not a high-order approximation to  $D^\alpha$ , but that in turn the  $K$ -operator depends on  $\alpha$ .

A basic step in the construction of the  $K$ -operator is the following in the one-dimensional case. The construction is given in the proof.

**Lemma 11.3.1.** *Let  $q \geq 1$ ,  $p \geq 1$ . There exists a unique trigonometric polynomial  $k_{q,p}(\sigma)$  of order  $p-1$  such that*

$$(11.3.2) \quad k_{q,p}(\sigma)(\hat{\chi}(\sigma))^q = 1 + O(|\sigma|^{2p}), \text{ as } |\sigma| \rightarrow 0.$$

Proof: We have, cf. (11.2.21), that  $\hat{\chi}(\sigma)^q = \left(\frac{\sin(\frac{1}{2}\sigma)}{\frac{1}{2}\sigma}\right)^q$ , or, setting  $\tau = \sin(\frac{1}{2}\sigma)$ , for small  $\tau$  or  $\sigma$ , Maclaurin expanding we have

$$(11.3.3) \quad \hat{\chi}(\sigma)^{-q} = \left(\frac{\arcsin \tau}{\tau}\right)^q = \sum_{j=0}^{\infty} \gamma_{q,j} \tau^{2j}.$$

We may therefore choose

$$(11.3.4) \quad k_{q,p}(\sigma) = \sum_{j=0}^{p-1} \gamma_{q,j} \left(\sin\left(\frac{1}{2}\sigma\right)\right)^{2j} = \sum_{j=0}^{p-1} \gamma_{q,j} 2^{-j} (1 - \cos \sigma)^j.$$

Uniqueness of  $k_{q,p}$  is immediate. For, if  $k_{q,p}^1$  and  $k_{q,p}^2$  both satisfy (11.3.2), then  $k_{q,p}^1 - k_{q,p}^2$  is a trigonometric polynomial of degree  $\leq p-1$  which vanishes to order  $2p$  at the origin. It is hence identically zero.  $\square$

For  $\beta = (\beta_1, \dots, \beta_n)$  with all  $\beta_j > 0$  and  $p \geq 1$ , we now define the  $n$ -dimensional trigonometric polynomial

$$(11.3.5) \quad k^{(\beta, p)}(\xi) = \prod_{j=1}^n k_{\beta_j, p}(\xi_j) = \sum_{\gamma} k_{\gamma}^{(\beta, p)} e^{-i\langle \xi, \gamma \rangle}.$$

Note that the sum on the right is finite.

Using the coefficient of this polynomial and the  $B$ -splines introduced in Section 11.2, we set

$$(11.3.6) \quad \mathcal{K}^{(\beta,p)}(x) = \sum_{\gamma} k_{\gamma}^{(\beta,p)} \psi_{(\beta)}(x - \gamma).$$

Then, from (11.2.5) and the above,

$$(11.3.7) \quad \begin{aligned} \hat{\mathcal{K}}^{(\beta,p)}(\xi) &= k^{(\beta,p)}(\xi) \hat{\psi}_{(\beta)}(\xi) = \prod_{j=1}^n k_{\beta_j,p}(\xi_j) \hat{\chi}(\xi_j)^{\beta_j} \\ &= 1 + O(|\xi|^{2p}), \quad \text{as } |\xi| \rightarrow 0. \end{aligned}$$

We also set

$$(11.3.8) \quad \mathcal{K}_h^{(\beta,p)}(x) = h^{-n} \mathcal{K}^{(\beta,p)}(h^{-1}x).$$

The relation (11.3.7) says that, in some sense, convolution with  $\mathcal{K}_h^{(\beta,p)}$  is an approximate identity of order  $h^{2p}$ . More precisely we have the following.

**Lemma 11.3.2.** *Let  $\beta = (\beta_1, \dots, \beta_n)$  with all  $\beta_j > 0$ , and let  $p \geq 1$ . Then for  $\Omega_1$ , with  $\Omega_2 = \Omega_1 + Mh$ ,  $M = M(\beta, p)$ ,*

$$(11.3.9) \quad \|v - \mathcal{K}_h^{(\beta,p)} * v\|_{L_{\infty}(\Omega_1)} \leq Ch^{2p} \|v\|_{W_{\infty}^{2p}(\Omega_2)}.$$

Proof: Set

$$(11.3.10) \quad \tilde{k}_j(\sigma) = k_{\beta_j,p}(\sigma) \hat{\chi}(\sigma)^{\beta_j}.$$

From (11.3.2) and since  $k_{\beta_j,p}(\sigma)$  and  $\hat{\chi}(\sigma)$  are bounded together with their first derivatives, we find using Lemma 11.2.1 that the analytic function  $(1 - \tilde{k}_j(\xi_j))\xi_j^{-2p} \in M_{\infty}(R^1)$ . Thus, by (11.2.4), using also (11.2.12) in one dimension, and (11.2.13),

$$(11.3.11) \quad \begin{aligned} &\|\mathcal{F}^{-1}(1 - \tilde{k}_j(h\xi_j))\widehat{v}\|_{L_{\infty}(R^n)} \\ &= h^{2p} \|\mathcal{F}^{-1}((1 - \tilde{k}_j(h\xi_j))(h\xi_j)^{-2p} \widehat{(D_j^{2p}v)})\|_{L_{\infty}(R^n)} \\ &\leq Ch^{2p} \|v\|_{W_{\infty}^{2p}(R^n)}. \end{aligned}$$

Now write by elementary algebra

$$(11.3.12) \quad 1 - \hat{\mathcal{K}}_h^{(\beta,p)}(h\xi) = 1 - \prod_{j=1}^n \tilde{k}_j(h\xi_j) = \sum_{j=1}^n \prod_{\ell < j} \tilde{k}_{\ell}(h\xi_{\ell})(1 - \tilde{k}_j(h\xi_j)).$$

It is easily seen (e.g., from (11.2.5), any trigonometric polynomial is in  $M_{\infty}$ , and  $\hat{\chi}$  is in  $M_{\infty}$  by (11.2.22)) that  $\tilde{k}_j \in M_{\infty}$ , and hence we have via (11.3.11) that

$$(11.3.13) \quad \|v - \mathcal{K}_h^{(\beta,p)} * v\|_{L_{\infty}(R^n)} \leq Ch^{2p} \|v\|_{W_{\infty}^{2p}(R^n)}.$$

If, as in (11.3.9), we are only interested in the norm over  $\Omega_1$ , since  $\mathcal{K}_h^{(\beta,p)} * v$  only involves the values of  $v$  over  $\Omega_2$ , we may extend  $v$  from  $\Omega_2$  in any manner whatsoever to  $R^n$  and obtain (11.3.9) from (11.3.13).  $\square$

We are now ready to define  $K_h^{(\alpha, \ell, p)}$ . With  $\mathbf{m} = (1, 1, \dots, 1)$  and  $\ell \geq 1$ , recall from (11.2.28) that  $D^\alpha \psi_{(\alpha+\ell\mathbf{m}), h} = \partial_h^\alpha \psi_{(\ell\mathbf{m}), h}$ . We hence have, cf. (11.3.6),

$$(11.3.14) \quad \begin{aligned} D^\alpha K_h^{(\alpha+\ell\mathbf{m}, p)} &= D^\alpha \left( \sum_\gamma k_\gamma^{(\alpha+\ell\mathbf{m}, p)} \psi_{(\alpha+\ell\mathbf{m}), h}(x - h\gamma) \right) \\ &= \sum_\gamma k_\gamma^{(\alpha+\ell\mathbf{m}, p)} \partial_h^\alpha \psi_{(\ell\mathbf{m}), h}(x - h\gamma) \\ &\equiv \partial_h^\alpha K_h^{(\alpha, \ell, p)} \end{aligned}$$

where thus

$$(11.3.15) \quad K_h^{(\alpha, \ell, p)}(x) = \sum_\gamma k_\gamma^{(\alpha+\ell\mathbf{m})} \psi_{(\ell\mathbf{m}), h}(x - \gamma h).$$

Having finally defined the operator occurring in Theorem 11.3.1, we shall now prove that theorem.

Proof of Theorem 11.3.1. We have

$$(11.3.16) \quad \begin{aligned} D^\alpha u - K_h^{(\alpha, \ell, p)} * (\partial_h^\alpha v) \\ &= [D^\alpha u - K_h^{(\alpha, \ell, p)} * (\partial_h^\alpha u)] + K_h^{(\alpha, \ell, p)} * \partial_h^\alpha (u - v) \\ &\equiv I_1 + I_2. \end{aligned}$$

Here, from (11.3.14), and by the nature of a convolution,

$$(11.3.17) \quad \begin{aligned} I_1 &= D^\alpha u - \partial_h^\alpha (K_h^{(\alpha, \ell, p)} * u) \\ &= D^\alpha u - (\partial_h^\alpha K_h^{(\alpha, \ell, p)}) * u \\ &= D^\alpha u - (D^\alpha K_h^{(\alpha+\ell\mathbf{m}, p)}) * u \\ &= D^\alpha u - K_h^{(\alpha+\ell\mathbf{m}, p)} * D^\alpha u. \end{aligned}$$

Thus from Lemma 11.3.2,

$$(11.3.18) \quad \|I_1\|_{L_\infty(\Omega_0)} \leq Ch^{2p} \|u\|_{W_\infty^{|\alpha|+2p}(\Omega_1)}.$$

We see that the correct estimate for  $I_2$ , and hence the theorem, would come from the following, cf. (11.3.15):

For  $\ell \geq 1$ ,  $w$  continuous,

$$(11.3.19) \quad \begin{aligned} &\|\psi_{(\ell\mathbf{m}), h} * w\|_{L_\infty(\Omega_0)} \\ &\leq C \left( \sum_{|\beta| \leq \ell+N_0} \|\partial_h^\beta w\|_{W_2^{-\ell}(\Omega_1)} + h^\ell \sum_{|\beta| \leq \ell} \|\partial_h^\beta w\|_{L_\infty(\Omega_1)} \right). \end{aligned}$$

To show (11.3.19), let  $\varphi \in C_0^\infty(-1, 1)$  with  $\varphi \equiv 1$  near the origin, and set  $\Phi(\xi) = \prod_{i=1}^n \varphi(\xi_j)$ . We then have (for  $w$  with compact support — we may change the function

at will outside of  $\Omega_2$ ),

$$\begin{aligned}
 (11.3.20) \quad \psi_{(\ell\mathbf{m}), h} * w &= \mathcal{F}^{-1}(\widehat{\psi}_{(\ell\mathbf{m})}(h\xi)\widehat{w}(\xi)) \\
 &= \mathcal{F}^{-1}(\widehat{\psi}_{(\ell\mathbf{m})}(h\xi)\Phi(h\xi)\widehat{w}(\xi)) \\
 &\quad + \mathcal{F}^{-1}(\widehat{\psi}_{(\ell\mathbf{m})}(h\xi)(1 - \Phi(h\xi))\widehat{w}(\xi)) \\
 &\equiv \mathcal{J}_1 + \mathcal{J}_2.
 \end{aligned}$$

For  $\mathcal{J}_1 = \mathcal{J}_1(x)$  we have from Cauchy–Schwarz’s inequality,

$$\begin{aligned}
 (11.3.21) \quad |\mathcal{J}_1| &= (2\pi)^{-n} \left| \int (1 + |\xi|^2)^{-N_0/2} e^{i\langle x, \xi \rangle} (1 + |\xi|^2)^{N_0/2} \right. \\
 &\quad \cdot \left. \widehat{\psi}_{(\ell\mathbf{m})}(h\xi)\Phi(h\xi)\widehat{w}(\xi) d\xi \right| \\
 &\leq C \|(1 + |\xi|^2)^{N_0/2} \widehat{\psi}_{(\ell\mathbf{m})}(h\xi)\Phi(h\xi)\widehat{w}(\xi)\|_{L_2(R^n)}.
 \end{aligned}$$

For  $h\xi$  in the support of  $\Phi(h\xi)$ ,

$$(11.3.22) \quad 1 + |\xi|^2 \leq C \left( 1 + \sum_{j=1}^n \left( \sin\left(\frac{1}{2}h\xi_j\right)/h \right)^2 \right).$$

Next note that the function  $\frac{a}{1+a} = 1 - \frac{1}{1+a}$  is increasing. Since  $\left(\frac{\sin(hy)}{h}\right)^2 \leq y^2$  it follows that

$$(11.3.23) \quad \left(\frac{\sin(hy)}{h}\right)^2 = \left(\frac{\sin(hy)}{h}\right)^2 \frac{1}{y^2} \leq \frac{1 + \left(\frac{\sin(hy)}{h}\right)^2}{1 + y^2}.$$

Thus, since  $\frac{\sin(\sigma)}{\sigma}$  is decreasing for  $0 \leq \sigma \leq 1$ , and bounded by 1, we have with  $\xi_0$  denoting the largest component of  $\xi$  (for  $|h\xi_0| < 1$ ),

$$\begin{aligned}
 (11.3.24) \quad \widehat{\psi}_{(\ell\mathbf{m})}(h\xi)^2 &\leq \left(\frac{\sin(\frac{1}{2}h\xi_0)}{\frac{1}{2}h\xi_0}\right)^{2\ell} \leq \left(\frac{1 + (\sin(\frac{1}{2}h\xi_0)/h)^2}{1 + \frac{1}{2}\xi_0^2}\right)^\ell \\
 &\leq C \frac{(1 + \sum_{j=1}^n (\sin(\frac{1}{2}h\xi_j)/h)^{2\ell})}{(1 + |\xi|^2)^\ell}.
 \end{aligned}$$

It follows from (11.3.22) and (11.3.24) that

$$\begin{aligned}
 (11.3.25) \quad &|(1 + |\xi|^2)^{N_0/2} \widehat{\psi}_{(\ell\mathbf{m})}(h\xi)\Phi(h\xi)| \\
 &\leq C \left( 1 + \sum_{j=1}^n \left( \sin\left(\frac{1}{2}h\xi_j\right)/h \right)^{2(N_0+\ell)} \right)^{1/2} (1 + |\xi|^2)^{-\ell/2}.
 \end{aligned}$$

By (11.2.9),  $\|w\|_{W_2^{-\ell}(R^n)} \simeq \|(1 + |\xi|^2)^{-\ell/2} \widehat{w}\|_{L_2(R^n)}$  and from (11.2.26), we find via

Parseval's relation that

$$\begin{aligned}
 (11.3.26) \quad |\mathcal{J}_1| &\leq C \left( \|w\|_{W_2^{-\ell}(R^n)} + \sum_{j=1}^n \|\partial_{h,j}^{N_0+\ell} w\|_{W_2^{-\ell}(R^n)} \right) \\
 &\leq C \sum_{|\beta| \leq \ell + N_0} \|\partial_h^\beta w\|_{W_2^{-\ell}(R^n)} \\
 &\leq C \sum_{|\beta| \leq \ell + N_0} \|\partial_h^\beta w\|_{W_2^{-\ell}(\Omega_1)}.
 \end{aligned}$$

(In the last step we again used the freedom to change  $w$  at will outside our basic domain.)

For  $\mathcal{J}_2$  (defined in (11.3.20)), using elementary algebra we write it as (cf. 11.3.12),

$$(11.3.27) \quad \mathcal{J}_2 = \mathcal{F}^{-1} \left( \left( \prod_j \widehat{\chi}(h\xi_j)^\ell \right) \left( \sum_j \prod_{i < j} \varphi(h\xi_i)(1 - \varphi(h\xi_j)) \widehat{w} \right) \right).$$

Now  $\widehat{\chi}$  and  $\varphi$  are in  $M_\infty$ , and likewise is  $(1 - \varphi(\xi_j))/\xi_j^\ell$ , by Lemma 11.2.1 and (11.2.13). Thus, cf. (11.2.26),

$$\begin{aligned}
 (11.3.28) \quad |\mathcal{J}_2| &\leq C \sum_{j=1}^n \|\mathcal{F}^{-1}((1 - \varphi(h\xi_j))\widehat{\chi}(h\xi_j)^\ell \widehat{w})\|_{L_\infty(R^n)} \\
 &= C \sum_{j=1}^n \left\| \mathcal{F}^{-1} \left( \left( \frac{1 - \varphi(h\xi_j)}{(\frac{1}{2}h\xi_j)^\ell} \right) \left( \sin \left( \frac{1}{2}h\xi_j \right) \right)^\ell \widehat{w} \right) \right\|_{L_\infty(R^n)} \\
 &= Ch^\ell \sum_{j=1}^n \left\| \mathcal{F}^{-1} \left( \left( \frac{1 - \varphi(h\xi_j)}{(h\xi_j)^\ell} \right) (\widehat{\partial_{h,j}^\ell w}) \right) \right\|_{L_\infty(R^n)} \\
 &\leq Ch^\ell \sum_{j=1}^n \|\partial_{h,j}^\ell w\|_{L_\infty(R^n)} \leq Ch^\ell \sum_{j=1}^n \|\partial_{h,j}^\ell w\|_{L_\infty(\Omega_1)}.
 \end{aligned}$$

Together, (11.3.26) and (11.3.28) show (11.3.19). As we have already noticed, this proves Theorem 11.3.1.  $\square$

#### 11.4. The $K$ -operator applied to finite element approximations in second order elliptic problems.

We apply Theorem 11.3.1 with  $v = u_h$ , where  $u_h \in S_h$  is given with  $A(u - u_h, \chi) = 0$  for  $\chi \in S_h^{comp}(\Omega_2)$ ,  $\Omega_0 \subset \subset \Omega_1 \subset \subset \Omega_2$ . Here  $S_h$  is translation invariant on  $\Omega_2$  in all the axes-directions  $x_j$ ,  $j = 1, \dots, n$ . (For simplicity we assume that the translation parameter is the same,  $h$ , in all directions.) From our investigations in Chapter 8 we have that

$$(11.4.1) \quad \|\partial_h^{\alpha+\beta}(u - u_h)\|_{L_\infty(\Omega_1)} \leq Ch^r$$

under various assumptions. (The logarithmic factor occurring in Chapter 8 may be avoided, cf. Remark 8.4.1.) Choosing  $2p \geq r + \ell$ , in order to obtain superconvergence of order  $h^{r+\ell}$ , it thus remains to estimate  $\|\partial_h^{\alpha+\beta}(u - u_h)\|_{W_2^{-\ell}(\Omega_1)}$ .

In the case of a basic operator with constant coefficients, we have

$$(11.4.2) \quad A(\partial_h^{\alpha+\beta}(u - u_h), \chi) = A(u - u_h, (\partial_h^{\alpha+\beta})\chi) = 0, \text{ for } \chi \in S_h^{comp}(\Omega_2).$$

We now need a local error estimate in a negative norm. We have not treated such estimates in these lectures, and we shall not give much detail here. Referring to [Nitsche and Schatz 1974, Lemma 4.2] we have

$$(11.4.3) \quad \begin{aligned} & \| \partial_h^{\alpha+\beta}(u - u_h) \|_{W_2^{-\ell}(\Omega_1)} \\ & \leq Ch^\gamma \| \partial_h^{\alpha+\beta}(u - u_h) \|_{W_2^1(\Omega_2)} + C \| \partial_h^{\alpha+\beta}(u - u_h) \|_{W_2^{-\ell-L}(\Omega_2)} \end{aligned}$$

for any  $L$ , with  $\gamma = \min(\ell+1, r-1)$ . The  $H^1$ -estimate for  $\partial_n^{\alpha+\beta}(u - u_h)$  is localized essentially as in Chapter 8 (again, details can be found in [Nitsche and Schatz 1974]), and the term involving  $W_2^{-\ell-L}$  on the right may easily be converted to  $\|u - u_h\|_{W_2^{-\ell}}$ , by taking  $L \geq |\alpha| + |\beta|$ . The upshot is that, if  $u$  is smooth enough,

$$(11.4.4) \quad \begin{aligned} & \| \partial_h^{\alpha+\beta}(u - u_h) \|_{W_2^{-\ell}(\Omega_1)} \\ & \leq Ch^\gamma \min_{\chi \in S_h} \| \partial_h^{\alpha+\beta}u - \chi \|_{W_2^1(\Omega_2)} + C \| u - u_h \|_{W_2^{-\ell}(\Omega_2)} \\ & \leq Ch^{r+\ell} \| u \|_{W_2^{r+|\alpha|+|\beta|}(\Omega_2)} + C \| u - u_h \|_{W_2^{-\ell}(\Omega_2)}, \end{aligned}$$

for  $\ell \leq r-2$ .

Thus, provided  $1 \leq \ell \leq r-2$  and  $\|u - u_h\|_{W_2^{-\ell}(\Omega_2)} \leq Ch^{r+\ell}$ , we have

$$(11.4.5) \quad \| D^\alpha u - K_h^{(\alpha, \ell, p)} * (\partial_h^\alpha u_h) \|_{L_\infty(\Omega_0)} \leq Ch^{r+\ell}.$$

The same estimate obtains in the case of variable coefficients, again using the ideas of [Nitsche and Schatz 1974] to prove (11.4.4). (There will be a lot of garbage terms when commuting the difference operator on to  $\chi$  in (11.4.2). These are handled by induction.) The reader will find details in [Bramble and Schatz 1977, Lemma 4.1].

## 11.5. Boundary integral equations and the $K$ -operator: an example.

A particularly dramatic use of the  $K$ -operator is given in [Tran 1993] in the context of boundary integral equations. We shall need to give a certain amount of background material in order to appreciate it. For deeper background, cf. e.g. [Sloan 1992].

We consider the Dirichlet problem on a disc in the plane,

$$(11.5.1) \quad \Delta u = 0 \quad \text{in } B_R = \{x : |x| < R\} \subset \subset \mathbb{R}^2,$$

$$(11.5.2) \quad u = g \quad \text{on } \Gamma_R = \partial B_R.$$

The so-called “direct” method of formulating this problem as an integral equation over  $\Gamma_R$  goes as follows.

Let  $\tilde{p}(y) = \frac{\partial u}{\partial n}(y)$ , the outward normal derivative for  $y \in \Gamma_R$ . By Green’s theorem, for  $x$  in  $\overset{\circ}{B}_R$ , the interior of  $B_R$  (and  $\log$  being the natural,  $e$ -based, logarithmic function, as is traditional in the present setting),

$$(11.5.3) \quad \begin{aligned} u(x) &= \frac{1}{2\pi} \int_{\Gamma_R} \left[ \frac{\partial}{\partial n_y} (\log|x-y|) u(y) - \log|x-y| \tilde{p}(y) \right] d\ell_y \\ &= \frac{1}{2\pi} \int_{\Gamma_R} \left[ \frac{\partial}{\partial n_y} (\log|x-y|) g(y) - \log|x-y| \tilde{p}(y) \right] d\ell_y, \quad x \in \overset{\circ}{B}_R. \end{aligned}$$

Letting  $x$  tend to  $\Gamma_R$  we have, in a standard way, the following integral equation over  $\Gamma_R$ ,

$$(11.5.4) \quad -\frac{1}{\pi} \int_{\Gamma_R} \log|x-y| \tilde{p}(y) d\ell_y \\ = g(x) - \frac{1}{\pi} \int_{\Gamma_R} \frac{\partial}{\partial n_y} (\log|x-y|) g(y) d\ell_y, \text{ for } x \in \Gamma_R.$$

(The method is called “direct” because we haven’t used reformulation in terms of double or single layer potentials, or, some such thing; in fact our unknown, the outflow derivative  $\tilde{p}(y)$ , is frequently of major interest in applications. Note that we have to know a fundamental solution to (11.5.1) in order to derive the boundary integral equation (11.5.4). This is the principal limitation of the method.)

It can be shown that (11.5.4) is equivalent with (11.5.1)–(11.5.2), provided  $R \neq 1$ . (If  $R = 1$ , since then  $\int_{\Gamma_1} \log|x-y| d\ell_y = 0$ , we are in trouble, at least we loose uniqueness in (11.5.4).)

Now parametrize the circle by

$$(11.5.5) \quad x(t) = R(\cos 2\pi t, \sin 2\pi t), \quad 0 \leq t \leq 1.$$

By use of elementary trigonometry, we then have

$$(11.5.6) \quad -\frac{1}{\pi} \int_{\Gamma_R} \log(|x-y|) \tilde{p}(y) d\ell_y \\ = \frac{-1}{2\pi} \int_0^1 \log(R^2(\cos 2\pi t - \cos 2\pi s)^2 \\ + (\sin 2\pi t - \sin 2\pi s)^2) \tilde{p}(y(s)) |y'(s)| ds \\ = -2 \int_0^1 \log(|2R \sin(\pi(t-s))|) \tilde{p}(y(s)) \frac{R}{2\pi} ds.$$

We also note that

$$(11.5.7) \quad \frac{\partial}{\partial n_y} \log|x-y| = \frac{n_y \cdot (y-x)}{|x-y|^2} = \frac{n_{y(s)} \cdot (y(s)-x(t))}{|x(t)-y(s)|^2}$$

is a smooth function of  $t$  and  $s$  if the boundary is smooth. Indeed, it equals the constant  $(2R)^{-1}$  for the circle. Letting then  $G(t)$  denote the right hand side of (11.5.4), and setting  $p(s) = \frac{R}{2\pi} \tilde{p}(y(s))$ , we thus obtain the equation

$$(11.5.8) \quad Ap(t) \equiv -2 \int_0^1 \log|2R \sin(\pi(t-s))| p(s) ds = G(t), \quad 0 \leq t \leq 1,$$

where  $G(t)$  is a smooth periodic function (provided  $g$  is smooth).

From [Gradshteyn and Ryzhik 1979, 1.441.2, p. 38] we have

$$(11.5.9) \quad -\frac{1}{2} \log(1 - \cos(2\pi t)) = \sum_{k=1}^{\infty} \frac{\cos(k2\pi t)}{k}$$

and so,

$$(11.5.10) \quad -2 \log(2R|\sin \pi t|) = -2 \log R + \sum_{k \neq 0} \frac{1}{|k|} e^{2\pi i k t}.$$

From this it follows that the effect of the operator  $A$  on  $p(s) = \sum c_j e^{2\pi i j s}$  is, essentially, to multiply the  $j^{\text{th}}$  Fourier coefficient  $c_j$ ,  $j \neq 0$ , by  $\frac{1}{|j|}$ . (We also smell a rat if  $R = 1$ !) Introducing the natural periodic Sobolev spaces  $H^m$  by the norms

$$(11.5.11) \quad \|p\|_{H^m} = \left( c_0^2 + \sum_{j \neq 0} |j|^{2m} |c_j|^2 \right)^{1/2}, \text{ for } -\infty < m < \infty,$$

we thus have

$$(11.5.12) \quad \|Ap\|_{H^{m+1}} \leq C \|p\|_{H^m},$$

i.e., the operator is “smoothing”.

Let us now further, for simplicity, assume that

$$(11.5.13) \quad R < 1.$$

Then clearly the operator  $A$  is coercive over  $H^{-1/2}$  in the sense that with a positive constant  $c$ , and the  $L_2$  inner product

$$(11.5.14) \quad (p, q) = \int_0^1 p(s) \overline{q(s)} ds,$$

we have

$$(11.5.15) \quad (Ap, p) \geq c \|p\|_{H^{-1/2}}^2.$$

It follows from Riesz' representation theorem that the equation  $Ap = G$  has a unique solution  $p$  in  $H^{-1/2}$ , for every  $G$  in  $H^{1/2}$ . Furthermore, it is easy to derive by Fourier analysis the regularity result that

$$(11.5.16) \quad \|p\|_{H^m} \simeq \|G\|_{H^{m+1}}.$$

(The particular case  $R = e^{-1/2}$  leads to an isometry in (11.5.16), cf. (11.5.10).)

Now for finite element solutions to  $Ap = G$ ! Let  $T_h$  be uniform subdivisions of length  $h$  of  $[0, 1]$ ,  $T_h = \{I_j\}_{j=1}^N$ , and set

$$(11.5.17) \quad S_h = \{\chi : \chi|_{I_j} \text{ is constant}\},$$

i.e.,  $S_h$  consists of discontinuous piecewise constants. The standard Galerkin finite element method is then to find  $p_h \in S_h$  by

$$(11.5.18) \quad (Ap_h, \chi) = (G, \chi), \text{ for } \chi \in S_h.$$

(Here the inner product is the  $L_2$ -inner product (11.5.14).) We assume that we can evaluate  $A$  exactly for  $q \in S_h$ , and that we can likewise evaluate exactly the inner product.

By the above,  $(Ap, p) \simeq \|p\|_{H^{-1/2}}^2$ , and so by standard techniques, and approximation theory,

$$(11.5.19) \quad \|p - p_h\|_{H^{-1/2}} \leq C \min_{\chi \in S_h} \|p - \chi\|_{H^{-1/2}} \leq Ch^{3/2},$$

if  $p \in H^1$  (which it does if  $G \in H^2$ ). An obvious duality argument then gives

$$(11.5.20) \quad \|p - p_h\|_{H^{-2}} \leq Ch^3.$$

Let us pause here to remark that negative norm error estimates are of interest for boundary integral equations for the following reason (among others): If we wish to evaluate  $u(x)$  at a point  $x$  inside  $B_R$ , we would naturally do it via (11.5.3), using  $\tilde{p}_h$ . Since  $\log|x-y|$  is smooth for  $x \in \overset{\circ}{B}_R$ ,  $y \in \Gamma_R$ , it is easy to see that the error in  $|u(x) - u_h(x)|$  would then be bounded by the error in any negative norm for  $p - p_h$ . Of course, constants deteriorate as  $x$  comes close to the boundary.

The so-called qualocation method (for “quadrature-modified collocation method”<sup>1)</sup>) was introduced to obtain even higher order negative norm estimates than in (11.5.20). (We continue to treat only the piecewise constant case.) Our presentation is based on [Sloan 1992a, Chapter 7] and [Sloan 1992b]; references to original work by Sloan and others such as [Sloan and Wendland 1989] and [Chandler and Sloan 1990] can be found in these expositions.

In the variant of the qualocation method that we shall give, it is still assumed that  $(Aq)(t)$  can be exactly evaluated for  $q \in S_h$ . Also, it will now be essential that the mesh is uniform (above, we could have considered quasi-uniform meshes, e.g.).

We now find  $p_h^Q \in S_h$  via

$$(11.5.21) \quad (Ap_h^Q, \chi)_h = (G, \chi)_h, \text{ for } \chi \in S_h,$$

where  $(p, q)_h$  is based on a composite quadrature rule; for  $I_j = (jh, (j+1)h)$ ,

$$(11.5.22) \quad \int_{I_j} f ds \cong \omega f(jh) + (1-\omega)f\left(\left(j + \frac{1}{2}\right)h\right).$$

(This form of the quadrature rule anticipates what is to come.) Since the functions in our space  $S_h$  are discontinuous at the nodes  $jh$ , we shall have to assign them a definite value there. We take that to be the mean from the left and right,

$$(11.5.23) \quad \chi(jh) = (\chi(jh - \varepsilon) + \chi(jh + \varepsilon))/2, \text{ for } \varepsilon < h, \chi \in S_h.$$

(We may note that  $A\chi \in H^1$ , for  $\chi \in S_h$ , and hence  $A\chi$  is continuous.)

It turns out to be possible to introduce a particular basis  $\psi_\mu$  for  $S_h$  which diagonalizes the matrix  $(A\psi_\mu, \psi_\nu)_h$  in (11.5.21). Thereafter, it turns out to be possible to derive a very precise expression for the error, in terms of the Fourier coefficients for  $p_h$  and  $p$ . Considering the leading term in that error expansion and choosing

$$(11.5.24) \quad \omega = \frac{3}{7}, \quad (1-\omega) = \frac{4}{7},$$

in the integration rule (11.5.22) to knock out that leading term, one then finds that

$$(11.5.25) \quad \|p - p_h^Q\|_{H^{-4}} \leq Ch^5,$$

for  $p \in H^4$ . The details are lengthy and will not be further commented on here.

The operator  $A$  is of convolution type, hence, translation invariant. So is our piecewise constant finite element space  $S_h$ , which is periodic so that we need not worry about boundaries. The analysis in [Tran 1993] now combines this with the  $K$ -operator, much in the spirit of Section 11.4. It also extends the analysis to the case when  $\Gamma$  is a general smooth curve in the plane and  $A$  a more general operator,

---

<sup>1)</sup>The method was introduced and, by and large, developed in Australia. It is sometimes referred to as the koalacation method.

not necessarily coming from the problem (11.5.1–2). (Both these extensions involve compact perturbations of our present type operators, not unlike going to variable coefficients in the second order partial differential equation case, cf. the remark in 11.6.iii below.)

Tran illustrates his analysis with a numerical example which I think is rather stunning, so, I'll reproduce it here.

The domain is the ellipse  $\frac{x_1^2}{4} + \frac{x_2^2}{9} \leq 1$  and the boundary function  $g$  is  $g(x_1, x_2) = \sin(x_1 - 0.1) \cosh(x_2 - 0.2)$ . The exact solution  $p$  is then known. He uses the qualocation method as above, and the particular  $K$ -operator is based on cubic (smoothest) splines, as Section 11.3 suggests ( $\ell = 4$ ). The function  $K$  is supported in  $[-4h, 4h]$ . The following is a sketch of it:




Figure 11.5.1

For the qualocation method used, superconvergence of order  $O(h^2)$  at midpoints is known. The following gives his numerical results. The numbers on the right in the second to fourth columns are the observed rates of convergence.

$N = 1/h$	$\ p - p_h^Q\ _{L_\infty(\Gamma)}$		$\max_i  p - p_h^Q (t_{i+1/2})$		$\ p - K_h * p_h^Q\ _{L_\infty(\Gamma)}$	
16	8.17		0.59		0.92	
32	4.22	0.95	0.24	1.28	0.426–1	4.43
64	2.08	1.02	0.61–1	2.00	0.957–3	5.48
128	1.05	0.99	0.154–1	1.99	0.197–4	5.60
256	0.52	1.00	0.385–2	2.00	0.435–6	5.50
512	0.26	1.00	0.962–3	2.00	0.107–7	5.34

Considering the pointwise global error, by trivial post-processing Tran, with a little help from friends, has managed to create a fifth order accurate approximation out of a first order accurate one. Not bad, or what?!

### 11.6. Remarks, including some other averaging methods.

i) The theory as given in Theorem 11.3.1 applies to the  $L_2$ -projection in locally translation invariant cases. Now, most favorably, we may have

$$(11.6.1) \quad \|u - u_h\|_{W_2^{-r}(\mathcal{D})} \leq Ch^{2r}$$

and thus pick up the same rate of convergence in the pointwise norm for approximations to any derivative of  $u$ .

ii) Likewise, Theorem 11.3.1 applies to time-continuous Galerkin approximations to parabolic problems, cf. e.g. [Bramble, Schatz, Thomée and Wahlbin 1977, Sections 6 and 7], [Thomée 1980a] and [Thomée 1984, Chapter 6].

iii) To put the result of [Tran 1993] in perspective, we offer the following references to previous work on integral equations. In [Richter 1978], “natural” (no averaging of any kind) superconvergence was considered for Fredholm integral equations of the second kind. [Sloan, Burn and Dattner 1975] introduced the idea of “iteration”, cf. [Nyström 1930], which we have seen in its two-point boundary value problem variant in Section 1.12.b. [Chandler 1980] gave an analysis for the  $K$ -operator as applied to the second kind Fredholm integral equation

$$(11.6.2) \quad (I - T)\varphi = f,$$

where  $T$  is a compact operator. Thus, for the Galerkin method, he treated the case of “the  $L_2$ -projection with compact perturbations”. In analogy, Tran treats the  $H^s$  projection, with compact perturbations, the case of  $s = -1/2$  being the one we have given some details about above.

iv) A minor variation on the  $K$ -operator in a special case in the plane occurs in [Oganesyan and Rukhovetz 1979, pp. 94 and 189]. It involves forming the average

$$(11.6.3) \quad \tilde{u}_h(x) = \frac{1}{4h^2} \int_{-h}^h \int_{-h}^h u_h(x + y) dy_1 dy_2.$$

For piecewise linears on a triangulation which is uniform in a neighborhood  $\Omega_0$  of  $\Omega_1$ , it is then not hard to show that

$$(11.6.4) \quad \|\nabla(u - \tilde{u}_h)\|_{L_\infty(\Omega_1)} \leq Ch^2 \ln 1/h,$$

provided  $u \in W_\infty^3(\Omega_0)$  and provided outside influences permit. Note that, from Chapter 8, we know that already applying simple centered difference quotients gives the same result.

v) The iteration technique in 1.12.a for finding the outflow derivative in a two-point boundary value problem generalizes, in a sense, to higher dimensions. Let

$$(11.6.5) \quad Lv = -\operatorname{div}(\alpha(x)\nabla v) - \sum_{i=1}^n \frac{\partial}{\partial x_i}(a_i(x)v) + a(x)v,$$

where  $\alpha(x)$  is a scalar function, and let  $u$  be the solution to the Dirichlet problem

$$(11.6.6) \quad \begin{aligned} Lu &= f \text{ in } \mathcal{D}, \\ u &= 0 \text{ on } \Gamma = \partial\mathcal{D}. \end{aligned}$$

For  $v$  smooth enough we then have by the divergence theorem,

$$(11.6.7) \quad \left\langle \alpha(x) \frac{\partial u}{\partial n}, v \right\rangle = A(u, v) - (f, v),$$

where  $\langle v, w \rangle = \int_{\Gamma} v w d\sigma$ ,  $A(v, w)$  is the natural bilinear form associated with  $L$ , and  $(v, w) = \int_{\mathcal{D}} v w dx$ .

Assume now for simplicity of description that  $\mathcal{D}$  is a polyhedral domain. On  $\mathcal{D}$  we have finite element spaces  $S_h(\mathcal{D})$ , which match  $\Gamma$  but which are *not* zero on  $\Gamma$ . Letting  $\overset{\circ}{S}_h(\mathcal{D})$  denote those functions in  $S_h(\mathcal{D})$  that *do* vanish on  $\Gamma$ , we find  $u_h \in \overset{\circ}{S}_h(\mathcal{D})$  by

$$(11.6.8) \quad A(u_h, \chi) = (f, \chi), \text{ for } \chi \in \overset{\circ}{S}_h(\mathcal{D}).$$

To form an approximation to the outflow derivative, or, rather, to  $p(x) = \alpha(x) \frac{\partial u}{\partial n}$ , let  $p_h \in S_h|_{\Gamma}$  be given by the analogue of (11.6.7),

$$(11.6.9) \quad \langle p_h, \chi \rangle = A(u_h, \tilde{\chi}) - (f, \tilde{\chi}), \text{ for } \chi \in S_h|_{\Gamma}$$

where  $\tilde{\chi}$  denotes any extension of  $\chi \in S_h|_{\Gamma}$  to  $S_h(\mathcal{D})$ .

By (11.6.8),  $p_h$  is independent of the particular extension  $\tilde{\chi}$  used. One may thus use the “simplest” one, cutting it down to be zero “as fast as possible”.

The extra work in computing  $p_h$  is commensurate to doing an  $L_2$ -projection over  $\Gamma$ .

In these lectures we shall not give an analysis of the error in  $p - p_h$ , but shall merely refer to [Douglas, Dupont and Wheeler 1974a], [Carey, Humprey and Wheeler 1981], [Carey 1982], [Carey, Chow and Seager 1985], [Chow and Lazarov 1989] and [Pehlivanov, Lazarov, Carey and Chow 1992] for some investigations.

A rather intriguing phenomenon is reported in some of the computational investigations above, see [Carey, Chow and Seager 1985]. To whit: Consider the polygonal situation in the plane.




Figure 11.6.1

Since  $u = 0$  on  $\Gamma$ , it follows that  $\nabla u(v_0) = 0$ , at the vertex  $v_0$ , if smooth. It appears that the accuracy in approximating the outflow derivative near  $v_0$  suffers unless one further imposes that the functions  $\chi \in S_h|_\Gamma$  in (11.6.9) also vanish at  $v_0$ .

vi) In a spirit somewhat similar to v), the Lagrange multiplier method (cf. [Babuška 1973]) incorporates a separate approximation to the outflow derivative in its basic formulation. We refer to [Bramble 1981] for some results about this approximation.

### 11.7. A superconvergent “global” averaging technique for function values.

This technique was introduced in [Louis 1979]. It applies to *any* mesh and *any* finite elements provided a suitable negative norm estimate for  $u - u_h$  is known. It involves, however, integration of  $u_h$  against a kernel in a unit size neighborhood of the point of interest, and, it also requires knowledge of an exact fundamental solution for the underlying partial differential equation.

As an example we consider the Dirichlet problem for Poisson's equation in the plane,

$$(11.7.1) \quad \begin{aligned} -\Delta u &= f \text{ in } \mathcal{D}, \\ u &= 0 \text{ on } \partial\mathcal{D}. \end{aligned}$$

Let  $x_0$  be a fixed point in  $\mathcal{D}$  and

$$(11.7.2) \quad B(x_0, R) = \{x : |x - x_0| \leq R\} \subseteq \mathcal{D},$$

where  $R$  is fixed, independent of  $h$  in particular. Let

$$(11.7.3) \quad \omega(x) = -\frac{1}{2\pi} \ln|x| + \psi(x)$$

where  $\psi(x) \in C^\infty$  is chosen so that

$$(11.7.4) \quad \omega(x) = \partial\omega(x)/\partial n = 0, \text{ for } |x| = R.$$

[Louis 1979] gives the example

$$(11.7.5) \quad \psi(x) = -\frac{1}{8\pi R^4}(|x|^4 - 4R^2|x|^2 + (3 - 4\ln R)R^4).$$

Note that it is not required that  $\omega(x) \in C^\infty$ , merely that  $\psi$  does.

By Green's second formula and (11.7.1), (11.7.4),

$$(11.7.6) \quad u(x_0) = \int_{B(x_0, R)} f(y)\omega(x_0 - y) + u(y)\Delta\psi(x_0 - y)dy.$$

With  $u_h$  any approximation to (11.7.1), we then define

$$(11.7.7) \quad \tilde{u}_h(x_0) = \int_{B(x_0, R)} f(y)\omega(x_0 - y) + u_h(y)\Delta\psi(x_0 - y)dy.$$

We assume that the integrals are exact (or, done to sufficiently high order of accuracy). We then have

$$(11.7.8) \quad (u - \tilde{u}_h)(x_0) = \int_{B(x_0, R)} (u - u_h)(y)\Delta\psi(x_0 - y)dy.$$

Since  $\psi \in \mathcal{C}^\infty$  and  $R$  is fixed, it is clear that the error in  $(u - \tilde{u}_h)(x_0)$  is governed by whatever negative norm accuracy  $u - u_h$  enjoys, cf. Section 6.2.

Finally, it is not hard to take into account the dependence on  $R$  of the error  $(u - \tilde{u}_h)(x_0)$ .


## Chapter 12. A computational investigation of superconvergence for first derivatives in the plane.

We shall review the paper [Babuška, Strouboulis, Upadhyay and Gangaraj 1993] and consider their results in light of our theories in Chapters 6 and 7.


The investigation reduces the problem of finding certain superconvergence points to that of finding the zeroes of certain polynomials. For each element pattern considered and each basic finite element, *only a finite number of polynomials are involved*. I.e., to find those superconvergence points, *there is no asymptotics w.r.t. h involved*. In this sense, the investigation may be called a “computational proof”.

### 12.1. Introduction.


Only so-called “natural” superconvergence points will be considered, i.e., no averaging, however simple, will be taken into account, cf. Remark 12.3.1 below. Only straight evaluation of the derivative of  $u_h$  at the point in question is allowed. Let us give details of the set-up for triangular elements. For each of four basic mesh-patterns considered later in the numerical examples we have a  $2h \times 2h$  “master cell” centered at  $x^0$  ( $= \bullet$ ).




Regular Pattern



Chevron Pattern



Union Jack Pattern



Criss Cross Pattern

Figure 12.1.1

(The shaded triangles will be used later, in Section 12.3.)

In a neighborhood  $\Omega_0$  of  $x^0$  the mesh is given by  $2h$ -periodic repetitions of one of these master cells, and so is translation invariant in each axis-direction by  $2h$  (or  $h$  in some directions and cases).

The finite element spaces considered will be continuous piecewise polynomials of total polynomial degree  $r - 1$  on each triangle. ( $C^1$  or higher finite elements will not be considered.)

Remark 12.1.1. What is important in the set-up of the master cells is that “certain things match up over opposite boundaries”. Expanding on this, in our  $C^0$ , total degree  $r - 1$ , triangular case we have a “standard” interpolation operator  $Int_h$  which interpolates the values of a function at the “principal lattice points”, illustrated here in the case of linears, quadratics, cubics and quartics.




Figure 12.1.2

(See [Nicolaides 1972 and 1973].) Note that the number of points on each edge are enough to determine completely the interpolant on that edge; hence  $C^0$  continuity is assured.

Furthermore, in all of our four pictorial examples, interpolation points on opposite sides in the master cell match up so that (taking  $x_0 = (0, 0)$ ),

$$Int_h(f(\cdot + 2h, \cdot))(-h, x_2) = Int_h(f(\cdot, \cdot))(h, x_2)$$

and correspondingly in the  $x_2$  direction. (In our technical work, this will enter in the proof of Lemma 12.2.1.)

We could thus also consider master cells of the form (still for  $C^0$  total degree polynomials, say)




Figure 12.1.3

but *not* of the form




Figure 12.1.4

□

Now assume that we have a basic finite element approximation  $u_h \in S_h(\Omega_0)$  to a function  $u$  which is sufficiently smooth on  $\Omega_0$ . Taking here only the case of the Poisson equation, we assume

$$(12.1.1) \quad D(u - u_h, \chi) = 0, \text{ for } \chi \in S_h^{\text{comp}}(\Omega_0),$$

where  $D(v, w) = \int \nabla v \cdot \nabla w$ . We shall further assume that

$$(12.1.2) \quad \|u - u_h\|_{L_\infty(\Omega_0)} \leq Ch^{r-L}.$$

Of course, this error is influenced from what happens outside of  $\Omega_0$ . Various pollution effects have been gotten rid of (at least beyond an order loss of  $h^L$ ). E.g., if the problem is over the basic domain  $\Omega_0 \subset\subset \mathcal{D}$  and  $\mathcal{D}$  has reentrant corners, the mesh may have been suitably refined at such corners, cf. Example 6.3.4.

We are now interested in superconvergence for either of the derivatives  $\frac{\partial}{\partial x_i}$ ,  $i = 1$  or 2. With

$$(12.1.3) \quad \Omega_1 = \{x : |x - x^0| \leq H\}, \quad \Omega_0 = \{x : |x - x^0| \leq 2H\}$$

and

$$(12.1.4) \quad H = h^\delta, \quad 0 < \delta < 1,$$

we shall show in Section 2 that

$$(12.1.5) \quad \frac{\partial}{\partial x_i}(u - u_h)(x) = \frac{\partial}{\partial x_i}\psi(x) + R_i(x), \quad i = 1, 2, \quad x \in \Omega_1,$$

where

$$(12.1.6) \quad \|R_i\|_{L_\infty(\Omega_1)} \leq C(h^{r-1+\delta} + h^{r-L-\delta}) \leq Ch^{r-1+\min(\delta, 1-L-\delta)}.$$

Provided  $L + \delta < 1$ , the term  $R_i(x)$  is thus “superconvergent” for first derivatives. Note that this is so even if the size  $H$  of the domain  $\Omega_0$  where the mesh is built up in a regular fashion from the master cell is very small,  $\delta \simeq 1$ ; then of course one has to be more stringent about pollution control,  $L \simeq 0$ . In other words, superconvergence is treated here as a local phenomenon.

As for the function  $\psi$ , it is based on computations involving *only the master cell*. In short, Taylor-expand  $u$  to polynomial degree  $r$  at  $x^0$  to get  $Q$ . Interpolate into  $S_h$  to form  $\rho = Q - \text{Int}_h Q$ . The effect of this is, as we shall prove in Section 12.2, among others to knock away the terms of order  $\leq r-1$  in  $Q$ , and to make  $\rho$  periodic of period  $2h$ . Hence, one needs to do it only over the master cell. Finally, over the master cell alone, find a  $2h$ -periodic finite element approximation to  $\rho$ . The function  $\psi$  will then be their difference, periodically extended. (See (12.2.25), (12.2.20,21) and (12.2.16) below for exact definition of  $\psi$ .)

From (12.1.5), (12.1.6) and the simple nature of  $\psi$ , cf. Remark 12.2.1, provided  $L + \delta < 1$ , we shall find that points  $x$  with a fixed relative position with respect to the master cell are superconvergent for  $\frac{\partial}{\partial x_i}$  on  $\Omega_1$  if and only if  $\frac{\partial\psi}{\partial x_i}(x) = 0$ . The computational experiments, to be reported in Section 3, then uses for  $Q$  any polynomial of degree  $r$  that may occur in a Taylor-expansion for  $u$ , that is, in general any monomial of degree  $p$ , and runs through these monomials and checks for  $\frac{\partial\psi}{\partial x_i} = 0$ . (Exactly how this check is done is not reported. It seems that interval arithmetic is not used.) Any such points found common to all monomials are then declared to be superconvergence points for that derivative. (In case one considers only, say, harmonic functions  $u$ , one need only consider harmonic polynomials  $Q$  of degree  $r$  in the computational experiments.)

## 12.2. Proof of (12.1.5), (12.1.6) and precise definition of the principal error term $\psi$ .

Recall that  $\Omega_1 \subset\subset \Omega_0$  are specific squares about the central point  $x^0$ , of side  $2H$ ,  $4H$ , respectively, with  $H = h^\delta$ . For simplicity we also assume that the  $2h$  periodic extensions of the master cell fit  $\Omega_1$  and  $\Omega_0$  exactly. We shall denote in general

$$(12.2.1) \quad c(x, h) = \{y : |y - x| \leq h\}.$$

Then  $c(x^0, h)$  is the master cell and, with  $x^{(i,j)} = (x_1^0 + 2ih, x_2^0 + 2jh)$ ,  $i, j$  integers,  $c(x^{(i,j)}, h)$  are the translated cells.

Let  $Q$  be the  $r^{\text{th}}$  order Taylor-expansion for  $u$  at  $x^0$ . Then

$$(12.2.2) \quad \|u - Q\|_{W_\infty^s(\Omega_0)} \leq CH^{r+1-s}, \text{ for } 0 \leq s \leq r+1.$$

As a tool for the present analysis (it will not enter into the final  $\psi$ ) we introduce the Neumann projection  $N_h Q$  of  $Q$  into  $S_h(\Omega_0)$ , i.e.,

$$(12.2.3) \quad D(Q - N_h Q, \chi) = 0, \text{ for } \chi \in S_h(\Omega_0),$$

$$(12.2.4) \quad \int_{\Omega_0} (Q - N_h Q) = 0.$$

We write

$$(12.2.5) \quad u - u_h = (Q - N_h Q) + [(u - Q) - (u_h - N_h Q)]$$

so that

$$(12.2.6) \quad \frac{\partial}{\partial x_i} (u - u_h)(x) = \frac{\partial}{\partial x_i} (Q - N_h Q)(x) + R_i^1(x), \text{ for } \chi \in \Omega_1,$$

where

$$(12.2.7) \quad \|R_i^1\|_{L_\infty(\Omega_1)} \leq \|(u - Q) - (u_h - N_h Q)\|_{W_\infty^1(\Omega_1)}.$$

From (12.1.1) and (12.2.3) we have

$$(12.2.8) \quad D((u - Q) - (u_h - N_h Q), \chi) = 0, \text{ for } \chi \in S_h^{\text{comp}}(\Omega_0)$$

so that by Theorem 5.5.1,

$$\begin{aligned} (12.2.9) \quad & \|R_i^1\|_{L_\infty(\Omega_1)} \\ & \leq C \min_{\chi \in S_h(\Omega_0)} \left( \|(u - Q) - \chi\|_{W_\infty^1(\Omega_0)} + H^{-1} \|(u - Q) - \chi\|_{L_\infty(\Omega_0)} \right) \\ & \quad + CH^{-2} \|(u - Q) - (u_h - N_h Q)\|_{L_2(\Omega_0)}. \end{aligned}$$

Here, using for  $\chi$  the interpolant of  $(u - Q)$  into  $S_h(\Omega_0)$ ,

$$(12.2.10) \quad \|(u - Q) - \chi\|_{W_\infty^1(\Omega_0)} \leq Ch^{r-1} \|u - Q\|_{W_\infty^r(\Omega_0)} \leq Ch^{r-1} H,$$

where we used (12.2.2) in the last step. Similarly,

$$(12.2.11) \quad H^{-1} \|(u - Q) - \chi\|_{L_\infty(\Omega_0)} \leq Ch^r \leq Ch^{r-1} H.$$

Thus,

$$(12.2.12) \quad \|R_i^1\|_{L_\infty(\Omega_1)} \leq Ch^{r-1} H + H^{-2} \|(u - Q) - (u_h - N_h Q)\|_{L_2(\Omega_0)}.$$

By our assumption (12.1.2),

$$(12.2.13) \quad \|u - u_h\|_{L_2(\Omega_0)} \leq CH \|u - u_h\|_{L_\infty(\Omega_0)} \leq CH h^{r-L}$$

and by standard  $L_2$  estimates (which are easily checked to be valid also on a small square),

$$(12.2.14) \quad \|Q - N_h Q\|_{L_2(\Omega_0)} \leq Ch^r \|Q\|_{W_2^r(\Omega_0)} \leq Ch^r H \|Q\|_{W_\infty^r(\Omega_0)} \leq CH h^{r-L}.$$

Hence (12.2.12) gives

$$(12.2.15) \quad \|R_i^1\|_{L_\infty(\Omega_1)} \leq C(h^{r-1} H + h^{r-L} H^{-1}).$$

We next prepare to attack  $\frac{\partial}{\partial x_i}(Q - N_h Q)(x)$  in (12.2.6).

Let  $Int_h$  denote the standard interpolation operator into  $S_h(\Omega_0)$  (cf. Remark 12.2.2 below), and let

$$(12.2.16) \quad \rho = Q - Int_h Q.$$

Let further  $H_{PER}^1(\Omega_0)$  denote the  $2h$ -periodic functions in  $W_2^1(\Omega_0)$ . A central observation is the following:

**Lemma 12.2.1.**  $\rho \in H_{PER}^1(\Omega_0)$ .

Proof: Certainly,  $\rho \in W_2^1(\Omega_0)$ . Since the situation is built up by translates of the master cell, the interpolant is translation invariant by  $2h$ , i.e.,

$$(12.2.17) \quad Int_h(f(\cdot + x^{(i,j)}))(x) = Int_h(f(\cdot))(x + x^{(i,j)}),$$

cf. Remark 12.1.1. Noting that for  $Q$  a polynomial of degree  $r$ ,

$$(12.2.18) \quad Q(x) = Q(x + x^{(i,j)}) - P_{ij}(x)$$

with  $P_{ij}(x)$  a polynomial of degree  $r - 1$ , which is reproduced by  $Int_h$ , we have

$$\begin{aligned} (12.2.19) \quad & Q(x) - Int_h(Q(\cdot))(x) \\ &= Q(x + x^{(i,j)}) - P_{ij}(x) - Int_h(Q(\cdot))(x) \\ &= Q(x + x^{(i,j)}) - Int_h((P_{ij} + Q)(\cdot))(x) \\ &= Q(x + x^{(i,j)}) - Int_h(Q(\cdot + x^{(i,j)}))(x) \\ &= Q(x + x^{(i,j)}) - Int_h(Q(\cdot))(x + x^{(i,j)}), \end{aligned}$$

which proves the lemma.  $\square$

Let now  $S_h^{PER}(c(x^0, h))$  ( $c(x^0, h)$  is the master cell, remember) denote the functions in  $S_h(c(x^0, h))$  which are  $2h$ -periodic, i.e., function values match up at boundaries. For any  $\varphi \in H_{PER}^1(\Omega_0)$  we define, *which is done on the master cell alone*,  $PP(\varphi) \in S_h^{PER}(c(x^0, h))$  ( $PP$  for periodic projection) by

$$(12.2.20) \quad D_{c(x^0, h)}(\varphi - PP(\varphi), \chi) = 0, \text{ for } \chi \in S_h^{PER}(c(x^0, h)),$$

$$(12.2.21) \quad \int_{c(x^0, h)} \varphi - PP(\varphi) = 0.$$

We also let  $PP(\varphi)$  denote its  $2h$ -periodic extension, which is then in  $H_{PER}^1(\Omega_0)$ . Another crucial result is the following.

**Lemma 12.2.2.** *For  $\varphi \in H_{PER}^1(\Omega_0)$ , we have*

$$(12.2.22) \quad D(\varphi - PP(\varphi), \chi) = 0, \text{ for all } \chi \in S_h^{comp}(\Omega_0).$$

Proof: For  $\gamma$  the relevant index set, we write in obvious notation,

$$(12.2.23) \quad D(PP(\varphi) - \varphi, \chi) = \sum_{(i,j) \in \gamma} D_{c(x^{(i,j)}, h)}(PP(\varphi) - \varphi, \chi).$$

By the periodicity of  $PP(\varphi)$  and  $\varphi$  then,

$$(12.2.24) \quad \begin{aligned} & \sum_{(i,j) \in \gamma} D_{c(x^{(i,j)}, h)}(PP(\varphi) - \varphi, \chi) \\ &= D_{c(x^0, h)}\left(PP(\varphi) - \varphi, \sum_{(i,j) \in \gamma} \chi(x + x^{(i,j)})\right). \end{aligned}$$

Since  $\sum_{(i,j) \in \gamma} \chi(x + x^{(i,j)}) \in S_h^{PER}(c(x^0, h))$ , we get (12.2.22) from (12.2.20).  $\square$

After these preparations, we attack  $Q - N_h Q$  in (12.2.6). We set

$$(12.2.25) \quad \psi = \rho - PP(\rho), \quad \rho \text{ given in (12.2.16),}$$

and write

$$(12.2.26) \quad Q - N_h Q = \psi + [Q - N_h Q - \psi].$$

Here, by (12.2.16),

$$(12.2.27) \quad Q - \psi = Q - (\rho - PP(\rho)) = Int_h Q + PP(\rho),$$

which shows that

$$(12.2.28) \quad Q - N_h Q - \psi = Int_h Q + PP(\rho) - N_h Q \in S_h(\Omega_0).$$

Further, from Lemma 12.2.2 and (12.2.3),

$$(12.2.29) \quad D(Q - N_h Q - \psi, \chi) = 0, \quad \text{for } \chi \in S_h^{comp}(\Omega_0).$$

From (12.2.28) and Theorem 5.5.1 then,

$$(12.2.30) \quad \begin{aligned} \|Q - N_h Q - \psi\|_{W_\infty^1(\Omega_1)} &\leq CH^{-2}\|Q - N_h Q - \psi\|_{L_2(\Omega_0)} \\ &\leq CH^{-2}(\|Q - N_h Q\|_{L_2(\Omega_0)} + \|\psi\|_{L_2(\Omega_0)}). \end{aligned}$$

As before, cf. (12.2.14),

$$(12.2.31) \quad H^{-2}\|Q - N_h Q\|_{L_2(\Omega_0)} \leq Ch^{r-L}H^{-1}.$$

Further, by a standard duality argument (albeit on a small region, periodic functions), since  $\psi = \rho - PP(\rho)$ , and  $\rho = Q - Int_h Q$ ,

$$(12.2.32) \quad \begin{aligned} H^{-2}\|\psi\|_{L_2(\Omega_0)} &\leq CH^{-2}h\|\psi\|_{W_2^1(\Omega_0)} \\ &\leq CH^{-2}h\|\rho\|_{W_2^1(\Omega_0)} \\ &\leq CH^{-2}hh^{r-1}\|Q\|_{W_2^r(\Omega_0)} \\ &\leq Ch^rH^{-1}\|Q\|_{W_\infty^r(\Omega_0)} \\ &\leq Ch^{r-L}H^{-1}. \end{aligned}$$

All in all from the above,

$$(12.2.33) \quad \|Q - N_h Q - \psi\|_{W_\infty^1(\Omega_1)} \leq Ch^{r-L}H^{-1}.$$

It follows from this and (12.2.6), (12.2.15) that

$$(12.2.34) \quad \frac{\partial}{\partial x_i}(u - u_h)(x) = \frac{\partial \psi}{\partial x_i}(x) + R_i(x), \quad \text{for } x \in \Omega,$$

where

$$(12.2.35) \quad \|R_i\|_{L_\infty(\Omega_1)} \leq C(h^{r-1}H + h^{r-L}H^{-1}) = C(h^{r-1+\delta} + h^{r-L-\delta}),$$

which is the desired result, already stated as (12.1.5), (12.1.6).

Remark 12.2.1. By scaling, one finds that  $\frac{\partial\psi}{\partial x_i}(x) = h^{r-1}\frac{\partial\widehat{\psi}}{\partial\widehat{x}}(\widehat{x})$  where  $\widehat{\cdot}$  denotes quantities on the master cell scaled to unit size. Then, a point  $x$  of fixed relative position in the master cell is superconvergent iff  $\frac{\partial\psi}{\partial x_i}(x) = 0$ .  $\square$

Remark 12.2.2. In our presentation above, we have, without commenting much on it, taken  $Int_h$  to be the “standard” interpolation operator, well known in the cases at hand, see Remark 12.1.1. Actually, the proof works the same with other interpolation operators, provided they respect Remark 12.1.1 on opposite boundaries. To illustrate, consider the analogous situation in one dimension and  $C^0$  quadratics: The “standard” would be to interpolate at mesh and midpoints, but one may equally well interpolate at meshpoints and a quarter point (or, whatever). In this connection, note that the application of the periodic projection  $PP$  in forming  $\psi$  removes the differences between interpolation operators so that the superconvergence points found do not depend on this ambiguity in defining the interpolation operator.

As it happens, some of the superconvergence points for derivatives found in the computational experiments do coincide with some of those for the “standard” interpolant. The theory here obviously (from the above comments in this remark) does not say so. The reader may wish to revisit Section 7.4.i in this regard.  $\square$


Remark 12.2.3. Although presented here only for the case of the Dirichlet form, the arguments in this section carry over without much essential change to the case of any constant coefficient operator. The case of smooth variable coefficient operators is somewhat messier: One may freeze the coefficients at  $x^0$  and then one has to consider, at quite a few places, what happens between the original form and the frozen one. (The case of  $-div(a(x_1, x_2)\nabla u) + \dots$  can also be handled analogously to Section 6.4, extended to a general size  $H$ , of course.)  $\square$

### 12.3. Results of computational studies, with comments.


We shall give the computational results of [Babuška, Strouboulis, Upadhyay and Gangaray 1993] and comment in some detail on their findings in light of our theory in these notes, mainly in light of our symmetry theory of Chapter 7 and also the tensor product theory of Chapter 6.

Considering the Poisson equation, running all four triangular pattern cases as depicted in Section 12.1, all cases  $2 \leq r \leq 8$ , running through all possible monomials of degree  $r$  for  $Q$ , and declaring a point superconvergent for  $\frac{\partial}{\partial x_1}$  for  $x$  lying in the shaded triangles in the figures of master cells pictured in Section 12.1 if  $\frac{\partial\psi}{\partial x_1}(x) = 0$  for *all* such monomials, they found:


For the regular pattern:



$r = 2$  : midpoint of the side parallel to  $x_1$ -axis




$r = 3$  : Gauss points




$r = 4$  : midpoint, as for  $r = 2$

$r = 5$  : no superconvergent points for  $\frac{\partial u}{\partial x_1}$



$r = 7$  : no superconvergence points for  $\frac{\partial u}{\partial x_1}$



For  $r$  even, these midpoints results are known by our symmetry arguments, since the average from both sides in the case of  $\frac{\partial}{\partial x_1}$  now coincides with  $\frac{\partial u_h}{\partial x_1}$ . Note that this is *not* so at the symmetric mesh points; there one must use averaging to obtain superconvergence also in  $\partial/\partial x_1$ . For  $r = 3$ , the result is in Remark 7.4.ii.

For the *Chevron pattern*, the midpoints of sides for  $r$  even stay as superconvergent points for  $\frac{\partial}{\partial x_1}$ . (The Gauss points for  $r = 3$  are lost.) Since the Chevron-pattern has the midpoints of  $x_1$ -parallel edges as symmetry points, our theory of symmetry explains these findings as well. (Since the mesh points are not symmetry points, the result for  $r = 3$  cannot be deduced as in Remark 7.4.ii. Indeed, according to these computations, it is lost.)

As for the *Union Jack mesh*, no superconvergence points for  $\frac{\partial}{\partial x_1}$  were found. Since midpoints are no longer symmetry points, we know that at the only symmetry points present, the meshpoints, we have to average. So, the results shouldn't show in the present numerical experiment, and they didn't.

As for the *Criss-Cross grid*, for the shaded triangle labelled 1 in the figure, the superconvergence was at the midpoints on the side parallel to  $x_1$  for  $\frac{\partial}{\partial x_1}$ , for  $r$  even. For  $r = 3$ , we have the same Gauss points as before, but for  $r$  odd,  $r \geq 5$ , no superconvergence points. Again, our symmetry theory has predicted them all (and predicts more if averaging is allowed).

For the triangle labelled 2 in the Criss-Cross grid, for  $\frac{\partial}{\partial x_1}$  our symmetry theory wouldn't predict any superconvergence points without averaging and, none were found in the numerical experiments.

The investigation in [Babuška, Strouboulis, Upadhyay and Gangaraj 1993] also considers rectangular elements. They treat  $C^0$ -elements of type:

(12.3.1) Tensor product elements; i.e., on each rectangle,

$$\chi(x_1, x_2) = \sum_{\substack{i,j \\ 0 \leq i,j \leq r-1}} c_{ij} x_1^i x_2^j.$$

(12.3.2) Serendipity (trunc) elements,

$$\chi(x_1, x_2) = \sum_{\substack{i,j \\ 0 \leq i+j \leq r-1}} c_{ij} x_1^i x_2^j + c_{r-1,1} x_1^{r-1} x_2 + c_{1,r-1} x_1 x_2^{r-1}.$$

(12.3.3) Intermediate degree  $r - 1$  (incomplete degree  $r$ )

$$\chi(x_1, x_2) = \sum_{\substack{i,j \\ 0 \leq i+j \leq r-1}} c_{ij} x_1^i x_2^j + \sum_{k=0}^{r-2} c_{r-1-k,k+1} x_1^{r-1-k} x_2^{k+1}.$$

(These elements are missing  $x_1^r$  and  $x_2^r$  from containing all polynomials of total degree  $r$ .)

For the *tensor-product elements*, they found the  $x_1$  Gauss lines as superconvergent lines. No surprise, exactly what our one-dimensional theory combined with the tensor product theory of Section 6.2 (and 6.4, for generalization) shows. (The theory is given in Chapter 6 only for  $H = 1$ . The generalization presently needed is not hard.) Note also that the meshes do not have to translation invariant for the theory in Chapter 6.

The very same superconvergence points as for tensor-products occurred for the *intermediate (incomplete) family* in (12.3.3), they reported (without much detail given). For  $r = 3$ , i.e., incomplete cubics, cf. Section 7.4.iii; otherwise our symmetry theory in Chapter 7 in these lectures is silent, except for the center of the square and midpoints of the  $x_1$ -parallel sides for  $r$  even, in the absence of averaging.

Finally, for the case of the *Serendipity family* in (12.3.2), for  $r = 2$  and 3, the superconvergence points are as for tensor products. (For  $r = 2$ , all three elements, tensor products, intermediate and Serendipity, are the same; for  $r = 3$ , Serendipity and intermediate are the same. So, this is expected, cf. Chapter 6 and Section 7.4.)

For the *Serendipity family with  $r = 4$* , for  $\frac{\partial}{\partial x_1}$ , a line of superconvergence and

four interior points were found:




Figure 12.3.1

The four points are at  $(\pm 0.7746 \dots, \pm 0.5774 \dots)$ . Note that, by elementary symmetry arguments, they *cannot* also be superconvergent for  $\frac{\partial}{\partial x_2}$  and so, being interior points, our symmetry theory will not predict them.

For the *Serendipity family* with  $r \geq 5$ , no superconvergence points for  $\frac{\partial}{\partial x_1}$  were reported for  $r$  odd. For  $r = 6$  and 8, the center of the square and the midpoints of the  $x_1$ -parallel sides were found, exactly as our symmetry theory says (in the absence of averaging).

In conclusion, our theories in these lectures have explained all superconvergence results found in [Babuška, Strouboulis, Upadhyay and Gangaraj 1993] for general solutions of Poisson's equation *except*

- a) Most of those found in the intermediate (incomplete) families on squares, and
- b) The results found in the Serendipity family on a square for  $r = 4$ , except for the center point and the midpoints of the  $x_1$ -parallel sides.

The paper also reported on the case when  $u$  in itself is a harmonic function. In that case, only the harmonic polynomials  $Q$  need be tested. Now plenty of superconvergence points for derivatives were found, for any  $r$  (not merely even  $r$ ), and the number of such points increasing with  $r$ . Of course, our theory in these lectures has not even thought about this very interesting case! Similarly, [Babuška, Strouboulis, Upadhyay and Ganagaraj 1994] investigates superconvergence in homogeneous solutions of the equations of elasticity.

Remark 12.3.1. As remarked in the beginning of Section 12.1, we have not taken into account averaging from two adjoining elements at a point on their common boundary, as in (7.2.12). It is trivial to do so in (12.1.5), and corresponding computational determination of all "averaged" superconvergence points could be performed. □

## References.

(For the guiding principles behind this list of references, see the Preface.)

- [Adeboye 1982] K. R. Adeboye, *Superconvergence results for Galerkin method for parabolic initial value problems via Laplace transforms*, Bull. Math. Soc. Sci. Math. R. S. Roumaine (N.S.) **26(74)** (1982), 115–127, {MR 84h:65095}.
- [Adjerid and Flaherty 1986] S. Adjerid and J. E. Flaherty, *A moving-mesh finite element method with local refinement for parabolic partial differential equations*, Comput. Methods Appl. Mech. Engrg. **55** (1986), 3–26, {MR 87h:65187}.
- [Adjerid and Flaherty 1988] S. Adjerid and J. E. Flaherty, *A local refinement finite-element method for two-dimensional parabolic systems*, SIAM J. Sci. Statist. Comput. **9** (1988), 792–811, {MR 89g:65136}.
- [Ainsworth and Craig 1992] M. Ainsworth and A. Craig, *A posteriori error estimates in the finite element method*, Numer. Math. **60** (1992), 429–463, {MR 92j:65163}.
- [Ainsworth, Zhu, Craig and Zienkiewicz 1989] M. Ainsworth, J. Z. Zhu, A. W. Craig and O. C. Zienkiewicz, *An analysis of the Zienkiewicz–Zhu a-posteriori error estimator in the finite element method*, Internat. J. Numer. Methods Engrg. **28** (1989), 2161–2174, {MR 90j:73053}.
- [Andreasyan 1987a] G. D. Andreasyan, *Superconvergence of averaged gradients on an irregular mesh in the finite element method*; (in Russian), Vestnik Moskov. Univ. Ser. XV Vychisl. Mat. Kibernet. (1987), 11–16, {MR 88j:65246}.
- [Andreasyan 1987b] G. D. Andreasyan, *Superconvergence of averaged gradients in the finite element method on rectangles*; (in Russian), Applied Mathematics 5, Erevan Univ. 130 (1987), 5–10, {MR 90e:65149}.
- [Andreasyan 1987c] G. D. Andreasyan, *Superconvergence of the finite element method for higher order one-dimensional boundary value problems*; (in Russian), Akad. Nauk Armyan. SSR Dokl. **84** (1987), 3–8, {MR 88e:65131}.
- [Andreev 1984a] A. B. Andreev, *Superconvergence of the gradient for linear triangle elements for elliptic and parabolic equations*, C. R. Acad. Bulgare Sci. **37** (1984), 293–296, {MR 85h:65230}.
- [Andreev 1984b] A. B. Andreev, *Error estimate of type superconvergence for quadratic triangular elements*, C. R. Acad. Bulgare Sci. **37** (1984), 1179–1182, {MR 86f:65186}.
- [Andreev 1985] A. B. Andreev, *Superconvergence du gradient des solutions approchées pour les équations paraboliques*, Serdica **11** (1985), 359–368, {MR 87j:65142}.
- [Andreev 1991] A. B. Andreev, *Superconvergence of the gradient of finite element eigenfunctions*, C. R. Acad. Bulgare Sci. **43** (1991), 9–11, {MR 92d:65185}.
- [Andreev and Andreasyan 1988] V. B. Andreev and G. D. Andreasyan, *Superconvergence of derivatives and their averagings in the finite element method for ordinary  $2m^{\text{th}}$ -order differential equations*; (in Russian), Computation Processes and Systems 6, Nauka, Moscow, 1988, pp. 31–39, {MR 90f:65147}.

- [Andreev and Lazarov 1988] A. B. Andreev and R. D. Lazarov, *Superconvergence of the gradient for quadratic triangular finite elements*, Numer. Methods Partial Differential Equations **4** (1988), 15–32, {MR 90m:65190}.
- [Arnold and Douglas 1979] D. N. Arnold and J. Douglas, Jr., *Superconvergence of the Galerkin approximation of a quasilinear parabolic equation in a single space variable*, Calcolo **16** (1979), 345–369, {MR 83b:65118}.
- [Arnold, Douglas and Thomée 1981] D. N. Arnold, J. Douglas, Jr. and V. Thomée, *Superconvergence of a finite element approximation to the solution of a Sobolev equation in a single space variable*, Math. Comp. **36** (1981), 55–63, {MR 82f:65108}.
- [Arnold and Winther 1982] D. N. Arnold and R. Winther, *A superconvergent finite element method for the Korteweg-de Vries equation*, Math. Comp. **38** (1982), 23–36, {MR 82m:65087}.
- [Atkinson 1976] K. E. Atkinson, *A Survey of Numerical Methods for the Solution of Fredholm Integral Equations of the Second Kind*, SIAM, 1976, {MR 58#3577}.
- [Aziz and Monk 1989] A. K. Aziz and P. Monk, *Continuous finite elements in space and time for the heat equation*, Math. Comp. **52** (1989), 255–274, {MR 90d:65189}.
- [Babuška 1970] I. Babuška, *Finite element methods for domains with corners*, Computing **6** (1970), 264–273, {MR 45#2934}.
- [Babuška 1973] I. Babuška, *The finite element method with Lagrangian multipliers*, Numer. Math. **20** (1973), 179–192, {MR 50#11806}.
- [Babuška and Aziz 1972] I. Babuška and A. K. Aziz, *Survey lectures on the mathematics foundations of the finite element method*, The Mathematical Foundations of the Finite Element Method (A. K. Aziz, ed.), Academic Press, 1972, pp. 1–359, {MR 54#9111}.
- [Babuška and Miller 1984] I. Babuška and A. Miller, *The post-processing in the finite element method, I-II*, Internat. J. Numer. Methods Engrg. **20** (1984), 1085–1109, 1111–1129.
- [Babuška and Osborn 1980] I. Babuška and J. Osborn, *Analysis of finite element methods for second order boundary value problems using mesh dependent norms*, Numer. Math. **34** (1980), 41–62, {MR 81g:65143}.
- [Babuška, Strouboulis, Gangaraj and Upadhyay 1994] I. Babuška, T. Strouboulis, S. K. Gangaraj and S. K. Upadhyay,  *$\eta\%$ -superconvergence in the interior of locally refined meshes of quadrilaterals: superconvergence of the gradient in finite element solutions of Laplace's and Poisson's equations* (to appear).
- [Babuška, Strouboulis and Upadhyay 1993a] I. Babuška, T. Strouboulis, C. S. Upadhyay,  *$\eta\%$ -superconvergence of finite element approximations in the interior of general meshes of triangles* (to appear).
- [Babuška, Strouboulis and Upadhyay 1993b] I. Babuška, T. Strouboulis and C. S. Upadhyay, *A model study of the quality of a-posteriori error estimates for linear elliptic problems. Part Ia: Error estimation in the interior of patchwise uniform grids of triangles* (to appear).

- [Babuška, Strouboulis, Upadhyay and Gangaraj 1993] I. Babuška, T. Strouboulis, C. S. Upadhyay and S. K. Gangaraj, *The study of superconvergence by a computer-based approach: Superconvergence of the gradient in finite element solutions of Laplace's and Poisson's equations* (to appear).
- [Babuška, Strouboulis, Upadhyay and Gangaraj 1994] I. Babuška, T. Strouboulis, C. S. Upadhyay and S. K. Gangaraj, *The study of superconvergence by a computer based approach: Superconvergence of the gradient of the displacement, the strain and stresses in finite element solutions for plane elasticity* (to appear).
- [Babuška, Strouboulis, Upadhyay, Gangaraj and Copps 1994] I. Babuška, T. Strouboulis, C. S. Upadhyay, S. K. Gangaraj and K. Copps, *Validation of a-posteriori error estimators by numerical approach*, Internat. J. Numer. Methods Engrg. **37** (1994), 1073–1124.
- [Bakker 1980] M. Bakker, *On the numerical solution of parabolic equations in a single space variable by the continuous time Galerkin method*, SIAM J. Numer. Anal. **17** (1980), 162–177, {MR 81m:65167}.
- [Bakker 1982] M. Bakker, *A note on  $C^0$  Galerkin methods for two-point boundary value problems*, Numer. Math. **38** (1982), 447–453, {MR 83e:65140}.
- [Bakker 1984] M. Bakker, *One-dimensional Galerkin methods and superconvergence at interior nodal points*, SIAM J. Numer. Anal. **21** (1984), 101–110, {MR 85f:65080}.
- [Barlow 1976] J. Barlow, *Optimal stress location in finite element models*, Internat. J. Numer. Methods. Engrg. **10** (1976), 243–251.
- [Barlow 1989] J. Barlow, *More on optimal stress points — reduced integration, element distortions and error estimation*, Internat. J. Numer. Methods Engrg. **28** (1989), 1487–1504, {MR 90h:73022a} Correction: {MR 90h:73022b}.
- [Barrett and Liu 1993] J. W. Barrett and W. B. Liu, *Finite element approximation of the  $p$ -Laplacian*, Math. Comp. **61** (1993), 523–537.
- [Barrett, Moore and Morton 1988] J. W. Barrett, G. Moore and K. W. Morton, *Optimal recovery in the finite-element method. I. Recovery from weighted  $L_2$  fits*, IMA J. Numer. Anal. **8** (1988), 149–184, {MR 90b:65167}.
- [Bennett and Sharpley 1988] C. Bennett and R. Sharpley, *Interpolation of Operators*, Academic Press, 1988, {MR 89e:46001}.
- [Birkhoff and Rota 1969] G. Birkhoff and G.-C. Rota, *Ordinary Differential Equations*, 2nd edition, Xerox College Publishing, 1969, {MR 38#4737}.
- [Blum, Lin and Rannacher 1986] H. Blum, Q. Lin and R. Rannacher, *Asymptotic error expansion and Richardson extrapolation for linear finite elements*, Numer. Math. **49** (1986), 11–38, {MR 87m:65172}.
- [Blum and Rannacher 1988] H. Blum and R. Rannacher, *Extrapolation techniques for reducing the pollution effect of reentrant corners in the finite element method*, Numer. Math. **52** (1988), 539–564, {MR 89e:65117}.

- [Blum and Rannacher 1990] H. Blum and R. Rannacher, *Finite-element eigenvalue computation on domains with reentrant corners using Richardson extrapolation*, J. Comput. Math. **8** (1990), 321–332, {MR 92i:65163}.
- [Bourlard 1988] M. Bourlard, *Superconvergence pour les méthodes de Galerkin raffinées résolvant les problèmes de Neumann et Dirichlet dans un polygone*, C. R. Acad. Sci. Paris Ser. I Math. **306** (1988), 211–216, {MR 88m:65171}.
- [Bramble 1970] J. H. Bramble, *On the convergence of difference approximations for second order uniformly elliptic operators*, in Numerical Solution of Field Problems in Continuum Physics, SIAM–AMS Proceedings, vol. 2, 1970, pp. 201–209, {MR 41#4828}.
- [Bramble 1981] J. H. Bramble, *The Lagrange multiplier method for Dirichlet's problem*, Math. Comp. **3** (1981), 1–11, {MR 83h:65119}.
- [Bramble and Hilbert 1970] J. H. Bramble and S. R. Hilbert, *Estimation of linear functionals on Sobolev spaces with application to Fourier transforms and spline interpolation*, SIAM J. Numer. Anal. **7** (1970), 113–124, {MR 41#7819}.
- [Bramble and Hilbert 1971] J. H. Bramble and S. R. Hilbert, *Bounds for a class of linear functionals with application to Hermite interpolation*, Numer. Math. **16** (1971), 362–369, {MR 44#7704}.
- [Bramble, Nitsche and Schatz 1975] J. H. Bramble, J. A. Nitsche and A. H. Schatz, *Maximum-norm interior estimates for Ritz-Galerkin methods*, Math. Comp. **29** (1975), 677–688, {MR 53#1975}.
- [Bramble and Schatz 1974] J. H. Bramble and A. H. Schatz, *Higher order local accuracy by averaging in the finite element method*, Mathematical Aspects of Finite Elements in Partial Differential Equations (C. deBoor, ed.), Academic Press, 1974, pp. 1–14, {MR 58#31903}.
- [Bramble and Schatz 1976] J. H. Bramble and A. H. Schatz, *Estimates for spline projections*, RAIRO Modél. Math. Anal. Numér. **10** (1976), 5–37, {MR 55#9563}.
- [Bramble and Schatz 1977] J. H. Bramble and A. H. Schatz, *Higher order local accuracy by averaging in the finite element method*, Math. Comp. **31** (1977), 94–111, {MR 55#4739}.
- [Bramble, Schatz, Thomée and Wahlbin 1977] J. H. Bramble, A. H. Schatz, V. Thomée and L. B. Wahlbin, *Some convergence estimates for semidiscrete Galerkin type approximations for parabolic equations*, SIAM J. Numer. Anal. **14** (1977), 218–240, {MR 56#7231}.
- [Bramble and Scott 1978] J. H. Bramble and R. Scott, *Simultaneous approximation in scales of Banach spaces*, Math. Comp. **32** (1978), 947–954, {MR 80a:65222}.
- [Bramble and Thomée 1974] J. H. Bramble and V. Thomée, *Interior maximum norm estimates for some simple finite element methods*, RAIRO Modél. Math. Anal. Numér. **8** (1974), 5–18, {MR 50#11808}.
- [Bramble and Xu 1989] J. H. Bramble and J. Xu, *A local post-processing technique for improving the accuracy in mixed finite element approximations*, SIAM J. Numer. Anal. **26** (1989), 1267–1275, {MR 90m:65193}.

- [Brenner, Thomée and Wahlbin 1975] P. Brenner, V. Thomée and L. B. Wahlbin, *Besov Spaces and Applications to Difference Methods for Initial Value Problems*, Lecture Notes in Mathematics 434, Springer, 1975, {MR 57#1106}.
- [Brezzi, Douglas and Fortin 1987] F. Brezzi, J. Douglas Jr. and M. Fortin, *Efficient rectangular mixed finite elements in two and three space variables*, RAIRO Model. Math. Anal. Numer. **21** (1987), 581–604, {MR 88j:65249}.
- [Cao and Zhu 1990] L. Q. Cao and Q.-D. Zhu, *A new kind of finite element and its superconvergence estimate*; (in Chinese), Natur. Sci. J. Xiangtan Univ. **12** (1990), 1–8, {MR 92j:65166}.
- [Carey 1982] G. F. Carey, *Derivative calculation from finite element solutions*, Comput. Methods Appl. Mech. Engrg. **35** (1982), 1–14, {MR 83m:73067}.
- [Carey, Chow and Seager 1985] G. F. Carey, S. S. Chow and M. K. Seager, *Approximate boundary-flux calculations*, Comput. Methods Appl. Mech. Engrg. **50** (1985), 107–120, {MR 87h:65191}.
- [Carey, Humphrey and Wheeler 1981] G. F. Carey, D. Humphrey and M. F. Wheeler, *Galerkin and collocation-Galerkin methods with superconvergence and optimal fluxes*, Internat. J. Numer. Methods Engrg. **17** (1981), 939–950, {MR 82g:80008}.
- [Carey and Seager 1985] G. F. Carey and M. Seager, *Projection and iteration in adaptive finite element refinement*, Internat. J. Numer. Methods Engrg. **21** (1985), 1681–1695, {MR 87b:65118}.
- [Cayco, Schatz and Wahlbin 1994] M.-E. Cayco, A. H. Schatz and L. B. Wahlbin (to appear).
- [Chandler 1980] G. A. Chandler, *Superconvergence for second kind integral equations*, The Application and Numerical Solution of Integral Equations, Sijthoff and Nordhoff, 1980, pp. 103–117, {MR 81h:45027}.
- [Chandler 1984] G. A. Chandler, *Mesh grading for boundary integral equations*, Computational Techniques and Applications (CTAC–83), North-Holland, 1984, pp. 289–296, {MR 86j:65149}.
- [Chandler 1984] G. A. Chandler, *Galerkin's method for boundary integral equations on polygonal domains*, J. Austral. Math. Soc. Ser. B **26** (1984), 1–13, {MR 86a:65116}.
- [Chandler and Sloan 1990] G. A. Chandler and I. H. Sloan, *Spline qualocation methods for boundary integral equations*, Numer. Math. **58** (1990), 537–567, {MR 91m:65323}; *Erratum*, Numer. Math. **62** (1992), 295.
- [Chatelin and Lebbar 1981] F. Chatelin and R. Lebbar, *The iterated projection solutions for the Fredholm integral equation of second kind*, J. Austral. Math. Soc. Ser. B **22** (1981), 439–451, {MR 82h:65096}.
- [Chatelin and Lebbar 1984] F. Chatelin and R. Lebbar, *Superconvergence results for the iterated projection method applied to a Fredholm integral equation of the second kind and the corresponding eigenvalue problem*, J. Integral Equations **6** (1984), 71–91, {MR 85i:65167}.

- [Chen 1979] C.-M. Chen, *Superconvergent points of Galerkin's method for two-point boundary value problems*; (in Chinese), Numer. Math. J. Chinese Univ. **1** (1979), 73–79, {MR 82e:65085}.
- [Chen 1980] C.-M. Chen, *Optimal points of the stresses for triangular linear element*; (in Chinese), Numer. Math. J. Chinese Univ. **2** (1980), 12–20, {MR 83d:65279}.
- [Chen 1981] C.-M. Chen, *Superconvergence of finite element solutions and their derivatives*; (in Chinese), Numer. Math. J. Chinese Univ. **3** (1981), 118–125, {MR 82m:65100}.
- [Chen 1982a] C.-M. Chen, *The Finite Element Method and Its Analysis in Improving Accuracy*; (in Chinese), Hunan Science Press, 1982.
- [Chen 1982b] C.-M. Chen, *Superconvergence of finite element for nonlinear problems*; (in Chinese), Numer. Math. J. Chinese Univ. **4** (1982), 222–228, {MR 83m:65083}.
- [Chen 1983a] C.-M. Chen, *Superconvergence of finite element approximations to nonlinear elliptic problems*, Proceedings of the China–France Symposium on Finite Element Methods (Beijing 1982), Science Press, Beijing, and Gordon and Breach 1983, 622–640, {MR 85h:65235}.
- [Chen 1983b] C.-M. Chen, *An estimate for elliptic boundary value problems and its applications to finite element methods*; (in Chinese), Numer. Math. J. Chinese Univ. **5** (1983), 215–223, {MR 86b:65128}.
- [Chen 1985a] C.-M. Chen,  *$W^{1,\infty}$  interior estimates for finite element methods on regular mesh*, J. Comput. Math. **3** (1985), 1–7, {MR 87b:65194}.
- [Chen 1985b] C.-M. Chen, *Superconvergence of Galerkin solutions for singular nonlinear two-point boundary value problems*; (in Chinese); Math. Numer. Sinica **7** (1985), 113–123, {MR 87b:65119}.
- [Chen 1985c] C.-M. Chen, *Superconvergence of finite element methods*; (in Chinese), Adv. in Math. **14** (1985), 39–51, {MR 87g:65134}.
- [Chen 1986] C.-M. Chen, *Superconvergence and extrapolation of the finite element approximations to quasilinear elliptic problems*, Dongbei Shuxue **2** (1986), 228–236, {MR 88g:65104}.
- [Chen 1988] C.-M. Chen, *High accuracy analysis of finite element methods — superconvergence and postprocessing*, Natur. Sci. J. Xiangtan Univ. **10** (1988), 114–123.
- [Chen 1990] C.-M. Chen, *Superconvergence in finite element in domain with reentrant corner*, Natur. Sci. J. Xiangtan Univ. **12** (1990), 134–141, {MR 92c:65124}.
- [Chen 1991] C.-M. Chen, *Correction of bilinear finite element*, Acta Math. Sci. (English ed.) **11** (1991), 13–19, {MR 92f:65132}.
- [Chen and Huang 1989] C.-M. Chen and Y. Q. Huang, *Extrapolation of triangular linear elements in general domain*, Numer. Math. J. Chinese Univ. **11** (1989), 1–16, {MR 91d:65169}.

- [Chen and Huang 1990] C.-M. Chen and Y. Q. Huang, *A correction method for triangular linear elements*; (in Chinese), Hunan Ann. Math. **10** (1990), 117–121, {MR 92i:65167}.
- [Chen and Jin 1987] C.-M. Chen and J. C. Jin, *Extrapolation of biquadratic finite element approximations in a rectangular domain*; (in Chinese), Natur. Sci. J. Xiangtan Univ. (1987), 18–28, {MR 88k:65109}.
- [Chen, Larsson and Zhang 1989] C.-M. Chen, S. Larsson and N. Y. Zhang, *Error estimates of optimal order for finite element methods with interpolated coefficients for the nonlinear heat equation*, IMA J. Numer. Anal. **9** (1989), 507–524, {MR 91k:65141}.
- [Chen and Lin 1989] C.-M. Chen and Q. Lin, *Extrapolation of finite element approximation in a rectangular domain*, J. Comput. Math. **7** (1989), 227–233, {MR 90i:65190}.
- [Chen and Liu 1987] C.-M. Chen and J. Liu, *Superconvergence of gradient of triangular linear element in general domain*, Natur. Sci. J. Xiantang Univ. **1** (1987), 114–127, {MR 88f:65193}.
- [Chen and Thomée 1985] C.-M. Chen and V. Thomée, *The lumped mass finite element method for a parabolic problem*, J. Austral. Math. Soc. Ser. B **26** (1985), 329–354, {MR 86m:65117}.
- [Chen and Zhang 1990] C.-M. Chen and Z. J. Zhang, *Correction of one-dimensional finite element method*, Natur. Sci. J. Xiangtan Univ. **12** (1990), 124–131, {MR 92c:65088}.
- [Chen, H.-S. 1988] H.-S. Chen, *A mixed finite element and superconvergence*; (in Chinese), Natural Sci. J. Xiangtan Univ. **10** (1988), {MR 90g:65140}.
- [Chen, H.-S. 1989] H.-S. Chen, *A rectangular finite element analysis*, Natur. Sci. J. Xiangtan Univ. **11** (1989), 1–11, {MR 92k:65160}.
- [Chen, H.-S. 1990] H.-S. Chen, *A superconvergence estimate for mixed finite element methods for plate bending problems*; (in Chinese), Math. Numer. Sinica **12** (1990), 28–32, {MR 91e:73075}.
- [Chen, H.-S. and Lin 1992] H.-S. Chen and Q. Lin, *Finite element approximation using a graded mesh on domains with reentrant corners*, Systems Sci. Math. Sci. **5** (1992), 127–140, {MR 94c:65115}.
- [Chen, H.-S. and Lin 1992] H.-S. Chen and Q. Lin, *A high accuracy analysis of the mixed finite element method for biharmonic equations*, Systems Sci. Math. Sci. **5** (1992), 352–364, {MR 93k:65091}.
- [Chen, Z.-X. 1992] Z.-X. Chen, *A quasi-projection analysis for elastic wave propagation in fluid-saturated porous media*, J. Comput. Math. **10** (1992), 366–375, {MR 93g:65121}.
- [Cheng 1984] S.-J. Cheng, *Superconvergence of finite element approximation for Navier-Stokes equations*, Tagungsband “Extrapolationsverfahren in der Methode der finiten Elemente”, Bonn. Math. Schrift. **158** (1984), 31–45, {MR 87b:65195}.

- [Chou and Li 1990] S. H. Chou and Q. Li, *Convergence of the nonconforming Wilson element for a class of nonlinear parabolic equations*, Math. Comp. **54** (1990), 509–524, {MR 90i:65192}.
- [Chow and Carey 1993] S.-S. Chow and G. F. Carey, *Superconvergence phenomena in nonlinear two-point boundary value problems*, Numer. Methods Partial Differential Equations **9** (1993), 561–577.
- [Chow, Carey and Lazarov 1991] S.-S. Chow, G. F. Carey and R. D. Lazarov, *Natural and postprocessed superconvergence in semilinear problems*, Numer. Methods Partial Differential Equations **7** (1991), 245–259, {MR 92m:65131}.
- [Chow and Lazarov 1989] S.-S. Chow and R. D. Lazarov, *Superconvergence of flux computations for nonlinear problems*, Bull. Austral. Math. Soc. **40** (1989), 465–479, {MR 91d:65149}.
- [Ciarlet 1991] P. G. Ciarlet, *Basic Error Estimates for Elliptic Problems*, Handbook of Numerical Analysis, vol. II, Finite Element Methods (Part 1) (P. G. Ciarlet and J. L. Lions, eds.), Elsevier, 1991, pp. 17–351, {MR 92f:65001}.
- [Ciavaldini and Crouzeix 1985] J. F. Ciavaldini and M. Crouzeix, *A finite element method scheme for one-dimensional elliptic equations with high super convergence of nodes*, Numer. Math. **46** (1985), 417–427, {MR 86g:65140}.
- [Clement 1975] P. Clement, *Approximation by finite element functions using local regularization*, RAIRO Modél. Math. Anal. Numér. **R-2** (1975), 77–84, {MR 53#4569}.
- [Cohen, Joly and Tordjman 1993] G. Cohen, P. Joly and N. Tordjman, *Construction and analysis of higher order finite elements with mass lumping for the wave equation*, Proceedings of the Second International Conference on Mathematical and Numerical Aspects of Wave Propagation, SIAM, 1993, pp. 152–160, {MR 94d:65058}.
- [Comodi 1989] M. I. Comodi, *The Hellan–Herrman–Johnson method: some new error estimates and postprocessing*, Math. Comp. **52** (1989), 17–29, {MR 89f:65120}.
- [Crouzeix and Thomée 1985] M. Crouzeix and V. Thomée, *The stability in  $L_p$  and  $W_p^1$  of the  $L_2$ -projection into finite element function spaces*, Math. Comp. **48** (1985), 521–532, {MR 88f:41016}.
- [Cullen 1982] M. J. P. Cullen, *The use of quadratic finite element methods and irregular grids in the solution of hyperbolic problems*, J. Comput. Phys. **45** (1982), 221–245, {MR 83f:65152}.
- [Dautov 1980] R. Z. Dautov, *Superconvergence of difference schemes for the third boundary value problem for quasi-linear elliptic equations*; (in Russian), Chisl. Metody Mekh. Sploshn. Sredy **11** (1980), 62–80, {MR 84b:65102}.
- [Dautov 1982] R. Z. Dautov, *Superconvergence of schemes of the finite element method with numerical integration for fourth-order quasilinear elliptic equations*, Differential Equations **18** (1982), 818–824, {MR 83k:65080}.

- [Dautov and Lapin 1979] R. Z. Dautov and A. V. Lapin, *Difference schemes of an arbitrary order of accuracy for quasilinear elliptic equations*; (in Russian), Izv. Vysš. Učebn. Zaved. Matematika **209** (1979), 24–37, {MR 81g:65128}.
- [Dautov and Lapin 1981] R. Z. Dautov and A. V. Lapin, *Investigation of the convergence, in mesh norms, of finite-element-method schemes with numerical integration for fourth-order elliptic equations*, Differential Equations **17** (1981), 807–817, {MR 83h:65122}.
- [Dautov, Lapin and Lyashko 1980] R. Z. Dautov, A. V. Lapin and A. D. Lyashko, *Some mesh schemes for quasilinear elliptic equations*, USSR Computational Math. and Math. Phys. **20** (1980), 62–78, {MR 81j:65121}.
- [deBoor 1978] C. deBoor, *A Practical Guide to Splines*, Applied Mathematical Sciences **27**, Springer, 1978, {MR 80a:65027}.
- [deBoor 1981] C. deBoor, *On a max-bound for the least squares spline approximant*, Approximation Theory and Function Spaces (Z. Ciesielski, ed.), North-Holland, 1981, pp. 163–175, {MR 84k:41012}.
- [deBoor and Fix 1973] C. deBoor and G. J. Fix, *Spline approximation by quasiinterpolants*, J. Approx. Theory **8** (1973), 19–45, {MR 49#5643}.
- [de Groen 1981] P. P. N. de Groen, *A finite element method with a large mesh-width for a stiff two-point boundary value problem*, J. Comput. Appl. Math. **7** (1981), 3–15, {MR 82d:65059}.
- [Deny and Lions 1953–54] J. Deny and J. L. Lions, *Les espaces du type de Beppo Levi*, Ann. Inst. Fourier (Grenoble) **5** (1953–54), 305–370, {MR 17#646}.
- [Descloux 1972] J. Descloux, *On finite element matrices*, SIAM J. Numer. Anal. **9** (1972), 260–265, {MR 46#8402}.
- [Ding and Lin 1989] Y. H. Ding and Q. Lin, *Finite element expansion for variable coefficient elliptic problems*, Systems Sci. Math. Sci. **2** (1989), 54–69, {MR 92d:65190}.
- [Dougalis 1979] V. A. Dougalis, *Multi-step Galerkin methods for hyperbolic equations*, Math. Comp. **33** (1979), 563–584, {MR 81b:65081}.
- [Dougalis and Serbin 1980] V. A. Dougalis and S. M. Serbin, *On the superconvergence of Galerkin approximations to second-order hyperbolic equations*, SIAM J. Numer. Anal. **17** (1980), 431–446, {MR 81j:65106}.
- [Douglas 1984] J. Douglas Jr., *Improved accuracy through superconvergence in the pressure in the simulation of miscible displacement*, Lecture Notes in Computer Science, North Holland, 1984, pp. 633–638, {MR 86f:65009}.
- [Douglas 1985] J. Douglas Jr., *Superconvergence in the pressure in the simulation of miscible displacement*, SIAM J. Numer. Anal. **22** (1985), 962–969, {MR 86j:65129}.
- [Douglas and Dupont 1973a] J. Douglas Jr. and T. Dupont, *Superconvergence for Galerkin methods for the two point boundary problem via local projections*, Numer. Math. **21** (1973), 270–278, {MR 48#10138}.

- [Douglas and Dupont 1973b] J. Douglas Jr. and T. Dupont, *Some superconvergence results for Galerkin methods for the approximate solution of two-point boundary value problems*, Topics in Numerical Analysis, Academic Press, 1973, pp. 89–92, {MR 51#2295}.
- [Douglas and Dupont 1974] J. Douglas Jr. and T. Dupont, *Galerkin approximations for the two point boundary problem using continuous, piecewise polynomial spaces*, Numer. Math. **22** (1974), 99–109, {MR 50#15360}.
- [Douglas and Dupont 1975] J. Douglas Jr. and T. Dupont, *A Galerkin method for a nonlinear Dirichlet problem*, Math. Comp. **29** (1975), 689–696, {MR 55#4742}.
- [Douglas, Dupont and Wahlbin 1975a] J. Douglas Jr., T. Dupont and L. B. Wahlbin, *Optimal  $L_\infty$  error estimates for Galerkin approximations to solutions of two-point boundary value problems*, Math. Comp. **29** (1975), 475–483, {MR 51#7298}.
- [Douglas, Dupont and Wahlbin 1975b] J. Douglas Jr., T. Dupont and L. B. Wahlbin, *The stability in  $L^q$  of the  $L^2$ -projection into finite element function spaces*, Numer. Math. **23** (1975), 193–197, {MR 52#4669}.
- [Douglas, Dupont and Wheeler 1974a] J. Douglas, Jr., T. Dupont and M. F. Wheeler, *A Galerkin procedure for approximating the flux on the boundary for elliptic and parabolic boundary value problems*, RAIRO Modél. Math. Anal. Numér. **8** (1974), 47–59, {MR 50#11811}.
- [Douglas, Dupont and Wheeler 1974b] J. Douglas, Jr., T. Dupont and M. F. Wheeler, *An  $L^\infty$  estimate and a superconvergence result for a Galerkin method for elliptic equations based on tensor products of piecewise polynomials*, RAIRO Modél. Math. Anal. Numér. **8** (1974), 61–66, {MR 50#11812}.
- [Douglas, Dupont and Wheeler 1974c] J. Douglas Jr., T. Dupont and W. F. Wheeler, *Some superconvergence results for an  $H^1$  Galerkin procedure for the heat equation*, Lecture Notes in Comput. Sci., vol. 10, Springer 1974, pp. 288–311, {MR 56#10056}.
- [Douglas and Gupta 1986] J. Douglas Jr. and C. P. Gupta, *Superconvergence for a mixed finite element method for elastic wave propagation in a plane domain*, Numer. Math. **49** (1986), 189–202, {MR 88c:65095}.
- [Douglas and Gupta 1988] J. Douglas Jr. and C. P. Gupta, *Superconvergence and interior estimates for a mixed finite element method for elastic waves in a planar domain*, Mat. Apl. Comput. **7** (1988), 75–99, {MR 90k:65172}.
- [Douglas, Gupta and Li 1985] J. Douglas, Jr., C. P. Gupta and G. Y. Li, *Interior and superconvergence esitmates for a primal hybrid finite element method for second order elliptic problems*, Calcolo **22** (1985), 187–207, {MR 87g:65137}.
- [Douglas and Milner 1985] J. Douglas Jr. and F. Milner, *Interior and superconvergence estimates for mixed methods for second order elliptic problems*, RAIRO Modél. Math. Anal. Numér. **19** (1985), 397–428, {MR 88a:65129}.
- [Douglas and Wang 1989] J. Douglas Jr. and J. Wang, *Superconvergence of mixed finite element methods on rectangular domains*, Calcolo **26** (1989), 121–133, {MR 92e:65147}.

- [Du and Liu 1984] M. S. Du and C. F. Liu, *Stability and convergence for the discrete-discontinuous finite element method*; (in Chinese), J. Numer. Methods Comput. Appl. **5** (1984), 219–231, {MR 86j:65131}.
- [Dunlap, Li, Taylor, Wahlbin and Warren] R. Dunlap, J. Li, A. Taylor, L. B. Wahlbin and J. Warren, *A computational investigation into superconvergence for second order two-point boundary value problems*, in preparation.
- [Dupont 1976] T. Dupont, *A unified theory of superconvergence for Galerkin methods for two-point boundary problems*, SIAM J. Numer. Anal. **13** (1976), 362–368, {MR53#12021}.
- [Dupont and Scott 1980] T. Dupont and R. Scott, *Polynomial approximation of functions in Sobolev spaces*, Math. Comp **34** (1980), 441–463, {MR 81h:65014}.
- [Duran 1990] R. Duran, *Superconvergence for rectangular mixed finite elements*, Numer. Math. **58** (1990), 287–298, {MR 92a:65289}.
- [Duran, Muschietti and Rodriguez 1992] R. Duran, M. A. Muschietti and R. Rodriguez, *Asymptotically exact error estimators for rectangular finite elements*, SIAM J. Numer. Anal. **29** (1992), 78–88, {MR 93a:65148}.
- [Dzhishkariani 1990] A. V. Dzhishkariani, *On the convergence of the finite element method with one iteration*; (in Russian), Zh. Vychisl. Mat. i Mat. Fiz. **30** (1990), 791–796, {MR 91h:65171}.
- [Dzhishkariani and Manveliani 1992] A. V. Dzhishkariani and G.S. Manveliani, *On the superconvergence of an approximate projection method with one iteration*; (in Russian), Georgian Academy of Science, Proceedings of A. Razmadze Mathematical Institut **100** (1992), 89–95.
- [El Hatri 1983a] M. El Hatri, *Superconvergence of the lumped mass approximation of parabolic equations*, C.R. Acad. Bulgare Sci. **36** (1983), 575–578, {MR85h:65187}.
- [El Hatri 1983b] M. El Hatri, *Superconvergence of axisymmetrical boundary value problem*, C. R. Acad. Bulgare Sci. **36** (1983), 1499–1502, {MR 86e:65153}.
- [El Hatri 1984] M. El Hatri, *Superconvergence in the finite element method for a degenerated boundary value problem*, Constructive Theory of Functions, Bulgarian Academy of Science, 1984, pp. 328–333.
- [El Hatri 1987] M. El Hatri, *Estimation d'erreur optimale et de type superconvergence de la methode des elements finis pour un probleme aux limites degeneres*, RAIRO Model. Math. Anal. Numer. **21** (1987), 27–61, {MR 89a:65168}.
- [Eriksson 1985] K. Eriksson, *Finite element methods of optimal order for problems with singular data*, Math. Comp. **44** (1985), 345–360, {MR 86j:65155b}.
- [Eriksson and Nie 1987] K. Eriksson and Y. Y. Nie, *Convergence analysis for a nonsymmetric Galerkin method for a class of singular boundary value problems in one space dimension*, Math. Comp. **49** (1987), 167–186, {MR 88h:65182}.
- [Ewing, Lazarov, Russell and Vassilevski 1990] R. E. Ewing, R. D. Lazarov, T. E. Russell and P. S. Vassilevsky, *Local refinement via domain decomposition*

- techniques for mixed finite element methods with rectangular Raviart–Thomas elements*, Third International Symposium on Domain Decomposition Methods for Partial Differential Equations, SIAM 1990, pp. 98–114, {MR 91f:65191}.
- [Ewing, Lazarov and Wang 1991] R. E. Ewing, R. D. Lazarov and J. Wang, *Superconvergence of the velocity along the Gauss lines in mixed finite element methods*, SIAM J. Numer. Anal. **18** (1991), 1015–1029, {MR 92e:65149}.
- [Ewing, Shen and Wang 1991] R. E. Ewing, J. Shen and J. Wang, *Application of superconvergence to problems in the simulation of miscible displacement*, Comput. Methods Appl. Mech. Engrg. **89** (1991), 73–84, {MR 92k:76078}.
- [Ewing and Wang 1992] R. E. Ewing and J. P. Wang, *Analysis of mixed finite element methods on locally refined grids*, Numer. Math. **63** (1992), 183–194, {MR 93h:65137}.
- [Fairweather 1978] G. Fairweather, *Finite Element Galerkin Methods for Differential Equations*, Lecture Notes in Pure and Applied Mathematics 34, Marcel Dekker, 1978, {MR 58#13781}.
- [Fan and Wang 1980] Y. X. Fan and S. J. Wang, *On the application of an interpolation matrix for computation of stresses in finite elements*, Numerical Methods in Geomechanics, Balkema, Rotterdam 1980, pp. 1273–1280, {MR 82e:73064}.
- [Filho 1968] F. V. Filho, *Comment on “Computation of stress resultants from the element stiffness matrices”* (= [Stricklin 1966]), AIAA Journal **6** (1968), 571–572.
- [Fish and Markolefas 1994] J. Fish and S. Markolefas, *Adaptive global-local refinement strategy based on the interior error estimates of the  $h$ -method*, Internat. J. Numer. Methods Engrg. **37** (1994), 827–838.
- [Fix and Nassif 1972] G. Fix and N. Nassif, *On finite element approximations to time-dependent problems*, Numer. Math. **19** (1972), 127–135, {MR 46#10218}.
- [Fletcher 1984] C. A. J. Fletcher, *Computational Galerkin Methods*, Springer Series in Computational Physics, 1984, {MR 85i:65003}.
- [Fortin and Mghazli 1992] M. Fortin and Z. Mghazli, *Analyse d’un élément mixte pour le problème de Stokes, II. Constructions et estimations d’erreur*, Numer. Math. **62** (1992), 161–188, {MR 93i:76054}.
- [Franca 1990] L. P. Franca, *On the superconvergence of the satisfying–Babuska–Brezzi method*, Internat. J. Numer. Methods Engrg. **29** (1990), 1715–1726, {MR 91g:65182}.
- [Frehse and Rannacher 1976] J. Frehse and R. Rannacher, *Eine  $L^1$ -Fehlerabschätzung diskreter Grundlösungen in der Methode der finiten Elemente*, Tagungsband “Finite Elemente”, Bonn. Math. Schrift. **89** (1975), 92–114, {MR 57#11104}.
- [Frehse and Rannacher 1978] J. Frehse and R. Rannacher, *Asymptotic  $L_\infty$ -error estimate for linear finite element approximations of quasilinear boundary value problems*, SIAM J. Numer. Anal. **15** (1978), 418–431, {MR 58#19224}.

- [Frey, Hall and Porsching 1978] A. E. Frey, C. A. Hall and T. A. Porsching, *Some results on the global inversion of bilinear and quadratic isoparametric finite element transformation*, Math. Comp. **32** (1978), 725–749, {MR 57#14508}.
- [Friedman 1969] A. Friedman, *Partial Differential Equations*, Holt, Rinehart and Winston, 1969, {MR 56#3433}.
- [Gastaldi and Nochetto 1987] L. Gastaldi and R. Nochetto, *Optimal  $L_\infty$ -error estimates for nonconforming and mixed finite element methods of lowest order*, Numer. Math. **50** (1987), 587–611, {MR 88f:65196}.
- [Gastaldi and Nochetto 1989] L. Gastaldi and R. Nochetto, *Sharp maximum norm error estimates for general mixed finite element approximations to second order elliptic equations*, RAIRO Model. Math. Anal. Numer. **23** (1989), 103–128, {MR 91b:65125}.
- [Genis 1984] A. Genis, *On finite element methods for the Euler–Poisson–Darboux equation*, SIAM J. Numer. Anal. **21** (1984), 1080–1106, {MR 86f:65171}.
- [Gilbarg and Trudinger 1983] D. Gilbarg and N. S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Second Edition, Grundlehren der mathematischen Wissenschaften, vol. 224, Springer, 1983, {MR 86c:35035}.
- [Goodsell and Whiteman 1989] G. Goodsell and J. R. Whiteman, *A unified treatment of superconvergent recovered gradient functions for piecewise linear finite element approximations*, Internat. J. Numer. Methods Engrg. **27** (1989), 469–481, {MR 91k:65144}.
- [Goodsell and Whiteman 1990] G. Goodsell and J. R. Whiteman, *Pointwise superconvergence of recovered gradients for piecewise linear finite element approximations to problems of planar linear elasticity*, Numer. Methods Partial Differential Equations **6** (1990), 59–74, {MR 91c:73081}.
- [Goodsell and Whiteman 1991] G. Goodsell and J. R. Whiteman, *Superconvergence of recovered gradients of piecewise quadratic finite element approximations, I, II*, Numer. Methods Partial Differential Equations **7** (1991), 61–83, 85–99, {MR 92e:65151 a,b}.
- [Gradshteyn and Ryzhik 1979] I. S. Gradshteyn and I. M. Ryzhik, *Tables of Integrals, Series and Products*, Corrected and Enlarged Edition, Academic Press, 1979, {MR 81g:33001}.
- [Grisvard 1985] P. Grisvard, *Elliptic Problems in Nonsmooth Domains*, Pitman, 1985, {MR 86m:35044}.
- [Gruber 1987] R. Gruber, *Finite elements in magnetohydrodynamics: ideal linear stability problem*, Finite Elements in Physics, North–Holland 1987, pp. 299–333, {MR 89b:65264}.
- [Haverkamp 1984] R. Haverkamp, *Eine Aussage zur  $L_\infty$ -Stabilität und zur genauen Konvergenzordnung der  $H_0^1$ -Projection*, Numer. Math. **44** (1984), 393–405, {MR 86f:65194}.

- [Heinrich 1987] B. Heinrich, *Finite Difference Methods on Irregular Networks: A Generalized Approach to Second Order Elliptic Problems*, International Series of Numerical Mathematics 82, Birkhäuser, 1987, {MR 88h:65004}.
- [Helfrich 1984] H.-P. Helfrich, *Asymptotic expansion for the finite element approximation of parabolic problems*, Tagungsband “Extrapolationsverfahren in der Methode der finiten Elemente”, Bonn. Math. Schrift. **158** (1984), 11–30, {MR 86k:65090}.
- [Hilbert 1973] S. Hilbert, *A mollifier useful for approximations in Sobolev spaces and some applications to approximating solutions of differential equations*, Math. Comp. **37** (1973), 81–89, {MR 48#10047}.
- [Hildebrandt and Wienholz 1964] S. Hildebrandt and E. Wienholz, *Constructive proofs of representation theorems in separable Hilbert space*, Comm. Pure Applied Math. **17** (1964), 369–373, {MR 29#3881}.
- [Hinton and Campbell 1974] E. Hinton and J. S. Campbell, *Local and global smoothing of discontinuous finite element functions using a least squares method*, Internat. J. Numer. Methods Engrg. **8** (1974), 461–480, {MR 53#14859}.
- [Hinton and Owen 1977] E. Hinton and D. R. J. Owen, *Finite Element Programming*, Academic Press, 1977.
- [Hinton, Scott and Ricketts 1975] E. Hinton, F. C. Scott and R. E. Ricketts, *Local least squares stress smoothing for parabolic isoparametric elements*, Internat. J. Numer. Methods Engrg. **9** (1975), 235–238.
- [Hlaváček and Křížek 1987] I. Hlaváček and M. Křížek, *On a superconvergent finite element scheme for elliptic systems, Parts I–III*, Apl. Mat. **32** (1987), 131–153, 200–213, 276–289, {MR 88c:65099, 88j:65251, 88j:65252}.
- [Höhn and Mittelmann 1981] W. Höhn and H. D. Mittelmann, *Some remarks on the discrete maximum principle for finite elements of higher order*, Computing **27** (1981), 145–154, {MR 83a:65109}.
- [Hörmander 1960] L. Hörmander, *Estimates for translation invariant operators in  $L_p$ -spaces*, Acta Math. **104** (1960), 93–140, {MR 22#12389}.
- [Huang and Wu 1982] D. Huang and D. Wu, *The superconvergence of the spline finite element solution and its second order derivative for the two-point boundary value problem of a fourth order differential equation*, J. Zhejiang Univ. **3** (1982), 92–99.
- [Huang, M. Y. and Wang 1984] M. Y. Huang and X. F. Wang, *A note on  $L_\infty$  estimates of semidiscrete Galerkin approximations*; (in Chinese), Numer. Math. J. Chinese Univ. **6** (1984), 376–379, {MR 87b:65162}.
- [Huang, X. F. 1986] X. F. Huang, *The optimal point of the gradient of finite element solution*, Appl. Math. Mech. **7** (1986), 785–794, {MR 88d:65164}.
- [Huang, Y. Q. and Lin 1991] Y. Q. Huang and Q. Lin, *Asymptotic expansions and superconvergence of finite element solutions to parabolic equations*; (in Chinese), J. Systems Sci. Math. Soci. **11** (1991), 327–335, {MR 92h:65170}.

- [Hughes 1987] T. J. R. Hughes, *The Finite Element Method*, Prentice-Hall, 1987, {MR 90i:65001}.
- [Hulme 1972] B. L. Hulme, *One-step piecewise polynomial Galerkin methods for initial value problems*, Math. Comp. **26** (1972), 415–426, {MR 47#9834}.
- [Irons and Razzaque 1972] B. M. Irons and A. Razzaque, *Experience with the patch test for convergence of finite elements*, Mathematical Foundations of the Finite Element Method with Applications to Partial Differential Equations, Academic Press, 1972, pp. 557–587, {MR 54#11813}.
- [Jamet 1976] P. Jamet, *Estimation d'erreur pour des éléments finis droits presque dégénérés*, RAIRO Modél. Math. Anal. Numér. **10** (1976), 43–61, {MR 56#13521}.
- [Jin and Huang 1993] J. C. Jin and Y. Q. Huang, *Superconvergence of derivatives of biquadratic finite element solutions*; (in Chinese), Natur. Sci. J. Xiangtan Univ. **15** (1993), 8–14.
- [Johnson and Pitkäranta 1982] C. Johnson and J. Pitkäranta, *Analysis of some mixed finite element methods related to reduced integration*, Math. Comp. **33** (1982), 375–400, {MR 83d:65287}.
- [Johnson and Thomée 1975] C. Johnson and V. Thomée, *Error estimates for a finite element approximation of a minimal surface*, Math. Comp. **29** (1975), 343–349, {MR 53#4571}.
- [Johnson, R. W. and MacKinnon 1992] R. W. Johnson and R. J. MacKinnon, *An auxiliary equation method for obtaining superconvergent finite-element approximations*, Comm. Appl. Numer. Methods **8** (1992), 99–107, {MR 92i:65168}.
- [Kantchev 1988] V. Kantchev, *Superconvergence of the gradient for linear finite elements for nonlinear elliptic problems*, Teubner-Texte Math. 107, Teubner 1988, 199–204.
- [Kendall and Wheeler 1976] R. P. Kendall and M. F. Wheeler, *A Crank-Nicolson  $H^{-1}$ -Galerkin procedure for parabolic problems in a single space variable*, SIAM J. Numer. Anal. **13** (1976), 861–876, {MR 58#13785}.
- [King and Serbin 1976] J. T. King and S. M. Serbin, *Boundary flux estimates for elliptic problems by the perturbed variational method*, Computing **16** (1976), 339–347, {MR 54#6524}.
- [Korneev 1982] V. G. Korneev, *Superconvergence of solutions of the finite element method in mesh norms*; (in Russian), Ž. Vyčisl. Mat. i. Mat. Fiz. **22** (1982), 1133–1148, {MR 84b:65114}.
- [Korneev 1983] V. G. Korneev, *Convergence of finite element solutions in difference norms*; (in Russian), Metody Vychisl. (1983), 3–45, {MR 85c:65138}.
- [Krasovskii 1967] J. P. Krasovskii, *Isolation of singularities of the Green's function*, Math. USSR-Izv. **1** (1967), 935–966, {MR 36#6788}.

- [Kreiss, Manteuffel, Swartz, Wendroff and White 1986] H. O. Kreiss, T. A. Manteuffel, B. K. Swartz, B. Wendroff and A. B. White Jr., *Supraconvergent schemes on irregular grids*, Math. Comp. **47** (1986), 537–554, {MR 88b:65082}.
- [Křížek 1986] M. Křížek, *Superconvergence results for linear triangular elements*, Lecture Notes in Mathematics 1192, Springer, 1986, pp. 315–320, {MR 88i:65132}.
- [Křížek 1994a] M. Křížek, *Higher order global accuracy of a weighted averaged gradient of the Courant element on irregular meshes*, Finite Element Methods: fifty years of the Courant element (M. Křížek, P. Neittaanmäki and R. Stenberg, eds.), Marcel Dekker, 1994, pp. 267–276.
- [Křížek 1994b] M. Křížek, *Supreconvergence phenomena in the finite element methods*, Comput. Methods Appl. Mech. Engrg. (to appear).
- [Křížek, Lin and Huang 1988] M. Křížek, Q. Lin and Y.-Q. Huang, *A nodal superconvergence arising from combination of linear and bilinear elements*, Systems Sci. Math. Sci. **1** (1988), 191–197, {MR 91f:65170}.
- [Křížek and Neittaanmäki 1984] M. Křížek and P. Neittaanmäki, *Superconvergence phenomenon in the finite element method arising from averaging gradients*, Numer. Math. **45** (1984), 105–116, {MR 86c:65135}.
- [Křížek and Neittaanmäki 1987a] M. Křížek and P. Neittaanmäki, *On superconvergence techniques*, Acta Appl. Math. **9** (1987), 175–198, {MR 88h:65208}.
- [Křížek and Neittaanmäki 1987b] M. Křížek and P. Neittaanmäki, *On a global superconvergence of the gradient of linear triangular elements*, J. Comput. Appl. Math. (1987), {MR 88h:65209}.
- [Křížek and Neittaanmäki 1989] M. Křížek and P. Neittaanmäki, *On  $O(h^4)$ -superconvergence of piecewise bilinear FE-approximations*, Mat. Apl. Comput. **8** (1989), 49–61, {MR 91g:65242}.
- [Křížek and Neittaanmäki 1990] M. Křížek and P. Neittaanmäki, *Finite Element Approximation of Variational Problems and Application*, John Wiley and Sons, 1990, {MR 91h:65174}.
- [Lapin 1987] A. V. Lapin, *Mesh schemes of higher order of accuracy for some classes of variational inequalities*, Soviet J. Numer. Anal. Math. Modelling **2** (1987), 37–55, {MR 88m:65176}.
- [Layton 1983] W. J. Layton, *The finite element method for a degenerate hyperbolic partial differential equation*, BIT **23** (1983), 231–238, {MR 84f:65073}.
- [Lazarov 1988] R. D. Lazarov, *Superconvergence of the gradient for triangular and tetrahedral finite elements of a solution of linear problems in elasticity theory*, (in Russian), Computational Processes and System **6**, Nakua Moscow 1988, pp.180–191, {MR 90f:65207}.
- [Lazarov, Andreev and El Hatri 1984] R. D. Lazarov, A. B. Andreev and M. El Hatri, *Superconvergence of the gradients in the finite element method for some elliptic and parabolic problems*, Variational–Difference Methods in Mathematical

- Physics, Part 2, (Proc. Fifth Int. Conf., Moscow 1983), Viniti, Moscow, 1984, pp. 13–25.
- [Lazarov and Pehlivanov 1991] R. D. Lazarov and A. I. Pehlivanov, *Local superconvergence analysis of 3-D linear finite elements for boundary flux calculations*, The Mathematics of Finite Elements and Applications VII (J. R. Whiteman, ed.), Academic Press, 1991, pp. 75–83, {MR 92d:65007}.
- [Levine 1985] N. Levine, *Superconvergent recovery of the gradient from piecewise linear finite-element approximations*, IMA J. Numer. Anal. **5** (1985), 407–427, {MR 87b:65206}.
- [Lesaint and Zlámal 1979] P. Lesaint and M. Zlámal, *Superconvergence of the gradient of finite element solutions*, RAIRO Modél. Math. Anal. Numér. **13** (1979), 139–166, {MR 80g:65112}.
- [Leyk 1982] Z. Leyk, *Superconvergence in the finite element method*; (in Polish), Mat. Stos. **20** (1982), 93–107, {MR 85k:65090}.
- [Li 1990] B. Li, *Superconvergence for higher-order triangular finite elements*, Chinese J. Numer. Math. Appl. **12** (1990), 75–79, {MR 92d:65196}.
- [Lin 1989] Q. Lin, *Finite element error expansion for nonuniform quadrilateral meshes*, Systems Sci. Math. Sci. **1** (1989), 275–282, {MR 92c:65137}.
- [Lin 1991] Q. Lin, *Superconvergence of FEM for singular solution*, J. Comput. Math. **9** (1991), 111–114, {MR 93b:65171}.
- [Lin 1992] Q. Lin, *Interpolated finite elements and global superconvergence*, Numerical Methods for Partial Differential Equations, World Sci. Publishing, 1992, pp. 91–95.
- [Lin and Lu 1984a] Q. Lin and T. Lu, *Asymptotic expansions for finite element approximation of elliptic problem on polygonal domains*, Computing Methods in Applied Sciences and Engineering VI (R. Glowinski and J.-L. Lions, eds.), Elsevier, 1984, pp. 317–321, {MR 86f:65009}.
- [Lin and Lu 1984b] Q. Lin and T. Lu, *Asymptotic expansions for finite element eigenvalues and finite element solution*, Tagungsband “Extrapolationsverfahren in der Methode der finiten Elemente”, Bonn. Math. Schrift. **158** (1984), 1–10, {MR 86j:65146}.
- [Lin, Lu and Shen 1983] Q. Lin, T. Lu and S.-M. Shen, *Maximum norm estimates, extrapolation and optimal points of stresses for the finite element methods on the strongly regular triangulation*, J. Comput. Math. **1** (1983), 376–383.
- [Lin, Sloan and Xie 1990] Q. Lin, I. H. Sloan and R. Xie, *Extrapolation of the iterated-collocation method for integral equations of the second kind*, SIAM J. Numer. Anal. **27** (1990), 1535–1541, {MR 91j:65199}.
- [Lin and Whiteman 1991] Q. Lin and J. R. Whiteman, *Superconvergence of recovered gradients of finite element approximations on nonuniform rectangular and quadrilateral meshes*, The Mathematics of Finite Elements and Applications VII (J. R. Whiteman, ed.), Academic Press, 1991, pp. 563–571, {MR 92d:65007}.

- [Lin and Xie 1986] Q. Lin and R.-F. Xie, *Some advances in the study of error expansion for finite elements*, J. Comput. Math. **4** (1986), 368–382, {MR 88j:65242}.
- [Lin and Xie 1987] Q. Lin and R.-F. Xie, *Error expansions for finite element approximations and their applications*, Lecture Notes in Mathematics, vol. 1297, Springer, 1987, pp. 98–112, {MR 89b:65272}.
- [Lin and Xie 1988] Q. Lin and R.-F. Xie, *How to recover the convergent rate for Richardson extrapolation on bounded domains*, J. Comput. Math. **6** (1988), 68–79, {MR 90b:65206}.
- [Lin and Xie 1989] Q. Lin and R.-F. Xie, *Error expansion for FEM and superconvergence under natural assumption*, J. Comput. Math. **7** (1989), 402–411.
- [Lin and Xie 1990] Q. Lin and R.-F. Xie, *Extrapolation of bilinear finite element solution with local refinement mesh*, Chinese Ann. Math. Ser. B. **11** (1990), 179–190, {MR 92a:65296}.
- [Lin and Xu 1985] Q. Lin and J. Xu, *Linear finite elements with high accuracy*, J. Comput. Math. **3** (1985), 115–133, {MR 87k:65141}.
- [Lin and Zhou 1992] Q. Lin and A. H. Zhou, *Defect correction for finite element gradient*, Systems Sci. Math. Sci. **5** (1992), 287–288, {MR 93h:65039}.
- [Lin and Zhou 1993] Q. Lin and A.-H. Zhou, *Notes on superconvergence and its related topics*, J. Comput. Math. **11** (1993), 211–214.
- [Lin and Zhu 1984] Q. Lin and Q.-D. Zhu, *Asymptotic expansion for the derivative of finite elements*, J. Comput. Math. **2** (1984), 361–363, {MR 87m:65185}.
- [Lin and Zhu 1986] Q. Lin and Q.-D. Zhu, *Local asymptotic expansion and extrapolation for finite elements*, J Comput. Math. **4** (1986), 263–265, {MR 87k:65142}.
- [Lions 1969] J. L. Lions, *Quelques Méthodes de Résolution des Problèmes aux Limites Non Linéaires*, Dunod, 1969, {MR 41#4326}.
- [Liu 1994a] X. Liu, *Interior error estimates for nonconforming finite element methods of the Stokes equation* (to appear).
- [Liu 1994b] X. Liu, *Lecture at the “Finite Element Circus”*, Penn State University, Fall 1994.
- [Long and Morton 1977] M. J. Long and K. W. Morton, *The use of divided differences in finite element calculations*, J. Inst. Maths. Applies. **19** (1977), 307–323, {MR 56#7116}.
- [Louis 1979] A. Louis, *Acceleration of convergence for finite element solutions of the Poisson equation*, Numer. Math. **33** (1979), 43–53, {MR 81i:65091}.
- [Luo 1992] P. Luo, *Superconvergence estimates for a nonconforming membrane element*, Northeast Math. J. **8** (1992), 477–482.
- [MacKinnon and Carey 1989] R. J. MacKinnon and G. F. Carey, *Superconvergent derivatives: a Taylor series analysis*, Internat. J. Numer. Methods Engrg. **28** (1989), 489–509, {MR 90e:73083}.

- [MacKinnon and Carey 1990] R. J. MacKinnon and G. F. Carey, *Nodal superconvergence and solution enhancement for a class of finite-element and finite-difference methods*, SIAM J. Sci. Statist. Comput. **11** (1990), 343–353, {MR 90m:65201}.
- [Makarenko 1987] E. N. Makarenko, *Superconvergence of schemes of the finite element method for the Poisson equation on regular nets*; (in Russian), Vestnik Moskov. Univ. Ser. XV Vychisl. Mat. Kibernet. (1987), 63–66, {MR 89a:65169}.
- [Mansfield 1981] L. Mansfield, *On mixed finite element methods for elliptic equations*, Comput. Math. Appl. **7** (1981), 59–66, {MR 82b:65133}.
- [Manteuffel and White 1986] T. A. Manteuffel and A. B. White Jr, *The numerical solution of second-order boundary value problems on nonuniform meshes*, Math. Comp. **47** (1986), 511–535, {MR 87m:65116}.
- [Marshall 1979] R. S. Marshall, *An alternative interpretation of superconvergence*, Internat. J. Numer. Methods. Engrg. **14** (1979), 1707–1709, {MR 80m:65079}.
- [Milner 1989] F. A. Milner,  *$L_\infty$ -error estimates for linear elasticity problems*, J. Comp. Appl. Math. **25** (1989), 305–313, {MR 90e:73023}.
- [Mock 1976] M. S. Mock, *Projection methods with different trial and test spaces*, Math. Comp. **30** (1976), 400–416, {MR 54#11814}.
- [Moore and Flaherty 1990] P. K. Moore and J. E. Flaherty, *A local refinement finite-element method for one-dimensional parabolic systems*, SIAM J. Numer. Anal. **27** (1990), 1422–1444, {MR 91h:65146}.
- [Morley 1989] M. E. Morley, *A family of mixed finite elements for linear elasticity*, Numer. Math. **55** (1989), 633–666, {MR 90f:73006}.
- [Nakao 1983] M. T. Nakao, *Interior estimates and superconvergence for  $H^{-1}$ -Galerkin method to elliptic equations*, Bull. Kyushu Inst. Tech. Math. Natur. Sci. **30** (1983), 19–30, {MR 85i:65154}.
- [Nakao 1984] M. T. Nakao, *Some superconvergence estimates for a Galerkin method for elliptic problems*, Bull. Kyushu Inst. Tech. Math. Natur. Sci. **31** (1984), 49–58, {MR 86e:65159}.
- [Nakao 1985a] M. T. Nakao, *Some superconvergence of Galerkin approximations for parabolic and hyperbolic problems in one space dimension*, Bull. Kyushu Inst. Tech. Math. Natur. Sci. **32** (1985), 1–14, {MR 86j:65123}.
- [Nakao 1985b] M. T. Nakao, *Error estimates of a Galerkin method for some nonlinear Sobolev equations in one space dimension*, Numer. Math. **47** (1985), 139–157, {MR 86j:65134}.
- [Nakao 1986] M. T. Nakao, *Some superconvergence for a Galerkin method by averaging gradients in one-dimensional problems*, J. Inform. Process. **9** (1986), 130–134, {MR 88e:65144}.
- [Nakao 1987a] M. T. Nakao, *Superconvergence of the gradient of Galerkin approximations for elliptic problems*, RAIRO Modél. Math. Anal. Numér. **21** (1987), 679–695, {MR 88m:65180}.

- [Nakao 1987b] M. T. Nakao, *Superconvergence of the gradient of Galerkin approximations for elliptic problems*, J. Comput. Appl. Math. **20** (1987), 341–348, {MR 89a:65172}.
- [Nakata and Wheeler 1986] M. Nakata and M. F. Wheeler, *Some superconvergence results for mixed finite element methods for linear parabolic equations*, Lecture Notes in Mathematics 1230, Springer, 1986, pp. 167–174, {MR 89b:65273}.
- [Nakata, Weiser and Wheeler 1985] M. Nakata, A. Weiser and M. F. Wheeler, *Some superconvergence results for mixed finite element methods for elliptic problems on rectangular domains*, The Mathematics of Finite Elements and Applications V (J. R. Whiteman, ed.), Academic Press, 1985, pp. 367–389, {MR 87g:65144}.
- [Natterer 1975] F. Natterer, *Über die punktweise Konvergenz finiter Elemente*, Numer. Math. **25** (1975), 67–77, {MR 57#14514}.
- [Nečas 1967] J. Nečas, *Les Méthodes Directes en Théorie des Équations Elliptiques*, Masson et C<sup>ie</sup>, 1967, {MR 37#3168}.
- [Neittaanmäki and Křížek 1988] R. Neittaanmäki and M. Křížek, *On  $0(h^4)$ -superconvergence of piecewise bilinear FE-approximations*, Teubner-Texte Math. 107, Teubner, 1988, pp. 250–255.
- [Neittaanmäki and Saranen 1982] P. Neittaanmäki and J. Saranen, *A modified least squares FE-element method for ideal fluid flow problems*, J. Comput. Appl. Math. **8** (1982), 165–169.
- [Ni 1986] P. Ni, *Superconvergence of generalized Galerkin method solutions and their derivatives*; (in Chinese), Dongbei Shida Xuebao **2** (1986), 13–17, {MR 87m:65189}.
- [Ni and Wu 1986] P. Ni and W. Wu, *Estimates of superconvergence for generalized Galerkin methods*; (in Chinese), Numer. Math. J. Chinese Univ. **8** (1986), 153–158, {MR 88b:65096}.
- [Nicolaides 1972] R. A. Nicolaides, *On a class of finite elements generated by Lagrange interpolation I*, SIAM J. Numer. Anal. **9** (1972), 435–445, {MR 47#6058}.
- [Nicolaides 1973] R. A. Nicolaides, *On a class of finite elements generated by Lagrange interpolation II*, SIAM J. Numer. Anal. **10** (1973), 182–189, {MR 47#6059}.
- [Nitsche 1975] J. A. Nitsche,  *$L_\infty$  convergence for finite element approximation*, Proceedings Second Conference on Finite Elements, Rennes, France (1975).
- [Nitsche 1976] J. A. Nitsche, *Über  $L_\infty$ -Abschätzungen von Projektionen auf finite Elemente*, Tagungsband “Finite Elemente”, Bonn. Math. Schrift. **89** (1976), 13–30, {MR 56#10062}.
- [Nitsche 1977a] J. A. Nitsche,  *$L_\infty$  convergence of finite element approximations*, Lecture Notes in Mathematics, vol. 606, Springer, 1977, pp. 261–274, {MR 58#8351}.

- [Nitsche 1977b] J. A. Nitsche, *On  $L_\infty$ -convergence of finite element approximations to the solution of a nonlinear boundary value problem*, Topics in Numerical Analysis III, Academic Press, 1987, pp. 317–325, {MR 81i:65094}.
- [Nitsche and Schatz 1972] J. Nitsche and A. H. Schatz, *On local approximation properties of  $L_2$ -projection on spline-subspaces*, Applicable Anal. **2** (1972), 161–168, {MR 53#1127}.
- [Nitsche and Schatz 1974] J. A. Nitsche and A. H. Schatz, *Interior estimates for Ritz–Galerkin methods*, Math. Comp. **28** (1974), 937–958, {MR 51#9525}.
- [Niu and Shephard 1993] Q. Niu and M. S. Shephard, *Superconvergent extraction techniques for finite element analysis*, Internat. J. Numer. Methods Engrg. **36** (1993), 811–836, {MR 93k:73080}.
- [Noor and Babuška 1987] A. K. Noor and I. Babuška, *Quality assessment and control of finite element solutions*, Finite Elem. Anal. Des. **3** (1987), 1–26.
- [Nyström 1930] E. J. Nyström, *Über die praktische Auflösung von Integral-Gleichungen mit Anwendungen auf Randwertaufgaben*, Acta Math. **54** (1930), 185–204.
- [Oden 1983] J. T Oden, *Finite Elements IV, Mathematical Aspects*, Prentice–Hall, 1983, {MR 86m:651001d}.
- [Oden and Brauchli 1971] J. T. Oden and J. Brauchli, *On the calculation of consistent stress distributions in finite element approximations*, Internat. J. Numer. Methods Engrg. **3** (1971), 317–325.
- [Oden and Demkowicz 1989] J. T. Oden and L. Demkowicz, *Advances in adaptive improvements: A survey of adaptive finite element methods in computational mechanics*, State-of-the-Art Surveys on Computational Mechanics, The American Society of Mechanical Engineers, 1989, pp. 441–467.
- [Oden and Reddy 1973] J. T. Oden and J. N. Reddy, *Note on an approximate method for computing consistent conjugate stresses in elastic finite elements*, Internat. J. Numer. Method Engrg. **6** (1973), 55–61.
- [Oganesyan, Rivkind and Rukhovetz 1973] L. A. Oganesyan, V. Ja. Rivkind and L. A. Rukhovetz, *Variation-difference methods for the solution of elliptic equations, Part I*; (in Russian), Proc. Sem Issue 5, Vilnius, 3–389, {MR 57#18146}.
- [Oganesyan and Rukhovetz 1969] L. A. Oganesyan and L. A. Rukhovetz, *Study of the rate of convergence of variational difference schemes for second order elliptic equations in a two-dimensional field with a smooth boundary*, U.S.S.R. Comput. Math. and Math. Phys. **9** (1969), 158–183, {MR 45#4665}.
- [Oganesyan and Rukhovetz 1979] L. A. Oganesyan and L. A. Rukhovetz, *Variational-difference Methods for the Solution of Elliptic Equations*; (in Russian), Izd. Akad. Nauk Armjanskoi SSR, Jerevan, 1979, {MR 82m:65105}.
- [O’Riordan and Stynes 1986] E. O’Riordan and M. Stynes, *An analysis of a superconvergence result for a singularly perturbed boundary value problem*, Math. Comp. **46** (1986), 81–92, {MR 87b:65107}.

- [Ortega and Rheinboldt 1970] J. M. Ortega and W. C. Rheinboldt, *Iterative Solution of Nonlinear Equations in Several Variables*, Academic Press, 1970, {MR 42#8686}.
- [Pehlivanov 1989] A. I. Pehlivanov, *Interior estimates of type superconvergence of the gradient in the finite element method*, C. R. Acad. Bulgare Sci. **42** (1989), 29–32, {MR 91e:65130}.
- [Pehlivanov, Lazarov, Carey and Chow 1992] A. I. Pehlivanov, R. D. Lazarov, G. F. Carey and S.-S. Chow, *Superconvergence analysis of approximate boundary-flux calculations*, Numer. Math. **63** (1992), 483–501.
- [Peng 1990] L. Peng, *A probabilistic algorithm for high accuracy finite elements*; (in Chinese), J. Southeast Univ. **20** (1990), 119–126, {MR 92h:65129}.
- [Powell 1981] M. J. D. Powell, *Approximation Theory and Methods*, Cambridge University Press, 1981, {MR 82f:41001}.
- [Prenter 1975] P. M. Prenter, *Splines and Variational Methods*, Wiley–Interscience 1975 (Wiley Classics Edition 1989), {MR 58#3287}.
- [Rachowicz and Oden 1989] W. Rachowicz and J. T. Oden, *On the accuracy and convergence of conjugate flux calculations*, Numer. Methods Partial Differential Equations **5** (1989), 143–156, {MR 90m:65202}.
- [Rannacher 1987] R. Rannacher, *Richardson extrapolation for a mixed finite element approximation of a plate bending problem*, Z. Angew. Math. Mech. **67** (1987), T381–T383, {MR 88g:65118}.
- [Rannacher 1991] R. Rannacher,  *$L_\infty$ -stability estimates and asymptotic error expansion for parabolic finite element equations*, Tagungsband “Extrapolation and defect correction”, Bonn. Math. Schrift. **228** (1991), 74–94, {MR 93i:65093}.
- [Rannacher and Scott 1982] R. Rannacher and R. Scott, *Some optimal error estimates for piecewise linear finite element approximations*, Math. Comp. **38** (1982), 437–445, {MR 83e:65180}.
- [Regińska 1986] T. Regińska, *Superconvergence of external approximation of eigenvalues of ordinary differential operators*, IMA J. Numer. Anal. **6** (1986), 309–323, {MR 89i:65083}.
- [Regińska 1987] T. Regińska, *The superconvergence effect in the finite element method for two-point boundary value problems*; (in Polish), Mat. Stos. **30** (1987), 97–111, {MR 89f:65085}.
- [Richter 1978] G. R. Richter, *Superconvergence of piecewise polynomial Galerkin approximations for Fredholm integral equations of the second kind*, Numer. Math. **31** (1978), 63–70, {MR 80a:65273}.
- [Rui 1992] H. X. Rui, *Superconvergence of finite element solutions to a class of Sobolev equations*; (in Chinese), Shandong Daxue Xuebao Ziran Kexue Ban **27** (1992), 301–308, {MR 93i:65094}.
- [Sanz-Serna and Abia 1984] J. M. Sanz-Serna and L. Abia, *Interpolation of the coefficients in nonlinear elliptic Galerkin procedures*, SIAM J. Numer. Anal. **21** (1984), 77–83, {MR 85c:65146}.

- [Saranen 1987] J. Saranen, *Local error estimates for some Petrov–Galerkin methods applied to strongly elliptic equations on curves*, Math. Comp. **48** (1987), 485–502, {MR 88m:65208}.
- [Schatz 1974] A. H. Schatz, *An observation concerning Ritz–Galerkin methods with indefinite bilinear forms*, Math. Comp. **28** (1974), 959–962, {MR 51#9526}.
- [Schatz 1980] A. H. Schatz, *A weak discrete maximum principle and stability of the finite element method in  $L^\infty$  on plane polygonal domains*, Math. Comp. **34** (1980), 77–91, {MR 81e:65063}.
- [Schatz 1990] A. H. Schatz, *An analysis of the finite element method for second order elliptic boundary value problems. An introduction*, Mathematical Theory of Finite Element and Boundary Element Methods (A. H. Schatz, V. Thomée and W. L. Wendland, eds.), Birkhäuser, 1990, pp. 9–133, {MR 92f:65004}.
- [Schatz 1994] A. H. Schatz, personal communication.
- [Schatz 1995] A. H. Schatz (to appear).
- [Schatz, Sloan and Wahlbin 1994] A. H. Schatz, I. H. Sloan and L. B. Wahlbin, *Superconvergence in finite element methods and meshes which are locally symmetric with respect to a point*, SIAM J. Numer. Anal. (to appear).
- [Schatz and Wahlbin 1977] A. H. Schatz and L. B. Wahlbin, *Interior maximum norm estimates for finite element methods*, Math. Comp. **31** (1977), 414–442, {MR 55#4748}.
- [Schatz and Wahlbin 1978] A. H. Schatz and L. B. Wahlbin, *Maximum norm estimates in the finite element method on plane polygonal domains, Part 1*, Math. Comp. **32** (1978), 73–109, {MR 58#19233a}.
- [Schatz and Wahlbin 1979] A. H. Schatz and L. B. Wahlbin, *Maximum norm estimates in the finite element method on plane polygonal domains, Part 2: Refinements*, Math. Comp. **33** (1979), 465–492, {MR 58#19233b}.
- [Schatz and Wahlbin 1981] A. H. Schatz and L. B. Wahlbin, *On a local asymptotic error estimate in finite elements and its use: numerical examples*, Advances in Computer Methods for Partial Differential Equations IV (R. Vichnevetsky and R. S. Stepleman, eds.), IMACS, 1981, pp. 14–17.
- [Schatz and Wahlbin 1982] A. H. Schatz and L. B. Wahlbin, *On the quasi-optimality in  $L_\infty$  of the  $\overset{\circ}{H}{}^1$ -projection into finite element spaces*, Math. Comp. **38** (1982), 1–21, {MR 82m:65106}.
- [Schatz and Wahlbin 1983] A. H. Schatz and L. B. Wahlbin, *On the finite element method for singularly perturbed reaction–diffusion problems in two and one dimensions*, Math. Comp. **40** (1983), 47–89, {MR 84c:65137}.
- [Schatz and Wahlbin 1994] A. H. Schatz and L. B. Wahlbin, *Interior maximum norm estimates for finite element methods Part II*, Math. Comp. (to appear).
- [Schatz and Wang 1994] A. H. Schatz and J. Wang, *Some new error estimates for the Ritz–Galerkin methods with minimal smoothness assumptions* (to appear).

- [Schellbach 1851] K. Schellbach, *Probleme der Variationsrechnung*, J. Reine Angew. Math. **41** (1851), 293–363.
- [Scott 1975] R. Scott, *Interpolated boundary conditions in the finite element method*, SIAM J. Numer. Anal. **12** (1975), 404–427, {MR 52#7162}.
- [Scott 1976] R. Scott, *Optimal  $L^\infty$  estimates for the finite element method on irregular meshes*, Math. Comp. **30** (1976), 681–697, {MR 55#9560}.
- [Scott and Zhang 1990] L. R. Scott and S. Zhang, *Finite element interpolation of nonsmooth functions satisfying boundary conditions*, Math. Comp. **54** (1990), 483–493, {MR 90j:65021}.
- [Shi 1992] J. Shi, *The extrapolation method for a class of nonsmooth Fredholm integral equations*, Systems Sci. Math. Sci. **5** (1992), 33–41, {MR 94d:45009}.
- [Shilov 1961] G. E. Shilov, *An Introduction to the Theory of Linear Spaces*, Prentice Hall, 1961, {MR 23#A3746}.
- [Sloan 1976] I. H. Sloan, *Improvement by iteration for compact operator equations*, Math. Comp. **30** (1976), 758–764, {MR 57#14435}.
- [Sloan 1992a] I. H. Sloan, *Error analysis for boundary integral methods*, Acta Numerica 1992 (A. Iserles, ed.), Cambridge University Press, 1992, pp. 287–339.
- [Sloan 1992b] I. H. Sloan, *Unconventional methods for boundary integral equations in the plane*, Proc. 1991 Dundee Conference on Numerical Analysis (D. F. Griffith and G. A. Watson, ed.), Longman Scientific and Technical, Harlow, 1992.
- [Sloan, Burn and Dattner 1975] I. H. Sloan, B. Burn and N. Dattner, *A new approach to the numerical solution of integral equations*, J. Comp. Physics **18** (1975), 92–105, {MR 53#1992}.
- [Sloan and Thomée 1985] I. H. Sloan and V. Thomée, *Superconvergence of the Galerkin iterates for integral equations of the second kind*, J. Integral Equations **9** (1985), 1–23, {MR 86j:65184}.
- [Sloan and Wendland 1989] I. H. Sloan and W. L. Wendland, *A quadrature based approach to improving the collocation method for splines of even degree*, Z. Anal. Anwendungen **8** (1989), 361–376, {MR 91a:65236}.
- [Squeff 1987] M. C. J. Squeff, *Superconvergence of mixed finite element methods for parabolic equations*, RAIRO Model. Math. Anal. Numer. **21** (1987), 327–352, {MR 88j:65207}.
- [Strang 1972] G. Strang, *Approximation in the finite element method*, Numer. Math. **19** (1972), 81–98, {MR 46#4677}.
- [Strang and Fix 1973] G. Strang and G. J. Fix, *An Analysis of the Finite Element Method*, Prentice Hall, 1973, {MR 56#1747}.
- [Stricklin 1966] J. A. Stricklin, *Computation of stress resultants from the element stiffness matrices*, AIAA Journal **4** (1966), 1095–1096.
- [Stricklin 1968] J. A. Stricklin, *Reply to F. V. Filho*, (= [Filho 1968]), AIAA Journal **6** (1968), 572.

- [Stynes and O'Riordan 1984] M. Stynes and E. O'Riordan, *A superconvergence result for a singularly perturbed boundary value problem*, Boole Press Conf. Ser. 6, pp. 309–319, {MR 86b:65084}.
- [Taylor and Zienkiewicz 1985] R. L. Taylor and O. C. Zienkiewicz, *A note on the “order of approximation”*, Internat. J. Solids and Structures **21** (1985), 793–797, {MR 88a:65093}.
- [Tessler, Riggs and Macy 1994] A. Tessler, H. R. Riggs and S. C. Macy, *A variational method for finite element stress recovery and error estimation*, Comput. Methods Appl. Mech. Engrg. **111** (1994), 369–382.
- [Thomée 1972] V. Thomée, *Spline approximation and difference schemes for the heat equation*, Mathematical Foundations of the Finite Element Method with Applications to Partial Differential Equations, Academic Press, 1972, pp. 711–746, {MR 49#11824}.
- [Thomée 1973] V. Thomée, *Convergence estimates for semi-discrete Galerkin methods for initial-value problems*, Lecture Notes in Mathematics 333, Springer, 1973, pp. 243–262, {MR 56#17147}.
- [Thomée 1977] V. Thomée, *High order local approximation to derivatives in the finite element method*, Math. Comp. **31** (1977), 652–660, {MR 55#11572}.
- [Thomée 1980a] V. Thomée, *Negative norm estimates and superconvergence for parabolic problems*, Math. Comp. **34** (1980), 93–113, {MR 81a:65092}.
- [Thomée 1980b] V. Thomée, *Galerkin-finite element methods for parabolic equations*, Acad. Sci. Fennica, Helsinki (1980), 943–952, {MR 81i:65100}.
- [Thomée 1984] V. Thomée, *Galerkin Finite Element Methods for Parabolic Problems*, Lecture Notes in Mathematics 1054, Springer, 1984, {MR 86k:65006}.
- [Thomée and Wendroff 1974] V. Thomée and B. Wendroff, *Convergence estimates for Galerkin method for variable coefficient initial value problems*, SIAM J. Numer. Anal. **11** (1974), 1059–1068, {MR 51#7309}.
- [Thomée and Westergren 1968] V. Thomée and B. Westergren, *Elliptic difference equations and interior regularity*, Numer. Math. **11** (1968), 196–210, {MR 36#7347}.
- [Thomée, Xu and Zhang 1989] V. Thomée, J.-C. Xu and N.-Y. Zhang, *Superconvergence of the gradient in piecewise linear finite element approximation to a parabolic problem*, SIAM J. Numer. Anal. **26** (1989), 553–573, {MR 90e:65165}.
- [Tong 1969] P. Tong, *Exact solutions of certain problems by finite-element method*, AIAA Journal **7** (1969), 178–180.
- [Tran 1993] T. Tran, *The K-operator and the qualocation method for strongly elliptic equations on smooth curves*, J. Integral Equations Appl. **5** (1993), 405–428.
- [Tran 1994] T. Tran, *The K-operator and the Galerkin method for strongly elliptic equations on smooth curves* (to appear).
- [Vacek 1976] J. Vacek, *Dual variational principle for an elliptic partial differential equation*, Apl. Mat. **21** (1976), 5–27, {MR 54#716}.

- [Volk 1986] W. Volk, *Globale Superkonvergenz bei der Lösung von linearen Randwertaufgaben*, Numer. Math. **48** (1986), 617–625, {MR 87i:65123}.
- [von Seggern 1981] R. von Seggern, *Ein Superkonvergenzresultat bei Anwendung der Methode der finiten Elemente auf lineare Integrodifferentialgleichungen*, Z. Angew. Math. Mech. **61** (1981), 320–321, {MR 84e:45009}.
- [Wahlbin 1984] L. B. Wahlbin, *On the sharpness of certain local estimates for  $\overset{\circ}{H}^1$  projections into finite element spaces: Influence of a reentrant corner*, Math. Comp. **42** (1984), 1–8, {MR 86b:65129}.
- [Wahlbin 1991] L. B. Wahlbin, *Local Behavior in Finite Element Methods*, Handbook of Numerical Analysis vol. II, Finite Element Methods (Part 1) (P. G. Ciarlet and J. L. Lions, eds.), Elsevier, 1991, pp. 355–522, {MR 92f:65001}.
- [Wahlbin 1992] L. B. Wahlbin, *On superconvergence up to boundaries in finite element methods: a counterexample*, SIAM J. Numer. Anal. **29** (1992), 937–946, {MR94d:65045}.
- [Wang 1985] H. Wang, *Superconvergence of finite element solutions and their derivatives for a class of quasilinear hyperbolic equations*; (in Chinese), Hunan Ann. Math. **5** (1985), {MR 88a:65135}.
- [Wang 1986] H. Wang, *Superconvergence of fully discretized finite element solutions and their derivatives for a class of quasilinear hyperbolic equations*; (in Chinese), Dongbei Shuxue **2** (1986), 205–214, {MR 88h:65217}.
- [Wang, J. 1991] J. Wang, *Superconvergence and extrapolation for mixed finite element methods on rectangular domains*, Math. Comp. **56** (1991), 477–503, {MR 91m:65280}.
- [Wendland 1990] W. L. Wendland, *Boundary element methods for elliptic problems*, Mathematical Theory of Finite Element and Boundary Element Methods (A. H. Schatz, V. Thomée and W. L. Wendland, eds.), Birkhäuser, 1990, pp. 219–276, {MR 92f:65004}.
- [Wheeler, J. A. 1973] J. A. Wheeler, *Simulation of heat transfer from a warm pipeline buried in permafrost*, in: Proceedings 74th National meeting of the American Institute of Chemical Engineers.
- [Wheeler, M. F. 1973] M. F. Wheeler, *An optimal  $L_\infty$  error estimate for Galerkin approximations to solutions of two point boundary value problems*, SIAM J. Numer. Anal. **10** (1973), 914–917, {MR 49#8399}.
- [Wheeler 1974] M. F. Wheeler, *Galerkin procedure for estimating the flux for two-point boundary value problems*, SIAM J. Numer. Anal. **11** (1974), 764–768, {MR 52#4644}.
- [Wheeler and Whiteman 1987a] M. F. Wheeler and J. R. Whiteman, *Superconvergent recovery of gradients on subdomains from piecewise linear finite-element approximations*, Numer. Methods Partial Differential Equations **3** (1987), 65–82, {MR 90h:65192}.
- [Wheeler and Whiteman 1987b] M. F. Wheeler and J. R. Whiteman, *Superconvergent recovery of gradients on subdomains from piecewise linear finite-element*

- approximations*, Numer. Methods Partial Differential Equations **3** (1987), 357–374, {MR 92a:65306}.
- [Whiteman, Beagles and Warby 1990] J. R. Whiteman, A. E. Beagles and M. K. Warby, *Theoretical and practical aspects of finite elements in the context of some problems of solid mechanics*, Jahresber. Deutsch. Math.–Verein **92** (1990), 77–88, {MR 91c:73084}.
- [Whiteman and Goodsell 1988] J. R. Whiteman and G. Goodsell, *Superconvergent recovery of stresses from finite element approximations on subdomains for planar problems of linear elasticity*, The Mathematics of Finite Elements and Applications VI (J. R. Whiteman, ed.), Academic Press, 1988, pp. 29–53, {MR 89h:65195}.
- [Whiteman and Goodsell 1989] J. R. Whiteman and G. Goodsell, *Some gradient superconvergence results in the finite element method*, Lecture Notes in Mathematics 1397, Springer, 1989, pp. 182–260, {MR 91a:65234}.
- [Whiteman and Goodsell 1991] J. R. Whiteman and G. Goodsell, *A survey of gradient superconvergence for finite element approximations to second order elliptic problems on triangular and tetrahedral meshes*, The Mathematics of Finite Elements and Applications VII (J. R. Whiteman, ed.), Academic Press, 1991, pp. 55–74, {MR 92d:65007}.
- [Wiberg and Abdulwahab 1992] N.–E. Wiberg and F. Abdulwahab, *An efficient postprocessing technique for stress problems based on superconvergent derivatives and equilibrium*, Numerical Methods in Engineering '92, Elsevier, 1992, pp. 25–32.
- [Wiberg and Abdulwahab 1993] N.–E. Wiberg and F. Abdulwahab, *Patch recovery based on superconvergent derivatives and equilibrium*, Internat. J. Numer. Methods Engrg. **36** (1993), 2703–2724, {MR 94c:73065}.
- [Wiberg and Li 1994] N.–E. Wiberg and X. D. Li, *Superconvergent patch recovery of finite-element solution and a posteriori  $L_2$  norm error estimate*, Comm. Numer. Methods Engrg. **10** (1994), 313–320.
- [Wielgosz 1982] C. Wielgosz, *Informations exactes données par des méthodes d'élément finis en mécanique*, J. Méc. Théor. Appl. **1** (1982), 323–329, {MR 85a:73061}.
- [Winther 1981] R. Winther, *A stable finite element method for initial-boundary value problems for first-order hyperbolic systems*, Math. Comp. **36** (1981), 65–86, {MR 81m:65181}.
- [Winther 1982] R. Winther, *A finite element method for a version of the Boussinesq equation*, SIAM J. Numer. Anal. **19** (1982), 561–570, {MR 83f:65184}.
- [Wu and Huang 1988] D. G. Wu and D. R. Huang, *Superconvergence of quadratic interpolation splines and quadratic spline finite element solutions*; (in Chinese), Numer. Math. J. Chinese Univ. **10** (1988), 224–229, {MR 90a:65032}.
- [Xu 1988] C. F. Xu, *The split iteration method at successive levels for solving finite element equations and its convergence rates*; (in Chinese), Math. Appl. **1** (1988), 59–66, {MR 90f:65212}.

- [Xu, J. 1985] J. Xu, *The error analysis and the improved algorithms for the infinite element method*, Proceedings of the 1984 Beijing Symposium on Differential Geometry and Differential Equations, Science Press, Beijing 1985, pp. 326–331, {MR 87b:65213}.
- [Xu, J. 1994] J. Xu, *Two-grid discretization techniques for linear and nonlinear partial differential equations*, SIAM J. Numer. Anal. (to appear).
- [Yang and Huang 1986] B. Yang and M. Y. Huang, *An asymptotic expression for the error of approximate solutions by the mixed finite element method*; (in Chinese), Lanzhou Daxue Xuebao **22** (1986), 5–9, {MR 88e:65151}.
- [Yang, Y. D. 1987] Y. D. Yang,  *$L_p$ -estimates and superconvergence of finite element approximations for eigenvalue problems*; (in Chinese), Numer. Math. J. Chinese Univ. **9** (1987), 83–87, {MR 88h:65197}.
- [Yang, Y. D. 1990] Y. D. Yang, *A theorem on superconvergence of shifts of finite element approximations for eigenvalue problems*; (in Chinese), J. Math. (Wuhan) **10** (1990), 229–234, {MR 91h:65166}.
- [Ženíšek 1990] A. Ženíšek, *Nonlinear Elliptic and Evolution Problems and Their Finite Element Approximation*, Academic Press, 1990, {MR 92c:65003}.
- [Zhang 1988] T. Zhang, *Superconvergence of a finite element method for a singular two-point boundary value problem*; (in Chinese), Numer. Math. J. Chinese Univ. **10** (1988), 185–188, {MR 90b:65162}.
- [Zhang, Z. 1992] Z. Zhang, *Arch beam models: finite element analysis and superconvergence*, Numer. Math. **61** (1992), 117–143, {MR 93b:65180}.
- [Zhou 1992] Q. H. Zhou, *A mathematical representation of the condensation method*, Natur. Sci. J. Xiangtan Univ. **14** (1992), 24–27, {MR 93g:65100}.
- [Zhu, J. Z. and Zienkiewicz 1990] J. Z. Zhu and O. C. Zienkiewicz, *Superconvergence recovery technique and a posteriori error estimates*, Internat. J. Numer. Methods. Engrg. **30** (1990), 1321–1339, {MR 91i:65141}.
- [Zhu 1981] Q.-D. Zhu, *A superconvergence result for the finite element method*; (in Chinese), Numer. Math. J. Chinese Univ. **3** (1981), 50–55, {MR 82f:65121}.
- [Zhu 1983a] Q.-D. Zhu, *Natural inner superconvergence for the finite element method*, Proceedings of the China–France Symposium on Finite Element Methods (Beijing 1982), Science Press, Beijing, and Gordon and Breach 1983, 935–960, {MR 85h:65253}.
- [Zhu 1983b] Q.-D. Zhu, *Uniform convergence estimates of derivatives for the finite element method*, Numer. Math. J. Chinese Univ. **5** (1983), 311–318, {MR 85k:65096}.
- [Zhu 1985] Q.-D. Zhu, *Uniform superconvergence estimates for the finite element method*; (in Chinese), Natur. Sci. J. Xiangtan Univ. (1985), 10–26, {MR 88h:65223}.
- [Zhu 1987] Q.-D. Zhu, *Some  $L_\infty$  estimates and interior superconvergence estimates for piecewise linear finite element approximations*; (in Chinese), Natur. Sci. J. Xiangtan Univ. (1987), 9–16, {MR 88g:65124}.

- [Zhu and Chen 1988] Q.-D. Zhu and H. S. Chen, *High accuracy error analysis of a linear finite element method for general boundary value problems*; (in Chinese), Natur. Sci. J. Xiangtan Univ. **10** (1988), 1–10, {MR 90j:65136}.
- [Zhu and Lin 1989] Q.-D. Zhu and Q. Lin, *The Hyperconvergence Theory of Finite Elements*; (in Chinese), Hunan Science and Technology Publishing House, Changsha, 1989, {MR 93j:65191}.
- [Zienkiewicz and Cheung 1967] O. C. Zienkiewicz and Y. K. Cheung, *The Finite Element Method in Structural and Continuum Mechanics*, McGraw-Hill, 1967, {MR 47#4518}.
- [Zienkiewicz, Taylor and Too 1971] O. C. Zienkiewicz, R. L. Taylor and J. M. Too, *Reduced integration techniques in general analysis of plates and shells*, Internat. J. Numer. Methods Engrg. **3** (1971), 275–290.
- [Zienkiewicz and Zhu 1987] O. C. Zienkiewicz and J. Z. Zhu, *A simple error estimator and adaptive procedure for practical engineering analysis*, Internat. J. Numer. Methods Engrg. **24** (1987), 337–357, {MR 87m:73055}.
- [Zienkiewicz and Zhu 1992a] O. C. Zienkiewicz and J. Z. Zhu, *The superconvergent patch recovery and a posteriori error estimates*, Internat. J. Numer. Methods Engrg. **33** (1992); Part 1, 1331–1364; Part 2, 1365–1382, {MR 93c:73098 and 73099}.
- [Zienkiewicz and Zhu 1992b] O. C. Zienkiewicz and J. Z. Zhu, *The superconvergent patch recovery (SPR) and adaptive finite element refinement*, Comput. Methods Appl. Mech. Engrg. **101** (1992), 207–224, {MR 93h:73042}.
- [Zlámal 1977] M. Zlámal, *Some superconvergence results in the finite element method*, Lecture Notes in Mathematics 606, Springer, 1977, pp. 353–362, {MR 58#8365}.
- [Zlámal 1978a] M. Zlámal, *Superconvergence and reduced integration in the finite element method*, Math. Comp. **32** (1978), 663–685, {MR 58#13794}.
- [Zlámal 1978b] M. Zlámal, *Superconvergence of gradients in the finite element method*; (in Russian), Akad. Nauk SSSR Sibirsk. Otdel. Vychisl. Tsentr (1978), 15–22, {MR 82e:65116}.

## **Subject Index.**

- Approximation theory 30
- Boundary element method 116
- Bramble–Hilbert lemma 37
- Computational domain 100
- Delta–function, discrete 10, 38
  - exponential decay in 10, 39
- Difference quotients 23, 87, 89, 101, 110
  - maximum–norm estimates via on translation invariant meshes 92
- Duality argument 31, 50
- Finite elements (cf. also "Meshes")
  - Hermite cubics 1, 16, 27
  - intermediate degree (incomplete) 134
  - isoparametric 98
  - Serendipity 134
  - smoohest splines 1
- Gauss points 81, 82, 133
- Hermite cubics 1, 16, 27
- Integral equation 25, 116, 121
- Inverse estimates 7, 28, 37
- Isoparametric elements 98
- $K$ –operator 107
- $L_2$ –projection 5, 36, 42
  - stability in  $L_p$  35, 39
  - superconvergence in 42
- Lagrange multiplier method 123
- Leibniz' rule, discrete 91
- Meshes (cf. also "Finite elements")
  - Chevron pattern 125, 133
  - criss–cross pattern 47, 125, 133
  - quasi–uniform 2
  - regular pattern 125, 133
  - symmetric w.r.t. a point 11, 44, 74
  - tensor product 43, 65, 96, 134
  - translation invariant 84, 98
  - uniform 12
  - Union Jack pattern 125, 133
- Minimal surface equation 95

- Monotone 95
- Multigrid 97
- Negative norm estimates
  - in elliptic projections 69
  - in  $L_2$ -projections 42
- Newton's method 96
- Outflow derivative 25, 121
- $p$ -Laplacian 95
- Physical domain 98
- Principal lattice points 126
- Quasiinterpolant 5
- Quasuniform meshes 2
- Smoothest splines 1
- Smoothing operator 32
- Superapproximation 5, 33, 36
- Superconvergence 3
  - nodal 3
- Symmetric meshes w.r.t. a point 11, 44, 74
- Tensor product elements 43, 65, 96, 134
- Translation invariant spaces 84, 98
  - maximum norm estimates via difference quotients 92
- Uniform meshes 12

- Vol. 1509: J. Aguadé, M. Castellet, F. R. Cohen (Eds.), Algebraic Topology - Homotopy and Group Cohomology. Proceedings, 1990. X, 330 pages. 1992.
- Vol. 1510: P. P. Kulish (Ed.), Quantum Groups. Proceedings, 1990. XII, 398 pages. 1992.
- Vol. 1511: B. S. Yadav, D. Singh (Eds.), Functional Analysis and Operator Theory. Proceedings, 1990. VIII, 223 pages. 1992.
- Vol. 1512: L. M. Adleman, M.-D. A. Huang, Primality Testing and Abelian Varieties Over Finite Fields. VII, 142 pages. 1992.
- Vol. 1513: L. S. Block, W. A. Coppel, Dynamics in One Dimension. VIII, 249 pages. 1992.
- Vol. 1514: U. Krengel, K. Richter, V. Warstat (Eds.), Ergodic Theory and Related Topics III, Proceedings, 1990. VIII, 236 pages. 1992.
- Vol. 1515: E. Ballico, F. Catanese, C. Ciliberto (Eds.), Classification of Irregular Varieties. Proceedings, 1990. VII, 149 pages. 1992.
- Vol. 1516: R. A. Lorentz, Multivariate Birkhoff Interpolation. IX, 192 pages. 1992.
- Vol. 1517: K. Keimel, W. Roth, Ordered Cones and Approximation. VI, 134 pages. 1992.
- Vol. 1518: H. Stichtenoth, M. A. Tsfasman (Eds.), Coding Theory and Algebraic Geometry. Proceedings, 1991. VIII, 223 pages. 1992.
- Vol. 1519: M. W. Short, The Primitive Soluble Permutation Groups of Degree less than 256. IX, 145 pages. 1992.
- Vol. 1520: Yu. G. Borisovich, Yu. E. Gliklikh (Eds.), Global Analysis – Studies and Applications V. VII, 284 pages. 1992.
- Vol. 1521: S. Busenberg, B. Forte, H. K. Kuiken, Mathematical Modelling of Industrial Process. Bari, 1990. Editors: V. Capasso, A. Fasano. VII, 162 pages. 1992.
- Vol. 1522: J.-M. Delort, F. B. I. Transformation. VII, 101 pages. 1992.
- Vol. 1523: W. Xue, Rings with Morita Duality. X, 168 pages. 1992.
- Vol. 1524: M. Coste, L. Mahé, M.-F. Roy (Eds.), Real Algebraic Geometry. Proceedings, 1991. VIII, 418 pages. 1992.
- Vol. 1525: C. Casacuberta, M. Castellet (Eds.), Mathematical Research Today and Tomorrow. VII, 112 pages. 1992.
- Vol. 1526: J. Azéma, P. A. Meyer, M. Yor (Eds.), Séminaire de Probabilités XXVI. X, 633 pages. 1992.
- Vol. 1527: M. I. Freidlin, J.-F. Le Gall, Ecole d'Eté de Probabilités de Saint-Flour XX – 1990. Editor: P. L. Hennequin. VIII, 244 pages. 1992.
- Vol. 1528: G. Isac, Complementarity Problems. VI, 297 pages. 1992.
- Vol. 1529: J. van Neerven, The Adjoint of a Semigroup of Linear Operators. X, 195 pages. 1992.
- Vol. 1530: J. G. Heywood, K. Masuda, R. Rautmann, S. A. Solonnikov (Eds.), The Navier-Stokes Equations II – Theory and Numerical Methods. IX, 322 pages. 1992.
- Vol. 1531: M. Stoer, Design of Survivable Networks. IV, 206 pages. 1992.
- Vol. 1532: J. F. Colombeau, Multiplication of Distributions. X, 184 pages. 1992.
- Vol. 1533: P. Jipsen, H. Rose, Varieties of Lattices. X, 162 pages. 1992.
- Vol. 1534: C. Greither, Cyclic Galois Extensions of Commutative Rings. X, 145 pages. 1992.
- Vol. 1535: A. B. Evans, Orthomorphism Graphs of Groups. VIII, 114 pages. 1992.
- Vol. 1536: M. K. Kwong, A. Zettl, Norm Inequalities for Derivatives and Differences. VII, 150 pages. 1992.
- Vol. 1537: P. Fitzpatrick, M. Martelli, J. Mawhin, R. Nussbaum, Topological Methods for Ordinary Differential Equations. Montecatini Terme, 1991. Editors: M. Furi, P. Zecca. VII, 218 pages. 1993.
- Vol. 1538: P.-A. Meyer, Quantum Probability for Probabilists. X, 287 pages. 1993.
- Vol. 1539: M. Coornaert, A. Papadopoulos, Symbolic Dynamics and Hyperbolic Groups. VIII, 138 pages. 1993.
- Vol. 1540: H. Komatsu (Ed.), Functional Analysis and Related Topics, 1991. Proceedings. XXI, 413 pages. 1993.
- Vol. 1541: D. A. Dawson, B. Maisonneuve, J. Spencer, Ecole d'Eté de Probabilités de Saint-Flour XXI - 1991. Editor: P. L. Hennequin. VIII, 356 pages. 1993.
- Vol. 1542: J. Fröhlich, Th. Kerler, Quantum Groups, Quantum Categories and Quantum Field Theory. VII, 431 pages. 1993.
- Vol. 1543: A. L. Dontchev, T. Zolezzi, Well-Posed Optimization Problems. XII, 421 pages. 1993.
- Vol. 1544: M. Schürmann, White Noise on Bialgebras. VII, 146 pages. 1993.
- Vol. 1545: J. Morgan, K. O'Grady, Differential Topology of Complex Surfaces. VIII, 224 pages. 1993.
- Vol. 1546: V. V. Kalashnikov, V. M. Zolotarev (Eds.), Stability Problems for Stochastic Models. Proceedings, 1991. VIII, 229 pages. 1993.
- Vol. 1547: P. Harmand, D. Werner, W. Werner, M-ideals in Banach Spaces and Banach Algebras. VIII, 387 pages. 1993.
- Vol. 1548: T. Urabe, Dynkin Graphs and Quadrilateral Singularities. VI, 233 pages. 1993.
- Vol. 1549: G. Vainikko, Multidimensional Weakly Singular Integral Equations. XI, 159 pages. 1993.
- Vol. 1550: A. A. Gonchar, E. B. Saff (Eds.), Methods of Approximation Theory in Complex Analysis and Mathematical Physics IV, 222 pages. 1993.
- Vol. 1551: L. Arkeryd, P. L. Lions, P.A. Markowich, S.R. S. Varadhan. Nonequilibrium Problems in Many-Particle Systems. Montecatini, 1992. Editors: C. Cercignani, M. Pulvirenti. VII, 158 pages 1993.
- Vol. 1552: J. Hilgert, K.-H. Neeb, Lie Semigroups and their Applications. XII, 315 pages. 1993.
- Vol. 1553: J.-L. Colliot-Thélène, J. Kato, P. Vojta. Arithmetic Algebraic Geometry. Trento, 1991. Editor: E. Ballico. VII, 223 pages. 1993.
- Vol. 1554: A. K. Lenstra, H. W. Lenstra, Jr. (Eds.), The Development of the Number Field Sieve. VIII, 131 pages. 1993.
- Vol. 1555: O. Liess, Conical Refraction and Higher Microlocalization. X, 389 pages. 1993.
- Vol. 1556: S. B. Kuksin, Nearly Integrable Infinite-Dimensional Hamiltonian Systems. XXVII, 101 pages. 1993.

- Vol. 1557: J. Azéma, P. A. Meyer, M. Yor (Eds.), Séminaire de Probabilités XXVII. VI, 327 pages. 1993.
- Vol. 1558: T. J. Bridges, J. E. Furter, Singularity Theory and Equivariant Symplectic Maps. VI, 226 pages. 1993.
- Vol. 1559: V. G. Sprindžuk, Classical Diophantine Equations. XII, 228 pages. 1993.
- Vol. 1560: T. Bartsch, Topological Methods for Variational Problems with Symmetries. X, 152 pages. 1993.
- Vol. 1561: I. S. Molchanov, Limit Theorems for Unions of Random Closed Sets. X, 157 pages. 1993.
- Vol. 1562: G. Harder, Eisensteinkohomologie und die Konstruktion gemischter Motive. XX, 184 pages. 1993.
- Vol. 1563: E. Fabes, M. Fukushima, L. Gross, C. Kenig, M. Röckner, D. W. Stroock, Dirichlet Forms. Varenna, 1992. Editors: G. Dell'Antonio, U. Mosco. VII, 245 pages. 1993.
- Vol. 1564: J. Jorgenson, S. Lang, Basic Analysis of Regularized Series and Products. IX, 122 pages. 1993.
- Vol. 1565: L. Boutet de Monvel, C. De Concini, C. Procesi, P. Schapira, M. Vergne. D-modules, Representation Theory, and Quantum Groups. Venezia, 1992. Editors: G. Zampieri, A. D'Agnolo. VII, 217 pages. 1993.
- Vol. 1566: B. Edixhoven, J.-H. Evertse (Eds.), Diophantine Approximation and Abelian Varieties. XIII, 127 pages. 1993.
- Vol. 1567: R. L. Dobrushin, S. Kusuoka, Statistical Mechanics and Fractals. VII, 98 pages. 1993.
- Vol. 1568: F. Weisz, Martingale Hardy Spaces and their Application in Fourier Analysis. VIII, 217 pages. 1994.
- Vol. 1569: V. Totik, Weighted Approximation with Varying Weight. VI, 117 pages. 1994.
- Vol. 1570: R. deLaubenfels, Existence Families, Functional Calculi and Evolution Equations. XV, 234 pages. 1994.
- Vol. 1571: S. Yu. Pilyugin, The Space of Dynamical Systems with the  $C^0$ -Topology. X, 188 pages. 1994.
- Vol. 1572: L. Götsche, Hilbert Schemes of Zero-Dimensional Subschemes of Smooth Varieties. IX, 196 pages. 1994.
- Vol. 1573: V. P. Havin, N. K. Nikolski (Eds.), Linear and Complex Analysis – Problem Book 3 – Part I. XXII, 489 pages. 1994.
- Vol. 1574: V. P. Havin, N. K. Nikolski (Eds.), Linear and Complex Analysis – Problem Book 3 – Part II. XXII, 507 pages. 1994.
- Vol. 1575: M. Mitrea, Clifford Wavelets, Singular Integrals, and Hardy Spaces. XI, 116 pages. 1994.
- Vol. 1576: K. Kitahara, Spaces of Approximating Functions with Haar-Like Conditions. X, 110 pages. 1994.
- Vol. 1577: N. Obata, White Noise Calculus and Fock Space. X, 183 pages. 1994.
- Vol. 1578: J. Bernstein, V. Lunts, Equivariant Sheaves and Functors. V, 139 pages. 1994.
- Vol. 1579: N. Kazamaki, Continuous Exponential Martingales and  $BMO$ . VII, 91 pages. 1994.
- Vol. 1580: M. Milman, Extrapolation and Optimal Decompositions with Applications to Analysis. XI, 161 pages. 1994.
- Vol. 1581: D. Bakry, R. D. Gill, S. A. Molchanov, Lectures on Probability Theory. Editor: P. Bernard. VIII, 420 pages. 1994.
- Vol. 1582: W. Balser, From Divergent Power Series to Analytic Functions. X, 108 pages. 1994.
- Vol. 1583: J. Azéma, P. A. Meyer, M. Yor (Eds.), Séminaire de Probabilités XXVIII. VI, 334 pages. 1994.
- Vol. 1584: M. Brokate, N. Kenmochi, I. Müller, J. F. Rodriguez, C. Verdi, Phase Transitions and Hysteresis. Montecatini Terme, 1993. Editor: A. Visintin. VII, 291 pages. 1994.
- Vol. 1585: G. Frey (Ed.), On Artin's Conjecture for Odd 2-dimensional Representations. VIII, 148 pages. 1994.
- Vol. 1586: R. Nillsen, Difference Spaces and Invariant Linear Forms. XII, 186 pages. 1994.
- Vol. 1587: N. Xi, Representations of Affine Hecke Algebras. VIII, 137 pages. 1994.
- Vol. 1588: C. Scheiderer, Real and Étale Cohomology. XXIV, 273 pages. 1994.
- Vol. 1589: J. Bellissard, M. Degli Esposti, G. Forni, S. Graffi, S. Isola, J. N. Mather, Transition to Chaos in Classical and Quantum Mechanics. Montecatini, 1991. Editor: S. Graffi. VII, 192 pages. 1994.
- Vol. 1590: P. M. Soardi, Potential Theory on Infinite Networks. VIII, 187 pages. 1994.
- Vol. 1591: M. Abate, G. Patrizio, Finsler Metrics – A Global Approach. IX, 180 pages. 1994.
- Vol. 1592: K. W. Breitung, Asymptotic Approximations for Probability Integrals. IX, 146 pages. 1994.
- Vol. 1593: J. Jorgenson & S. Lang, D. Goldfeld, Explicit Formulas for Regularized Products and Series. VIII, 154 pages. 1994.
- Vol. 1594: M. Green, J. Murre, C. Voisin, Algebraic Cycles and Hodge Theory. Torino, 1993. Editors: A. Albano, F. Bardelli. VII, 275 pages. 1994.
- Vol. 1595: R.D.M. Accola, Topics in the Theory of Riemann Surfaces. IX, 105 pages. 1994.
- Vol. 1596: L. Heindorf, L. B. Shapiro, Nearly Projective Boolean Algebras. X, 202 pages. 1994.
- Vol. 1597: B. Herzog, Kodaira-Spencer Maps in Local Algebra. XVII, 176 pages. 1994.
- Vol. 1598: J. Berndt, F. Tricerri, L. Vanhecke, Generalized Heisenberg Groups and Damek-Ricci Harmonic Spaces. VIII, 125 pages. 1995.
- Vol. 1599: K. Johannson, Topology and Combinatorics of 3-Manifolds. XVIII, 446 pages. 1995.
- Vol. 1600: W. Narkiewicz, Polynomial Mappings. VII, 130 pages. 1995.
- Vol. 1601: A. Pott, Finite Geometry and Character Theory. VII, 181 pages. 1995.
- Vol. 1602: J. Winkelmann, The Classification of Three-dimensional Homogeneous Complex Manifolds. XI, 230 pages. 1995.
- Vol. 1603: V. Ene, Real Functions. Current Topics. XIII, 310 pages. 1995.
- Vol. 1604: A. Huber, Mixed Motives and their Realization in Derived Categories. XV, 207 pages. 1995.
- Vol. 1605: L. B. Wahlbin, Superconvergence in Galerkin Finite Element Methods. XI, 166 pages. 1995.