

Notes for the talk

Lagrangian fibrations on Nikulin orbifolds

Giacomo Nanni

giacomonanni.github.io

Università di Bologna & Ruhr-Universität Bochum

2 October 2026

These are notes for the talk and may contain inaccuracies.
The content follows my handwritten notes; the $\text{\LaTeX}$ source was mainly written by an AI tool (Claude Opus 5.5).

Lagrangian fibrations on Nikulin orbifolds

Theorem (Nanni 2025)

There are exactly 2 deformation families of Lagrangian fibrations on Nikulin-type orbifolds.

Corollary

The SYZ conjecture holds for Nikulin-type orbifolds.

Outline

- 1 IHS manifolds and the SYZ conjecture
- 2 Nikulin orbifolds and their Lagrangian fibrations
- 3 Deforming Lagrangian fibrations
- 4 Monodromy orbits and classification
- 5 References

K3 surfaces

Definition

A **K3 surface** is a compact complex Kähler surface S such that

- $\pi_1(S) = 0$;
- $H^0(S, \Omega_S^2) = \mathbb{C}\sigma$, with σ symplectic.

Example: $V(f_4) \subset \mathbb{P}^3$, a smooth quartic. **Fact:** $(H^2(S, \mathbb{Z}), \cap) \stackrel{\varphi}{\cong} \Lambda$ (marking).

Theorem (Torelli; Pyatetskii-Shapiro–Shafarevich 1971, Burns–Rapoport 1975)

There is a moduli space $\mathcal{M}_\Lambda$ of marked K3 surfaces, and the period map

$$\begin{array}{ccc} \mathcal{P}: & \mathcal{M}_\Lambda & \longrightarrow D \subset \mathbb{P}(\Lambda_{\mathbb{C}}) \\ & (S, \varphi) & \longmapsto \varphi[\mathbb{C}\sigma] \end{array}$$

induces an isomorphism on each connected component of the Hausdorff reduction of $\mathcal{M}_\Lambda$.

$$D = \{x : x^2 = 0, \ x\bar{x} > 0\}.$$

Irreducible holomorphic symplectic manifolds

Definition

An **irreducible holomorphic symplectic (IHS) manifold** is a compact complex Kähler manifold X such that

- $\pi_1(X) = 0$;
- $H^0(X, \Omega_X^2) = \mathbb{C}\sigma$, with σ symplectic.

Def/Prop (Beauville 1983, Fujiki 1987). The **Beauville–Bogomolov–Fujiki form** is the “unique” quadratic form q_X on $H^2(X, \mathbb{Z})$ such that there is $c_X \in \mathbb{R}_{>0}$ with

$$c_X q_X(\alpha)^n = \int_X \alpha^{2n} \quad \forall \alpha \in H^2(X, \mathbb{Z}), \quad 2n = \dim_{\mathbb{C}} X.$$

Theorem (Verbitsky 2013; Huybrechts 2012)

Fix a lattice Λ . The period map $\mathcal{P}: \mathcal{M}_\Lambda \rightarrow D \subset \mathbb{P}(\Lambda_{\mathbb{C}})$ of marked IHS manifolds of lattice Λ (defined as for K3 surfaces) induces an isomorphism on each connected component of the Hausdorff reduction of $\mathcal{M}_\Lambda$.

Fibrations of IHS manifolds

Theorem (Matsushita 1999)

A fibration $\psi: X \rightarrow B$ (surjective, connected fibres, $0 < \dim B < 2n$) satisfies:

- $\dim B = n$;
- $\rho(B) = 1$: $\text{Pic}(B)_{\mathbb{Q}} = \mathbb{Q}H$, H ample;
- the general fibre is a Lagrangian abelian variety.

$\ell = \psi^* c_1(H)$ is **algebraic, nef, isotropic**:

$$c_X q_X(\ell)^n = \int_X \psi^* c_1(H)^{2n} = \int_B H^{2n} = 0 \quad \Rightarrow \quad q_X(\ell) = 0.$$

Corollary

The very general IHS does **not** admit a fibration. Idea: $\sigma \in \bigcup_{\ell^2=0} \ell^\perp$, $\ell \in \Lambda$ (countable union).

The SYZ conjecture

Conjecture (SYZ; Verbitsky 2010)

If X admits an algebraic nef isotropic class ℓ , X admits a Lagrangian fibration with class a multiple of ℓ .

SYZ = Strominger–Yau–Zaslow (*Mirror symmetry is T-duality*, 1996)

Known: $K3^{[n]}$ [Bayer–Macrì 2014], Kum_n [Yoshioka 2016], OG6 [Mongardi–Rapagnetta 2021], OG10 [Mongardi–Onorati 2022].

Examples of IHS [Beauville 1983, Fujiki 1983]

$$\begin{array}{ccc} S \times S & & \text{too many 2-forms: } \lambda_1 \sigma \boxplus \lambda_2 \sigma \\ \downarrow / \mathfrak{S}_2 & & \downarrow \text{wavy} \\ S^{[2]} \xrightarrow{\text{Bl}} S^{(2)} & & \text{only } \lambda(\sigma \boxplus \sigma) \text{ descends} \end{array}$$

Symplectic orbifolds

Definition

An **orbifold** is a complex space locally isomorphic to quotients V/G , $V \subseteq \mathbb{C}^{2n}$ open, G a finite group.

Definition

An **irreducible symplectic orbifold** is a compact complex Kähler orbifold Y such that

- $\pi_1(Y_{\text{reg}}) = 0$;
- $H^0(Y, \Omega^2) = \mathbb{C}\sigma$, σ symplectic;
- $\text{codim Sing } Y > 2$.

Torelli theorem [Menet 2020]

Lagrangian fibrations [Schwald 2020]

Nikulin orbifolds

Let S be a K3 surface and ι a symplectic involution on S .

$$\begin{array}{ccccc}
 & & S^{[2]} & & \iota^{[2]} \text{ symplectic involution} \\
 & & \downarrow & & \\
 \text{Sing } Y: 28 \text{ points} & Y \xrightarrow{\text{Bl}_\Sigma} & S^{[2]}/\iota^{[2]} & \text{Sing}(S^{[2]}/\iota^{[2]}) : & \begin{cases} \Sigma \text{ K3 surface} \\ 28 = \binom{8}{2} \text{ points} \end{cases}
 \end{array}$$

8 fixed points of ι on S [Nikulin 1979]; general case [Mongardi 2012]

Definition

Such Y are called **Nikulin orbifolds**; their (locally trivial) deformations are of **Nikulin type**.

BBF lattice: $H^2(Y, \mathbb{Z}) \cong U(2)^3 \oplus E_8(-1) \oplus \langle -2 \rangle^2$, $c_Y = 6$ [Menet 2015].

Examples of fibrations

$\varepsilon: S \rightarrow \mathbb{P}^1$ elliptic, ι_+, ι_- symplectic involutions (e.g. on the Fermat quartic)

$$\varepsilon \circ \iota_+ = \varepsilon$$

$$\begin{array}{ccc} S^{[2]} & \longrightarrow & \mathbb{P}^2 \\ \downarrow & \nearrow & \\ Y_+ & \longrightarrow & S^{[2]}/\iota_+ \end{array}$$

fibre of type (1,2) [Barth 1987]; $\text{div } \ell = 1$

$$\varepsilon \circ \iota_- = \tau \circ \varepsilon$$

$$\begin{array}{ccc} S^{[2]} & \longrightarrow & \mathbb{P}^2 \\ \downarrow & & \downarrow \\ Y_- & \longrightarrow & S^{[2]}/\iota_- \longrightarrow \mathbb{P}^2/\tau \end{array}$$

fibre of type (1,1) [Wieneck 2016]; $\text{div } \ell = 2$

Different invariants $\Rightarrow$ not deformation equivalent.

Deforming: monodromy

A Lagrangian fibration $\psi: X \rightarrow B$ is the same as a semiample (isotropic) line bundle L on X .

Take a deformation (locally trivial) of X :

$$\begin{array}{ccc} X & \hookrightarrow & \mathcal{X} \\ \downarrow & & \downarrow \\ 0 & \in & T \end{array}$$

Definition

A **deformation** of ψ is such a deformation together with a line bundle $\mathcal{L}$ on $\mathcal{X}$ with $\mathcal{L}|_X = L$, such that $L_t := \mathcal{L}|_{X_t}$ is semiample for all $t \in T$.

Introduce markings. Assume $\pi_1(T) = 0$ and fix a marking $\varphi: H^2(X, \mathbb{Z}) \rightarrow \Lambda$.

- For $t \in T$ choose a path γ_t from 0 to t : parallel transport $P_{\gamma_t}: H^2(X, \mathbb{Z}) \rightarrow H^2(X_t, \mathbb{Z})$, and set $\varphi_t := \varphi \circ P_{\gamma_t}^{-1}$.
- φ_t does not depend on γ_t : $\varphi_t^{-1} \circ \varphi'_t = P_{\gamma_t} P_{\gamma'_t}^{-1} = P_{\gamma_t(\gamma'_t)^{-1}} = \text{id}$, since $\gamma_t(\gamma'_t)^{-1} \in \pi_1(T) = 0$.

Deforming along paths

Remark. If $(\mathcal{X}, \mathcal{L})$ deforms (X, L) , then $\varphi_t(c_1(L_t)) = \ell \in \Lambda$
 $\forall t \in T \rightsquigarrow \mathcal{P}(T) \subseteq \ell^\perp$.


Theorem (Matsushita 2016;
Engel–Filipazzi–Greer–Mauri–Svaldi 2025)

A Lagrangian fibration (X, L) deforms over the locus where ℓ stays algebraic $(\mathcal{P}(T) \cap \ell^\perp)$, for T a small neighbourhood of 0.

What if $\pi_1(T) \neq 0$? Then the choice of γ_t matters! So fix a γ_t .

Theorem (smooth: Markman 2014, Matsushita 2017; singular: Onorati–Ríos Ortiz 2025)


If ℓ stays algebraic and nef along γ_t , (X, L) deforms along γ_t . Moreover, if ℓ is algebraic and nef at the extremes, then there is a deformation of (X, L) containing X_t .



The monodromy group

How to reach other components?

We still have freedom on the marking.



Definition (Markman 2011)

The **monodromy group** $\text{Mon}^2(X)$ of X is the group of parallel transport operators P_γ , $\gamma \in \pi_1(T)$, for some deformation $\mathcal{X} \rightarrow T$.

Fact. If (X, φ_1) , (X, φ_2) are in the same connected component of $\mathcal{M}_\Lambda$, then

$$\varphi_2^{-1} \circ \varphi_1 \in \text{Mon}^2(X).$$

$\Rightarrow$ the class of a fibration can be moved anywhere in its Mon^2 -orbit.

How many orbits?

Isotropic classes up to monodromy: at most 3 orbits [Menet–Riess 2022]

Theorem (Brandhorst–Menet–Muller 2024; independently Nanni 2025)

Mon²(Y) is maximal: $\text{Mon}^2(Y) = O^+(H^2(Y, \mathbb{Z}))$.

Corollary

Exactly 2 orbits of primitive isotropic classes, distinguished by the divisibility (1 or 2).

Both are realised: Y_+ (div = 1) and Y_- (div = 2).

Theorem (Nanni 2025)

There are 2 deformation families of Lagrangian fibrations on Nikulin-type orbifolds.

Sketch of proof

- Take any (Y, L) . Take $(\hat{Y}, \hat{L})$ the one example with the same invariants (div of the primitive class).
- Fix a marking φ on Y , deform it to a marking $\hat{\varphi}$ on $\hat{Y}$.
- $\exists P_\gamma \in \text{Mon}^2(\hat{Y})$ such that $\hat{\varphi} \circ P_\gamma(\hat{L}) = \varphi(L)$.
- $\Rightarrow$ by [Onorati–Ríos Ortiz 2025], (Y, L) and $(\hat{Y}, \hat{L})$ are deformation equivalent as fibrations.

Corollary (of the proof)

SYZ: start from a nef isotropic algebraic class instead of L ; the same argument shows it is semiample.

References I

- **W. Barth** (1987). *Abelian surfaces with $(1, 2)$ -polarization*. Adv. Stud. Pure Math. 10, 41–84. [doi:10.2969/aspm/01010041](https://doi.org/10.2969/aspm/01010041)
- **A. Bayer, E. Macrì** (2014). *MMP for moduli of sheaves on K3s via wall-crossing: nef and movable cones, Lagrangian fibrations*. Invent. Math. 198, 505–590. [doi:10.1007/s00222-014-0501-8](https://doi.org/10.1007/s00222-014-0501-8)
- **A. Beauville** (1983). *Variétés kählériennes dont la première classe de Chern est nulle*. J. Differential Geom. 18, 755–782. [doi:10.4310/jdg/1214438181](https://doi.org/10.4310/jdg/1214438181)
- **S. Brandhorst, G. Menet, S. Müller** (2024). *Automorphisms of Nikulin-type orbifolds*. [arXiv:2411.04668](https://arxiv.org/abs/2411.04668), to appear in Rev. Mat. Iberoam.
- **D. Burns, M. Rapoport** (1975). *On the Torelli problem for kählerian K3 surfaces*. Ann. Sci. ÉNS 8, 235–273. [doi:10.24033/asens.1287](https://doi.org/10.24033/asens.1287)
- **P. Engel, S. Filipazzi, F. Greer, M. Mauri, R. Svaldi** (2025). *Boundedness of some fibered K-trivial varieties*. [arXiv:2507.00973](https://arxiv.org/abs/2507.00973)
- **A. Fujiki** (1983). *On primitively symplectic compact Kähler V-manifolds of dimension four*. Progr. Math. 39, 71–250
- **A. Fujiki** (1987). *On the de Rham cohomology group of a compact Kähler symplectic manifold*. Adv. Stud. Pure Math. 10, 105–165. [doi:10.2969/aspm/01010105](https://doi.org/10.2969/aspm/01010105)
- **D. Huybrechts** (2012). *A global Torelli theorem for hyperkähler manifolds [after M. Verbitsky]*. Séminaire Bourbaki, Exp. 1040, Astérisque 348, 375–403. [arXiv:1106.5573](https://arxiv.org/abs/1106.5573)

References II

- E. Markman (2011). *A survey of Torelli and monodromy results for holomorphic-symplectic varieties*. Springer Proc. Math. 8, 257–322. [doi:10.1007/978-3-642-20300-8_15](https://doi.org/10.1007/978-3-642-20300-8_15)
- E. Markman (2014). *Lagrangian fibrations of holomorphic-symplectic varieties of $K3^{[n]}$ -type*. Springer Proc. Math. Stat. 71, 241–283. [doi:10.1007/978-3-319-05404-9_10](https://doi.org/10.1007/978-3-319-05404-9_10)
- D. Matsushita (1999). *On fibre space structures of a projective irreducible symplectic manifold*. Topology 38, 79–83. [doi:10.1016/S0040-9383\(98\)00003-2](https://doi.org/10.1016/S0040-9383(98)00003-2)
- D. Matsushita (2016). *On deformations of Lagrangian fibrations*. Progr. Math. 315, 237–243. [doi:10.1007/978-3-319-29959-4_9](https://doi.org/10.1007/978-3-319-29959-4_9)
- D. Matsushita (2017). *On isotropic divisors on irreducible symplectic manifolds*. Adv. Stud. Pure Math. 74, 291–312. [doi:10.2969/aspm/07410291](https://doi.org/10.2969/aspm/07410291)
- G. Menet (2015). *Beauville–Bogomolov lattice for a singular symplectic variety of dimension 4*. J. Pure Appl. Algebra 219, 1455–1495. [doi:10.1016/j.jpaa.2014.06.012](https://doi.org/10.1016/j.jpaa.2014.06.012)
- G. Menet (2020). *Global Torelli theorem for irreducible symplectic orbifolds*. J. Math. Pures Appl. 137, 213–237. [doi:10.1016/j.matpur.2020.03.010](https://doi.org/10.1016/j.matpur.2020.03.010)
- G. Menet, U. Riess (2022). *Wall divisors on irreducible symplectic orbifolds of Nikulin-type*. [arXiv:2209.03667](https://arxiv.org/abs/2209.03667)
- G. Mongardi (2012). *Symplectic involutions on deformations of $K3^{[2]}$* . Cent. Eur. J. Math. 10, 1472–1485. [doi:10.2478/s11533-012-0073-z](https://doi.org/10.2478/s11533-012-0073-z)

References III

- **G. Mongardi, C. Onorati** (2022). *Birational geometry of irreducible holomorphic symplectic tenfolds of O'Grady type*. Math. Z. 300, 3497–3526. [doi:10.1007/s00209-021-02966-6](https://doi.org/10.1007/s00209-021-02966-6)
- **G. Mongardi, A. Rapagnetta** (2021). *Monodromy and birational geometry of O'Grady's sixfolds*. J. Math. Pures Appl. 146, 31–68. [doi:10.1016/j.matpur.2020.12.006](https://doi.org/10.1016/j.matpur.2020.12.006)
- **G. Nanni** (2025). *Monodromy of Nikulin orbifolds*. [arXiv:2503.14441](https://arxiv.org/abs/2503.14441), to appear in Kyoto J. Math.
- **G. Nanni** (2025). *Lagrangian fibrations on Nikulin-type orbifolds*. [arXiv:2512.19244](https://arxiv.org/abs/2512.19244)
- **V. V. Nikulin** (1979). *Finite groups of automorphisms of Kählerian K3 surfaces*. Trudy Moskov. Mat. Obshch. 38, 75–137
- **C. Onorati, Á. D. Ríos Ortiz** (2025). *The SYZ conjecture for singular moduli spaces of sheaves on K3 surfaces*. [arXiv:2510.01005](https://arxiv.org/abs/2510.01005)
- **I. I. Pyatetskii-Shapiro, I. R. Shafarevich** (1971). *A Torelli theorem for algebraic surfaces of type K3*. Izv. Akad. Nauk SSSR Ser. Mat. 35, 530–572; Engl. transl. Math. USSR-Izv. 5, 547–588. [doi:10.1070/IM1971v005n03ABEH001075](https://doi.org/10.1070/IM1971v005n03ABEH001075)
- **M. Schwald** (2020). *Fujiki relations and fibrations of irreducible symplectic varieties*. Épijournal Géom. Algébrique 4, Art. 7. [doi:10.46298/epiga.2020.volume4.4557](https://doi.org/10.46298/epiga.2020.volume4.4557)
- **A. Strominger, S.-T. Yau, E. Zaslow** (1996). *Mirror symmetry is T-duality*. Nuclear Phys. B 479, 243–259. [doi:10.1016/0550-3213\(96\)00434-8](https://doi.org/10.1016/0550-3213(96)00434-8)

- **M. Verbitsky** (2010). *Hyperkähler SYZ conjecture and semipositive line bundles*. Geom. Funct. Anal. 19, 1481–1493. [doi:10.1007/s00039-009-0037-z](https://doi.org/10.1007/s00039-009-0037-z)
- **M. Verbitsky** (2013). *Mapping class group and a global Torelli theorem for hyperkähler manifolds*. Duke Math. J. 162, 2929–2986. [doi:10.1215/00127094-2382680](https://doi.org/10.1215/00127094-2382680), [arXiv:0908.4121](https://arxiv.org/abs/0908.4121)
- **B. Wieneck** (2016). *On polarization types of Lagrangian fibrations*. Manuscripta Math. 151, 305–327. [doi:10.1007/s00229-016-0845-z](https://doi.org/10.1007/s00229-016-0845-z)
- **K. Yoshioka** (2016). *Bridgeland's stability and the positive cone of the moduli spaces of stable objects on an abelian surface*. Adv. Stud. Pure Math. 69, 473–537. [doi:10.2969/aspm/06910473](https://doi.org/10.2969/aspm/06910473)