

A SUPERSPACE APPROACH TO SUPERGRAVITY

SUPERSYMMETRY EXAM

Giancarlo Oancia

February 17, 2025

University of Bologna

MOTIVATION

Local susy implies gravity.

Local susy implies gravity.
Global susy in terms of superfields.

Local susy implies gravity.

Global susy in terms of superfields.

Can we study local susy in superspace formalism?

JUST SAY 'SUPER'

CURVED SUPERSPACE

Conventions

- Einstein indices: $\hat{M} = (\hat{\mu}, \hat{\alpha}, \hat{\dot{\alpha}})$. Lorentz indices: $M = (\mu, \alpha, \dot{\alpha})$;

Conventions

- Einstein indices: $\hat{M} = (\hat{\mu}, \hat{\alpha}, \hat{\dot{\alpha}})$. Lorentz indices: $M = (\mu, \alpha, \dot{\alpha})$;
- Superdiffeomorphisms: $z'^{\hat{\mu}} = z^{\hat{\mu}} + \xi^{\hat{\mu}}(z)$;

Conventions

- Einstein indices: $\hat{M} = (\hat{\mu}, \hat{\alpha}, \hat{\dot{\alpha}})$. Lorentz indices: $M = (\mu, \alpha, \dot{\alpha})$;
- Superdiffeomorphisms: $z'^{\hat{\mu}} = z^{\hat{\mu}} + \xi^{\hat{\mu}}(z)$;
- Local Lorentz: $x'^{\mu} = x^{\nu} \Lambda_{\nu}^{\mu}(z)$, $\theta'^{\alpha} = \theta^{\beta} \Lambda_{\beta}^{\alpha}(z)$, $\bar{\theta}'_{\dot{\alpha}} = \bar{\theta}_{\dot{\beta}} \Lambda^{\dot{\beta}}_{\dot{\alpha}}(z)$;

Conventions

- Einstein indices: $\hat{M} = (\hat{\mu}, \hat{\alpha}, \hat{\dot{\alpha}})$. Lorentz indices: $M = (\mu, \alpha, \dot{\alpha})$;
- Superdiffeomorphisms: $z'^{\hat{\mu}} = z^{\hat{\mu}} + \xi^{\hat{\mu}}(z)$;
- Local Lorentz: $x'^{\mu} = x^{\nu} \Lambda_{\nu}^{\mu}(z)$, $\theta'^{\alpha} = \theta^{\beta} \Lambda_{\beta}^{\alpha}(z)$, $\bar{\theta}'_{\dot{\alpha}} = \bar{\theta}_{\dot{\beta}} \Lambda^{\dot{\beta}}_{\dot{\alpha}}(z)$;
- Supervielbein: $E_{\hat{M}}^N(z)$, $g_{\hat{M}\hat{N}} = (-)^{|\hat{M}|(|N|+|\hat{N}|)} E_{\hat{N}}^N E_{\hat{M}}^M \eta_{MN}$;

Conventions

- Einstein indices: $\hat{M} = (\hat{\mu}, \hat{\alpha}, \hat{\dot{\alpha}})$. Lorentz indices: $M = (\mu, \alpha, \dot{\alpha})$;
- Superdiffeomorphisms: $z'^{\hat{\mu}} = z^{\hat{\mu}} + \xi^{\hat{\mu}}(z)$;
- Local Lorentz: $x'^{\mu} = x^{\nu} \Lambda_{\nu}^{\mu}(z)$, $\theta'^{\alpha} = \theta^{\beta} \Lambda_{\beta}^{\alpha}(z)$, $\bar{\theta}'_{\dot{\alpha}} = \bar{\theta}_{\dot{\beta}} \Lambda^{\dot{\beta}}_{\dot{\alpha}}(z)$;
- Supervielbein: $E_{\hat{M}}^N(z)$, $g_{\hat{M}\hat{N}} = (-)^{|\hat{M}|(|N|+|\hat{N}|)} E_{\hat{N}}^N E_{\hat{M}}^M \eta_{MN}$;
- Superconnection: $\Omega_{\hat{M}MN}$;

Conventions

- Einstein indices: $\hat{M} = (\hat{\mu}, \hat{\alpha}, \hat{\dot{\alpha}})$. Lorentz indices: $M = (\mu, \alpha, \dot{\alpha})$;
- Superdiffeomorphisms: $z'^{\hat{\mu}} = z^{\hat{\mu}} + \xi^{\hat{\mu}}(z)$;
- Local Lorentz: $x'^{\mu} = x^{\nu} \Lambda_{\nu}^{\mu}(z)$, $\theta'^{\alpha} = \theta^{\beta} \Lambda_{\beta}^{\alpha}(z)$, $\bar{\theta}'_{\dot{\alpha}} = \bar{\theta}_{\dot{\beta}} \Lambda^{\dot{\beta}}_{\dot{\alpha}}(z)$;
- Supervielbein: $E_{\hat{M}}^N(z)$, $g_{\hat{M}\hat{N}} = (-)^{|\hat{M}|(|N|+|\hat{N}|)} E_{\hat{N}}^N E_{\hat{M}}^M \eta_{MN}$;
- Superconnection: $\Omega_{\hat{M}MN}$;
- Covariant derivatives:

$$\mathfrak{D}_{\hat{M}} X^M = \partial_{\hat{M}} X^M + (-)^{|\hat{M}||N|} X^N \Omega_{\hat{M}N}^M,$$

$$\mathfrak{D}_N X^M = E_N^{\hat{M}} \mathfrak{D}_{\hat{M}} X^M;$$

Conventions

- Einstein indices: $\hat{M} = (\hat{\mu}, \hat{\alpha}, \hat{\dot{\alpha}})$. Lorentz indices: $M = (\mu, \alpha, \dot{\alpha})$;
- Superdiffeomorphisms: $z'^{\hat{\mu}} = z^{\hat{\mu}} + \xi^{\hat{\mu}}(z)$;
- Local Lorentz: $x'^{\mu} = x^{\nu} \Lambda_{\nu}^{\mu}(z)$, $\theta'^{\alpha} = \theta^{\beta} \Lambda_{\beta}^{\alpha}(z)$, $\bar{\theta}'_{\dot{\alpha}} = \bar{\theta}_{\dot{\beta}} \Lambda^{\dot{\beta}}_{\dot{\alpha}}(z)$;
- Supervielbein: $E_{\hat{M}}^N(z)$, $g_{\hat{M}\hat{N}} = (-)^{|\hat{M}|(|N|+|\hat{N}|)} E_{\hat{N}}^N E_{\hat{M}}^M \eta_{MN}$;
- Superconnection: $\Omega_{\hat{M}MN}$;
- Covariant derivatives:

$$\mathfrak{D}_{\hat{M}} X^M = \partial_{\hat{M}} X^M + (-)^{|\hat{M}||N|} X^N \Omega_{\hat{M}N}^M,$$

$$\mathfrak{D}_N X^M = E_N^{\hat{M}} \mathfrak{D}_{\hat{M}} X^M;$$

- Torsion: $T_{\hat{M}\hat{N}}^N$. Curvature: $R_{\hat{M}\hat{N}M}^N$;

Conventions

- Einstein indices: $\hat{M} = (\hat{\mu}, \hat{\alpha}, \hat{\dot{\alpha}})$. Lorentz indices: $M = (\mu, \alpha, \dot{\alpha})$;
- Superdiffeomorphisms: $z'^{\hat{\mu}} = z^{\hat{\mu}} + \xi^{\hat{\mu}}(z)$;
- Local Lorentz: $x'^{\mu} = x^{\nu} \Lambda_{\nu}^{\mu}(z)$, $\theta'^{\alpha} = \theta^{\beta} \Lambda_{\beta}^{\alpha}(z)$, $\bar{\theta}'_{\dot{\alpha}} = \bar{\theta}_{\dot{\beta}} \Lambda^{\dot{\beta}}_{\dot{\alpha}}(z)$;
- Supervielbein: $E_{\hat{M}}^N(z)$, $g_{\hat{M}\hat{N}} = (-)^{|\hat{M}|(|N|+|\hat{N}|)} E_{\hat{N}}^N E_{\hat{M}}^M \eta_{MN}$;
- Superconnection: $\Omega_{\hat{M}MN}$;
- Covariant derivatives:

$$\mathfrak{D}_{\hat{M}} X^M = \partial_{\hat{M}} X^M + (-)^{|\hat{M}||N|} X^N \Omega_{\hat{M}N}^M,$$

$$\mathfrak{D}_N X^M = E_N^{\hat{M}} \mathfrak{D}_{\hat{M}} X^M;$$

- Torsion: $T_{\hat{M}\hat{N}}^N$. Curvature: $R_{\hat{M}\hat{N}M}^N$;
- $[\mathfrak{D}_M, \mathfrak{D}_N]_{|M||N|} = T_{MN}^P \mathfrak{D}_P - 1/2 R_{MN\underline{\alpha}\underline{\beta}} J^{\underline{\alpha}\underline{\beta}}$.

Flat Limit

- $\Omega_{\hat{M}\hat{M}}{}^P = 0;$
- $\mathfrak{D}_{\hat{M}} = \partial_{\hat{M}};$

Flat Limit

- $\Omega_{\hat{M}\hat{M}}{}^P = 0;$
- $\mathfrak{D}_{\hat{M}} = \partial_{\hat{M}};$
- $\mathcal{D}_\mu = \partial_\mu = E_\mu{}^{\hat{M}}\partial_{\hat{M}}, \mathcal{D}_\alpha = E_\alpha{}^{\hat{M}}\partial_{\hat{M}}, \bar{\mathcal{D}}^{\dot{\alpha}} = E^{\dot{\alpha}\hat{M}}\partial_{\hat{M}};$

Flat Limit

- $\Omega_{\hat{M}\hat{M}}^P = 0;$
- $\mathfrak{D}_{\hat{M}} = \partial_{\hat{M}};$
- $\mathcal{D}_\mu = \partial_\mu = E_\mu^{\hat{M}} \partial_{\hat{M}}, \mathcal{D}_\alpha = E_\alpha^{\hat{M}} \partial_{\hat{M}}, \bar{D}^{\dot{\alpha}} = E^{\dot{\alpha}\hat{M}} \partial_{\hat{M}};$
- $\mathcal{D}_\alpha = \partial_\alpha + i\sigma^\mu_{\alpha\dot{\alpha}} \bar{\theta}^{\dot{\alpha}} \partial_\mu, \bar{D}_{\dot{\alpha}} = \bar{\partial}_{\dot{\alpha}} - i\theta^\alpha \sigma^\mu_{\alpha\dot{\alpha}} \partial_\mu;$

Flat Limit

- $\Omega_{\hat{M}\hat{M}}^P = 0;$
- $\mathfrak{D}_{\hat{M}} = \partial_{\hat{M}};$
- $\mathcal{D}_\mu = \partial_\mu = E_\mu^{\hat{M}} \partial_{\hat{M}}, \mathcal{D}_\alpha = E_\alpha^{\hat{M}} \partial_{\hat{M}}, \bar{\mathcal{D}}^{\dot{\alpha}} = E^{\dot{\alpha}\hat{M}} \partial_{\hat{M}};$
- $\mathcal{D}_\alpha = \partial_\alpha + i\sigma^\mu_{\alpha\dot{\alpha}} \bar{\theta}^{\dot{\alpha}} \partial_\mu, \bar{\mathcal{D}}_{\dot{\alpha}} = \bar{\partial}_{\dot{\alpha}} - i\theta^\alpha \sigma^\mu_{\alpha\dot{\alpha}} \partial_\mu;$
- $E_M^{\hat{M}} = \begin{pmatrix} \delta_\mu^{\hat{\mu}} & 0 & 0 \\ i\delta_\mu^{\hat{\mu}} (\sigma^\mu \bar{\theta})_\alpha & \delta_\alpha^{\hat{\alpha}} & 0 \\ -i\delta_\mu^{\hat{\mu}} (\theta \sigma^\mu)_{\dot{\beta}} \varepsilon^{\dot{\alpha}\dot{\beta}} & 0 & \delta_{\hat{\alpha}}^{\dot{\alpha}} \end{pmatrix};$

Flat Limit

- $\Omega_{\hat{M}\hat{M}}^P = 0;$
- $\mathfrak{D}_{\hat{M}} = \partial_{\hat{M}};$
- $\mathcal{D}_\mu = \partial_\mu = E_\mu^{\hat{M}} \partial_{\hat{M}}, \mathcal{D}_\alpha = E_\alpha^{\hat{M}} \partial_{\hat{M}}, \bar{\mathcal{D}}^{\dot{\alpha}} = E^{\dot{\alpha}\hat{M}} \partial_{\hat{M}};$
- $\mathcal{D}_\alpha = \partial_\alpha + i\sigma^\mu_{\alpha\dot{\alpha}} \bar{\theta}^{\dot{\alpha}} \partial_\mu, \bar{\mathcal{D}}_{\dot{\alpha}} = \bar{\partial}_{\dot{\alpha}} - i\theta^\alpha \sigma^\mu_{\alpha\dot{\alpha}} \partial_\mu;$
- $E_M^{\hat{M}} = \begin{pmatrix} \delta_{\mu}^{\hat{\mu}} & 0 & 0 \\ i\delta_{\mu}^{\hat{\mu}}(\sigma^\mu \bar{\theta})_{\alpha} & \delta_{\alpha}^{\hat{\alpha}} & 0 \\ -i\delta_{\mu}^{\hat{\mu}}(\theta \sigma^\mu)_{\dot{\beta}} \varepsilon^{\dot{\alpha}\dot{\beta}} & 0 & \delta_{\dot{\alpha}}^{\hat{\alpha}} \end{pmatrix};$
- $\{\mathcal{D}_\alpha, \bar{\mathcal{D}}_{\dot{\alpha}}\} = -2i\sigma^\mu_{\alpha\dot{\alpha}} \partial_\mu.$

Bianchi identities.

Bianchi identities.

Dragon theorem.

Bianchi identities.

Dragon theorem.

Sugra algebra is Susy algebra in flat limit.

Bianchi identities.

Dragon theorem.

Sugra algebra is Susy algebra in flat limit.

No spin higher than two.

Bianchi identities.

Dragon theorem.

Sugra algebra is Susy algebra in flat limit.

No spin higher than two.

Constraints on torsion tensor.

FOR YOUR SAFETY
LET'S SKIP THE (SUPER)COMPUTATION

Supergravity Algebra

$$\{\mathfrak{D}_\alpha, \mathfrak{D}_\beta\} = -1/2 R_{\alpha\beta\gamma\delta} J^{\delta\gamma};$$

$$\{\bar{\mathfrak{D}}_{\dot{\alpha}}, \bar{\mathfrak{D}}_{\dot{\beta}}\} = -1/2 R_{\dot{\alpha}\dot{\beta}\dot{\gamma}\dot{\delta}} J^{\dot{\delta}\dot{\gamma}};$$

$$\{\mathfrak{D}_\alpha, \bar{\mathfrak{D}}_{\dot{\alpha}}\} = T_{\alpha\dot{\alpha}}{}^\mu \mathfrak{D}_\mu - 1/2 R_{\alpha\dot{\alpha}\beta\gamma} J^{\gamma\beta} - 1/2 R_{\alpha\dot{\alpha}\dot{\beta}\dot{\gamma}} J^{\dot{\gamma}\dot{\beta}};$$

$$[\mathfrak{D}_\alpha, \mathfrak{D}_\mu] = T_{\alpha\mu}{}^\beta \mathfrak{D}_\beta - T_{\alpha\mu}{}^{\dot{\beta}} \bar{\mathfrak{D}}_{\dot{\beta}} - 1/2 R_{\alpha\mu\beta\gamma} J^{\gamma\beta} - 1/2 R_{\alpha\mu\dot{\beta}\dot{\gamma}} J^{\dot{\gamma}\dot{\beta}};$$

$$[\bar{\mathfrak{D}}_{\dot{\alpha}}, \mathfrak{D}_\mu] = T_{\dot{\alpha}\mu}{}^\beta \mathfrak{D}_\beta - T_{\dot{\alpha}\mu}{}^{\dot{\beta}} \bar{\mathfrak{D}}_{\dot{\beta}} - 1/2 R_{\dot{\alpha}\mu\beta\gamma} J^{\gamma\beta} - 1/2 R_{\dot{\alpha}\mu\dot{\beta}\dot{\gamma}} J^{\dot{\gamma}\dot{\beta}};$$

$$[\mathfrak{D}_\mu, \mathfrak{D}_\nu] = T_{\mu\nu}{}^\beta \mathfrak{D}_\beta - T_{\mu\nu}{}^{\dot{\beta}} \bar{\mathfrak{D}}_{\dot{\beta}} - 1/2 R_{\mu\nu\beta\gamma} J^{\gamma\beta} - 1/2 R_{\mu\nu\dot{\beta}\dot{\gamma}} J^{\dot{\gamma}\dot{\beta}}.$$

Supergravity Algebra

$$\{\mathfrak{D}_\alpha, \mathfrak{D}_\beta\} = -1/2 R_{\alpha\beta\gamma\delta} J^{\delta\gamma};$$

$$\{\bar{\mathfrak{D}}_{\dot{\alpha}}, \bar{\mathfrak{D}}_{\dot{\beta}}\} = -1/2 R_{\dot{\alpha}\dot{\beta}\dot{\gamma}\dot{\delta}} J^{\dot{\delta}\dot{\gamma}};$$

$$\{\mathfrak{D}_\alpha, \bar{\mathfrak{D}}_{\dot{\alpha}}\} = T_{\alpha\dot{\alpha}}{}^\mu \mathfrak{D}_\mu - 1/2 R_{\alpha\dot{\alpha}\beta\gamma} J^{\gamma\beta} - 1/2 R_{\alpha\dot{\alpha}\dot{\beta}\dot{\gamma}} J^{\dot{\gamma}\dot{\beta}};$$

$$[\mathfrak{D}_\alpha, \mathfrak{D}_\mu] = T_{\alpha\mu}{}^\beta \mathfrak{D}_\beta - T_{\alpha\mu}{}^{\dot{\beta}} \bar{\mathfrak{D}}_{\dot{\beta}} - 1/2 R_{\alpha\mu\beta\gamma} J^{\gamma\beta} - 1/2 R_{\alpha\mu\dot{\beta}\dot{\gamma}} J^{\dot{\gamma}\dot{\beta}};$$

$$[\bar{\mathfrak{D}}_{\dot{\alpha}}, \mathfrak{D}_\mu] = T_{\dot{\alpha}\mu}{}^\beta \mathfrak{D}_\beta - T_{\dot{\alpha}\mu}{}^{\dot{\beta}} \bar{\mathfrak{D}}_{\dot{\beta}} - 1/2 R_{\dot{\alpha}\mu\beta\gamma} J^{\gamma\beta} - 1/2 R_{\dot{\alpha}\mu\dot{\beta}\dot{\gamma}} J^{\dot{\gamma}\dot{\beta}};$$

$$[\mathfrak{D}_\mu, \mathfrak{D}_\nu] = T_{\mu\nu}{}^\beta \mathfrak{D}_\beta - T_{\mu\nu}{}^{\dot{\beta}} \bar{\mathfrak{D}}_{\dot{\beta}} - 1/2 R_{\mu\nu\beta\gamma} J^{\gamma\beta} - 1/2 R_{\mu\nu\dot{\beta}\dot{\gamma}} J^{\dot{\gamma}\dot{\beta}}.$$

Super-Weyl Transformation

$$\delta_\Sigma E_{\hat{M}}{}^\mu = (\Sigma + \Sigma^\dagger) E_{\hat{M}}{}^\mu;$$

$$\delta_\Sigma E_{\hat{M}}{}^\alpha = (2\Sigma^\dagger - \Sigma) E_{\hat{M}}{}^\alpha + i/2 E_{\hat{M}}{}^\mu (\bar{\mathfrak{D}}\Sigma^\dagger \bar{\sigma}_\mu)^\alpha;$$

$$\delta_\Sigma E_{\hat{M}\dot{\alpha}} = (2\Sigma - \Sigma^\dagger) E_{\hat{M}\dot{\alpha}} + i/2 E_{\hat{M}}{}^\mu (\mathfrak{D}\Sigma \sigma_\mu)_{\dot{\alpha}}.$$

Independent Superfields

- Chiral superfield $\mathcal{R}$;
- Real vector superfield $\mathcal{G}_\mu$;
- Symmetric chiral superfield $\mathcal{W}_{(\alpha\beta\gamma)}$.

Independent Superfields

- Chiral superfield $\mathcal{R}$;
- Real vector superfield $\mathcal{G}_\mu$;
- Symmetric chiral superfield $\mathcal{W}_{(\alpha\beta\gamma)}$.

Supergravity Multiplet

- Massless graviton $e_{\hat{\mu}}^{\mu}$;
- Massless gravitino $(\psi_{\hat{\mu}}^{\alpha}, \bar{\psi}_{\hat{\mu}\dot{\alpha}})$;
- Auxiliary complex scalar M ;
- Auxiliary real vector b_{μ} .

CURVED SUPERSPACE: GAUGE FIXING

Using local Lorentz and superdiffeomorphisms, we can fix the gauge

$$\Omega_{\hat{\mu}M}{}^N| = \omega_{\hat{\mu}M}{}^N(x), \quad \Omega_{\hat{\alpha}M}{}^N| = 0, \quad \Omega^{\hat{\alpha}}{}_{\hat{M}}{}^N| = 0;$$

$$E_{\hat{M}}{}^M| = \begin{pmatrix} e_{\hat{\mu}}{}^{\mu}(x) & 1/2 \psi_{\hat{\mu}}{}^{\alpha}(x) & 1/2 \bar{\psi}_{\hat{\mu}\dot{\alpha}}(x) \\ 0 & \delta_{\hat{\alpha}}^{\alpha} & 0 \\ 0 & 0 & \delta_{\hat{\alpha}}^{\dot{\alpha}} \end{pmatrix};$$

$$E_M{}^{\hat{M}}| = \begin{pmatrix} e_{\mu}{}^{\hat{\mu}}(x) & -1/2 \psi_{\mu}{}^{\hat{\alpha}}(x) & -1/2 \bar{\psi}_{\mu\hat{\alpha}}(x) \\ 0 & \delta_{\alpha}^{\hat{\alpha}} & 0 \\ 0 & 0 & \delta_{\dot{\alpha}}^{\hat{\dot{\alpha}}} \end{pmatrix};$$

$$\mathcal{R}(z)| = -1/6 M(x);$$

$$\mathcal{G}_{\mu}(z)| = -1/3 b_{\mu}(x).$$

CURVED SUPERSPACE: SUPERGRAVITY TRANSFORMATION

Defining $\tilde{\theta}_\alpha = \theta^{\hat{\alpha}} E_{\hat{\alpha}}{}^\beta \varepsilon_{\alpha\beta} \neq \theta_\alpha$, the supergravity transformation is

$$\begin{aligned}\xi^\mu(z) &= 2i(\tilde{\theta}\sigma^\mu\bar{\varepsilon} - \varepsilon\sigma^\mu\tilde{\bar{\theta}}), \quad \xi^\alpha(z) = \varepsilon^\alpha, \quad \bar{\xi}_{\dot{\alpha}}(z) = \bar{\varepsilon}_{\dot{\alpha}}, \\ L_{\alpha\beta}(z) &= 1/3 \left[\tilde{\theta}_\alpha(2\varepsilon_\beta M^* + b_{\beta\dot{\gamma}}\bar{\varepsilon}^{\dot{\gamma}}) + \tilde{\theta}_\beta(2\varepsilon_\alpha M^* + b_{\alpha\dot{\gamma}}\bar{\varepsilon}^{\dot{\gamma}}) \right], \\ L_{\dot{\alpha}\dot{\beta}}(z) &= 1/3 \left[\tilde{\bar{\theta}}_{\dot{\alpha}}(2\bar{\varepsilon}_{\dot{\beta}} M + \varepsilon^\gamma b_{\gamma\dot{\beta}}) + \tilde{\bar{\theta}}_{\dot{\beta}}(2\bar{\varepsilon}_{\dot{\alpha}} M + \varepsilon^\gamma b_{\gamma\dot{\alpha}}) \right], \\ L_{\mu\nu}(z) &= -1/2 \left[(\bar{\sigma}_\mu\sigma_\nu)^{\dot{\alpha}}{}_{\dot{\beta}} L_{\dot{\alpha}}{}^{\dot{\beta}} + (\sigma_\mu\bar{\sigma}_\nu)_\alpha{}^\beta L^\alpha{}_\beta \right],\end{aligned}$$

CURVED SUPERSPACE: SUPERGRAVITY TRANSFORMATION

Defining $\tilde{\theta}_\alpha = \theta^{\hat{\alpha}} E_{\hat{\alpha}}^\beta \varepsilon_{\alpha\beta} \neq \theta_\alpha$, the supergravity transformation is

$$\begin{aligned}\xi^\mu(z) &= 2i(\tilde{\theta}\sigma^\mu\bar{\varepsilon} - \varepsilon\sigma^\mu\tilde{\bar{\theta}}), \quad \xi^\alpha(z) = \varepsilon^\alpha, \quad \bar{\xi}_{\dot{\alpha}}(z) = \bar{\varepsilon}_{\dot{\alpha}}, \\ L_{\alpha\beta}(z) &= 1/3 \left[\tilde{\theta}_\alpha(2\varepsilon_\beta M^* + b_{\beta\dot{\gamma}}\bar{\varepsilon}^{\dot{\gamma}}) + \tilde{\theta}_\beta(2\varepsilon_\alpha M^* + b_{\alpha\dot{\gamma}}\bar{\varepsilon}^{\dot{\gamma}}) \right], \\ L_{\dot{\alpha}\dot{\beta}}(z) &= 1/3 \left[\tilde{\bar{\theta}}_{\dot{\alpha}}(2\bar{\varepsilon}_{\dot{\beta}} M + \varepsilon^\gamma b_{\gamma\dot{\beta}}) + \tilde{\bar{\theta}}_{\dot{\beta}}(2\bar{\varepsilon}_{\dot{\alpha}} M + \varepsilon^\gamma b_{\gamma\dot{\alpha}}) \right], \\ L_{\mu\nu}(z) &= -1/2 \left[(\bar{\sigma}_\mu\sigma_\nu)^{\dot{\alpha}}_{\dot{\beta}} L_{\dot{\alpha}}^{\dot{\beta}} + (\sigma_\mu\bar{\sigma}_\nu)_\alpha^\beta L^\alpha_\beta \right],\end{aligned}$$

The transformation of the supergravity multiplet is

$$\begin{aligned}\delta e_{\hat{\mu}}^\mu &= \delta E_{\hat{\mu}}^\mu \Big| = -i \left[\varepsilon(x) \sigma^\mu \bar{\psi}_{\hat{\mu}} - \psi_{\hat{\mu}} \sigma^\mu \bar{\varepsilon}(x) \right] \\ \delta \psi_{\hat{\mu}}^\alpha &= 2\delta E_{\hat{\mu}}^\alpha \Big| = -2\mathfrak{D}_{\hat{\mu}}\varepsilon^\alpha(x) + \frac{i}{3} e_{\hat{\mu}}^\nu (\bar{\varepsilon}(x) \bar{\sigma}_\nu)^\alpha M \\ &\quad + \frac{i}{3} e_{\hat{\mu}}^\nu [(\varepsilon(x) \sigma_\rho \bar{\sigma}_\nu)^\alpha b^\rho - 3\varepsilon^\alpha(x) b_\nu] \\ \delta M &= -6\delta\mathcal{R} \Big| = \dots \quad \delta b^\mu = -3\delta\mathcal{G}^\mu \Big| = \dots\end{aligned}$$

where $\mathfrak{D}_{\hat{\mu}}\psi_{\hat{\nu}}^\alpha = \partial_{\hat{\mu}}\psi_{\hat{\nu}}^\alpha + \psi_{\hat{\nu}}^\beta \omega_{\hat{\mu}\beta}^\alpha$.

SUPERFIELDS IN CURVED SPACETIME

Chiral Superfield: Definition

$$\bullet \bar{\mathcal{D}}_{\dot{\alpha}} \Phi = 0, \mathcal{D}_{\alpha} \Phi^{\dagger} = 0;$$

Chiral Superfield: Definition

- $\bar{\mathcal{D}}_{\dot{\alpha}}\Phi = 0, \mathcal{D}_{\alpha}\Phi^{\dagger} = 0;$
- $\Phi = \Phi(z^{\hat{M}}) = \Phi(x^{\hat{\mu}}, \theta^{\hat{\alpha}}, \bar{\theta}_{\hat{\dot{\alpha}}});$
- Derivatives have Lorentz indices. Expansion has Einstein indices.

A NIGHTMARE

Chiral Superfield: Definition

$$\bullet \bar{\mathcal{D}}_{\dot{\alpha}}\Phi = 0, \mathcal{D}_{\alpha}\Phi^{\dagger} = 0;$$

Chiral Superfield: Definition

- $\bar{\mathcal{D}}_{\dot{\alpha}}\Phi = 0, \mathcal{D}_{\alpha}\Phi^{\dagger} = 0;$
- Use: $\int d^2\theta d^2\bar{\theta} S(x, \theta, \bar{\theta}) = -\frac{1}{4} \int d^2\theta \bar{\mathcal{D}} \cdot \bar{\mathcal{D}} S(x, \theta, \bar{\theta});$

Chiral Superfield: Definition

- $\bar{\mathfrak{D}}_{\dot{\alpha}}\Phi = 0, \mathfrak{D}_{\alpha}\Phi^{\dagger} = 0;$
- Use: $\int d^2\theta d^2\bar{\theta} S(x, \theta, \bar{\theta}) = -\frac{1}{4} \int d^2\theta \bar{\mathfrak{D}} \cdot \bar{\mathfrak{D}} S(x, \theta, \bar{\theta});$
- Components with Lorentz indices:

$$\begin{aligned}\phi &= \Phi|, & \xi_{\alpha} &= \frac{1}{\sqrt{2}} \mathfrak{D}_{\alpha} \Phi|, & F &= \frac{1}{4} \mathfrak{D} \cdot \mathfrak{D} \Phi|, \\ \phi^{\dagger} &= \Phi^{\dagger}|, & \bar{\xi}^{\dot{\alpha}} &= \frac{1}{\sqrt{2}} \bar{\mathfrak{D}}^{\dot{\alpha}} \Phi^{\dagger}|, & F^{\dagger} &= \frac{1}{4} \bar{\mathfrak{D}} \cdot \bar{\mathfrak{D}} \Phi^{\dagger}|;\end{aligned}$$

Chiral Superfield: Definition

- $\bar{\mathcal{D}}_{\dot{\alpha}} \Phi = 0, \mathcal{D}_{\alpha} \Phi^{\dagger} = 0;$
- Use: $\int d^2\theta d^2\bar{\theta} S(x, \theta, \bar{\theta}) = -\frac{1}{4} \int d^2\theta \bar{\mathcal{D}} \cdot \bar{\mathcal{D}} S(x, \theta, \bar{\theta});$
- Components with Lorentz indices:

$$\begin{aligned} \phi &= \Phi|, & \xi_{\alpha} &= \frac{1}{\sqrt{2}} \mathcal{D}_{\alpha} \Phi|, & F &= \frac{1}{4} \mathcal{D} \cdot \mathcal{D} \Phi|, \\ \phi^{\dagger} &= \Phi^{\dagger}|, & \bar{\xi}^{\dot{\alpha}} &= \frac{1}{\sqrt{2}} \bar{\mathcal{D}}^{\dot{\alpha}} \Phi^{\dagger}|, & F^{\dagger} &= \frac{1}{4} \bar{\mathcal{D}} \cdot \bar{\mathcal{D}} \Phi^{\dagger}|; \end{aligned}$$

- $\Xi^{\dagger} = [\mathcal{D} \cdot \mathcal{D} - 8\mathcal{R}^{\dagger}] \Phi, \quad \Xi = [\bar{\mathcal{D}} \cdot \bar{\mathcal{D}} - 8\mathcal{R}] \Phi^{\dagger}.$

Chiral Superfield: Definition

- $\bar{\mathcal{D}}_{\dot{\alpha}} \Phi = 0, \mathcal{D}_{\alpha} \Phi^{\dagger} = 0;$
- Use: $\int d^2\theta d^2\bar{\theta} S(x, \theta, \bar{\theta}) = -\frac{1}{4} \int d^2\theta \bar{\mathcal{D}} \cdot \bar{\mathcal{D}} S(x, \theta, \bar{\theta});$
- Components with Lorentz indices:

$$\begin{aligned} \phi &= \Phi|, \quad \xi_{\alpha} = \frac{1}{\sqrt{2}} \mathcal{D}_{\alpha} \Phi|, \quad F = \frac{1}{4} \mathcal{D} \cdot \mathcal{D} \Phi|, \\ \phi^{\dagger} &= \Phi^{\dagger}|, \quad \bar{\xi}^{\dot{\alpha}} = \frac{1}{\sqrt{2}} \bar{\mathcal{D}}^{\dot{\alpha}} \Phi^{\dagger}|, \quad F^{\dagger} = \frac{1}{4} \bar{\mathcal{D}} \cdot \bar{\mathcal{D}} \Phi^{\dagger}|; \end{aligned}$$

- $\Xi^{\dagger} = [\mathcal{D} \cdot \mathcal{D} - 8\mathcal{R}^{\dagger}] \Phi, \quad \Xi = [\bar{\mathcal{D}} \cdot \bar{\mathcal{D}} - 8\mathcal{R}] \Phi^{\dagger}.$
- Supergravity transformation:

$$\delta \Phi = -\xi^M \mathcal{D}_M \Phi = -\xi^{\alpha} \mathcal{D}_{\alpha} \Phi - \xi^{\mu} \mathcal{D}_{\mu} \Phi;$$

Vector Superfield: Definition

- Definition: $V = V^\dagger$;

Vector Superfield: Definition

- Definition: $V = V^\dagger$;
- Gauge transformation: $e^{2gV} \rightarrow e^{-2ig\Lambda} e^{2gV} e^{2ig\Lambda^\dagger}$;

Vector Superfield: Definition

- Definition: $V = V^\dagger$;
- Gauge transformation: $e^{2gV} \rightarrow e^{-2ig\Lambda} e^{2gV} e^{2ig\Lambda^\dagger}$;
- Wess-Zumino Gauge:

$$V| = 0, \quad \mathfrak{D}_\alpha V| = 0, \quad \bar{\mathfrak{D}}_{\dot{\alpha}} V| = 0, \quad \mathfrak{D} \cdot \mathfrak{D} V| = 0, \quad \bar{\mathfrak{D}} \cdot \bar{\mathfrak{D}} V| = 0;$$

Vector Superfield: Definition

- Definition: $V = V^\dagger$;
- Gauge transformation: $e^{2gV} \rightarrow e^{-2ig\Lambda} e^{2gV} e^{2ig\Lambda^\dagger}$;
- Wess-Zumino Gauge:

$$V| = 0, \quad \mathfrak{D}_\alpha V| = 0, \quad \bar{\mathfrak{D}}_{\dot{\alpha}} V| = 0, \quad \mathfrak{D} \cdot \mathfrak{D} V| = 0, \quad \bar{\mathfrak{D}} \cdot \bar{\mathfrak{D}} V| = 0;$$

- Field-strength superfield:

$$W_\alpha = -\frac{1}{4}(\bar{\mathfrak{D}} \cdot \bar{\mathfrak{D}} - 8\mathcal{R})e^{2gV}\mathfrak{D}_\alpha e^{-2gV}, \quad \bar{W}_{\dot{\alpha}} = \frac{1}{4}(\mathfrak{D} \cdot \mathfrak{D} - 8\mathcal{R}^\dagger)e^{-2gV}\bar{\mathfrak{D}}_{\dot{\alpha}} e^{2gV};$$

Vector Superfield: Definition

- Definition: $V = V^\dagger$;
- Gauge transformation: $e^{2gV} \rightarrow e^{-2ig\Lambda} e^{2gV} e^{2ig\Lambda^\dagger}$;
- Wess-Zumino Gauge:

$$V| = 0, \quad \mathfrak{D}_\alpha V| = 0, \quad \bar{\mathfrak{D}}_{\dot{\alpha}} V| = 0, \quad \mathfrak{D} \cdot \mathfrak{D} V| = 0, \quad \bar{\mathfrak{D}} \cdot \bar{\mathfrak{D}} V| = 0;$$

- Field-strength superfield:

$$W_\alpha = -\frac{1}{4}(\bar{\mathfrak{D}} \cdot \bar{\mathfrak{D}} - 8\mathcal{R})e^{2gV}\mathfrak{D}_\alpha e^{-2gV}, \quad \bar{W}_{\dot{\alpha}} = \frac{1}{4}(\mathfrak{D} \cdot \mathfrak{D} - 8\mathcal{R}^\dagger)e^{-2gV}\bar{\mathfrak{D}}_{\dot{\alpha}} e^{2gV};$$

- Under gauge: $W_\alpha \rightarrow e^{-2ig\Lambda} W_\alpha e^{2ig\Lambda}$;

Vector Superfield: Definition

- Definition: $V = V^\dagger$;
- Gauge transformation: $e^{2gV} \rightarrow e^{-2ig\Lambda} e^{2gV} e^{2ig\Lambda^\dagger}$;
- Wess-Zumino Gauge:

$$V| = 0, \quad \mathfrak{D}_\alpha V| = 0, \quad \bar{\mathfrak{D}}_{\dot{\alpha}} V| = 0, \quad \mathfrak{D} \cdot \mathfrak{D} V| = 0, \quad \bar{\mathfrak{D}} \cdot \bar{\mathfrak{D}} V| = 0;$$

- Field-strength superfield:

$$W_\alpha = -\frac{1}{4}(\bar{\mathfrak{D}} \cdot \bar{\mathfrak{D}} - 8\mathcal{R})e^{2gV} \mathfrak{D}_\alpha e^{-2gV}, \quad \bar{W}_{\dot{\alpha}} = \frac{1}{4}(\mathfrak{D} \cdot \mathfrak{D} - 8\mathcal{R}^\dagger)e^{-2gV} \bar{\mathfrak{D}}_{\dot{\alpha}} e^{2gV};$$

- Under gauge: $W_\alpha \rightarrow e^{-2ig\Lambda} W_\alpha e^{2ig\Lambda}$;
- Physical components:

$$V_\mu = \frac{1}{4} \bar{\sigma}_\mu^{\dot{\alpha}\alpha} [\mathfrak{D}_\alpha, \bar{\mathfrak{D}}_{\dot{\alpha}}] V|, \quad \lambda_\alpha = \frac{i}{2g} W_\alpha|, \quad \bar{\lambda}_{\dot{\alpha}} = -\frac{i}{2g} \bar{W}_{\dot{\alpha}}|,$$

$$D = \frac{1}{4g} \mathfrak{D}^\alpha W_\alpha| = \frac{1}{4g} \bar{\mathfrak{D}}_{\dot{\alpha}} \bar{W}^{\dot{\alpha}}|.$$

Gauge Interactions

- Gauge transformation: $\Phi \rightarrow e^{-2ig\Lambda}\Phi$;

Gauge Interactions

- Gauge transformation: $\Phi \rightarrow e^{-2ig\Lambda}\Phi$;
- Gauge invariant term: $\Phi^\dagger e^{-2igV} \Phi$;

Gauge Interactions

- Gauge transformation: $\Phi \rightarrow e^{-2ig\Lambda}\Phi$;
- Gauge invariant term: $\Phi^\dagger e^{-2igV} \Phi$;

Gauge Interactions

- Gauge transformation: $\Phi \rightarrow e^{-2ig\Lambda}\Phi$;
- Gauge invariant term: $\Phi^\dagger e^{-2igV} \Phi$;
- Relevant superfield:

$$\begin{aligned}\Upsilon &= (\bar{\mathfrak{D}} \cdot \bar{\mathfrak{D}} - 8\mathcal{R})\Phi^\dagger e^{-2gV} \\ &= \Xi - 2g(\bar{\mathfrak{D}} \cdot \bar{\mathfrak{D}} - 8\mathcal{R})\Phi^\dagger V + 2g^2(\bar{\mathfrak{D}} \cdot \bar{\mathfrak{D}} - 8\mathcal{R})\Phi^\dagger V^2\end{aligned}$$

Gauge Interactions

- Gauge transformation: $\Phi \rightarrow e^{-2ig\Lambda}\Phi$;
- Gauge invariant term: $\Phi^\dagger e^{-2igV} \Phi$;
- Relevant superfield:

$$\begin{aligned}\Upsilon &= (\bar{\mathfrak{D}} \cdot \bar{\mathfrak{D}} - 8\mathcal{R})\Phi^\dagger e^{-2gV} \\ &= \Xi - 2g(\bar{\mathfrak{D}} \cdot \bar{\mathfrak{D}} - 8\mathcal{R})\Phi^\dagger V + 2g^2(\bar{\mathfrak{D}} \cdot \bar{\mathfrak{D}} - 8\mathcal{R})\Phi^\dagger V^2\end{aligned}$$

- Supergravity transformation: $\delta V = -\xi^M \mathfrak{D}_M V$.

SUPERGRAVITY INVARIANT ACTION

Integration on Superspace

- Invariant measure of integration?

Integration on Superspace

- Invariant measure of integration?
- Superdeterminant: $d^4x d^4\theta = \text{sdet}(E_{\hat{M}}^M) d^4\hat{x} d^4\hat{\theta} \equiv E d^4\hat{x} d^4\hat{\theta};$

Integration on Superspace

- Invariant measure of integration?
- Superdeterminant: $d^4x d^4\theta = \text{sdet}(E_{\hat{M}}^M) d^4\hat{x} d^4\hat{\theta} \equiv E d^4\hat{x} d^4\hat{\theta};$
- For a chiral superfield Φ : derivatives have Lorentz indices.
Expansion has Einstein indices;

Integration on Superspace

- Invariant measure of integration?
- Superdeterminant: $d^4x d^4\theta = \text{sdet}(E_{\hat{M}}^M) d^4\hat{x} d^4\hat{\theta} \equiv E d^4\hat{x} d^4\hat{\theta}$;
- For a chiral superfield Φ : derivatives have Lorentz indices.
Expansion has Einstein indices;
- Define $z^{\bar{M}} = (x^{\hat{\mu}}, \Theta^\alpha, \bar{\Theta}_{\dot{\alpha}})$.

WESS: 'A TOUR DE FORCE'

Defining Hybrid Variables

- Define $z^{\bar{M}} = (x^{\hat{\mu}}, \Theta^\alpha, \bar{\Theta}_{\dot{\alpha}})$ such that:

$$\begin{aligned}\Phi(x, \Theta) &= \Phi| + \sqrt{2}\Theta \cdot (\mathfrak{D}\Phi) \Big| - \frac{1}{4}\Theta \cdot \Theta (\mathfrak{D} \cdot \mathfrak{D}\Phi) \Big| \\ &= \phi + \sqrt{2}\Theta \cdot \chi - \Theta \cdot \Theta F;\end{aligned}$$

Defining Hybrid Variables

- Define $z^{\bar{M}} = (x^{\hat{\mu}}, \Theta^\alpha, \bar{\Theta}_{\dot{\alpha}})$ such that:

$$\begin{aligned}\Phi(x, \Theta) &= \Phi| + \sqrt{2}\Theta \cdot (\mathfrak{D}\Phi) \Big| - \frac{1}{4}\Theta \cdot \Theta (\mathfrak{D} \cdot \mathfrak{D}\Phi) \Big| \\ &= \phi + \sqrt{2}\Theta \cdot \chi - \Theta \cdot \Theta F;\end{aligned}$$

- $S_{\text{curved}} = \int d^4x d^2\Theta \Delta f(\Phi) + h.c.;$

Defining Hybrid Variables

- Define $z^{\bar{M}} = (x^{\hat{\mu}}, \Theta^\alpha, \bar{\Theta}_{\dot{\alpha}})$ such that:

$$\begin{aligned}\Phi(x, \Theta) &= \Phi| + \sqrt{2}\Theta \cdot (\mathfrak{D}\Phi) \Big| - \frac{1}{4}\Theta \cdot \Theta (\mathfrak{D} \cdot \mathfrak{D}\Phi) \Big| \\ &= \phi + \sqrt{2}\Theta \cdot \chi - \Theta \cdot \Theta F;\end{aligned}$$

- $S_{\text{curved}} = \int d^4x d^2\Theta \Delta f(\Phi) + h.c.$;
- Chiral density Δ , supergravity invariant, such that:

$$\delta(\Delta\Phi) = -(-)^{|\bar{M}|} \partial_{\bar{M}}(\eta^{\bar{M}} \Delta\Phi);$$

Defining Hybrid Variables

- Define $z^{\bar{M}} = (x^{\hat{\mu}}, \Theta^\alpha, \bar{\Theta}_{\dot{\alpha}})$ such that:

$$\begin{aligned}\Phi(x, \Theta) &= \Phi| + \sqrt{2}\Theta \cdot (\mathfrak{D}\Phi) \Big| - \frac{1}{4}\Theta \cdot \Theta (\mathfrak{D} \cdot \mathfrak{D}\Phi) \Big| \\ &= \phi + \sqrt{2}\Theta \cdot \chi - \Theta \cdot \Theta F;\end{aligned}$$

- $S_{\text{curved}} = \int d^4x d^2\Theta \Delta f(\Phi) + h.c.$;
- Chiral density Δ , supergravity invariant, such that:

$$\delta(\Delta\Phi) = -(-)^{|\bar{M}|} \partial_{\bar{M}}(\eta^{\bar{M}} \Delta\Phi);$$

- Choose $\mathcal{E} = e [1 + i\Theta\sigma^{\hat{\mu}}\bar{\psi}_{\hat{\mu}} - \Theta \cdot \Theta (M^* + \bar{\psi}_{\hat{\mu}}\bar{\sigma}^{\hat{\mu}\hat{\nu}}\psi_{\hat{\nu}})]$;

Defining Hybrid Variables

- Define $z^{\bar{M}} = (x^{\hat{\mu}}, \Theta^\alpha, \bar{\Theta}_{\dot{\alpha}})$ such that:

$$\begin{aligned}\Phi(x, \Theta) &= \Phi \Big| + \sqrt{2} \Theta \cdot (\mathfrak{D} \Phi) \Big| - \frac{1}{4} \Theta \cdot \Theta (\mathfrak{D} \cdot \mathfrak{D} \Phi) \Big| \\ &= \phi + \sqrt{2} \Theta \cdot \chi - \Theta \cdot \Theta F;\end{aligned}$$

- $S_{\text{curved}} = \int d^4x d^2\Theta \Delta f(\Phi) + h.c.$;
- Chiral density Δ , supergravity invariant, such that:

$$\delta(\Delta \Phi) = -(-)^{|\bar{M}|} \partial_{\bar{M}} (\eta^{\bar{M}} \Delta \Phi);$$

- Choose $\mathcal{E} = e [1 + i \Theta \sigma^{\hat{\mu}} \bar{\psi}_{\hat{\mu}} - \Theta \cdot \Theta (M^* + \bar{\psi}_{\hat{\mu}} \bar{\sigma}^{\hat{\mu}\hat{\nu}} \bar{\psi}_{\hat{\nu}})]$;
- We can expand in Θ the superfields $\Xi, \Upsilon, W_\alpha, \mathcal{R}$.

Global Susy Action

$$\begin{aligned} S = & \int d^4x \, d^2\theta \, d^2\bar{\theta} \, \frac{1}{2} [K(\Phi, \Phi^\dagger e^{-2gV}) + K(e^{-2gV} \Phi, \Phi^\dagger)] \\ & + \int d^4x \, d^2\theta \, W(\Phi) + \int d^4x \, d^2\bar{\theta} \, W^*(\Phi^\dagger) \\ & + \frac{1}{16g^2} \int d^4x \, d^2\theta \, h_{ab}(\Phi) W^{a\alpha} W_\alpha^b + \frac{1}{16g^2} \int d^4x \, d^2\bar{\theta} \, h_{ab}^*(\Phi^\dagger) \bar{W}^{\dot{a}}_\alpha \bar{W}^{b\dot{\alpha}} \end{aligned}$$

Global Susy Action

$$\begin{aligned} S = & \int d^4x \, d^2\theta \, d^2\bar{\theta} \, \frac{1}{2} [K(\Phi, \Phi^\dagger e^{-2gV}) + K(e^{-2gV} \Phi, \Phi^\dagger)] \\ & + \int d^4x \, d^2\theta \, W(\Phi) + \int d^4x \, d^2\bar{\theta} \, W^*(\Phi^\dagger) \\ & + \frac{1}{16g^2} \int d^4x \, d^2\theta h_{ab}(\Phi) W^{a\alpha} W_\alpha^b + \frac{1}{16g^2} \int d^4x \, d^2\bar{\theta} h_{ab}^*(\Phi^\dagger) \bar{W}^{\dot{a}}_\alpha \bar{W}^{b\dot{\alpha}} \end{aligned}$$

Principle of Covariance

- $\theta \rightarrow \Theta$;
- $d^2\theta \rightarrow d^2\Theta \mathcal{E}$;
- $\bar{\mathcal{D}} \cdot \bar{\mathcal{D}} \rightarrow (\bar{\mathfrak{D}} \cdot \bar{\mathfrak{D}} - 8\mathcal{R})$.

SUPERGRAVITY LAGRANGIAN: RESULT

Supergravity Lagrangian

$$\mathcal{L} = \int d^2\Theta \mathcal{E} \left\{ \frac{3}{8} (\bar{\mathfrak{D}} \cdot \bar{\mathfrak{D}} - 8\mathcal{R}) e^{-\frac{1}{3}K(\Phi, \Phi^\dagger e^{-2gV})} + W(\Phi) \right. \\ \left. + \frac{1}{16g^2} h_{ab}(\Phi) W^{a\alpha} W^b_{\alpha} \right\} + h.c.$$

SUPERGRAVITY LAGRANGIAN: RESULT

Supergravity Lagrangian

$$\mathcal{L} = \int d^2\Theta \mathcal{E} \left\{ \frac{3}{8} (\bar{\mathfrak{D}} \cdot \bar{\mathfrak{D}} - 8\mathcal{R}) e^{-\frac{1}{3}K(\Phi, \Phi^\dagger e^{-2gV})} + W(\Phi) \right. \\ \left. + \frac{1}{16g^2} h_{ab}(\Phi) W^{a\alpha} W^b_{\alpha} \right\} + h.c.$$

Units:

$$[\Theta] = [\theta] = -\frac{1}{2}, [\mathcal{E}] = [e] = 0, [\mathfrak{D}] = [\mathcal{D}] = \frac{1}{2}, [K] = 2.$$

Expanding:

$$\mathcal{L} \supset \frac{3}{8} M_P^2 \int d^2\Theta \mathcal{E} (\bar{\mathfrak{D}} \cdot \bar{\mathfrak{D}} - 8\mathcal{R}) e^{-\frac{1}{3M_P^2} K(\Phi, \Phi^\dagger e^{-2gV})} + h.c.$$

SUPERGRAVITY LAGRANGIAN: RESULT

Supergravity Lagrangian

$$\mathcal{L} = \int d^2\Theta \mathcal{E} \left\{ \frac{3}{8} (\bar{\mathfrak{D}} \cdot \bar{\mathfrak{D}} - 8\mathcal{R}) e^{-\frac{1}{3}K(\Phi, \Phi^\dagger e^{-2gV})} + W(\Phi) \right. \\ \left. + \frac{1}{16g^2} h_{ab}(\Phi) W^{a\alpha} W^b_{\alpha} \right\} + h.c.$$

Units:

$$[\Theta] = [\theta] = -\frac{1}{2}, [\mathcal{E}] = [e] = 0, [\mathfrak{D}] = [\mathcal{D}] = \frac{1}{2}, [K] = 2.$$

Expanding:

$$\mathcal{L} \supset \frac{3}{8} M_P^2 \int d^2\Theta \mathcal{E} (\bar{\mathfrak{D}} \cdot \bar{\mathfrak{D}} - 8\mathcal{R}) e^{-\frac{1}{3M_P^2} K(\Phi, \Phi^\dagger e^{-2gV})} + h.c. \\ = -6M_P^2 \int d^2\Theta \mathcal{E} \mathcal{R} + \left[-\frac{1}{8} \int d^2\Theta \mathcal{E} (\bar{\mathfrak{D}} \cdot \bar{\mathfrak{D}} - 8\mathcal{R}) K(\Phi, \Phi^\dagger e^{-2gV}) + h.c. \right] \\ + \mathcal{O}\left(\frac{1}{M_P^2}\right).$$

Super-Weyl Transformation

$$\begin{aligned}\mathcal{L} \rightarrow \mathcal{L}' = & \frac{3}{8} \int d^2\Theta \mathcal{E}(\bar{\mathfrak{D}} \cdot \bar{\mathfrak{D}} - 8\mathcal{R}) e^{2(\Sigma + \Sigma^\dagger)} \exp \left\{ -\frac{1}{3} K(\Phi, \Phi^\dagger e^{-2gV}) \right\} \\ & + \int d^2\Theta \mathcal{E} e^{6\Sigma} W(\Phi) + \frac{1}{16g^2} \int d^2\Theta \mathcal{E} h_{ab}(\Phi) W^{a\alpha} W_\alpha^b + h.c.\end{aligned}$$

Super-Weyl Transformation

$$\begin{aligned}\mathcal{L} \rightarrow \mathcal{L}' = & \frac{3}{8} \int d^2\Theta \mathcal{E}(\bar{\mathfrak{D}} \cdot \bar{\mathfrak{D}} - 8\mathcal{R}) e^{2(\Sigma + \Sigma^\dagger)} \exp \left\{ -\frac{1}{3} K(\Phi, \Phi^\dagger e^{-2gV}) \right\} \\ & + \int d^2\Theta \mathcal{E} e^{6\Sigma} W(\Phi) + \frac{1}{16g^2} \int d^2\Theta \mathcal{E} h_{ab}(\Phi) W^{a\alpha} W_\alpha^b + h.c.\end{aligned}$$

Kahler Transformation

To have an invariant scalar potential:

$$\begin{aligned}K(\Phi, \Phi^\dagger e^{-2gV}) & \rightarrow K(\Phi, \Phi^\dagger e^{-2gV}) + F(\Phi) + F^*(\Phi^\dagger), \\ W(\Phi) & \rightarrow e^{-F(\Phi)} W(\Phi).\end{aligned}$$

SUPERGRAVITY LAGRANGIAN: SYMMETRIES

Super-Weyl Transformation

$$\mathcal{L} \rightarrow \mathcal{L}' = \frac{3}{8} \int d^2\Theta \mathcal{E}(\bar{\mathfrak{D}} \cdot \bar{\mathfrak{D}} - 8\mathcal{R}) e^{2(\Sigma + \Sigma^\dagger)} \exp \left\{ -\frac{1}{3} K(\Phi, \Phi^\dagger e^{-2gV}) \right\} \\ + \int d^2\Theta \mathcal{E} e^{6\Sigma} W(\Phi) + \frac{1}{16g^2} \int d^2\Theta \mathcal{E} h_{ab}(\Phi) W^{a\alpha} W_\alpha^b + h.c.$$

Kahler Transformation

To have an invariant scalar potential:

$$K(\Phi, \Phi^\dagger e^{-2gV}) \rightarrow K(\Phi, \Phi^\dagger e^{-2gV}) + F(\Phi) + F^*(\Phi^\dagger), \\ W(\Phi) \rightarrow e^{-F(\Phi)} W(\Phi).$$

Then:

$$\mathcal{L}' \rightarrow \mathcal{L}'' = \frac{3}{8} \int d^2\Theta \mathcal{E}(\bar{\mathfrak{D}} \cdot \bar{\mathfrak{D}} - 8\mathcal{R}) e^{2(\Sigma + \Sigma^\dagger)} e^{-\frac{1}{3}K - \frac{1}{3}(F + F^*)} \\ + \int d^2\Theta \mathcal{E} e^{-F + 6\Sigma} W + \frac{1}{16g^2} \int d^2\Theta \mathcal{E} h_{ab}(\Phi) W^{a\alpha} W_\alpha^b + h.c.$$

Generalised Kahler Potential

- For $F = \ln(W)$:

$$\begin{aligned}\mathcal{L} = & \frac{3}{8} \int d^2\Theta \mathcal{E} (\bar{\mathfrak{D}} \cdot \bar{\mathfrak{D}} - 8\mathcal{R}) e^{-\frac{1}{3}\mathcal{G}(\Phi, \Phi^\dagger e^{-2gV})} + \int d^2\Theta \mathcal{E} \\ & + \frac{1}{16g^2} \int d^2\Theta \mathcal{E} h_{ab}(\Phi) W^{a\alpha} W_\alpha^b + h.c.\end{aligned}$$

Generalised Kahler Potential

- For $F = \ln(W)$:

$$\mathcal{L} = \frac{3}{8} \int d^2\Theta \mathcal{E} (\bar{\mathfrak{D}} \cdot \bar{\mathfrak{D}} - 8\mathcal{R}) e^{-\frac{1}{3}\mathcal{G}(\Phi, \Phi^\dagger e^{-2gV})} + \int d^2\Theta \mathcal{E} \\ + \frac{1}{16g^2} \int d^2\Theta \mathcal{E} h_{ab}(\Phi) W^{a\alpha} W_\alpha^b + h.c.$$

- Generalised Kahler potential: $\mathcal{G} = K + \ln |W|^2$

Final Result

$$\begin{aligned}\mathcal{L} = & \frac{3}{8} \int d^2\Theta \mathcal{E}(\bar{\mathfrak{D}} \cdot \bar{\mathfrak{D}} - 8\mathcal{R})e^{-\frac{1}{3}\mathcal{G}(\Phi, \Phi^\dagger e^{-2gV})} + \int d^2\Theta \mathcal{E} \\ & + \frac{1}{16g^2} \int d^2\Theta \mathcal{E} h_{ab}(\Phi) W^{a\alpha} W_\alpha^b + h.c.\end{aligned}$$

EXPLICIT COMPUTATION

HAVE FUN
COMPUTING THE COMPONENTS