

CIRCLE COMPACTIFICATION AND T-DUALITY

STRING THEORY EXAM

Giancarlo Oancia

January 28, 2025

University of Bologna

MOTIVATION

Superstring lives in 10d.

Superstring lives in 10d.
We live in 4d.

Superstring lives in 10d.

We live in 4d.

What to do with those extra-dimensions?

WHAT REALLY IS SPACETIME?

KALUZA-KLEIN COMPACTIFICATION

KALUZA-KLEIN COMPACTIFICATION: MAIN RESULTS

Compactification of a scalar field on $\mathcal{M}_D \rightarrow \mathcal{M}_{D-1} \times S^1$:

Kaluza-Klein Masses

$$m_s^2 = M^2 + \frac{s^2}{R^2}.$$

Negligible for

$$E \ll \frac{1}{R}.$$

KALUZA-KLEIN COMPACTIFICATION: MAIN RESULTS

Compactification of a **scalar field** on $\mathcal{M}_D \rightarrow \mathcal{M}_{D-1} \times S^1$:

Kaluza-Klein Masses

$$m_s^2 = M^2 + \frac{s^2}{R^2}.$$

Negligible for

$$E \ll \frac{1}{R}.$$

Compactification of **Einstein gravity** on $\mathcal{M}_D \rightarrow \mathcal{M}_{D-1} \times S^1$:

Result

- Decomposition $SO(1, D-1) \rightarrow SO(1, D-2)$;
- Gauge symmetry;
- $\text{Vol}(S^1) = \int_0^{2\pi R} dy \sqrt{G_{yy}^{(0)}} = e^{-(D-3)\alpha_{D-1}\phi} \cdot 2\pi R.$

SUPERSTRING

Closed Critical String in 10d

- Critical setting \mathcal{M}_{10} . Closed string;
- Bosonic X^μ and fermionic ψ^μ fields;

Closed Critical String in 10d

- Critical setting \mathcal{M}_{10} . Closed string;
- Bosonic X^μ and fermionic ψ^μ fields;
- Lightcone gauge: $X^\mu \rightarrow (X^\pm, X^i)$, $\psi^\mu \rightarrow (\psi^\pm, \psi^i)$;
- $X^i = X_L^i(\tau + \sigma) + X_R^i(\tau, \sigma)$, $\psi^i = \psi_L^i(\tau + \sigma) + \psi_R^i(\tau - \sigma)$;

Closed Critical String in 10d

- Critical setting \mathcal{M}_{10} . Closed string;
- Bosonic X^μ and fermionic ψ^μ fields;
- Lightcone gauge: $X^\mu \rightarrow (X^\pm, X^i)$, $\psi^\mu \rightarrow (\psi^\pm, \psi^i)$;
- $X^i = X_L^i(\tau + \sigma) + X_R^i(\tau, \sigma)$, $\psi^i = \psi_L^i(\tau + \sigma) + \psi_R^i(\tau - \sigma)$;
- Mode expansion: $\psi_{L/R}^i(\xi^\pm) = \sqrt{\frac{2\pi}{l}} \sum_{r \in \mathbb{Z} + \phi} b_r^{i(\sim)} e^{-\frac{2\pi i}{l} r \xi^\pm}$;

Closed Critical String in 10d

- Critical setting \mathcal{M}_{10} . Closed string;
- Bosonic X^μ and fermionic ψ^μ fields;
- Lightcone gauge: $X^\mu \rightarrow (X^\pm, X^i)$, $\psi^\mu \rightarrow (\psi^\pm, \psi^i)$;
- $X^i = X_L^i(\tau + \sigma) + X_R^i(\tau, \sigma)$, $\psi^i = \psi_L^i(\tau + \sigma) + \psi_R^i(\tau - \sigma)$;
- Mode expansion: $\psi_{L/R}^i(\xi^\pm) = \sqrt{\frac{2\pi}{l}} \sum_{r \in \mathbb{Z} + \phi} b_r^{i(\sim)} e^{-\frac{2\pi i}{l} r \xi^\pm}$;
- NS sector: $\phi = 1/2$, $a^{(\sim)} = 1/2$;
- R sector: $\phi = 0$, $a^{(\sim)} = 0$;

Closed Critical String in 10d

- Critical setting \mathcal{M}_{10} . Closed string;
- Bosonic X^μ and fermionic ψ^μ fields;
- Lightcone gauge: $X^\mu \rightarrow (X^\pm, X^i)$, $\psi^\mu \rightarrow (\psi^\pm, \psi^i)$;
- $X^i = X_L^i(\tau + \sigma) + X_R^i(\tau, \sigma)$, $\psi^i = \psi_L^i(\tau + \sigma) + \psi_R^i(\tau - \sigma)$;
- Mode expansion: $\psi_{L/R}^i(\xi^\pm) = \sqrt{\frac{2\pi}{l}} \sum_{r \in \mathbb{Z} + \phi} b_r^{(\sim)i} e^{-\frac{2\pi i}{l} r \xi^\pm}$;
- NS sector: $\phi = 1/2$, $\tilde{a} = 1/2$;
- R sector: $\phi = 0$, $\tilde{a} = 0$;
- Level-matching condition: $M_L^2 = M_R^2$;
- Mass-shell condition: $M_{L/R}^2 = \frac{2}{\alpha'} \left(N_\perp - \tilde{a} \right)$.

SUPERSTRING: THE SECTORS

sector	G-parity	state	little group rep.	$\alpha' M_R^2/2$	statistics
NS	—	$ 0\rangle_{NS}$	$SO(9) : \mathbf{1}$	$-1/2$	boson
NS	+	$b_{-1/2}^i 0\rangle_{NS}$	$SO(8) : \mathbf{8}_v$	0	boson
R	+	$ 0\rangle_R, B_{a_1}^+ B_{a_2}^+ 0\rangle_R,$ $B_1^+ B_2^+ B_3^+ B_4^+ 0\rangle_R$	$SO(8) : \mathbf{8}_s$	0	fermion
R	—	$B_{a_1}^+ 0\rangle_R,$ $B_{a_1}^+ B_{a_2}^+ B_{a_3}^+ 0\rangle_R$	$SO(8) : \mathbf{8}_c$	0	fermion

SUPERSTRING: TYPE II THEORIES

Type IIA			Type IIB		
sector	fields	$SO(8)$	sector	fields	$SO(8)$
(NS_+, NS_+)	$\Phi, B_{[\mu\nu]}, G_{(\mu\nu)}$	$\mathbf{8}_v \otimes \mathbf{8}_v$	(NS_+, NS_+)	$\Phi, B_{[\mu\nu]}, G_{(\mu\nu)}$	$\mathbf{8}_v \otimes \mathbf{8}_v$
(R_+, R_-)	C_1, C_3	$\mathbf{8}_s \otimes \mathbf{8}_c$	(R_+, R_+)	C_0, C_2, C_4^+	$\mathbf{8}_s \otimes \mathbf{8}_s$
(NS_+, R_-)	$\tilde{\lambda}_a, \tilde{\psi}_a^\mu$	$\mathbf{8}_v \otimes \mathbf{8}_c$	(NS_+, R_+)	$\lambda_a^{(1)}, \psi_a^{(1)\mu}$	$\mathbf{8}_v \otimes \mathbf{8}_s$
(R_+, NS_+)	λ_a, ψ_a^μ	$\mathbf{8}_s \otimes \mathbf{8}_v$	(R_+, NS_+)	$\lambda_a^{(2)}, \psi_a^{(2)\mu}$	$\mathbf{8}_s \otimes \mathbf{8}_v$

TYPE II STRING: COMPACTIFICATION

TYPE II STRING: FIELD THEORETIC COMPACTIFICATION

Type IIA					
sector	$SO(8)$	10d fields	$SO(7)$	$SO(7)$ irrep	9d fields
$(\text{NS}_+, \text{NS}_+)$	$\mathbf{8}_v \otimes \mathbf{8}_v$	$\Phi, B_{[\hat{\mu}\hat{\nu}]}, G_{(\hat{\mu}\hat{\nu})}$	$\mathbf{7} \otimes \mathbf{7}$	$\mathbf{1} \oplus \mathbf{21} \oplus \mathbf{27}$	$\phi, B_{[\mu\nu]}, G_{(\mu\nu)}$
			$(\mathbf{7} \otimes \mathbf{1}) \oplus (\mathbf{1} \otimes \mathbf{7})$	$\mathbf{7} \oplus \mathbf{7}$	$G_{\mu 9}, B_{\mu 9}$
			$\mathbf{1} \otimes \mathbf{1}$	$\mathbf{1}$	G_{99}
(R_+, R_-)	$\mathbf{8}_s \otimes \mathbf{8}_c$	$C_{\hat{\mu}}, C_{[\hat{\mu}\hat{\nu}\hat{\rho}]}$	$\mathbf{8} \otimes \mathbf{8}$	$\mathbf{1} \oplus \mathbf{7} \oplus \mathbf{21} \oplus \mathbf{35}$	$A_9, A_\mu, C_{9\mu\nu}, C_{\mu\nu\rho}$
$(\text{NS}_+, \text{R}_-)$	$\mathbf{8}_v \otimes \mathbf{8}_c$	$\tilde{\lambda}_a, \tilde{\psi}_a^{\hat{\mu}}$	$\mathbf{7} \otimes \mathbf{8}$	$\mathbf{8} \oplus \mathbf{48}$	$\tilde{\psi}_a^\mu, \tilde{\psi}_a^9$
			$\mathbf{1} \otimes \mathbf{8}$	$\mathbf{8}$	$\tilde{\lambda}_a$
$(\text{R}_+, \text{NS}_+)$	$\mathbf{8}_s \otimes \mathbf{8}_v$	$\lambda_a, \psi_a^{\hat{\mu}}$	$\mathbf{8} \otimes \mathbf{7}$	$\mathbf{8} \oplus \mathbf{48}$	ψ_a^μ, ψ_a^9
			$\mathbf{8} \otimes \mathbf{1}$	$\mathbf{8}$	λ_a
Type IIB					
sector	$SO(8)$	10d fields	$SO(7)$	$SO(7)$ irrep	9d fields
$(\text{NS}_+, \text{NS}_+)$	$\mathbf{8}_v \otimes \mathbf{8}_v$	$\Phi, B_{[\hat{\mu}\hat{\nu}]}, G_{(\hat{\mu}\hat{\nu})}$	$\mathbf{7} \otimes \mathbf{7}$	$\mathbf{1} \oplus \mathbf{21} \oplus \mathbf{27}$	$\phi, B_{[\mu\nu]}, G_{(\mu\nu)}$
			$(\mathbf{7} \otimes \mathbf{1}) \oplus (\mathbf{1} \otimes \mathbf{7})$	$\mathbf{7} \oplus \mathbf{7}$	$G_{\mu 9}, B_{\mu 9}$
			$\mathbf{1} \otimes \mathbf{1}$	$\mathbf{1}$	G_{99}
(R_+, R_+)	$\mathbf{8}_s \otimes \mathbf{8}_s$	$C_0, C_{[\hat{\mu}\hat{\nu}]}, C_{[\hat{\mu}\hat{\nu}\hat{\rho}\hat{\sigma}]}^+$	$\mathbf{8} \otimes \mathbf{8}$	$\mathbf{1} \oplus \mathbf{7} \oplus \mathbf{21} \oplus \mathbf{35}$	$a, C_{\mu 9}, C_{\mu\nu}, C_{\mu\nu\rho 9}$
$(\text{NS}_+, \text{R}_+)$	$\mathbf{8}_v \otimes \mathbf{8}_s$	$\lambda_a^{(1)}, \psi_a^{(1)\hat{\mu}}$	$\mathbf{7} \otimes \mathbf{8}$	$\mathbf{8} \oplus \mathbf{48}$	$\psi_a^{(1)9}, \psi_a^{(1)\mu}$
			$\mathbf{1} \otimes \mathbf{8}$	$\mathbf{8}$	$\lambda_a^{(1)}$
$(\text{R}_+, \text{NS}_+)$	$\mathbf{8}_s \otimes \mathbf{8}_v$	$\lambda_a^{(2)}, \psi_a^{(2)\hat{\mu}}$	$\mathbf{8} \otimes \mathbf{7}$	$\mathbf{8} \oplus \mathbf{48}$	$\psi_a^{(2)9}, \psi_a^{(2)\mu}$
			$\mathbf{8} \otimes \mathbf{1}$	$\mathbf{8}$	$\lambda_a^{(2)}$

TYPE II STRING: FIELD THEORETIC COMPACTIFICATION

Type IIA					
sector	$SO(8)$	10d fields	$SO(7)$	$SO(7)$ irrep	9d fields
(NS_+, NS_+)	$\mathbf{8}_v \otimes \mathbf{8}_v$	$\Phi, B_{[\hat{\mu}\hat{\nu}]}, G_{(\hat{\mu}\hat{\nu})}$	$\mathbf{7} \otimes \mathbf{7}$	$\mathbf{1} \oplus \mathbf{21} \oplus \mathbf{27}$	$\phi, B_{[\mu\nu]}, G_{(\mu\nu)}$
			$(\mathbf{7} \otimes \mathbf{1}) \oplus (\mathbf{1} \otimes \mathbf{7})$	$\mathbf{7} \oplus \mathbf{7}$	$G_{\mu 9}, B_{\mu 9}$
			$\mathbf{1} \otimes \mathbf{1}$	$\mathbf{1}$	G_{99}
(R_+, R_-)	$\mathbf{8}_s \otimes \mathbf{8}_c$	$C_{\hat{\mu}}, C_{[\hat{\mu}\hat{\nu}\hat{\rho}]}$	$\mathbf{8} \otimes \mathbf{8}$	$\mathbf{1} \oplus \mathbf{7} \oplus \mathbf{21} \oplus \mathbf{35}$	$A_9, A_\mu, C_{9\mu\nu}, C_{\mu\nu\rho}$
(NS_+, R_-)	$\mathbf{8}_v \otimes \mathbf{8}_c$	$\tilde{\lambda}_a, \tilde{\psi}_a^{\hat{\mu}}$	$\mathbf{7} \otimes \mathbf{8}$	$\mathbf{8} \oplus \mathbf{48}$	$\tilde{\psi}_a^\mu, \tilde{\psi}_a^9$
			$\mathbf{1} \otimes \mathbf{8}$	$\mathbf{8}$	$\tilde{\lambda}_a$
(R_+, NS_+)	$\mathbf{8}_s \otimes \mathbf{8}_v$	$\lambda_a, \psi_a^{\hat{\mu}}$	$\mathbf{8} \otimes \mathbf{7}$	$\mathbf{8} \oplus \mathbf{48}$	ψ_a^μ, ψ_a^9
			$\mathbf{8} \otimes \mathbf{1}$	$\mathbf{8}$	λ_a
Type IIB					
sector	$SO(8)$	10d fields	$SO(7)$	$SO(7)$ irrep	9d fields
(NS_+, NS_+)	$\mathbf{8}_v \otimes \mathbf{8}_v$	$\Phi, B_{[\hat{\mu}\hat{\nu}]}, G_{(\hat{\mu}\hat{\nu})}$	$\mathbf{7} \otimes \mathbf{7}$	$\mathbf{1} \oplus \mathbf{21} \oplus \mathbf{27}$	$\phi, B_{[\mu\nu]}, G_{(\mu\nu)}$
			$(\mathbf{7} \otimes \mathbf{1}) \oplus (\mathbf{1} \otimes \mathbf{7})$	$\mathbf{7} \oplus \mathbf{7}$	$G_{\mu 9}, B_{\mu 9}$
			$\mathbf{1} \otimes \mathbf{1}$	$\mathbf{1}$	G_{99}
(R_+, R_+)	$\mathbf{8}_s \otimes \mathbf{8}_s$	$C_0, C_{[\hat{\mu}\hat{\nu}]}, C_{[\hat{\mu}\hat{\nu}\hat{\rho}\hat{\sigma}]}^+$	$\mathbf{8} \otimes \mathbf{8}$	$\mathbf{1} \oplus \mathbf{7} \oplus \mathbf{21} \oplus \mathbf{35}$	$a, C_{\mu 9}, C_{\mu\nu}, C_{\mu\nu\rho 9}$
(NS_+, R_+)	$\mathbf{8}_v \otimes \mathbf{8}_s$	$\lambda_a^{(1)}, \psi_a^{(1)\hat{\mu}}$	$\mathbf{7} \otimes \mathbf{8}$	$\mathbf{8} \oplus \mathbf{48}$	$\psi_a^{(1)9}, \psi_a^{(1)\mu}$
			$\mathbf{1} \otimes \mathbf{8}$	$\mathbf{8}$	$\lambda_a^{(1)}$
(R_+, NS_+)	$\mathbf{8}_s \otimes \mathbf{8}_v$	$\lambda_a^{(2)}, \psi_a^{(2)\hat{\mu}}$	$\mathbf{8} \otimes \mathbf{7}$	$\mathbf{8} \oplus \mathbf{48}$	$\psi_a^{(2)9}, \psi_a^{(2)\mu}$
			$\mathbf{8} \otimes \mathbf{1}$	$\mathbf{8}$	$\lambda_a^{(2)}$

sector	$SO(8)$	10d fields
(NS_+, NS_+)	$\mathbf{8}_v \otimes \mathbf{8}_v$	$\Phi, B_{[\hat{\mu}\hat{\nu}]}, G_{(\hat{\mu}\hat{\nu})}$
(R_+, R_-)	$\mathbf{8}_s \otimes \mathbf{8}_c$	$C_{\hat{\mu}}, C_{[\hat{\mu}\hat{\nu}\hat{\rho}]}$
(NS_+, R_-)	$\mathbf{8}_v \otimes \mathbf{8}_c$	$\tilde{\lambda}_a, \tilde{\psi}_a^{\hat{\mu}}$
(R_+, NS_+)	$\mathbf{8}_s \otimes \mathbf{8}_v$	$\lambda_a, \psi_a^{\hat{\mu}}$

TYPE II STRING: FIELD THEORETIC COMPACTIFICATION

Type IIA					
sector	$SO(8)$	10d fields	$SO(7)$	$SO(7)$ irrep	9d fields
(NS_+, NS_+)	$\mathbf{8}_v \otimes \mathbf{8}_v$	$\Phi, B_{[\hat{\mu}\hat{\nu}]}, G_{(\hat{\mu}\hat{\nu})}$	$\mathbf{7} \otimes \mathbf{7}$	$\mathbf{1} \oplus \mathbf{21} \oplus \mathbf{27}$	$\phi, B_{[\mu\nu]}, G_{(\mu\nu)}$
			$(\mathbf{7} \otimes \mathbf{1}) \oplus (\mathbf{1} \otimes \mathbf{7})$	$\mathbf{7} \oplus \mathbf{7}$	$G_{\mu 9}, B_{\mu 9}$
			$\mathbf{1} \otimes \mathbf{1}$	$\mathbf{1}$	G_{99}
(R_+, R_-)	$\mathbf{8}_s \otimes \mathbf{8}_c$	$C_{\hat{\mu}}, C_{[\hat{\mu}\hat{\nu}\hat{\rho}]}$	$\mathbf{8} \otimes \mathbf{8}$	$\mathbf{1} \oplus \mathbf{7} \oplus \mathbf{21} \oplus \mathbf{35}$	$A_9, A_\mu, C_{9\mu\nu}, C_{\mu\nu\rho}$
(NS_+, R_-)	$\mathbf{8}_v \otimes \mathbf{8}_c$	$\tilde{\lambda}_a, \tilde{\psi}_a^{\hat{\mu}}$	$\mathbf{7} \otimes \mathbf{8}$	$\mathbf{8} \oplus \mathbf{48}$	$\tilde{\psi}_a^\mu, \tilde{\psi}_a^9$
			$\mathbf{1} \otimes \mathbf{8}$	$\mathbf{8}$	$\tilde{\lambda}_a$
(R_+, NS_+)	$\mathbf{8}_s \otimes \mathbf{8}_v$	$\lambda_a, \psi_a^{\hat{\mu}}$	$\mathbf{8} \otimes \mathbf{7}$	$\mathbf{8} \oplus \mathbf{48}$	ψ_a^μ, ψ_a^9
			$\mathbf{8} \otimes \mathbf{1}$	$\mathbf{8}$	λ_a
Type IIB					
sector	$SO(8)$	10d fields	$SO(7)$	$SO(7)$ irrep	9d fields
(NS_+, NS_+)	$\mathbf{8}_v \otimes \mathbf{8}_v$	$\Phi, B_{[\hat{\mu}\hat{\nu}]}, G_{(\hat{\mu}\hat{\nu})}$	$\mathbf{7} \otimes \mathbf{7}$	$\mathbf{1} \oplus \mathbf{21} \oplus \mathbf{27}$	$\phi, B_{[\mu\nu]}, G_{(\mu\nu)}$
			$(\mathbf{7} \otimes \mathbf{1}) \oplus (\mathbf{1} \otimes \mathbf{7})$	$\mathbf{7} \oplus \mathbf{7}$	$G_{\mu 9}, B_{\mu 9}$
			$\mathbf{1} \otimes \mathbf{1}$	$\mathbf{1}$	G_{99}
(R_+, R_+)	$\mathbf{8}_s \otimes \mathbf{8}_s$	$C_0, C_{[\hat{\mu}\hat{\nu}]}, C_{[\hat{\mu}\hat{\nu}\hat{\rho}\hat{\sigma}]}^+$	$\mathbf{8} \otimes \mathbf{8}$	$\mathbf{1} \oplus \mathbf{7} \oplus \mathbf{21} \oplus \mathbf{35}$	$a, C_{\mu 9}, C_{\mu\nu}, C_{\mu\nu\rho 9}$
(NS_+, R_+)	$\mathbf{8}_v \otimes \mathbf{8}_s$	$\lambda_a^{(1)}, \psi_a^{(1)\hat{\mu}}$	$\mathbf{7} \otimes \mathbf{8}$	$\mathbf{8} \oplus \mathbf{48}$	$\psi_a^{(1)9}, \psi_a^{(1)\mu}$
			$\mathbf{1} \otimes \mathbf{8}$	$\mathbf{8}$	$\lambda_a^{(1)}$
(R_+, NS_+)	$\mathbf{8}_s \otimes \mathbf{8}_v$	$\lambda_a^{(2)}, \psi_a^{(2)\hat{\mu}}$	$\mathbf{8} \otimes \mathbf{7}$	$\mathbf{8} \oplus \mathbf{48}$	$\psi_a^{(2)9}, \psi_a^{(2)\mu}$
			$\mathbf{8} \otimes \mathbf{1}$	$\mathbf{8}$	$\lambda_a^{(2)}$

TYPE II STRING: FIELD THEORETIC COMPACTIFICATION

Type IIA					
sector	$SO(8)$	10d fields	$SO(7)$	$SO(7)$ irrep	9d fields
(NS_+, NS_+)	$\mathbf{8}_v \otimes \mathbf{8}_v$	$\Phi, B_{[\hat{\mu}\hat{\nu}]}, G_{(\hat{\mu}\hat{\nu})}$	$\mathbf{7} \otimes \mathbf{7}$	$\mathbf{1} \oplus \mathbf{21} \oplus \mathbf{27}$	$\phi, B_{[\mu\nu]}, G_{(\mu\nu)}$
			$(\mathbf{7} \otimes \mathbf{1}) \oplus (\mathbf{1} \otimes \mathbf{7})$	$\mathbf{7} \oplus \mathbf{7}$	$G_{\mu 9}, B_{\mu 9}$
			$\mathbf{1} \otimes \mathbf{1}$	$\mathbf{1}$	G_{99}
(R_+, R_-)	$\mathbf{8}_s \otimes \mathbf{8}_c$	$C_{\hat{\mu}}, C_{[\hat{\mu}\hat{\nu}\hat{\rho}]}$	$\mathbf{8} \otimes \mathbf{8}$	$\mathbf{1} \oplus \mathbf{7} \oplus \mathbf{21} \oplus \mathbf{35}$	$A_9, A_\mu, C_{9\mu\nu}, C_{\mu\nu\rho}$
(NS_+, R_-)	$\mathbf{8}_v \otimes \mathbf{8}_c$	$\tilde{\lambda}_a, \tilde{\psi}_a^{\hat{\mu}}$	$\mathbf{7} \otimes \mathbf{8}$	$\mathbf{8} \oplus \mathbf{48}$	$\tilde{\psi}_a^\mu, \tilde{\psi}_a^9$
			$\mathbf{1} \otimes \mathbf{8}$	$\mathbf{8}$	$\tilde{\lambda}_a$
(R_+, NS_+)	$\mathbf{8}_s \otimes \mathbf{8}_v$	$\lambda_a, \psi_a^{\hat{\mu}}$	$\mathbf{8} \otimes \mathbf{7}$	$\mathbf{8} \oplus \mathbf{48}$	ψ_a^μ, ψ_a^9
			$\mathbf{8} \otimes \mathbf{1}$	$\mathbf{8}$	λ_a
Type IIB					
sector	$SO(8)$	10d fields	$SO(7)$	$SO(7)$ irrep	9d fields
(NS_+, NS_+)	$\mathbf{8}_v \otimes \mathbf{8}_v$	$\Phi, B_{[\hat{\mu}\hat{\nu}]}, G_{(\hat{\mu}\hat{\nu})}$	$\mathbf{7} \otimes \mathbf{7}$	$\mathbf{1} \oplus \mathbf{21} \oplus \mathbf{27}$	$\phi, B_{[\mu\nu]}, G_{(\mu\nu)}$
			$(\mathbf{7} \otimes \mathbf{1}) \oplus (\mathbf{1} \otimes \mathbf{7})$	$\mathbf{7} \oplus \mathbf{7}$	$G_{\mu 9}, B_{\mu 9}$
			$\mathbf{1} \otimes \mathbf{1}$	$\mathbf{1}$	G_{99}
(R_+, R_+)	$\mathbf{8}_s \otimes \mathbf{8}_s$	$C_0, C_{[\hat{\mu}\hat{\nu}]}, C_{[\hat{\mu}\hat{\nu}\hat{\rho}\hat{\sigma}]}^+$	$\mathbf{8} \otimes \mathbf{8}$	$\mathbf{1} \oplus \mathbf{7} \oplus \mathbf{21} \oplus \mathbf{35}$	$a, C_{\mu 9}, C_{\mu\nu}, C_{\mu\nu\rho 9}$
(NS_+, R_+)	$\mathbf{8}_v \otimes \mathbf{8}_s$	$\lambda_a^{(1)}, \psi_a^{(1)\hat{\mu}}$	$\mathbf{7} \otimes \mathbf{8}$	$\mathbf{8} \oplus \mathbf{48}$	$\psi_a^{(1)9}, \psi_a^{(1)\mu}$
			$\mathbf{1} \otimes \mathbf{8}$	$\mathbf{8}$	$\lambda_a^{(1)}$
(R_+, NS_+)	$\mathbf{8}_s \otimes \mathbf{8}_v$	$\lambda_a^{(2)}, \psi_a^{(2)\hat{\mu}}$	$\mathbf{8} \otimes \mathbf{7}$	$\mathbf{8} \oplus \mathbf{48}$	$\psi_a^{(2)9}, \psi_a^{(2)\mu}$
			$\mathbf{8} \otimes \mathbf{1}$	$\mathbf{8}$	$\lambda_a^{(2)}$

Type IIA

$SO(7)$	$SO(7)$ irrep	9d fields
$\mathbf{7} \otimes \mathbf{7}$	$\mathbf{1} \oplus \mathbf{21} \oplus \mathbf{27}$	$\phi, B_{[\mu\nu]}, G_{(\mu\nu)}$
$(\mathbf{7} \otimes \mathbf{1}) \oplus (\mathbf{1} \otimes \mathbf{7})$	$\mathbf{7} \oplus \mathbf{7}$	$G_{\mu 9}, B_{\mu 9}$
$\mathbf{1} \otimes \mathbf{1}$	$\mathbf{1}$	G_{99}
$\mathbf{8} \otimes \mathbf{8}$	$\mathbf{1} \oplus \mathbf{7} \oplus \mathbf{21} \oplus \mathbf{35}$	$A_9, A_\mu, C_{9\mu\nu}, C_{\mu\nu\rho}$
$\mathbf{7} \otimes \mathbf{8}$	$\mathbf{8} \oplus \mathbf{48}$	$\tilde{\psi}_a^\mu, \tilde{\psi}_a^9$
$\mathbf{1} \otimes \mathbf{8}$	$\mathbf{8}$	$\tilde{\lambda}_a$
$\mathbf{8} \otimes \mathbf{7}$	$\mathbf{8} \oplus \mathbf{48}$	ψ_a^μ, ψ_a^9
$\mathbf{8} \otimes \mathbf{1}$	$\mathbf{8}$	λ_a

TYPE II STRING: FIELD THEORETIC COMPACTIFICATION

Type IIA					
sector	$SO(8)$	10d fields	$SO(7)$	$SO(7)$ irrep	9d fields
(NS_+, NS_+)	$\mathbf{8}_v \otimes \mathbf{8}_v$	$\Phi, B_{[\hat{\mu}\hat{\nu}]}, G_{(\hat{\mu}\hat{\nu})}$	$\mathbf{7} \otimes \mathbf{7}$	$\mathbf{1} \oplus \mathbf{21} \oplus \mathbf{27}$	$\phi, B_{[\mu\nu]}, G_{(\mu\nu)}$
			$(\mathbf{7} \otimes \mathbf{1}) \oplus (\mathbf{1} \otimes \mathbf{7})$	$\mathbf{7} \oplus \mathbf{7}$	$G_{\mu 9}, B_{\mu 9}$
			$\mathbf{1} \otimes \mathbf{1}$	$\mathbf{1}$	G_{99}
(R_+, R_-)	$\mathbf{8}_s \otimes \mathbf{8}_c$	$C_{\hat{\mu}}, C_{[\hat{\mu}\hat{\nu}\hat{\rho}]}$	$\mathbf{8} \otimes \mathbf{8}$	$\mathbf{1} \oplus \mathbf{7} \oplus \mathbf{21} \oplus \mathbf{35}$	$A_9, A_\mu, C_{9\mu\nu}, C_{\mu\nu\rho}$
(NS_+, R_-)	$\mathbf{8}_v \otimes \mathbf{8}_c$	$\tilde{\lambda}_a, \tilde{\psi}_a^{\hat{\mu}}$	$\mathbf{7} \otimes \mathbf{8}$	$\mathbf{8} \oplus \mathbf{48}$	$\tilde{\psi}_a^\mu, \tilde{\psi}_a^9$
			$\mathbf{1} \otimes \mathbf{8}$	$\mathbf{8}$	$\tilde{\lambda}_a$
(R_+, NS_+)	$\mathbf{8}_s \otimes \mathbf{8}_v$	$\lambda_a, \psi_a^{\hat{\mu}}$	$\mathbf{8} \otimes \mathbf{7}$	$\mathbf{8} \oplus \mathbf{48}$	ψ_a^μ, ψ_a^9
			$\mathbf{8} \otimes \mathbf{1}$	$\mathbf{8}$	λ_a
Type IIB					
sector	$SO(8)$	10d fields	$SO(7)$	$SO(7)$ irrep	9d fields
(NS_+, NS_+)	$\mathbf{8}_v \otimes \mathbf{8}_v$	$\Phi, B_{[\hat{\mu}\hat{\nu}]}, G_{(\hat{\mu}\hat{\nu})}$	$\mathbf{7} \otimes \mathbf{7}$	$\mathbf{1} \oplus \mathbf{21} \oplus \mathbf{27}$	$\phi, B_{[\mu\nu]}, G_{(\mu\nu)}$
			$(\mathbf{7} \otimes \mathbf{1}) \oplus (\mathbf{1} \otimes \mathbf{7})$	$\mathbf{7} \oplus \mathbf{7}$	$G_{\mu 9}, B_{\mu 9}$
			$\mathbf{1} \otimes \mathbf{1}$	$\mathbf{1}$	G_{99}
(R_+, R_+)	$\mathbf{8}_s \otimes \mathbf{8}_s$	$C_0, C_{[\hat{\mu}\hat{\nu}]}, C_{[\hat{\mu}\hat{\nu}\hat{\rho}\hat{\sigma}]}^+$	$\mathbf{8} \otimes \mathbf{8}$	$\mathbf{1} \oplus \mathbf{7} \oplus \mathbf{21} \oplus \mathbf{35}$	$a, C_{\mu 9}, C_{\mu\nu}, C_{\mu\nu\rho 9}$
(NS_+, R_+)	$\mathbf{8}_v \otimes \mathbf{8}_s$	$\lambda_a^{(1)}, \psi_a^{(1)\hat{\mu}}$	$\mathbf{7} \otimes \mathbf{8}$	$\mathbf{8} \oplus \mathbf{48}$	$\psi_a^{(1)9}, \psi_a^{(1)\mu}$
			$\mathbf{1} \otimes \mathbf{8}$	$\mathbf{8}$	$\lambda_a^{(1)}$
(R_+, NS_+)	$\mathbf{8}_s \otimes \mathbf{8}_v$	$\lambda_a^{(2)}, \psi_a^{(2)\hat{\mu}}$	$\mathbf{8} \otimes \mathbf{7}$	$\mathbf{8} \oplus \mathbf{48}$	$\psi_a^{(2)9}, \psi_a^{(2)\mu}$
			$\mathbf{8} \otimes \mathbf{1}$	$\mathbf{8}$	$\lambda_a^{(2)}$

TYPE II STRING: FIELD THEORETIC COMPACTIFICATION

Type IIA					
sector	$SO(8)$	10d fields	$SO(7)$	$SO(7)$ irrep	9d fields
(NS_+, NS_+)	$\mathbf{8}_v \otimes \mathbf{8}_v$	$\Phi, B_{[\hat{\mu}\hat{\nu}]}, G_{(\hat{\mu}\hat{\nu})}$	$\mathbf{7} \otimes \mathbf{7}$	$\mathbf{1} \oplus \mathbf{21} \oplus \mathbf{27}$	$\phi, B_{[\mu\nu]}, G_{(\mu\nu)}$
			$(\mathbf{7} \otimes \mathbf{1}) \oplus (\mathbf{1} \otimes \mathbf{7})$	$\mathbf{7} \oplus \mathbf{7}$	$G_{\mu 9}, B_{\mu 9}$
			$\mathbf{1} \otimes \mathbf{1}$	$\mathbf{1}$	G_{99}
(R_+, R_-)	$\mathbf{8}_s \otimes \mathbf{8}_c$	$C_{\hat{\mu}}, C_{[\hat{\mu}\hat{\nu}\hat{\rho}]}$	$\mathbf{8} \otimes \mathbf{8}$	$\mathbf{1} \oplus \mathbf{7} \oplus \mathbf{21} \oplus \mathbf{35}$	$A_9, A_\mu, C_{9\mu\nu}, C_{\mu\nu\rho}$
(NS_+, R_-)	$\mathbf{8}_v \otimes \mathbf{8}_c$	$\tilde{\lambda}_a, \tilde{\psi}_a^{\hat{\mu}}$	$\mathbf{7} \otimes \mathbf{8}$	$\mathbf{8} \oplus \mathbf{48}$	$\tilde{\psi}_a^\mu, \tilde{\psi}_a^9$
			$\mathbf{1} \otimes \mathbf{8}$	$\mathbf{8}$	$\tilde{\lambda}_a$
(R_+, NS_+)	$\mathbf{8}_s \otimes \mathbf{8}_v$	$\lambda_a, \psi_a^{\hat{\mu}}$	$\mathbf{8} \otimes \mathbf{7}$	$\mathbf{8} \oplus \mathbf{48}$	ψ_a^μ, ψ_a^9
			$\mathbf{8} \otimes \mathbf{1}$	$\mathbf{8}$	λ_a
Type IIB					
sector	$SO(8)$	10d fields	$SO(7)$	$SO(7)$ irrep	9d fields
(NS_+, NS_+)	$\mathbf{8}_v \otimes \mathbf{8}_v$	$\Phi, B_{[\hat{\mu}\hat{\nu}]}, G_{(\hat{\mu}\hat{\nu})}$	$\mathbf{7} \otimes \mathbf{7}$	$\mathbf{1} \oplus \mathbf{21} \oplus \mathbf{27}$	$\phi, B_{[\mu\nu]}, G_{(\mu\nu)}$
			$(\mathbf{7} \otimes \mathbf{1}) \oplus (\mathbf{1} \otimes \mathbf{7})$	$\mathbf{7} \oplus \mathbf{7}$	$G_{\mu 9}, B_{\mu 9}$
			$\mathbf{1} \otimes \mathbf{1}$	$\mathbf{1}$	G_{99}
(R_+, R_+)	$\mathbf{8}_s \otimes \mathbf{8}_s$	$C_0, C_{[\hat{\mu}\hat{\nu}]}, C_{[\hat{\mu}\hat{\nu}\hat{\rho}\hat{\sigma}]}^+$	$\mathbf{8} \otimes \mathbf{8}$	$\mathbf{1} \oplus \mathbf{7} \oplus \mathbf{21} \oplus \mathbf{35}$	$a, C_{\mu 9}, C_{\mu\nu}, C_{\mu\nu\rho 9}$
(NS_+, R_+)	$\mathbf{8}_v \otimes \mathbf{8}_s$	$\lambda_a^{(1)}, \psi_a^{(1)\hat{\mu}}$	$\mathbf{7} \otimes \mathbf{8}$	$\mathbf{8} \oplus \mathbf{48}$	$\psi_a^{(1)9}, \psi_a^{(1)\mu}$
			$\mathbf{1} \otimes \mathbf{8}$	$\mathbf{8}$	$\lambda_a^{(1)}$
(R_+, NS_+)	$\mathbf{8}_s \otimes \mathbf{8}_v$	$\lambda_a^{(2)}, \psi_a^{(2)\hat{\mu}}$	$\mathbf{8} \otimes \mathbf{7}$	$\mathbf{8} \oplus \mathbf{48}$	$\psi_a^{(2)9}, \psi_a^{(2)\mu}$
			$\mathbf{8} \otimes \mathbf{1}$	$\mathbf{8}$	$\lambda_a^{(2)}$

sector	$SO(8)$	10d fields
(NS_+, NS_+)	$\mathbf{8}_v \otimes \mathbf{8}_v$	$\Phi, B_{[\hat{\mu}\hat{\nu}]}, G_{(\hat{\mu}\hat{\nu})}$
(R_+, R_+)	$\mathbf{8}_s \otimes \mathbf{8}_s$	$C_0, C_{[\hat{\mu}\hat{\nu}]}, C_{[\hat{\mu}\hat{\nu}\hat{\rho}\hat{\sigma}]}^+$
(NS_+, R_+)	$\mathbf{8}_v \otimes \mathbf{8}_s$	$\lambda_a^{(1)}, \psi_a^{(1)\hat{\mu}}$
(R_+, NS_+)	$\mathbf{8}_s \otimes \mathbf{8}_v$	$\lambda_a^{(2)}, \psi_a^{(2)\hat{\mu}}$

TYPE II STRING: FIELD THEORETIC COMPACTIFICATION

Type IIA					
sector	$SO(8)$	10d fields	$SO(7)$	$SO(7)$ irrep	9d fields
(NS_+, NS_+)	$\mathbf{8}_v \otimes \mathbf{8}_v$	$\Phi, B_{[\hat{\mu}\hat{\nu}]}, G_{(\hat{\mu}\hat{\nu})}$	$\mathbf{7} \otimes \mathbf{7}$	$\mathbf{1} \oplus \mathbf{21} \oplus \mathbf{27}$	$\phi, B_{[\mu\nu]}, G_{(\mu\nu)}$
			$(\mathbf{7} \otimes \mathbf{1}) \oplus (\mathbf{1} \otimes \mathbf{7})$	$\mathbf{7} \oplus \mathbf{7}$	$G_{\mu 9}, B_{\mu 9}$
			$\mathbf{1} \otimes \mathbf{1}$	$\mathbf{1}$	G_{99}
(R_+, R_-)	$\mathbf{8}_s \otimes \mathbf{8}_c$	$C_{\hat{\mu}}, C_{[\hat{\mu}\hat{\nu}\hat{\rho}]}$	$\mathbf{8} \otimes \mathbf{8}$	$\mathbf{1} \oplus \mathbf{7} \oplus \mathbf{21} \oplus \mathbf{35}$	$A_9, A_\mu, C_{9\mu\nu}, C_{\mu\nu\rho}$
(NS_+, R_-)	$\mathbf{8}_v \otimes \mathbf{8}_c$	$\tilde{\lambda}_a, \tilde{\psi}_a^{\hat{\mu}}$	$\mathbf{7} \otimes \mathbf{8}$	$\mathbf{8} \oplus \mathbf{48}$	$\tilde{\psi}_a^\mu, \tilde{\psi}_a^9$
			$\mathbf{1} \otimes \mathbf{8}$	$\mathbf{8}$	$\tilde{\lambda}_a$
(R_+, NS_+)	$\mathbf{8}_s \otimes \mathbf{8}_v$	$\lambda_a, \psi_a^{\hat{\mu}}$	$\mathbf{8} \otimes \mathbf{7}$	$\mathbf{8} \oplus \mathbf{48}$	ψ_a^μ, ψ_a^9
			$\mathbf{8} \otimes \mathbf{1}$	$\mathbf{8}$	λ_a
Type IIB					
sector	$SO(8)$	10d fields	$SO(7)$	$SO(7)$ irrep	9d fields
(NS_+, NS_+)	$\mathbf{8}_v \otimes \mathbf{8}_v$	$\Phi, B_{[\hat{\mu}\hat{\nu}]}, G_{(\hat{\mu}\hat{\nu})}$	$\mathbf{7} \otimes \mathbf{7}$	$\mathbf{1} \oplus \mathbf{21} \oplus \mathbf{27}$	$\phi, B_{[\mu\nu]}, G_{(\mu\nu)}$
			$(\mathbf{7} \otimes \mathbf{1}) \oplus (\mathbf{1} \otimes \mathbf{7})$	$\mathbf{7} \oplus \mathbf{7}$	$G_{\mu 9}, B_{\mu 9}$
			$\mathbf{1} \otimes \mathbf{1}$	$\mathbf{1}$	G_{99}
(R_+, R_+)	$\mathbf{8}_s \otimes \mathbf{8}_s$	$C_0, C_{[\hat{\mu}\hat{\nu}]}, C_{[\hat{\mu}\hat{\nu}\hat{\rho}\hat{\sigma}]}^+$	$\mathbf{8} \otimes \mathbf{8}$	$\mathbf{1} \oplus \mathbf{7} \oplus \mathbf{21} \oplus \mathbf{35}$	$a, C_{\mu 9}, C_{\mu\nu}, C_{\mu\nu\rho 9}$
(NS_+, R_+)	$\mathbf{8}_v \otimes \mathbf{8}_s$	$\lambda_a^{(1)}, \psi_a^{(1)\hat{\mu}}$	$\mathbf{7} \otimes \mathbf{8}$	$\mathbf{8} \oplus \mathbf{48}$	$\psi_a^{(1)9}, \psi_a^{(1)\mu}$
			$\mathbf{1} \otimes \mathbf{8}$	$\mathbf{8}$	$\lambda_a^{(1)}$
(R_+, NS_+)	$\mathbf{8}_s \otimes \mathbf{8}_v$	$\lambda_a^{(2)}, \psi_a^{(2)\hat{\mu}}$	$\mathbf{8} \otimes \mathbf{7}$	$\mathbf{8} \oplus \mathbf{48}$	$\psi_a^{(2)9}, \psi_a^{(2)\mu}$
			$\mathbf{8} \otimes \mathbf{1}$	$\mathbf{8}$	$\lambda_a^{(2)}$

TYPE II STRING: FIELD THEORETIC COMPACTIFICATION

Type IIA					
sector	$SO(8)$	10d fields	$SO(7)$	$SO(7)$ irrep	9d fields
(NS_+, NS_+)	$\mathbf{8}_v \otimes \mathbf{8}_v$	$\Phi, B_{[\hat{\mu}\hat{\nu}]}, G_{(\hat{\mu}\hat{\nu})}$	$\mathbf{7} \otimes \mathbf{7}$	$\mathbf{1} \oplus \mathbf{21} \oplus \mathbf{27}$	$\phi, B_{[\mu\nu]}, G_{(\mu\nu)}$
			$(\mathbf{7} \otimes \mathbf{1}) \oplus (\mathbf{1} \otimes \mathbf{7})$	$\mathbf{7} \oplus \mathbf{7}$	$G_{\mu 9}, B_{\mu 9}$
			$\mathbf{1} \otimes \mathbf{1}$	$\mathbf{1}$	G_{99}
(R_+, R_-)	$\mathbf{8}_s \otimes \mathbf{8}_c$	$C_{\hat{\mu}}, C_{[\hat{\mu}\hat{\nu}\hat{\rho}]}$	$\mathbf{8} \otimes \mathbf{8}$	$\mathbf{1} \oplus \mathbf{7} \oplus \mathbf{21} \oplus \mathbf{35}$	$A_9, A_\mu, C_{9\mu\nu}, C_{\mu\nu\rho}$
(NS_+, R_-)	$\mathbf{8}_v \otimes \mathbf{8}_c$	$\tilde{\lambda}_a, \tilde{\psi}_a^{\hat{\mu}}$	$\mathbf{7} \otimes \mathbf{8}$	$\mathbf{8} \oplus \mathbf{48}$	$\tilde{\psi}_a^\mu, \tilde{\psi}_a^9$
			$\mathbf{1} \otimes \mathbf{8}$	$\mathbf{8}$	$\tilde{\lambda}_a$
(R_+, NS_+)	$\mathbf{8}_s \otimes \mathbf{8}_v$	$\lambda_a, \psi_a^{\hat{\mu}}$	$\mathbf{8} \otimes \mathbf{7}$	$\mathbf{8} \oplus \mathbf{48}$	ψ_a^μ, ψ_a^9
			$\mathbf{8} \otimes \mathbf{1}$	$\mathbf{8}$	λ_a
Type IIB					
sector	$SO(8)$	10d fields	$SO(7)$	$SO(7)$ irrep	9d fields
(NS_+, NS_+)	$\mathbf{8}_v \otimes \mathbf{8}_v$	$\Phi, B_{[\hat{\mu}\hat{\nu}]}, G_{(\hat{\mu}\hat{\nu})}$	$\mathbf{7} \otimes \mathbf{7}$	$\mathbf{1} \oplus \mathbf{21} \oplus \mathbf{27}$	$\phi, B_{[\mu\nu]}, G_{(\mu\nu)}$
			$(\mathbf{7} \otimes \mathbf{1}) \oplus (\mathbf{1} \otimes \mathbf{7})$	$\mathbf{7} \oplus \mathbf{7}$	$G_{\mu 9}, B_{\mu 9}$
			$\mathbf{1} \otimes \mathbf{1}$	$\mathbf{1}$	G_{99}
(R_+, R_+)	$\mathbf{8}_s \otimes \mathbf{8}_s$	$C_0, C_{[\hat{\mu}\hat{\nu}]}, C_{[\hat{\mu}\hat{\nu}\hat{\rho}\hat{\sigma}]}^+$	$\mathbf{8} \otimes \mathbf{8}$	$\mathbf{1} \oplus \mathbf{7} \oplus \mathbf{21} \oplus \mathbf{35}$	$a, C_{\mu 9}, C_{\mu\nu}, C_{\mu\nu\rho 9}$
(NS_+, R_+)	$\mathbf{8}_v \otimes \mathbf{8}_s$	$\lambda_a^{(1)}, \psi_a^{(1)\hat{\mu}}$	$\mathbf{7} \otimes \mathbf{8}$	$\mathbf{8} \oplus \mathbf{48}$	$\psi_a^{(1)9}, \psi_a^{(1)\mu}$
			$\mathbf{1} \otimes \mathbf{8}$	$\mathbf{8}$	$\lambda_a^{(1)}$
(R_+, NS_+)	$\mathbf{8}_s \otimes \mathbf{8}_v$	$\lambda_a^{(2)}, \psi_a^{(2)\hat{\mu}}$	$\mathbf{8} \otimes \mathbf{7}$	$\mathbf{8} \oplus \mathbf{48}$	$\psi_a^{(2)9}, \psi_a^{(2)\mu}$
			$\mathbf{8} \otimes \mathbf{1}$	$\mathbf{8}$	$\lambda_a^{(2)}$

Type IIB

$SO(7)$	$SO(7)$ irrep	9d fields
$\mathbf{7} \otimes \mathbf{7}$	$\mathbf{1} \oplus \mathbf{21} \oplus \mathbf{27}$	$\phi, B_{[\mu\nu]}, G_{(\mu\nu)}$
$(\mathbf{7} \otimes \mathbf{1}) \oplus (\mathbf{1} \otimes \mathbf{7})$	$\mathbf{7} \oplus \mathbf{7}$	$G_{\mu 9}, B_{\mu 9}$
$\mathbf{1} \otimes \mathbf{1}$	$\mathbf{1}$	G_{99}
$\mathbf{8} \otimes \mathbf{8}$	$\mathbf{1} \oplus \mathbf{7} \oplus \mathbf{21} \oplus \mathbf{35}$	$a, C_{\mu 9}, C_{\mu\nu}, C_{\mu\nu\rho 9}$
$\mathbf{7} \otimes \mathbf{8}$	$\mathbf{8} \oplus \mathbf{48}$	$\psi_a^{(1)9}, \psi_a^{(1)\mu}$
$\mathbf{1} \otimes \mathbf{8}$	$\mathbf{8}$	$\lambda_a^{(1)}$
$\mathbf{8} \otimes \mathbf{7}$	$\mathbf{8} \oplus \mathbf{48}$	$\psi_a^{(2)9}, \psi_a^{(2)\mu}$
$\mathbf{8} \otimes \mathbf{1}$	$\mathbf{8}$	$\lambda_a^{(2)}$

TYPE II STRING: FIELD THEORETIC COMPACTIFICATION

Type IIA					
sector	$SO(8)$	10d fields	$SO(7)$	$SO(7)$ irrep	9d fields
$(\text{NS}_+, \text{NS}_+)$	$\mathbf{8}_v \otimes \mathbf{8}_v$	$\Phi, B_{[\hat{\mu}\hat{\nu}]}, G_{(\hat{\mu}\hat{\nu})}$	$\mathbf{7} \otimes \mathbf{7}$	$\mathbf{1} \oplus \mathbf{21} \oplus \mathbf{27}$	$\phi, B_{[\mu\nu]}, G_{(\mu\nu)}$
			$(\mathbf{7} \otimes \mathbf{1}) \oplus (\mathbf{1} \otimes \mathbf{7})$	$\mathbf{7} \oplus \mathbf{7}$	$G_{\mu 9}, B_{\mu 9}$
			$\mathbf{1} \otimes \mathbf{1}$	$\mathbf{1}$	G_{99}
(R_+, R_-)	$\mathbf{8}_s \otimes \mathbf{8}_c$	$C_{\hat{\mu}}, C_{[\hat{\mu}\hat{\nu}\hat{\rho}]}$	$\mathbf{8} \otimes \mathbf{8}$	$\mathbf{1} \oplus \mathbf{7} \oplus \mathbf{21} \oplus \mathbf{35}$	$A_9, A_\mu, C_{9\mu\nu}, C_{\mu\nu\rho}$
$(\text{NS}_+, \text{R}_-)$	$\mathbf{8}_v \otimes \mathbf{8}_c$	$\tilde{\lambda}_a, \tilde{\psi}_a^{\hat{\mu}}$	$\mathbf{7} \otimes \mathbf{8}$	$\mathbf{8} \oplus \mathbf{48}$	$\tilde{\psi}_a^\mu, \tilde{\psi}_a^9$
			$\mathbf{1} \otimes \mathbf{8}$	$\mathbf{8}$	$\tilde{\lambda}_a$
$(\text{R}_+, \text{NS}_+)$	$\mathbf{8}_s \otimes \mathbf{8}_v$	$\lambda_a, \psi_a^{\hat{\mu}}$	$\mathbf{8} \otimes \mathbf{7}$	$\mathbf{8} \oplus \mathbf{48}$	ψ_a^μ, ψ_a^9
			$\mathbf{8} \otimes \mathbf{1}$	$\mathbf{8}$	λ_a
Type IIB					
sector	$SO(8)$	10d fields	$SO(7)$	$SO(7)$ irrep	9d fields
$(\text{NS}_+, \text{NS}_+)$	$\mathbf{8}_v \otimes \mathbf{8}_v$	$\Phi, B_{[\hat{\mu}\hat{\nu}]}, G_{(\hat{\mu}\hat{\nu})}$	$\mathbf{7} \otimes \mathbf{7}$	$\mathbf{1} \oplus \mathbf{21} \oplus \mathbf{27}$	$\phi, B_{[\mu\nu]}, G_{(\mu\nu)}$
			$(\mathbf{7} \otimes \mathbf{1}) \oplus (\mathbf{1} \otimes \mathbf{7})$	$\mathbf{7} \oplus \mathbf{7}$	$G_{\mu 9}, B_{\mu 9}$
			$\mathbf{1} \otimes \mathbf{1}$	$\mathbf{1}$	G_{99}
(R_+, R_+)	$\mathbf{8}_s \otimes \mathbf{8}_s$	$C_0, C_{[\hat{\mu}\hat{\nu}]}, C_{[\hat{\mu}\hat{\nu}\hat{\rho}\hat{\sigma}]}^+$	$\mathbf{8} \otimes \mathbf{8}$	$\mathbf{1} \oplus \mathbf{7} \oplus \mathbf{21} \oplus \mathbf{35}$	$a, C_{\mu 9}, C_{\mu\nu}, C_{\mu\nu\rho 9}$
$(\text{NS}_+, \text{R}_+)$	$\mathbf{8}_v \otimes \mathbf{8}_s$	$\lambda_a^{(1)}, \psi_a^{(1)\hat{\mu}}$	$\mathbf{7} \otimes \mathbf{8}$	$\mathbf{8} \oplus \mathbf{48}$	$\psi_a^{(1)9}, \psi_a^{(1)\mu}$
			$\mathbf{1} \otimes \mathbf{8}$	$\mathbf{8}$	$\lambda_a^{(1)}$
$(\text{R}_+, \text{NS}_+)$	$\mathbf{8}_s \otimes \mathbf{8}_v$	$\lambda_a^{(2)}, \psi_a^{(2)\hat{\mu}}$	$\mathbf{8} \otimes \mathbf{7}$	$\mathbf{8} \oplus \mathbf{48}$	$\psi_a^{(2)9}, \psi_a^{(2)\mu}$
			$\mathbf{8} \otimes \mathbf{1}$	$\mathbf{8}$	$\lambda_a^{(2)}$

IS THIS A COINCIDENCE?

IS THIS A COINCIDENCE?
IS THIS A FEATURE OF THE MASSLESS SPECTRUM?

Setting and Main Results

Lightcone quantization:

- $\mathcal{M}_{10} \rightarrow \mathcal{M}_9 \times S^1, X^9 \simeq X^9 + 2\pi R;$
- $X^i(\tau, \sigma + l) = X^i(\tau, \sigma);$

Setting and Main Results

Lightcone quantization:

- $\mathcal{M}_{10} \rightarrow \mathcal{M}_9 \times S^1, X^9 \simeq X^9 + 2\pi R;$
- $X^i(\tau, \sigma + l) = X^i(\tau, \sigma);$
- $X^9(\tau, \sigma + l) = X^9(\tau, \sigma) + 2\pi R\omega;$
- $p_9 = s/R, s \in \mathbb{Z}.$

Setting and Main Results

Lightcone quantization:

- $\mathcal{M}_{10} \rightarrow \mathcal{M}_9 \times S^1, X^9 \simeq X^9 + 2\pi R;$
- $X^i(\tau, \sigma + l) = X^i(\tau, \sigma);$
- $X^9(\tau, \sigma + l) = X^9(\tau, \sigma) + 2\pi R\omega;$
- $p_9 = s/R, s \in \mathbb{Z}.$

Mode decomposition:

- $X_{L/R}^9(\xi^\pm) = \frac{x^9}{2} + \frac{\alpha'\pi}{l} p_{L/R} \xi^\pm + i\sqrt{\frac{\alpha'}{2}} \sum_{n \neq 0} \frac{\alpha_n^9}{n} e^{-\frac{2\pi i}{l} n \xi^\pm};$
- $p_L = \left(\frac{s}{R} + \frac{\omega R}{\alpha'}\right), p_R = \left(\frac{s}{R} - \frac{\omega R}{\alpha'}\right);$

Setting and Main Results

Lightcone quantization:

- $\mathcal{M}_{10} \rightarrow \mathcal{M}_9 \times S^1, X^9 \simeq X^9 + 2\pi R;$
- $X^i(\tau, \sigma + l) = X^i(\tau, \sigma);$
- $X^9(\tau, \sigma + l) = X^9(\tau, \sigma) + 2\pi R\omega;$
- $p_9 = s/R, s \in \mathbb{Z}.$

Mode decomposition:

- $X_{L/R}^9(\xi^\pm) = \frac{x^9}{2} + \frac{\alpha'\pi}{l} p_{L/R} \xi^\pm + i\sqrt{\frac{\alpha'}{2}} \sum_{n \neq 0} \frac{\alpha_n^9}{n} e^{-\frac{2\pi i}{l} n \xi^\pm};$
- $p_L = \left(\frac{s}{R} + \frac{\omega R}{\alpha'}\right), p_R = \left(\frac{s}{R} - \frac{\omega R}{\alpha'}\right);$
- $M_L^2 = \frac{p_L^2}{2} + \frac{2}{\alpha'} \left(\tilde{N}_\perp - \tilde{a}\right), \quad M_R^2 = \frac{p_R^2}{2} + \frac{2}{\alpha'} (N_\perp - a);$
- $M_L^2 = M_R^2, M^2 = M_L^2 + M_R^2.$

T-DUALITY: INTRODUCTION

Mass-Shell Condition

$$M^2 = \frac{s^2}{R^2} + \frac{\omega^2 R^2}{\alpha'^2} + \frac{2}{\alpha'}(\tilde{N}_\perp + N_\perp - a - \tilde{a}),$$
$$p_L = \left(\frac{s}{R} + \frac{\omega R}{\alpha'} \right), \quad p_R = \left(\frac{s}{R} - \frac{\omega R}{\alpha'} \right).$$

Mass-Shell Condition

$$M^2 = \frac{s^2}{R^2} + \frac{\omega^2 R^2}{\alpha'^2} + \frac{2}{\alpha'}(\tilde{N}_\perp + N_\perp - a - \tilde{a}),$$
$$p_L = \left(\frac{s}{R} + \frac{\omega R}{\alpha'} \right), \quad p_R = \left(\frac{s}{R} - \frac{\omega R}{\alpha'} \right).$$

From Spectrum

$$R \rightarrow R' = \alpha' / R,$$
$$(s, \omega) \rightarrow (s', \omega') = (\omega, s).$$

Mass-Shell Condition

$$M^2 = \frac{s^2}{R^2} + \frac{\omega^2 R^2}{\alpha'^2} + \frac{2}{\alpha'}(\tilde{N}_\perp + N_\perp - a - \tilde{a}),$$
$$p_L = \left(\frac{s}{R} + \frac{\omega R}{\alpha'} \right), \quad p_R = \left(\frac{s}{R} - \frac{\omega R}{\alpha'} \right).$$

From Spectrum

$$R \rightarrow R' = \alpha'/R,$$
$$(s, \omega) \rightarrow (s', \omega') = (\omega, s).$$

From Full Theory

$$p_L \rightarrow p_L, \quad p_R \rightarrow -p_R,$$
$$X_L^9 \rightarrow X_L^9, \quad X_R^9 \rightarrow -X_R^9.$$

Mass-Shell Condition

$$M^2 = \frac{s^2}{R^2} + \frac{\omega^2 R^2}{\alpha'^2} + \frac{2}{\alpha'}(\tilde{N}_\perp + N_\perp - a - \tilde{a}),$$
$$p_L = \left(\frac{s}{R} + \frac{\omega R}{\alpha'} \right), \quad p_R = \left(\frac{s}{R} - \frac{\omega R}{\alpha'} \right).$$

From Spectrum

$$R \rightarrow R' = \alpha'/R,$$
$$(s, \omega) \rightarrow (s', \omega') = (\omega, s).$$

From Full Theory

$$p_L \rightarrow p_L, \quad p_R \rightarrow -p_R,$$
$$X_L^9 \rightarrow X_L^9, \quad X_R^9 \rightarrow -X_R^9.$$

from Worldsheet Supersymmetry

$$\psi_L^9 \rightarrow \psi_L^9, \quad \psi_R^9 \rightarrow -\psi_R^9.$$

T-Duality

Transformation:

$$\begin{aligned}X^9(\tau, \sigma) &= X_L^9(\xi^+) + X_R^9(\xi^-) \rightarrow X'^9(\tau, \sigma) = X_L^9(\xi^+) - X_R^9(\xi^-), \\ \psi^9(\tau, \sigma) &= \psi_L^9(\xi^+) + \psi_R^9(\xi^-) \rightarrow \psi'^9(\tau, \sigma) = \psi_L^9(\xi^+) - \psi_R^9(\xi^-).\end{aligned}$$

T-Duality

Transformation:

$$\begin{aligned}X^9(\tau, \sigma) &= X_L^9(\xi^+) + X_R^9(\xi^-) \rightarrow X'^9(\tau, \sigma) = X_L^9(\xi^+) - X_R^9(\xi^-), \\ \psi^9(\tau, \sigma) &= \psi_L^9(\xi^+) + \psi_R^9(\xi^-) \rightarrow \psi'^9(\tau, \sigma) = \psi_L^9(\xi^+) - \psi_R^9(\xi^-).\end{aligned}$$

On the modes:

$$\begin{aligned}\tilde{b}_r^9 &\rightarrow \tilde{b}'^9_r = \tilde{b}_r^9, & b_r^9 &\rightarrow b'^9_r = -b_r^9, \\ B_4^\pm &= \frac{1}{\sqrt{2}} (b_0^8 \pm i b_0^9) \rightarrow \frac{1}{\sqrt{2}} (b_0^8 \mp i b_0^9) = B_4^\mp.\end{aligned}$$

T-Duality

Transformation:

$$\begin{aligned} X^9(\tau, \sigma) &= X_L^9(\xi^+) + X_R^9(\xi^-) \rightarrow X'^9(\tau, \sigma) = X_L^9(\xi^+) - X_R^9(\xi^-), \\ \psi^9(\tau, \sigma) &= \psi_L^9(\xi^+) + \psi_R^9(\xi^-) \rightarrow \psi'^9(\tau, \sigma) = \psi_L^9(\xi^+) - \psi_R^9(\xi^-). \end{aligned}$$

On the modes:

$$\begin{aligned} \tilde{b}_r^9 &\rightarrow \tilde{b}'^9_r = \tilde{b}_r^9, & b_r^9 &\rightarrow b'^9_r = -b_r^9, \\ B_4^\pm &= \frac{1}{\sqrt{2}} (b_0^8 \pm i b_0^9) \rightarrow \frac{1}{\sqrt{2}} (b_0^8 \mp i b_0^9) = B_4^\mp. \end{aligned}$$

On the sectors:

$$\begin{aligned} (R_+, R_\pm) &\rightarrow (R_+, R_\mp), \\ (NS_+, R_\pm) &\rightarrow (NS_+, R_\mp). \end{aligned}$$

Type IIB on S^1 with $R \cong$ Type IIA on S'^1 with $R' = \frac{\alpha'}{R}$.

$$\begin{aligned}X^9(\tau, \sigma) &= X_L^9(\xi^+) + X_R^9(\xi^-), \\X'^9(\tau, \sigma) &= X_L^9(\xi^+) - X_R^9(\xi^-).\end{aligned}$$

THEN, WHAT REALLY IS SPACETIME?