



HIGH-PROTEIN

RECIPE COLLECTION

Clinical & behavioural systems for satiety, muscle preservation and practical kitchen mastery.

Revised & expanded edition · 2026

J.C. STORGANAU

Contents

Foreword: The 2026 Dietary Reset 3

1. The Biology of Satiety 5

2. Decoupling Muscle Protein Synthesis from the “30-Gram Limit” 7

3. The Brain Connection: Leidy’s fMRI Findings and the Breakfast Anchor 9

4. Navigating 2026’s Biggest Misconceptions and Friction Points 11

5. Comparative Chemistry of Whey and Plant-Based Proteins 13

6. The “Ingredient Prep” Kitchen System and Rapid Recipes 15

7. Your 5-Stage Transformation Roadmap 19

Appendix: Key Takeaways, Quick Start and References 21

HOW TO USE THIS BOOK

New to high-protein eating? Read the Foreword, then jump to Chapter 6 and cook one recipe from each section. **Want the science first?** Read Chapters 1 to 3 in order. **Short on time?** The “In short” box that closes each chapter gives the whole idea in four bullets, and the Appendix has a one-page 7-day quick start.

Important: the information in this book is based on scientific research and is provided for informational purposes only. It is not medical advice, diagnosis or treatment. Please consult a qualified healthcare professional before making health-related changes, particularly if you have a medical condition, are pregnant, take medication (including GLP-1 therapies), or have a history of disordered eating. The author and publisher disclaim liability for outcomes resulting from use of this information.

FOREWORD

Foreword: The 2026 Dietary Reset


Why the protein conversation changed, what the new guidance says, and how to turn it into dinner.

In January 2026, US federal health officials released the **Dietary Guidelines for Americans, 2025–2030**. The headline change was visual: the familiar MyPlate graphic gave way to a food pyramid turned upside down. Whole, nutrient-dense protein foods and healthy fats now sit at the broad top of the graphic, vegetables and fruit alongside them, while refined carbohydrates, added sugars and highly processed foods are pushed toward the narrow tip.

1.2–1.6
g protein per kg body weight per day now suggested for adults

0.8
g/kg, the long-standing RDA: the minimum to avoid deficiency, not an optimal target

Jan 2026
release of the 2025–2030 Dietary Guidelines



The 2025–2030 “inverted pyramid”, simplified. Consult the official guidelines for the full graphic and serving guidance.

The reasoning is that the old 0.8 g/kg figure was set to prevent deficiency in sedentary adults. It was never designed to protect muscle, support healthy ageing, or help manage appetite and blood sugar. A higher intake range of 1.2 to 1.6 g/kg/day is now suggested, and the guidelines encourage getting that protein mainly from whole foods.

THE DEBATE, FAIRLY STATED

Not everyone agrees on the details. Cardiometabolic organisations welcome the focus on whole foods but caution that a rapid rise in red meat and saturated fat could raise cardiovascular risk unless it is balanced with fibre, fish, legumes and unsaturated fats. This book takes the middle path: **protein-forward, plant-rich, and fibre-aware**. You will see fish, poultry, dairy and plant proteins far more often than red meat.

Find your daily target

Multiply your body weight in kilograms by 1.2 (lower end) and 1.6 (upper end). If your weight is well above a healthy range, a dietitian may suggest basing the calculation on a goal weight instead.

Body weight	Daily target (1.2–1.6 g/kg)	Per meal across 4 eating occasions (≈0.4 g/kg)
60 kg (132 lb)	72–96 g	≈ 24 g
70 kg (154 lb)	84–112 g	≈ 28 g
80 kg (176 lb)	96–128 g	≈ 32 g
90 kg (198 lb)	108–144 g	≈ 36 g
100 kg (220 lb)	120–160 g	≈ 40 g

Many adults land comfortably at 25 to 40 g per meal, which is the range this book is built around. Between the rise of appetite-suppressing GLP-1 medications and conflicting social media advice, hitting that range without spending hours in the kitchen or eating dry, joyless food is the real challenge. That is what the rest of this manual solves.


CHAPTER 1

The Biology of Satiety

Hunger is a hormonal signal, not a character flaw. Protein is the macronutrient that speaks to that system most clearly.

The endocrine control loop

Many weight-management plans fail because they treat hunger as a willpower problem. In reality, satiety is run by a coordinated network of gut hormones and brain centres, and it responds strongly to what is on your plate. When you eat protein, amino acids and small peptides act on specialised cells lining the gut, which send three kinds of signal to the brainstem and hypothalamus.



How dietary protein shapes appetite signalling (simplified).

Signal	Where it comes from	What it does
GLP-1 and PYY	L-cells in the lower gut	Slow gastric emptying and prolong the physical feeling of fullness.
CCK (cholecystokinin)	I-cells in the upper small intestine	Rapid “I’m full” message to the brain via the vagus nerve.
Ghrelin (reduced)	Oxyntic cells in the stomach	Protein tends to lower this main hunger hormone, quietening “food noise” between meals.


WHY THIS MATTERS IF YOU TAKE A GLP-1 MEDICATION

GLP-1-based medicines work on these same pathways, which is why they are so effective at reducing appetite. Approved options now include oral semaglutide, orforglipron and tirzepatide, while newer agents such as retatrutide and CagriSema are in late-stage trials or under regulatory review (check current status). When appetite drops sharply, **protein becomes the nutrient most worth protecting**: a protein-first approach helps limit the loss of lean muscle that can accompany rapid weight loss. Agree your targets with the clinician who prescribes your medication.

The thermic effect of food

Protein has a second, quieter advantage. Digesting, absorbing and processing food costs energy, a cost known as the **Thermic Effect of Food (TEF)**, or diet-induced thermogenesis.

$$TEF\ (\%) = (\text{energy spent processing a food} \div \text{energy the food provides}) \times 100$$



Typical thermic effect by macronutrient. Ranges vary between studies and individuals.

So for every 100 calories of protein eaten, roughly 20 to 30 are spent on processing, against 5 to 10 for carbohydrate and 0 to 3 for fat. The liver does much of that work: breaking peptide bonds, converting surplus amino acids to urea, and driving gluconeogenesis, all of which cost ATP.

KEEP IT IN PROPORTION

Across a mixed diet TEF accounts for only about 10% of daily energy use, so a protein-rich diet adds a modest boost, not a metabolic miracle. Its value is cumulative: a small energy edge, plus better fullness and better preservation of resting energy expenditure while you eat less.

IN SHORT

- Protein triggers GLP-1, PYY and CCK and tends to lower ghrelin, so it keeps you full longer.
- Processing protein costs 20–30% of its calories, versus 5–10% for carbs and 0–3% for fat.
- The thermic effect is a modest bonus, not a weight-loss trick.
- If you use appetite-suppressing medication, protein is the nutrient most worth protecting.

CHAPTER 2

Decoupling Muscle Protein Synthesis from the “30-Gram Limit”


Your gut can absorb far more than 30 g in one meal. What really matters is how often your muscles get a strong signal to build.

A persistent fitness-culture claim says your body can use only 30 g of protein per meal and wastes the rest. The claim confuses two separate processes: **digestive absorption** and **muscle protein synthesis (MPS)**.

	The myth	The reality
Absorption	Anything beyond 30 g is excreted.	Your gut absorbs almost everything. A big meal is simply digested more slowly, over several hours.
Muscle building	30 g is a hard ceiling.	Each meal has a diminishing-returns point for switching on MPS. Extra amino acids are still used, mostly for energy and other tissues.
Strategy	Eat as much as you like at dinner.	Spreading protein across the day gives repeated muscle-building signals.

Even beats skewed

The strongest evidence comes from a crossover trial by Mamerow and colleagues, published in *The Journal of Nutrition*. Volunteers ate the same 90 g of protein and the same calories per day, either **skewed** toward dinner (roughly 10, 16 and 63 g) or **evenly spread** (roughly 30 g at each meal). Using tracer infusions and muscle biopsies, the researchers found 24-hour muscle protein synthesis was about **25% higher** on the even pattern.



Same 90 g/day in both patterns. Even distribution gave ≈25% higher 24-hour muscle protein synthesis (Mamerow et al., 2014).

READ THE STUDY IN CONTEXT

The participants were healthy adults eating a controlled 90 g/day, not athletes or older adults, and the trial was short. It shows that *distribution* can matter, not that every meal must be exactly 30 g. Treat it as a strong reason to avoid the “skip breakfast, feast at dinner” pattern.

The leucine trigger

Muscle-building machinery (the mTORC1 pathway) switches on when the amino acid **leucine** reaches a threshold inside muscle cells. Roughly 2 to 3 g of leucine does it, which many people reach with about 25 to 40 g of high-quality protein. A breakfast of toast, cereal or coffee delivers perhaps 10 g, so the switch never flips, and a large dinner cannot retroactively make up for it.

A practical framework

Schoenfeld and Aragon proposed a simple rule: aim for about **0.4 g of protein per kg of body weight per meal**, spread over at least four eating occasions. That gives 1.6 g/kg/day, and going up to roughly 0.55 g/kg per meal reaches about 2.2 g/kg/day, the upper end used in muscle-building research. For most readers of this book, 1.2 to 1.6 g/kg is plenty.

EATING PLANT-BASED?

Most plant proteins contain a little less leucine and are slightly less digestible than animal proteins. Aim for the upper end of the per-meal range (30 to 40 g), or combine sources such as soy, pea and wheat proteins.

IN SHORT

- Absorption has no 30 g cap; muscle-building signalling per meal does taper off.
- Even distribution produced about 25% higher 24-hour MPS than a dinner-heavy pattern in the same-calorie trial.
- Aim for roughly 25–40 g (or ≈0.4 g/kg) at each of three to four eating occasions.
- Breakfast is the easiest place to fix, which leads directly to Chapter 3.

CHAPTER 3

The Brain Connection: Leidy's fMRI Findings and the Breakfast Anchor

The hardest part of a diet is often the evening. New research suggests the evening is shaped by the morning.

Dr. Heather Leidy and colleagues studied how breakfast shapes appetite and brain responses to food. In a small study of overweight, "breakfast-skipping" late-adolescent girls (around 20 participants), each person tried three morning patterns in turn: **skipping breakfast**, a **normal-protein breakfast** (13 g), or a **high-protein breakfast** (35 g, from eggs and beef). Before lunch, participants viewed food images while undergoing functional MRI (fMRI) scans.




Illustration of the reward-related activation reported after a skipped or low-protein breakfast versus a 35 g protein breakfast.

Morning pattern	Brain response to food cues	Behaviour
Skipped breakfast	Highest activity in reward and craving regions (amygdala, insula, hippocampus and related areas).	Strongest drive toward high-fat, high-sugar food.
Normal breakfast (13 g protein)	Some reduction compared with skipping.	Modest effect on later eating.
High-protein breakfast (35 g)	Clearly lower reward-region activation, with changes in prefrontal and parahippocampal areas.	Less evening snacking on high-fat and sugary foods; steadier appetite.

WHAT THE SCANS DO AND DO NOT SHOW

fMRI measures blood-flow changes in brain regions linked to reward; it does not measure dopamine directly, and the study was small, short-term and in one group of young women. The result is best read as a consistent, plausible signal that a protein-rich morning reduces cravings, supported by further trials, not as proof that breakfast "switches off" cravings.

Beyond cravings

- **Less evening snacking.** A protein-rich breakfast reduced later snacking on high-fat, sugary convenience foods in these trials.
- **Steadier blood sugar.** About 35 g of protein at breakfast has been linked to more stable glucose through the day.
- **Focus and energy.** Some short-term studies report better concentration before lunch after a high-protein breakfast, though results are mixed and individual.

Build your Morning Anchor

The practical takeaway is the **30g Morning Anchor**: aim for 30 to 35 g of protein within an hour or two of waking. Five easy ways to get there:

Anchor breakfast	How	Protein
Savoury cottage cheese bowl	1½ cups cottage cheese, cherry tomatoes, cucumber, everything-bagel seasoning, olive oil.	≈ 34 g
Microwave egg bites	4 eggs, ⅓ cup cottage cheese, spinach and a little bacon, 3 minutes in a silicone mould.	≈ 32 g
Greek yogurt and quinoa bowl	1 cup nonfat Greek yogurt, ½ cup cooked quinoa, 2 tbsp hemp seeds, berries.	≈ 35 g
Omelette plus cottage cheese	3-egg omelette with spinach, ½ cup cottage cheese on the side.	≈ 30 g
Whey-and-milk smoothie	1 scoop whey (≈24 g), 1 cup milk, frozen berries, oats.	≈ 32 g

IN SHORT

- A 35 g protein breakfast reduced reward-region responses and evening snacking compared with skipping or a 13 g breakfast.
- Evidence is promising but comes from small, short-term studies.
- Aim for 30–35 g of protein within an hour or two of waking.
- If you are not hungry in the morning, start with a smaller protein serving and build up.

CHAPTER 4

Navigating 2026’s Biggest Misconceptions and Friction Points

Moving to a higher-protein diet brings real-world snags. Knowing them in advance is what keeps you consistent.

“Mouth satisfaction” versus “stomach fullness”

A common beginner complaint is feeling physically full yet still wanting to eat. It happens when a meal has volume but no sensory variety. Dry, unseasoned chicken with plain steamed broccoli stretches the stomach, but without fat, texture or bright flavours the brain’s reward centres stay unsatisfied, and the cravings arrive later. The fix is to make meals satisfying to eat, not just macro-friendly.

> **THE SATISFACTION FORMULA**
>
> **Protein** (25–40 g) + **Fat or fibre** (avocado, olive oil, hemp seeds, beans) + **Texture** (crunch, creaminess) + **Brightness** (lemon, pickles, herbs, heat). Hit all four and you feel done.

The ultra-processed food “gray area”

Public concern about ultra-processed foods (UPFs) is high in 2026, and it sometimes drifts into fear-based messaging. Many beginners assume that anything in a package is a UPF and must be avoided. Dietitians point out that plain Greek yogurt, canned tuna, packaged firm tofu and high-fibre wraps come in packets, yet are nutritious and make consistency far easier. Focus on the overall pattern, and use a quick label check:

- **Short ingredient list** you recognise (tuna, water, salt) beats a long list of additives.
- **Protein per serving:** look for at least 15 g, or roughly 10 g of protein per 100 kcal.
- **Check sodium** on deli meats, canned fish, pickles and stock; these are the saltiest convenience staples.
- **Limit added sugar** in flavoured yogurts, bars and sauces.

Kidney and bone myths

One persistent myth says high-protein diets cause acidosis, kidney damage and bone loss through calcium excretion. Trials lasting 6 to 12 months have repeatedly found no harmful effects on kidney function or bone mineral density in **healthy adults**, and healthy kidneys adapt to higher intakes. Adequate calcium, vitamin D and fruit and vegetables remain important for bones.


> **CHECK WITH A CLINICIAN FIRST IF…**
>
> you have chronic kidney disease or reduced kidney function, you are pregnant or breastfeeding, you have a history of disordered eating, you are under 18, or you are managing diabetes or other conditions with medication. Protein targets for these groups are individual.

Do you need to rotate protein sources?

An online claim says too much choline from eggs and chicken causes a “mood flatline”. The clinical evidence for this is thin, and ordinary food intakes sit well below the established upper limit. If you feel flat after changing your diet, the likelier explanations are simply too few total calories or carbohydrates, poor sleep, or not drinking enough. Variety is still good practice, so rotate poultry, fish, dairy, lean beef, tofu, tempeh and legumes to broaden your vitamins, minerals and fats.

The protein–fibre synergy

Raising protein while cutting carbohydrates often causes bloating and constipation, because the fibre from grains, fruit and legumes drops at the same time. The current “fibremaxxing” trend has a sound core: pair protein with fibre. Jumping from 10 g to 40 g overnight backfires, so use the **+5 g weekly ramp** and raise fluids alongside it.



The +5 g weekly fibre ramp with matching fluid increases. Stop climbing when you reach a comfortable level; 30 g/day is the UK guideline.

Week	Fibre (g/day)	Fluids (L/day)	Easy additions
Start	≈ 10	1.5	Your current baseline
1	15	1.8	Add a vegetable to two meals; swap to a wholemeal wrap
2	20	2.1	Add ½ cup beans or lentils daily
3	25	2.4	Add berries and 1 tbsp chia or ground flax
4	30	2.7	Choose edamame or chickpea pasta once or twice a week
5	35	3.0	Hold here; go to 40 only if comfortable

Fluid needs vary with body size, climate and activity. If you have kidney or heart conditions, follow your clinician’s fluid advice.

IN SHORT

- Make meals satisfying: protein, fat or fibre, texture and brightness.
- Packaged does not mean bad; read labels for protein, ingredients and sodium.
- Healthy adults do not need to fear higher protein for kidneys or bones; certain medical groups do need guidance.
- Raise fibre by about 5 g and fluids by about 0.3 L each week.

CHAPTER 5

Comparative Chemistry of Whey and Plant-Based Proteins

Choosing a protein powder or plant staple comes down to budget, digestion and dietary preference.

Whey, from cheese vat to shaker

Whey is a fast-digesting, highly bioavailable by-product of cheesemaking, naturally rich in leucine. Increasing levels of filtration and processing produce three main forms with distinct trade-offs.

Liquid whey
Cheesemaking by-product
→
Concentrate (WPC)
Ultrafiltration, 70–80% protein
→
Isolate (WPI)
Microfiltration, 90%+ protein
→
Hydrolysate (WPH)
Enzymes break chains into peptides

	Concentrate (WPC)	Isolate (WPI)	Hydrolysate (WPH)
Protein by weight	70–80%	90%+	Typically 90%+
Digestion speed	Fast	Faster	Fastest
Lactose and fat	Around 7–8% lactose and fat (varies by product)	Under about 3% lactose; trace fat	Trace
Bioactives	Retains immunoglobulins and lactoferrin	Most removed in processing	Largely broken down
Taste and texture	Creamy, rich	Thin, clean, mixes easily	Thin, can taste bitter
Cost	Economical baseline	Moderate premium (about 30–50%)	Highest
Best for	Budget-minded beginners with normal dairy tolerance	Lactose-sensitive or low-carb eaters	Elite athletes, very sensitive digestion

BUYING TIPS

Choose products with a short ingredient list and a third-party testing mark (such as Informed Sport) where possible. Remember that fast digestion is not essential for most people; **total daily protein and its spread across meals matter more than the type of whey.**

Debunking the “incomplete plant protein” myth

Plant-based eaters are often told to pair foods carefully at every meal to get all nine essential amino acids. Biochemistry has retired this idea. **All plant proteins contain all nine essential amino acids.** Some are relatively lower in one (grains in lysine, legumes in methionine), but the body keeps a pool of free amino acids and balances the books over the day. Eat a variety of sources across 24 hours and aim for slightly higher totals, and you are covered.

High-density plant protein staples

Food	Serving	Protein	Notes
Seitan (vital wheat gluten)	4 oz (≈115 g)	≈ 30 g	Around 200 kcal; low in lysine, so pair with legumes. Not suitable for coeliac disease.
Textured vegetable protein (TVP)	½ cup rehydrated	≈ 12 g	Soy crumbles; replaces mince in tacos, chilli and bolognese.
Edamame or lentil pasta	2 oz (≈56 g) dry	≈ 20–25 g	Higher protein than chickpea pasta (about 11–14 g per 2 oz); check the packet.
Nutritional yeast	¼ cup	≈ 8 g	Savoury, cheesy topping; many brands are fortified with B-vitamins.
Hemp seeds	3 tbsp	≈ 10 g	Provide magnesium and omega-3 fats; low in lysine, so vary your sources.
Tempeh	4 oz (≈115 g)	≈ 20 g	Fermented soy; firm, nutty and high in fibre.

IN SHORT


- Concentrate is the affordable default; choose isolate if lactose or carbs are a concern.
- Digestion speed matters far less than your total daily protein and its spread.
- Every plant protein is complete in the sense that matters; variety over a day covers the gaps.
- Plant eaters should aim for the upper end of per-meal targets.

CHAPTER 6

The “Ingredient Prep” Kitchen System and Rapid Recipes

Prep components, not meals. You get fresh, varied dinners in minutes without the Wednesday food fatigue.

Traditional meal prep produces identical boxes of dried-out protein and soggy vegetables, and by midweek nobody wants them. **Ingredient Prep** flips the approach: spend about 30 minutes twice a week (for example Sunday and Wednesday) cooking *components* and storing them separately.



Bulk meal prep (left) versus modular Ingredient Prep (right).

Component	Prep ideas	Keeps (fridge)
Versatile proteins	Shredded rotisserie chicken, seasoned lean beef or turkey mince, hard-boiled eggs	3–4 days (eggs in shell up to 7)
Prepared vegetables	Sliced bell peppers, washed baby spinach, steamed broccoli, diced cucumber	3–4 days
Carbohydrate bases	Quinoa, brown rice or high-protein pasta	See food-safety note below

FOOD SAFETY FOR PREPPED FOOD

Cool cooked rice and grains quickly (within about an hour) and refrigerate. UK NHS guidance says to eat chilled rice within one day, while US guidance allows 3–4 days; when in doubt, go shorter, which is another reason to prep twice a week. Keep cooked meat and poultry 3–4 days at most, reheat until piping hot, and keep the fridge at 5°C or below.

Your 30-minute prep session

1. **Minutes 0–5: start the slow things**
   Put grains on to cook, boil eggs for 10 to 12 minutes, and preheat the oven or pan.
2. **Minutes 5–15: cook the proteins**
   Sear seasoned turkey or beef mince; shred rotisserie chicken; season and rest.
3. **Minutes 15–25: vegetables**
   Steam broccoli, slice peppers, dice cucumber, wash spinach.
4. **Minutes 25–30: cool and store**
   Spread grains on a tray to cool fast, then store every component in its own container.

The Power Bowl (“Trash Salad”) blueprint

With components ready, any bowl takes under five minutes. Follow the five layers:

Layer	Choose	Amount
1. Base	Fresh greens, or pre-cooked quinoa, brown rice or high-protein pasta	2 cups greens or 1 cup grains
2. Protein	Shredded chicken, seared turkey mince, cubed tofu, hard-boiled eggs	4–6 oz (115–170 g)
3. Colour and crunch	Cucumber, peppers, tomatoes, pickled onion	A generous handful
4. Satiety booster	Avocado, hemp seeds, hummus	¼ avocado, a sprinkle, or a spoonful
5. Flavour	Nonfat Greek yogurt, lemon juice, fresh herbs and spices	2–3 tbsp

Under-30-minute recipe blueprints

Protein values are estimates per serving and vary by brand, so check labels. Recipes serve one unless noted.

1. Quick no-cook and cold prep

Recipe	Time	Protein	Key ingredients and kitchen hack
Cottage Cheese Veggie Pack	2 min	≈ 34 g	Mix 1½ cups cottage cheese with halved cherry tomatoes, cucumber and everything-bagel seasoning.
Turkey Pickle Roll-Ups	3 min	≈ 32 g	Roll 6 oz deli turkey and 2 oz low-fat cheese around pickle spears with a swipe of mustard.
Greek Yogurt Chicken Salad	4 min	≈ 40 g	Mix 4 oz shredded chicken with ¼ cup nonfat Greek yogurt, diced celery and lemon juice.
Tuna Avocado Salad Bowl	5 min	≈ 31 g	Mash 1 can tuna in water with ½ avocado, 2 tbsp Greek yogurt, lemon juice and black pepper in the container.
Canned Salmon Avocado Bowl	5 min	≈ 36 g	Mix 1 can salmon with ½ avocado, 2 tbsp Greek yogurt, capers, lemon zest and olive oil.
Smoked Salmon Quinoa Bowl	5 min	≈ 30 g	Top pre-cooked quinoa with 4 oz smoked salmon, cucumber and a Greek yogurt and dill dressing.
Tuna Cucumber Boats	6 min	≈ 38 g	Scoop tuna salad (1½ cans) into hollowed cucumber halves in place of bread.
Dill Pickle Chicken Salad	7 min	≈ 37 g	Mix 1 can chicken breast with ½ cup nonfat Greek yogurt, diced dill pickles and a splash of pickle juice.
Egg Salad Supergreen Wraps	8 min	≈ 33 g	Mash 4 hard-boiled eggs with ⅓ cup Greek yogurt; wrap in briefly blanched collard leaves.

2. Fast air-fryer and microwave

Recipe	Time	Protein	Key ingredients and kitchen hack
Philly Cheesesteak Bowl	8 min	≈ 33 g	Microwave pre-cooked steak, peppers and mushrooms; top with melted provolone over cauliflower rice.
Microwave Egg Bites	8 min	≈ 32 g	Whisk 4 eggs, ⅓ cup cottage cheese, spinach and a little bacon in a silicone mould; microwave about 3 minutes.
Air Fryer Tilapia Peri-Peri	10 min	≈ 41 g	Coat 200 g tilapia with olive oil, garlic paste and peri-peri seasoning; air fry 10 minutes, flipping halfway.
Crispy Air Fryer Walleye	10 min	≈ 42 g	Lightly crumb walleye fillets (cod or haddock work well in the UK); air fry at 200°C / 400°F for 10 minutes until golden.
Turkey Taco Bowls	15 min	≈ 35 g	Sear lean turkey mince with taco seasoning; serve over microwave brown rice with black beans and salsa.
Buffalo Chicken Smash Tacos	20 min	≈ 38 g	Press shredded buffalo chicken into corn tortillas on a hot skillet; top with light Greek yogurt ranch.



Seared salmon with brown rice and roasted broccoli: a complete high-protein plate in about 25 minutes.

3. One-pot and sheet-pan dinners

Recipe	Time	Protein	Key ingredients and kitchen hack
Ground Turkey Sweet Potato Skillet	9 min	≈ 35 g	Sauté 6 oz ground turkey with pre-diced sweet potato, baby spinach and garlic in one skillet.
Teriyaki Turkey Skillet	15 min	≈ 35 g	Brown 6 oz turkey mince; add broccoli and a low-sugar teriyaki sauce; serve over cauliflower rice or quinoa.
Bone Broth Mac and Cheese	20 min	≈ 33 g	Boil chickpea pasta directly in beef bone broth; stir in sharp cheddar and Greek yogurt until creamy.
Bone Broth Spaghetti	30 min	≈ 29 g	Cook pasta in beef bone broth; top with lean turkey mince and your favourite marinara.
Creamy Salmon Alfredo Pasta	30 min	≈ 41 g	Toss chickpea pasta with seared salmon, light cream cheese, parmesan, garlic and baby spinach.
High-Protein Marry Me Chicken	30 min	≈ 61 g	Sear an 8 oz chicken breast with sun-dried tomatoes and broth; finish with whipped cottage cheese. Use 5 oz chicken for ≈ 45 g.

One day, about 130 g of protein

Meal	Example	Protein
Breakfast	Savoury cottage cheese bowl	≈ 34 g
Lunch	Greek Yogurt Chicken Salad over greens	≈ 40 g
Snack	Nonfat Greek yogurt with berries (¾ cup)	≈ 18 g
Dinner	Air Fryer Tilapia with roasted broccoli and quinoa	≈ 41 g
Total		≈ 133 g

Quick conversions: 1 oz ≈ 28 g; 4 oz ≈ 115 g; 1 cup ≈ 240 ml; 400°F ≈ 200°C; 1 lb ≈ 450 g.

IN SHORT

- Prep components, not meals: two 30-minute sessions a week.
- Build bowls from five layers: base, protein, colour, satiety booster, flavour.
- Cool and store prepped food promptly, and use the food-safety timings.
- Hit 25–40 g at each meal using the recipe tables as a menu.

CHAPTER 7

Your 5-Stage Transformation Roadmap

A structured path from food stress to food freedom: one stage per week, each with one system and one recipe.

Information alone rarely changes habits. This roadmap pairs each lesson with a single kitchen system and a matching emotional shift, so that by the end of five weeks you are not following rules; you are running a routine that fits your life.



Stage	You learn	System and key recipe	How it should feel
1. The Satiety Breakthrough	How hunger hormones (GLP-1, PYY, CCK, ghrelin) and the thermic effect of food work.	**System:** the 30g Morning Anchor. **Recipe:** Savoury Cottage Cheese Bowl.	Frustrated → hopeful. Past struggles were biology, not weak will.
2. The Power Pantry Blueprint	How to navigate the supermarket, read labels and choose good convenience foods.	**System:** the Power Pantry checklist (below). **Recipe:** Tuna Avocado Salad Bowl.	Fearful → capable. Convenience foods without guilt.
3. The Modular Ingredient Prep	A component-based alternative to traditional meal prep.	**System:** the 30-minute prep session. **Recipe:** Teriyaki Turkey Skillet.	Stressed → organised. Healthy eating fits a busy week.
4. The Protein–Fibre Synergy	Why pairing protein with fibre supports gut comfort.	**System:** the weekly +5 g fibre ramp. **Recipe:** Greek Yogurt and Quinoa Bowl.	Bloated → comfortable and energised.
5. Sustainable Autonomy	Flexibility, social eating, and adjusting protein for activity.	**System:** the Power Bowl assembly method. **Recipe:** High-Protein Marry Me Chicken.	Unsure → confident. You can adapt on your own.

The Power Pantry checklist

PROTEINS

- ☐ Eggs
- ☐ Rotisserie or canned chicken
- ☐ Canned tuna, salmon, sardines
- ☐ Lean turkey or beef mince
- ☐ Firm tofu or tempeh
- ☐ Smoked salmon

DAIRY AND ALTERNATIVES

- ☐ Nonfat Greek yogurt
- ☐ Cottage cheese
- ☐ Sharp cheddar or parmesan
- ☐ Milk or fortified soy milk
- ☐ Whey or plant protein powder

FIBRE AND CARBS

- ☐ Quinoa and brown rice
- ☐ Edamame or chickpea pasta
- ☐ Black beans and lentils
- ☐ High-fibre wraps
- ☐ Oats and berries

FLAVOUR BOOSTERS

- ☐ Lemons and fresh herbs
- ☐ Pickles and capers
- ☐ Everything-bagel seasoning
- ☐ Hot sauce and mustard
- ☐ Hemp seeds, olive oil

WHEN LIFE GETS IN THE WAY

Restaurant meal? Pick the dish built around a protein, add a vegetable side and don't sweat the rest. Missed your prep day? A rotisserie chicken, a bag of salad and a tub of cottage cheese is a perfectly good plan B. Consistency over weeks matters far more than any single meal.

IN SHORT

- Work through one stage per week: one system, one recipe.
- Use the Power Pantry list to make shopping and prep automatic.
- Plan B meals keep you consistent when the week falls apart.
- Adjust protein for your activity, age and health, ideally with professional input.

APPENDIX

Appendix: Key Takeaways, Quick Start and References

Everything above, condensed into one page of recommendations and a first week to try.

Synthesised recommendations

1. **Establish a morning protein anchor**
   Aim for 30–35 g of high-quality protein within an hour or two of waking to help curb afternoon brain fog, steady blood sugar and reduce late-night cravings.
2. **Spread protein through the day**
   Target about 0.4 g/kg (roughly 25–40 g) per meal across three to four meals, within a daily range of 1.2–1.6 g/kg.
3. **Ramp fibre and fluids together**
   Add about 5 g of fibre and 0.3 L of fluid per week until you reach 30–40 g and 2.5–3 L, if that suits your health.
4. **Prep components, not meals**
   Two short sessions a week: two proteins, three prepared vegetables and one carbohydrate base, stored separately.
5. **Choose supplements by tolerance**
   Whey concentrate is the affordable default; switch to isolate if lactose or carbs are an issue; consider soy or pea blends if plant-based.
6. **Use healthy convenience foods**
   Canned fish, pre-cooked grains, high-fibre wraps and pre-chopped vegetables keep prep under 15 minutes. Watch sodium.

Your 7-day quick start

Day	Do this
1	Calculate your protein range (Foreword). Stock the Power Pantry.
2	Eat a 30–35 g breakfast. Note how your evening cravings feel.
3	Do your first 30-minute ingredient prep session.
4	Build two Power Bowls from your components. Add 5 g of fibre.
5	Cook one recipe from each section of Chapter 6.
6	Do your second prep session; try a new protein source.
7	Review: what was easy, what was boring, what will you change?

Selected references

- Dietary Guidelines for Americans, 2025–2030. US Department of Health and Human Services and US Department of Agriculture, January 2026.

- Mamerow MM, et al. Dietary protein distribution positively influences 24-h muscle protein synthesis in healthy adults. *J Nutr.* 2014;144(6):876–880.
- Schoenfeld BJ, Aragon AA. How much protein can the body use in a single meal for muscle-building? *J Int Soc Sports Nutr.* 2018;15:10.
- Leidy HJ, et al. Beneficial effects of a higher-protein breakfast on the appetitive, hormonal, and neural signals controlling energy intake regulation in overweight/obese, “breakfast-skipping,” late-adolescent girls. *Am J Clin Nutr.* 2013;97(4):677–688.
- Halton TL, Hu FB. The effects of high protein diets on thermogenesis, satiety and weight loss: a critical review. *J Am Coll Nutr.* 2004;23(5):373–385.

Please verify citations against the original publications before reuse. Product, regulatory and guideline details reflect information available in October 2026.