

GASTROENTEROLOGY CONSULTANTS, P.C.

BRIAN D. CLARKE, M.D.
(317) 621-2100

You are scheduled for a colonoscopy on _____, _____ 200__.

Report to outpatient registration at _____ a.m./p.m. at

- ☐ Community Hospital North at 7150 Clearvista Pkwy, Phone (317) 621-5193
- ☐ Carmel Endoscopy Center at 13421 Old Meridian St, Phone(317) 706-1600

You need to be accompanied by a driver for this exam. (A taxi cab or bus is **not** acceptable.)

**** This prep is not recommended for patients with constipation – Ask for Gallon Prep ****

YOU WILL NEED TO PURCHASE:

HalfLyte[®] and Bisacodyl Tablets Bowel Prep Kit (Prescription required)

Prepare **HalfLyte[®] solution** day before prep and refrigerate overnight if you prefer cold liquids.

MEDICATION RESTRICTIONS

You may continue to take regular medications on day before procedure except medication that would prevent your colon from being cleaned out. Example: Anti-diarrheal medications, fiber supplements.

Heart or blood pressure medications: Take these on the morning of the colon exam.

Diabetics: Take half of your evening dose of **insulin** on the day before the procedure but do not take oral diabetes medication or insulin on the day of the colonoscopy until completed. (If any questions, please contact the doctor that prescribes your insulin.)

Iron supplements (Ferrous sulfate): Stop 1 week before colonoscopy.

Blood thinner: Please discontinue **Coumadin (warfarin)**, **Lovenox (heparin)**, **Aspirin** 3 days before colonoscopy and **Plavix (clopidogrel)**, **Aggrenox (aspirin/dipyridamole)**, **Persantine (dipyridamole)**, **Pletal (cilostazol)**, **Ticlid (ticlopidine)** 5 days before the colonoscopy. Discuss with your prescribing physician for their approval before you stop any medicine that affects blood clotting.

COLON PREP INSTRUCTIONS – The Day Before Your Test

1. **Clear Liquid Diet** (see list below) beginning at breakfast. **No solid food allowed.**
2. **2:00 p.m. or immediately after work** Take 4 bisacodyl delayed release tablets with water (do NOT chew or crush). First bowel movement should occur in about 1-4 hours after taking the laxative pills.
3. Wait for a bowel movement (or **maximum of 4 hours**) then drink **HalfLyte[®]** solution, 1 (8 oz) glass every 10 minutes (approximately 8 glasses). **Drink ALL of the solution.** Rapid drinking of each portion is better than drinking small amounts continuously. A watery bowel movement should occur in approximately 1 hour after drinking the solution. You may experience some abdominal bloating and distention before the bowels start to move. If severe discomfort or distention occurs, stop drinking the solution temporarily or drink each portion at longer intervals until these symptoms disappear.
4. **Midnight Please have nothing to eat until colonoscopy is completed.** Water is permitted.
5. If colonoscopy is in afternoon, clear liquids are permitted up to 3 hours before your arrival time.

CLEAR LIQUID DIET

(No red or purple artificial colors)

Jell-O[®] or gelatin (plain without fruit), coffee or tea, low-salt bouillon/broth, juices without pulp, popsicles, soda-pop, Kool-aid[®], Gatorade[®] & hard candy. No milk products or fruit fiber.

FINAL INSTRUCTIONS

- Bring current medication list with dosages
- If you use a CPAP machine at night for sleep apnea, please bring this with you.

*****Please note: There is a \$75.00 fee for procedures cancelled less than 48 hours (2 business days) before scheduled appointment time*****