

GASTROENTEROLOGY CONSULTANTS, P.C.

BRIAN D. CLARKE, M.D.
(317) 621-2100

You are scheduled for a colonoscopy on _____, _____ 200__.

Report to outpatient registration at _____ a.m./p.m. at

- ☐ Community Hospital North at 7150 Clearvista Pkwy, Phone (317) 621-5193
- ☐ Carmel Endoscopy Center at 13421 Old Meridian St, Phone(317) 706-1600

You need to be accompanied by a driver for this exam. (A taxi cab or bus is **not** acceptable.)

**** If you have heart failure, kidney disease, high blood pressure or salt restriction, do not use this prep.**

YOU WILL NEED TO PURCHASE:

OsmoPrep™ Tablets (Prescription required) – You will receive a bottle of 32 tablets

MEDICATION RESTRICTIONS

You may continue to take regular medications on day before procedure except medication that would prevent your colon from being cleaned out. Example: Anti-diarrheal medications, fiber supplements.

Heart or blood pressure medications: Take these on the morning of the colon exam.

Diabetics: Take half of your evening dose of **insulin** on the day before the procedure but do not take oral diabetes medication or insulin on the day of the colonoscopy until completed. (If any questions, please contact the doctor that prescribes your insulin.)

Iron supplements (Ferrous sulfate): Stop 1 week before colonoscopy.

Blood thinner: Please discontinue **Coumadin (warfarin)**, **Lovenox (heparin)**, **Aspirin** 3 days before colonoscopy and **Plavix (clopidogrel)**, **Aggrenox (aspirin/dipyridamole)**, **Persantine (dipyridamole)**, **Pletal (cilostazol)**, **Ticlid (ticlopidine)** 5 days before the colonoscopy. Discuss with your prescribing physician for their approval before you stop any medicine that affects blood clotting.

COLON PREP INSTRUCTIONS – Begin Day Before Your Test

1. Clear Liquid Diet (see list below) beginning at breakfast. **No solid food allowed.**
2. **6 p.m.** Take 4 **OsmoPrep™ Tablets** with at least 8 oz. of clear liquid every 15 minutes for a total of 5 doses in an hour. **Do not exceed 20 tablets.** Initially, you may feel slightly bloated, but will become more comfortable as you have bowel movements. You will begin having bowel movements 45 minutes to 1 hour after taking the Osmo Prep tablets so stay close to a toilet facility.

Morning of your colonoscopy at least 3 hours before arrival time (You may need to set an alarm!)

3. **Early morning** (Example: 4 a.m. if arrival time at 7 a.m.) or **8 a.m.** if afternoon colonoscopy
Take 4 **OsmoPrep™ Tablets** with at least 8 oz. of clear liquid every 15 minutes for a total of 3 doses in 30 minutes. **Do not exceed 12 tablets.**
4. If colonoscopy in afternoon, clear liquids are permitted up to 3 hours before your arrival time.

CLEAR LIQUID DIET

(No red or purple artificial colors)

Jell-O® or gelatin (plain without fruit), coffee or tea, low-salt bouillon/broth, juices without pulp, popsicles, soda-pop, Kool-aid®, Gatorade® & hard candy. No milk products or fruit fiber.

FINAL INSTRUCTIONS

- Bring current medication list with dosages
- If you use a CPAP machine at night for sleep apnea, please bring this with you.

*****Please note: There is a \$75.00 fee for procedures cancelled less than 48 hours (2 business days) before scheduled appointment time*****