

GASTROENTEROLOGY CONSULTANTS, P.C.

BRIAN D. CLARKE, M.D.
(317) 621-2100

You are scheduled for a colonoscopy on _____, _____ 200__.

Report to outpatient registration at _____ a.m./p.m. at

- ☐ Community Hospital North at 7150 Clearvista Pkwy, Phone (317) 621-5193
- ☐ Carmel Endoscopy Center at 13421 Old Meridian St, Phone(317) 706-1600

You need to be accompanied by a driver for this exam. (A taxi cab or bus is **not** acceptable.)

**** If you have heart failure, kidney disease, high blood pressure or salt restriction, do not use this prep.**

YOU WILL NEED TO PURCHASE:

Fleet® Phospho-soda® EZ-Prep™ (No prescription required)

Or buy Fleet® Phospho-soda® Ginger-Lemon Flavored – two 1.5 oz. bottles if EZ-Prep not available

MEDICATION RESTRICTIONS

You may continue to take regular medications on day before procedure except medication that would prevent your colon from being cleaned out. Example: Anti-diarrheal medications, fiber supplements.

Heart or blood pressure medications: Take these on the morning of the colon exam.

Diabetics: Take half of your evening dose of **insulin** on the day before the procedure but do not take oral diabetes medication or insulin on the day of the colonoscopy until completed. (If any questions, please contact the doctor that prescribes your insulin.)

Iron supplements (Ferrous sulfate): Stop 1 week before colonoscopy.

Blood thinner: Please discontinue **Coumadin (warfarin)**, **Lovenox (heparin)**, **Aspirin** 3 days before colonoscopy and **Plavix (clopidogrel)**, **Aggrenox (aspirin/dipyridamole)**, **Persantine (dipyridamole)**, **Pletal (cilostazol)**, **Ticlid (ticlopidine)** 5 days before the colonoscopy. Discuss with your prescribing physician for their approval before you stop any medicine that affects blood clotting.

COLON PREP INSTRUCTIONS – The Day Before Your Test

1. Clear Liquid Diet (see list below) beginning at breakfast. **No solid food allowed.**
2. **Late morning or immediately after work*** Mix **Fleet® Phospho-soda®** 1.5 fl. oz. with 12 oz of cold water, mix well and drink all of it. Immediately follow with at least 12 oz of clear liquids.
3. **Between Doses** Drink at least 2 more glasses of clear liquids (at least 12 oz each). More is better.
4. **12 hours from first dose (at least 3 hours before arrival time)** Repeat step 2. with remaining bottle of **Fleet® Phospho-soda®** and immediately drink at least 12 oz of clear liquid afterward.
5. If colonoscopy in afternoon, clear liquids are permitted up to 3 hours before your arrival time.
 - * The prep start time may be later if your colonoscopy is scheduled in the afternoon but you should allow 12 hours between doses of **Fleet® Phospho-soda®** and at least 3 hours before leaving home.

CLEAR LIQUID DIET

(No red or purple artificial colors)

Jell-O® or gelatin (plain without fruit), coffee or tea, low-salt bouillon/broth, juices without pulp, popsicles, soda-pop, Kool-aid®, Gatorade® & hard candy. No milk products or fruit fiber.

FINAL INSTRUCTIONS

- Bring current medication list with dosages
- If you use a CPAP machine at night for sleep apnea, please bring this with you.

*****Please note: There is a \$75.00 fee for procedures cancelled less than 48 hours (2 business days) before scheduled appointment time*****