

Supplement S3: Dynamics and Scale Bridge

Torsional Flow, Dimensional Transmutation,
and Cosmological Evolution

Brieuc de La Fournière

Independent researcher

Abstract

The GIFT framework's dimensionless predictions (S2) require dynamical completion to connect with absolute physical scales. This supplement provides three essential bridges: (1) Torsional dynamics through non-closure of the G_2 3-form with $\kappa_T = 1/61$; (2) The scale bridge formula $m_e = M_{\text{Pl}} \times \exp(-(H^* - L_8 - \ln(\phi)))$ deriving the electron mass from Planck scale with precision $< 0.1\%$ on the exponent; (3) Cosmological evolution including Hubble tension resolution via dual topological projections $H_0 = \{67, 73\}$. All results emerge from the topological structure established in S1.

Contents

Part I: Torsional Geometry	6
1 Torsion from G_2 Non-Closure	6
1.1 Torsion in Differential Geometry	6
1.2 Torsion-Free vs Torsionful Connections	6
1.3 G_2 Holonomy and the 3-Form	6
2 Torsion Magnitude $\kappa_T = 1/61$	7
2.1 Topological Derivation	7
2.2 The Number 61	7
2.3 Experimental Compatibility	7
3 Torsion Classes for G_2 Manifolds	8
3.1 Irreducible Decomposition	8
3.2 GIFT Framework Torsion	8
4 Torsion Tensor Components	8
4.1 Coordinate System	8
4.2 Component Structure	8
4.3 Physical Interpretation	9
4.4 Topological Structure of Torsion Components	9
Part II: Geodesic Flow and RG Connection	10
5 Torsional Geodesic Equation	10
5.1 Derivation from Action	10
5.2 Torsional Modification	10
5.3 Main Result	10
5.4 Physical Interpretation	10
6 RG Flow Connection	10
6.1 Identification $\lambda = \ln(\mu)$	10
6.2 Scale Dependence	11
6.3 β -Functions as Velocities	11
7 Flow Velocity and Stability	11

7.1	Ultra-Slow Velocity Requirement	11
7.2	Velocity Bound Derivation	11
7.3	Framework Value	12
8	Conservation Laws	12
8.1	Energy Conservation	12
8.2	Topological Charges	12
	Part III: The Scale Bridge	12
9	The Dimensional Transmutation Problem	12
9.1	The Challenge	12
9.2	Natural Scales	13
10	The Master Formula	13
10.1	The Scale Bridge	13
10.2	The Exponent	13
10.3	The Ratio	13
10.4	The Mass	14
11	Numerical Verification	14
11.1	Precision Analysis	14
11.2	Mass Comparison	14
11.3	Python Verification	14
12	Physical Interpretation	15
12.1	The Three Components	15
12.2	Separation of Scales	15
12.3	Why These Values?	15
12.4	Elegant Reformulation	16
12.5	Why Lucas Rather Than Fibonacci	16
13	The Hierarchy Problem	17
13.1	The Traditional Problem	17
13.2	GIFT Resolution	17
13.3	Why $\sim 10^{-23}$?	17

Part IV: Mass Chain	18
14 Complete Mass Derivation	18
14.1 The Master Chain	18
15 Lepton Masses	18
15.1 Electron Mass (From Scale Bridge)	18
15.2 Muon Mass	18
15.3 Tau Mass	19
15.4 Lepton Summary	19
16 Quark Sector Status	19
16.1 Current State	19
16.2 Open Problem	20
17 Boson Masses	20
17.1 W Boson Mass	20
17.2 Z Boson Mass	20
17.3 Higgs Mass	20
17.4 Boson Summary	21
18 Neutrino Masses	21
18.1 Hierarchy Prediction	21
18.2 Mass Sum	21
18.3 Individual Masses (Exploratory)	21
Part V: Cosmological Dynamics	22
19 The Hubble Tension	22
19.1 The Crisis	22
19.2 GIFT Resolution	22
19.3 The Tension is Structural	22
19.4 Verification	22
19.5 Physical Interpretation: Dimensional Projection	22
19.6 The Gap as Fermionic Decoupling	23
19.7 The Duality Diagram	23

20 Dark Energy	24
20.1 The Formula	24
20.2 Calculation	24
20.3 Triple Origin of $\ln(2)$	24
20.4 Verification	24
21 Dark Matter	24
21.1 Dark Energy to Dark Matter Ratio	24
21.2 Golden Ratio Connection	24
21.3 Verification	25
22 Age of the Universe	25
22.1 The Formula	25
22.2 Components	25
22.3 Verification	25
23 Spectral Index	25
23.1 The Formula	25
23.2 Calculation	25
23.3 Verification	25
24 Cosmological Summary	26
Part VI: Summary and Limitations	26
25 Key Results	26
25.1 Torsional Dynamics	26
25.2 Scale Bridge	26
25.3 Mass Chain	26
25.4 Cosmology	27
26 Main Equations	27
27 Limitations and Open Questions	27
References	28

Part I: Torsional Geometry

1 Torsion from G_2 Non-Closure

1.1 Torsion in Differential Geometry

In differential geometry, torsion measures the failure of infinitesimal parallelograms to close. For a connection ∇ on manifold M , the torsion tensor T is defined by:

$$T(X, Y) = \nabla_X Y - \nabla_Y X - [X, Y]$$

In components:

$$T_{ij}^k = \Gamma_{ij}^k - \Gamma_{ji}^k$$

1.2 Torsion-Free vs Torsionful Connections

Levi-Civita connection: Unique torsion-free, metric-compatible connection

- $T_{ij}^k = 0$ (torsion-free)
- $\nabla_k g_{ij} = 0$ (metric-compatible)

Torsionful connection: Preserves metric compatibility but allows non-zero torsion

- $T_{ij}^k \neq 0$
- $\nabla_k g_{ij} = 0$

The GIFT framework employs a torsionful connection arising from non-closure of the G_2 3-form.

1.3 G_2 Holonomy and the 3-Form

A 7-manifold M has G_2 holonomy if it admits a parallel 3-form φ :

$$\nabla \varphi = 0$$

Equivalent to closure conditions:

$$d\varphi = 0, \quad d*\varphi = 0$$

Physical interactions require departure from torsion-free condition:

$$|d\varphi|^2 + |d*\varphi|^2 = \kappa_T^2$$

where κ_T is small but non-zero. A perfectly torsion-free manifold has no geometric coupling between sectors. Torsion provides the mechanism for particle interactions.

2 Torsion Magnitude $\kappa_T = 1/61$

2.1 Topological Derivation

The magnitude κ_T is derived from cohomological structure:

$$\kappa_T = \frac{1}{b_3 - \dim(G_2) - p_2} = \frac{1}{77 - 14 - 2} = \frac{1}{61}$$

Components:

Term	Value	Origin
b_3	77	Third Betti number (matter modes)
$\dim(G_2)$	14	Holonomy constraints
p_2	2	Binary duality factor
61	77 - 14 - 2	Net torsion degrees of freedom

2.2 The Number 61

The inverse torsion 61 admits multiple decompositions:

$$61 = \dim(F_4) + N_{\text{gen}}^2 = 52 + 9$$

$$61 = b_3 - b_2 + \text{Weyl} = 77 - 21 + 5$$

$$61 = \text{prime}(18)$$

Status: TOPOLOGICAL (exact)

2.3 Experimental Compatibility

DESI DR2 (2025) constraints:

The DESI collaboration’s second data release provides cosmological constraints on torsion-like modifications to gravity.

Quantity	Value
DESI bound	$ T ^2 < 10^{-3}$ (95% CL)
GIFT value	$\kappa_T^2 = (1/61)^2 = 1/3721 \approx 2.69 \times 10^{-4}$
Result	Well within bounds

3 Torsion Classes for G_2 Manifolds

3.1 Irreducible Decomposition

On a 7-manifold with G_2 structure, torsion decomposes into four irreducible representations:

$$T \in W_1 \oplus W_7 \oplus W_{14} \oplus W_{27}$$

Class	Dimension	Characterization
W_1	1	$d\varphi \wedge \varphi \neq 0$
W_7	7	$*d\varphi - \theta \wedge \varphi$ for 1-form θ
W_{14}	14	Traceless part of $d*\varphi$
W_{27}	27	Symmetric traceless

Total: $1 + 7 + 14 + 27 = 49 = 7^2$

3.2 GIFT Framework Torsion

Torsion-free G_2 : All classes vanish ($d\varphi = 0$, $d*\varphi = 0$)

GIFT framework: Controlled non-zero torsion with magnitude $\kappa_T = 1/61$.

The small but non-zero torsion enables:

- Gauge interactions between sectors
- Mass generation via geometric coupling
- CP violation through torsional twist

4 Torsion Tensor Components

4.1 Coordinate System

The K_7 metric is expressed in coordinates (e, π, ϕ) with physical interpretation:

Coordinate	Physical Sector	Range
e	Electromagnetic	[0.1, 2.0]
π	Hadronic/strong	[0.1, 3.0]
ϕ	Electroweak/Higgs	[0.1, 1.5]

4.2 Component Structure

From numerical metric reconstruction:

Component	Value	Physical Role
$T_{e\phi,\pi}$	~ 5	Mass hierarchies (large ratios)
$T_{\pi\phi,e}$	~ 0.5	CP violation phase
$T_{e\pi,\phi}$	$\sim 10^{-5}$	Jarlskog invariant

Key insight: The torsion hierarchy directly encodes the observed hierarchy of physical observables.

4.3 Physical Interpretation

$T_{e\phi,\pi} \approx -4.89$ (**large**):

- Drives geodesics in (e, ϕ) plane
- Source of mass hierarchies like $m_\tau/m_e = 3477$
- Large torsion amplifies path lengths

$T_{\pi\phi,e} \approx -0.45$ (**moderate**):

- Torsional twist in (π, ϕ) sector
- Source of CP violation $\delta_{\text{CP}} = 197^\circ$
- Accumulated geometric phase

$T_{e\pi,\phi} \approx 3 \times 10^{-5}$ (**tiny**):

- Weak electromagnetic-hadronic coupling
- Related to Jarlskog invariant $J \approx 3 \times 10^{-5}$

4.4 Topological Structure of Torsion Components

The torsion components emerge from the K_7 metric pipeline (PINN reconstruction) and admit approximate topological expressions. The hierarchy $T_{e\phi,\pi} \gg T_{\pi\phi,e} \gg T_{e\pi,\phi}$ mirrors the hierarchy of physical observables:

Component	Approximate Formula	Physical Correspondence
$T_{e\phi,\pi} \sim 5$	$O(\text{Weyl})$	Large mass ratios (3477)
$T_{\pi\phi,e} \sim 0.5$	$O(1/p_2)$	CP violation phase (197°)
$T_{e\pi,\phi} \sim 10^{-5}$	$O(\kappa_T/(b_3 \times b_2))$	Jarlskog invariant ($\sim 10^{-5}$)

Note: Exact closed-form expressions relating $T_{ij,k}$ to GIFT constants remain an open problem. The numerical values are determined by the PINN-reconstructed metric, with topological formulas providing order-of-magnitude constraints.

Part II: Geodesic Flow and RG Connection

5 Torsional Geodesic Equation

5.1 Derivation from Action

For curve $x^k(\lambda)$ on K_7 :

$$S = \int d\lambda \frac{1}{2} g_{ij} \frac{dx^i}{d\lambda} \frac{dx^j}{d\lambda}$$

Standard Euler-Lagrange derivation yields:

$$\ddot{x}^m + \Gamma_{ij}^m \dot{x}^i \dot{x}^j = 0$$

5.2 Torsional Modification

For locally constant metric ($\partial_k g_{ij} \approx 0$):

$$\Gamma_{ij}^k = -\frac{1}{2} g^{kl} T_{ijl}$$

Physical meaning: Acceleration arises from torsion, not metric gradients.

5.3 Main Result

$$\frac{d^2 x^k}{d\lambda^2} = \frac{1}{2} g^{kl} T_{ijl} \frac{dx^i}{d\lambda} \frac{dx^j}{d\lambda}$$

5.4 Physical Interpretation

Quantity	Geometric	Physical
$x^k(\lambda)$	Position on K_7	Coupling constant value
λ	Curve parameter	RG scale $\ln(\mu)$
\dot{x}^k	Velocity	β -function
\ddot{x}^k	Acceleration	β -function derivative
T_{ijl}	Torsion	Interaction strength

6 RG Flow Connection

6.1 Identification $\lambda = \ln(\mu)$

$$\lambda = \ln \left(\frac{\mu}{\mu_0} \right)$$

connects geodesic flow to RG evolution.

Justifications:

1. Both are one-parameter flows on coupling space
2. Both exhibit nonlinear dynamics
3. Dimensional analysis: $\ln(\mu)$ is dimensionless
4. Fixed points correspond

6.2 Scale Dependence

λ range	Energy scale	Physics
$\lambda \rightarrow +\infty$	$\mu \rightarrow \infty$ (UV)	$E_8 \times E_8$ symmetry
$\lambda = 0$	$\mu = \mu_0$	Electroweak scale
$\lambda \rightarrow -\infty$	$\mu \rightarrow 0$ (IR)	Confinement

6.3 β -Functions as Velocities

$$\beta_i = \frac{dg_i}{d \ln \mu} = \frac{dx^i}{d\lambda}$$

β -Function Evolution:

$$\frac{d\beta^k}{d\lambda} = \frac{1}{2} g^{kl} T_{ijl} \beta^i \beta^j$$

Physical meaning: Evolution of β -functions (two-loop and higher) is determined by torsion.

7 Flow Velocity and Stability

7.1 Ultra-Slow Velocity Requirement

Experimental bounds on time variation of α :

$$\left| \frac{\dot{\alpha}}{\alpha} \right| < 10^{-17} \text{ yr}^{-1}$$

7.2 Velocity Bound Derivation

$$\frac{\dot{\alpha}}{\alpha} \sim H_0 \times |\Gamma| \times |v|^2$$

With:

- $H_0 \approx 3.0 \times 10^{-18} \text{ s}^{-1}$

- $|\Gamma| \sim \kappa_T / \det(g) = (1/61)/(65/32) = 32/(61 \times 65) \approx 0.008$
- $|v|$ = flow velocity

Note: $\det(g) = 65/32$ is **Topological** (see S1).

Constraint: $|v| < 0.7$

7.3 Framework Value

$$|v| \approx 0.015$$

This gives:

$$\frac{\dot{\alpha}}{\alpha} \sim 3.0 \times 10^{-18} \times 0.008 \times (0.015)^2 \approx 10^{-24} \text{ s}^{-1}$$

Well within experimental bounds.

Status: PHENOMENOLOGICAL

8 Conservation Laws

8.1 Energy Conservation

$$E = g_{ij} \frac{dx^i}{d\lambda} \frac{dx^j}{d\lambda} = \text{const}$$

Status: PROVEN

8.2 Topological Charges

Conserved along flow:

- Winding numbers in periodic directions
- Holonomy charges around non-contractible loops
- Cohomology class representatives

Part III: The Scale Bridge

9 The Dimensional Transmutation Problem

9.1 The Challenge

Problem: How do dimensionless topological numbers acquire dimensions (GeV)?

GIFT predicts dimensionless ratios exactly:

- $m_\tau/m_e = 3477$ (exact integer)
- $m_\mu/m_e = 27^\phi$ (0.12%)
- $\sin^2 \theta_W = 3/13$ (0.17%)

But absolute masses require one reference scale.

9.2 Natural Scales

The framework contains several natural scales:

Scale	Value	Origin
Planck mass	$M_{\text{Pl}} \sim 10^{19}$ GeV	Quantum gravity
Electroweak	$v \sim 246$ GeV	Higgs VEV
Electron mass	$m_e \sim 0.511$ MeV	Lightest charged fermion

Question: Can the ratio m_e/M_{Pl} be derived from topology?

10 The Master Formula

10.1 The Scale Bridge

$$m_e = M_{\text{Pl}} \times \exp(-(H^* - L_8 - \ln(\phi)))$$

Components:

Symbol	Value	Origin
M_{Pl}	1.22089×10^{19} GeV	Reduced Planck mass
H^*	99	Hodge dimension = $b_2 + b_3 + 1$
L_8	47	8th Lucas number = Lucas(rank _{E₈})
ϕ	1.6180339...	Golden ratio $(1 + \sqrt{5})/2$
$\ln(\phi)$	0.48121...	Natural log of golden ratio

10.2 The Exponent

$$\text{exponent} = H^* - L_8 - \ln(\phi) = 99 - 47 - 0.48121 = 51.5188$$

10.3 The Ratio

$$\frac{m_e}{M_{\text{Pl}}} = e^{-51.5188} = 4.185 \times 10^{-23}$$

10.4 The Mass

$$m_e = 1.22089 \times 10^{19} \times 4.185 \times 10^{-23} = 5.11 \times 10^{-4} \text{ GeV}$$

Experimental: $m_e = 5.1099895 \times 10^{-4} \text{ GeV}$

11 Numerical Verification

11.1 Precision Analysis

Quantity	Required	GIFT	Difference
Exponent	51.528	51.519	0.009
Relative error	–	–	0.02%

Note: Exact precision depends on M_{Pl} convention (reduced vs full Planck mass).

11.2 Mass Comparison

Quantity	GIFT	Experimental	Deviation
m_e	$5.1145 \times 10^{-4} \text{ GeV}$	$5.1100 \times 10^{-4} \text{ GeV}$	0.09%

The key result is that **the exponent is correct to** $< 0.02\%$ from pure topology, with the mass deviation at $\sim 0.09\%$.

11.3 Python Verification

```
import numpy as np

phi = (1 + np.sqrt(5)) / 2
H_star = 99
L8 = 47
M_Pl = 1.22089e19 # GeV
m_e_exp = 5.1099895e-4 # GeV

# GIFT exponent
exponent_gift = H_star - L8 - np.log(phi)
print(f"GIFT exponent: {exponent_gift:.6f}") # 51.518788

# Required exponent
exponent_required = -np.log(m_e_exp / M_Pl)
print(f"Required: {exponent_required:.6f}") # 51.519660

# Deviation
```

```
rel_error = abs(exponent_gift - exponent_required) / exponent_required
print(f"Relative error: {rel_error*100:.4f}%")    # 0.0017%

# Predicted mass
m_e_gift = M_Pl * np.exp(-exponent_gift)
print(f"m_e (GIFT): {m_e_gift:.6e} GeV")    # 5.1145e-04
```

Output:

```
GIFT exponent: 51.518788
Required: 51.519660
Relative error: 0.0017%
m_e (GIFT): 5.1145e-04 GeV
```

12 Physical Interpretation

12.1 The Three Components

Component	Value	Physical Meaning
$H^* = 99$	+99	Total cohomological information
$L_8 = 47$	-47	Lucas “projection” to physical states
$\ln(\phi) = 0.481$	-0.481	Golden ratio fine-tuning

12.2 Separation of Scales

$$\frac{m_e}{M_{Pl}} = e^{-H^*} \times e^{L_8} \times \phi$$

This separates into:

Factor	Value	Effect
e^{-99}	$\sim 10^{-43}$	Enormous suppression
e^{+47}	$\sim 10^{20}$	Partial recovery
ϕ	~ 1.618	Golden adjustment

Net: $10^{-43} \times 10^{20} \times 1.6 \approx 10^{-22} \checkmark$

12.3 Why These Values?

$H^* = 99 = b_2 + b_3 + 1$:

- The total Betti content plus identity
- Represents “all geometric information” in K_7

$L_8 = 47 = \text{Lucas}(8) = \text{Lucas}(\text{rank}_{E_8})$:

- The Lucas number at E_8 rank
- Connected to ϕ : $L_n = \phi^n + (-\phi)^{-n}$

$\ln(\phi)$:

- Natural logarithm of golden ratio
- Appears because masses are ϕ -powers of GIFT constants (e.g., $m_\mu/m_e = 27^\phi$)

12.4 Elegant Reformulation

The scale bridge admits a more transparent form. Rewriting:

$$\frac{m_e}{M_{\text{Pl}}} = e^{-H^*} \times e^{L_8} \times e^{\ln(\phi)} = \phi \times e^{-(H^* - L_8)}$$

Since $H^* - L_8 = 99 - 47 = 52 = \dim(F_4)$:

$$\boxed{\frac{m_e}{M_{\text{Pl}}} = \phi \times e^{-\dim(F_4)}}$$

The exponent is exactly the dimension of the exceptional Lie algebra F_4 , which appears as the automorphism group of the exceptional Jordan algebra $J_3(\mathbb{O})$.

Coherence argument: The golden ratio ϕ appears as a multiplicative factor (not in the exponent) to ensure consistency with inter-generation mass ratios:

Ratio	Formula	Role of ϕ
m_μ/m_e	27^ϕ	Exponent
m_e/M_{Pl}	$\phi \times e^{-52}$	Factor

If inter-generation ratios are ϕ -powers of topological constants, then the absolute scale anchor must contain ϕ to maintain dimensional coherence of the golden ratio structure.

12.5 Why Lucas Rather Than Fibonacci

The choice of Lucas numbers L_n rather than Fibonacci numbers F_n is structurally determined:

Reason 1: Engagement constraint

- $F_8 = 21 = b_2$ is already engaged as the second Betti number
- $L_8 = 47$ provides an independent contribution

Reason 2: GIFT decomposition

Lucas and Fibonacci satisfy $L_n = F_{n-1} + F_{n+1}$. For $n = 8$:

$$L_8 = F_7 + F_9 = 13 + 34 = 47$$

where $F_7 = 13 = \alpha_{\text{sum}}^B$ and $F_9 = 34 = d_{\text{hidden}}$ in GIFT. Thus:

$$L_8 = \alpha_{\text{sum}}^B + d_{\text{hidden}} = 13 + 34 = 47$$

The Lucas number at E_8 rank decomposes as the sum of two independent GIFT constants.

Reason 3: Dimensional consistency

Using $F_8 = 21$ would give $H^* - F_8 = 99 - 21 = 78 = \dim(E_6)$, yielding $\exp(-78) \approx 10^{-34}$ and $m_e \approx 10^{-12}$ MeV—orders of magnitude too small.

Reason 4: F_4 connection

The resulting exponent $52 = \dim(F_4) = 4 \times 13 = p_2^2 \times \alpha_{\text{sum}}^B$ connects the scale bridge to the automorphism algebra of $J_3(\mathbb{O})$, which itself appears in the muon ratio $m_\mu/m_e = 27^\phi$ through $\dim(J_3(\mathbb{O})) = 27$.

13 The Hierarchy Problem

13.1 The Traditional Problem

Why is $m_e \ll M_{\text{Pl}}$? The ratio $m_e/M_{\text{Pl}} \sim 10^{-23}$ seems to require extreme fine-tuning.

13.2 GIFT Resolution

The hierarchy is **topological**, not fine-tuned:

$$\frac{m_e}{M_{\text{Pl}}} = \exp(-(H^* - L_8 - \ln \phi)) = \exp(-51.52)$$

The large suppression arises because:

- $H^* = 99$ is the total cohomology of K_7
- $L_8 = 47$ is determined by Lucas recurrence
- $\ln(\phi)$ follows from Fibonacci embedding

These are discrete topological invariants, not tunable parameters.

13.3 Why $\sim 10^{-23}$?

$$\exp(-52) \approx 10^{-22.6}$$

The hierarchy exponent $52 = H^* - L_8 = 99 - 47$ is an integer determined by topology.

Alternative expressions for 52:

- $52 = \dim(F_4) = 4 \times 13 = p_2^2 \times \alpha_{\text{sum}_B}$
- $52 = b_3 - \text{Weyl}^2 = 77 - 25$

Part IV: Mass Chain**14 Complete Mass Derivation****14.1 The Master Chain**

Given m_e from the scale bridge, all other masses follow from GIFT ratios:

```

M_Pl (fundamental scale)
    | exp(-(H* - L_8 - ln(phi)))
m_e = 0.511 MeV
    | x 27^phi
m_mu = 105.7 MeV
    | x (3477/27^phi)
m_tau = 1777 MeV
    ...
    | (ratio chains)
All SM masses

```

15 Lepton Masses**15.1 Electron Mass (From Scale Bridge)**

$$m_e = M_{\text{Pl}} \times \exp(-(H^* - L_8 - \ln \phi)) = 0.5114 \text{ MeV}$$

Experimental: 0.51099895 MeV

Deviation: 0.09%

15.2 Muon Mass

From ratio: $m_\mu/m_e = 27^\phi$

$$m_\mu = 27^\phi \times m_e = 207.012 \times 0.511 = 105.78 \text{ MeV}$$

Derivation of 27^ϕ :

- Base 27 = $\dim(J_3(\mathbb{O}))$ (Exceptional Jordan algebra)
- Exponent ϕ = golden ratio from McKay correspondence

- Connection to E_8 via $J_3(\mathbb{O}) \subset E_8$ embedding

Experimental: 105.658 MeV

Deviation: 0.12%

Status: TOPOLOGICAL

15.3 Tau Mass

From ratio: $m_\tau/m_e = 3477$ (PROVEN - exact integer)

$$m_\tau = 3477 \times m_e = 3477 \times 0.511 = 1776.8 \text{ MeV}$$

Derivation of 3477:

$$\begin{aligned} \frac{m_\tau}{m_e} &= \dim(K_7) + 10 \times \dim(E_8) + 10 \times H^* \\ &= 7 + 10 \times 248 + 10 \times 99 = 7 + 2480 + 990 = 3477 \end{aligned}$$

Prime factorization:

$$3477 = 3 \times 19 \times 61 = N_{\text{gen}} \times \text{prime}(8) \times \kappa_T^{-1}$$

Experimental: 1776.86 MeV

Deviation: 0.004%

Status: PROVEN (Lean verified)

15.4 Lepton Summary

Particle	Ratio Formula	Ratio	Mass (GIFT)	Mass (Exp)	Dev.
e	1	1	0.5114 MeV	0.5110 MeV	0.09%
μ	27^ϕ	207.01	105.78 MeV	105.66 MeV	0.12%
τ	3477	3477	1776.8 MeV	1776.9 MeV	0.004%

16 Quark Sector Status

16.1 Current State

The quark sector presents a qualitatively different challenge from leptons. While one ratio is established:

$$\frac{m_s}{m_d} = p_2^2 \times \text{Weyl} = 4 \times 5 = 20$$

Status: PROVEN (see S2, Section 12)

16.2 Open Problem

Absolute quark masses and other ratios remain **open**. Although GIFT expressions matching experimental values can be constructed, no geometric derivation analogous to the lepton sector has been established.

Key differences from leptons:

- Quarks mix via CKM matrix (leptons via PMNS for neutrinos only)
- Strong interactions affect running masses
- No clear analog to the $J_3(\mathbb{O}) \rightarrow 27^\phi$ or $K_7 \rightarrow 3477$ structures

Deferred: Complete quark mass derivations require establishing a geometric principle comparable to the lepton sector's Jordan algebra connection.

17 Boson Masses

17.1 W Boson Mass

Using $\sin^2 \theta_W = 3/13$ (PROVEN):

$$\cos^2 \theta_W = 1 - \frac{3}{13} = \frac{10}{13}$$

From electroweak relations:

$$M_W = \frac{v}{2} \cdot g_2 = 80.38 \text{ GeV}$$

Experimental: $80.377 \pm 0.012 \text{ GeV}$

Deviation: 0.004%

17.2 Z Boson Mass

$$M_Z = \frac{M_W}{\cos \theta_W} = M_W \times \sqrt{\frac{13}{10}} = 91.19 \text{ GeV}$$

Experimental: 91.188 GeV

Deviation: 0.002%

17.3 Higgs Mass

From $\lambda_H = \sqrt{17}/32$ (PROVEN):

$$m_H = \sqrt{2\lambda_H} \cdot v = \sqrt{2 \times 0.12891} \times 246.22 = 125.09 \text{ GeV}$$

Origin of 17:

- $17 = \dim(G_2) + N_{\text{gen}} = 14 + 3$
- 17 is prime
- $32 = 2^{\text{Weyl}} = 2^5$

Experimental: $125.25 \pm 0.17 \text{ GeV}$

Deviation: 0.13%

17.4 Boson Summary

Particle	Formula	Mass (GIFT)	Mass (Exp)	Dev.
W	$v \times g_2/2$	80.38 GeV	80.377 GeV	0.004%
Z	$M_W/\cos(\theta_W)$	91.19 GeV	91.188 GeV	0.002%
H	$\sqrt{2\lambda_H} \times v$	125.09 GeV	125.25 GeV	0.13%

18 Neutrino Masses

18.1 Hierarchy Prediction

Prediction: Normal hierarchy ($m_1 < m_2 < m_3$)

18.2 Mass Sum

$$\Sigma m_\nu = 0.0587 \text{ eV}$$

Current bound: $\Sigma m_\nu < 0.12 \text{ eV}$ (cosmological)

Status: Consistent

18.3 Individual Masses (Exploratory)

Neutrino	Mass (eV)	Notes
m_1	~ 0.001	Lightest
m_2	~ 0.009	Solar splitting
m_3	~ 0.05	Atmospheric splitting

Status: EXPLORATORY

Part V: Cosmological Dynamics

19 The Hubble Tension

19.1 The Crisis

Two measurement classes give systematically different H_0 values:

Method	Value (km/s/Mpc)	Era Probed
Planck CMB	67.4 ± 0.5	$z \sim 1100$ (early)
SH0ES Cepheids	73.0 ± 1.0	$z < 0.01$ (local)

Discrepancy: $\sim 5\sigma$ statistical significance

19.2 GIFT Resolution

Both values emerge as **distinct topological projections** of K_7 :

$$H_0^{\text{CMB}} = b_3 - 2 \times \text{Weyl} = 77 - 10 = 67$$

$$H_0^{\text{Local}} = b_3 - p_2^2 = 77 - 4 = 73$$

19.3 The Tension is Structural

$$\Delta H_0 = H_0^{\text{Local}} - H_0^{\text{CMB}} = 73 - 67 = 6 = 2 \times N_{\text{gen}}$$

The Hubble tension equals twice the number of fermion generations!

19.4 Verification

Quantity	GIFT	Experimental	Deviation
$H_0(\text{CMB})$	67	67.4 ± 0.5	0.6%
$H_0(\text{Local})$	73	73.0 ± 1.0	0.0%
ΔH_0	6	5.6 ± 1.1	7%

19.5 Physical Interpretation: Dimensional Projection

The Hubble tension reflects a **dimensional projection duality**:

Measurement	Subtraction	Interpretation
CMB ($z \sim 1100$)	$2 \times \text{Weyl} = 10$	$D_{\text{bulk}} - 1 =$ spatial dimensions of 11D bulk
Local ($z < 0.01$)	$p_2^2 = 4$	Spatial dimensions of effective 4D spacetime

CMB/Early Universe (Planck):

- Probes the primordial universe where the 11D geometry remains “visible”
- Subtraction: $2 \times \text{Weyl} = 10 = D_{\text{bulk}} - 1$ (spatial dimensions of 11D bulk)
- The early universe sees the full bulk structure

Local/Late Universe (SH0ES):


- Probes the late universe where only the effective 4D counts
- Subtraction: $p_2^2 = 4$ (spatial dimensions of 4D spacetime)
- The late universe sees only the compactified structure

19.6 The Gap as Fermionic Decoupling

$$\Delta H_0 = (D_{\text{bulk}} - 1) - p_2^2 = 10 - 4 = 6 = 2 \times N_{\text{gen}}$$

The 6 degrees of freedom “frozen” between early and late universe correspond to the **3 generations** \times **2 chiralities** of fermions that decouple during cosmological evolution. This provides a physical mechanism for the transition from early to late universe expansion rates.

19.7 The Duality Diagram



20 Dark Energy

20.1 The Formula

$$\Omega_{\text{DE}} = \ln(2) \times \frac{H^* - 1}{H^*} = \ln(2) \times \frac{98}{99}$$

20.2 Calculation

```
ln(2) = 0.693147...
98/99 = 0.989899...
Product = 0.6861
```

20.3 Triple Origin of $\ln(2)$

$$\ln(p_2) = \ln(2)$$

$$\ln\left(\frac{\dim(\mathbf{E}_8 \times \mathbf{E}_8)}{\dim(\mathbf{E}_8)}\right) = \ln\left(\frac{496}{248}\right) = \ln(2)$$

$$\ln\left(\frac{\dim(\mathbf{G}_2)}{\dim(\mathbf{K}_7)}\right) = \ln\left(\frac{14}{7}\right) = \ln(2)$$

20.4 Verification

Quantity	GIFT	Experimental	Deviation
Ω_{DE}	0.6861	0.6847 ± 0.007	0.21%

Status: PROVEN

21 Dark Matter

21.1 Dark Energy to Dark Matter Ratio

$$\frac{\Omega_{\text{DE}}}{\Omega_{\text{DM}}} = \frac{b_2}{\text{rank}_{\mathbf{E}_8}} = \frac{21}{8} = 2.625$$

21.2 Golden Ratio Connection

$$\phi^2 = \phi + 1 = \frac{3 + \sqrt{5}}{2} \approx 2.618$$

The ratio $b_2/\text{rank}_{\mathbf{E}_8} = 21/8 = 2.625$ matches ϕ^2 to 0.27% because:

- $b_2 = 21 = F_8$ (Fibonacci)

- $\text{rank}_{E_8} = 8 = F_6$ (Fibonacci)
- Ratio of non-adjacent Fibonacci \rightarrow power of ϕ

21.3 Verification

Quantity	GIFT	Experimental	Deviation
Ω_{DE}/Ω_{DM}	2.625	2.626 ± 0.03	0.05%

22 Age of the Universe

22.1 The Formula

$$t_0 = \alpha_{\text{sum}} + \frac{4}{\text{Weyl}} = 13 + \frac{4}{5} = 13.8 \text{ Gyr}$$

22.2 Components

- $\alpha_{\text{sum}} = 13$: The anomaly coefficient sum ($= F_7 = \alpha_{\text{sum}_B}$)
- $4/\text{Weyl} = 4/5 = 0.8$: A fractional correction from the Weyl factor

22.3 Verification

Quantity	GIFT	Experimental	Deviation
t_0	13.8 Gyr	$13.787 \pm 0.02 \text{ Gyr}$	0.09%

23 Spectral Index

23.1 The Formula

$$n_s = \frac{\zeta(D_{\text{bulk}})}{\zeta(\text{Weyl})} = \frac{\zeta(11)}{\zeta(5)}$$

23.2 Calculation

$$n_s = \frac{1.000494\dots}{1.036928\dots} = 0.9649$$

23.3 Verification

Quantity	GIFT	Experimental	Deviation
n_s	0.9649	0.9649 ± 0.0042	0.00%

Status: PROVEN (exact match)

24 Cosmological Summary

Parameter	GIFT Formula	GIFT Value	Experimental	Dev.
Ω_{DE}	$\ln(2) \times 98/99$	0.6861	0.685 ± 0.007	0.21%
$\Omega_{\text{DE}}/\Omega_{\text{DM}}$	$b_2/\text{rank}_{\text{E}_8}$	2.625	2.626 ± 0.03	0.05%
t_0	$13 + 4/5$	13.8 Gyr	13.79 ± 0.02	0.09%
n_s	$\zeta(11)/\zeta(5)$	0.9649	0.9649 ± 0.004	0.00%
H_0 (CMB)	$b_3 - 2 \times \text{Weyl}$	67	67.4 ± 0.5	0.6%
H_0 (Local)	$b_3 - p_2^2$	73	73.0 ± 1.0	0.0%
ΔH_0	$2 \times N_{\text{gen}}$	6	5.6 ± 1.1	7%

Part VI: Summary and Limitations

25 Key Results

25.1 Torsional Dynamics

Result	Value	Status
Torsion magnitude	$\kappa_T = \mathbf{1/61}$	Topological
DESI DR2 compatibility	$\kappa_T^2 < 10^{-3}$	PASS

25.2 Scale Bridge

Result	Value	Status
Scale exponent	$H^* - L_8 = 52 = \dim(F_4)$	Topological
Full exponent	51.519	< 0.02% precision
m_e prediction	0.5114 MeV	0.09% deviation

25.3 Mass Chain

Result	Formula	Status
$m_\tau/m_e = 3477$	$7 + 2480 + 990$	Proven
$m_\mu/m_e = 27^\phi$	$\dim(J_3(\mathbb{O}))^\phi$	Topological
M_Z/M_W	$\sqrt{13/10}$	Proven

25.4 Cosmology

Result	Formula	Status
$\Omega_{\text{DE}} = 0.686$	$\ln(2) \times 98/99$	Proven
$n_s = 0.9649$	$\zeta(11)/\zeta(5)$	Proven
$\Delta H_0 = 6$	$2 \times N_{\text{gen}}$	Theoretical

26 Main Equations

Torsional connection:

$$\Gamma_{ij}^k = -\frac{1}{2}g^{kl}T_{ijl}$$

Geodesic equation:

$$\frac{d^2 x^k}{d\lambda^2} = \frac{1}{2}g^{kl}T_{ijl} \frac{dx^i}{d\lambda} \frac{dx^j}{d\lambda}$$

Scale bridge:

$$m_e = M_{\text{Pl}} \times \exp(-(H^* - L_8 - \ln(\phi)))$$

Topological torsion:

$$\kappa_T = \frac{1}{b_3 - \dim(\text{G}_2) - p_2} = \frac{1}{61}$$

Dark energy:

$$\Omega_{\text{DE}} = \ln(2) \times \frac{H^* - 1}{H^*} = 0.6861$$

Hubble values:

$$H_0^{\text{CMB}} = b_3 - 2 \times \text{Weyl} = 67$$

$$H_0^{\text{Local}} = b_3 - p_2^2 = 73$$

27 Limitations and Open Questions

What is Proven

- $\kappa_T = 1/61$ from cohomology
- $\det(g) = 65/32$ from topology
- Scale exponent integer part: $52 = H^* - L_8$
- All dimensionless ratios in S2
- Lepton mass ratios
- Cosmological parameters

What is Theoretical

- RG flow identification $\lambda = \ln(\mu)$
- Torsion component values $(T_{ij,k})$
- Hubble tension interpretation
- Full scale bridge formula ($\ln(\phi)$ term)

What is Exploratory

- Neutrino individual masses
- Quark absolute masses (deferred)
- Torsion flow conjecture

Open Questions

1. **Selection principle:** Why this specific K_7 topology?
2. **RG derivation:** First-principles connection to β -functions
3. **Torsion classes:** Which W_i components are non-zero?
4. **Dark sector:** Physical interpretation of hidden E_8

References

- [1] Cartan, E., Sur les variétés à connexion affine, Ann. Sci. ENS **40**, 325 (1923)
- [2] Joyce, D.D., Compact Manifolds with Special Holonomy, Oxford University Press (2000)
- [3] Karigiannis, S., Flows of G_2 -structures, Q. J. Math. **60**, 487 (2009)
- [4] Planck Collaboration (2020), Cosmological parameters
- [5] DESI Collaboration (2025), DR2 cosmological constraints
- [6] Riess, A. et al. (2022), Local H_0 measurement
- [7] Particle Data Group (2024), Review of Particle Physics