

Supplement S1: Mathematical Foundations

E_8 Exceptional Lie Algebra, G_2 Holonomy Manifolds, and K_7 Construction

Version: 3.2

Author: Briec de La Fournière

Independent researcher

Abstract

This supplement presents the mathematical architecture underlying GIFT. Part I develops E_8 exceptional Lie algebra with the Exceptional Chain theorem. Part II introduces G_2 holonomy manifolds. Part III establishes K_7 manifold construction via twisted connected sum, building compact G_2 manifolds by gluing asymptotically cylindrical building blocks. Part IV establishes the algebraic reference form $\varphi_{\text{ref}} = (65/32)^{1/14} \times \varphi_0$ with exact $\det(g) = 65/32$; Joyce's theorem ensures a torsion-free metric exists. Core algebraic relations are formally verified in Lean 4 (v3.2.0).

Contents

1	The Octonionic Foundation	3
1.1	Why This Framework Exists	3
1.2	The Division Algebra Chain	3
1.3	G_2 as Octonionic Automorphisms	3
1.4	Why $\dim(K_7) = 7$	4
2	E_8 Exceptional Lie Algebra	4
2.1	Root System and Dynkin Diagram	4
2.2	Basic Data	4
2.3	Root System Construction	4
2.4	Cartan Matrix	5
2.5	Weyl Group	5
2.6	Order and Factorization	5
2.7	Topological Factorization Theorem	5
2.8	Exceptional Chain	5
2.9	The Pattern	5
2.10	Exceptional Chain Theorem	6
2.11	$E_8 \times E_8$ Product Structure	6
2.12	Direct Sum	6
2.13	τ Numerator Connection	6
2.14	Binary Duality Parameter	6
2.15	Exceptional Algebras from Octonions	6
2.16	Exceptional Jordan Algebra $J_3(\mathbb{O})$	7
2.17	F_4 Connection	7
2.18	Exceptional Differences	7
3	G_2 Holonomy Manifolds	7
3.1	Definition and Properties	7
3.2	G_2 as Exceptional Holonomy	7
3.3	Holonomy Classification (Berger)	7
3.4	Torsion: Definition and GIFT Interpretation	8
3.5	Topological Invariants	8
3.6	Derived Constants	8

3.7	The 61 Decomposition	8
4	K_7 Manifold Construction	9
4.1	Twisted Connected Sum Framework	9
4.2	TCS Construction	9
4.3	Asymptotically Cylindrical G_2 Manifolds	9
4.4	Building Blocks	9
4.5	Cohomological Structure	10
4.6	Mayer-Vietoris Analysis	10
4.7	Betti Number Derivation	10
4.8	Complete Betti Spectrum	10
4.9	Third Betti Number Decomposition	10
5	Metric Structure and Verification	11
5.1	Structural Metric Invariants	11
5.2	The Zero-Parameter Paradigm	11
5.3	Torsion Capacity $\kappa_T = 1/61$	11
5.4	Metric Determinant $\det(g) = 65/32$	12
5.5	Formal Certification	12
5.6	Algebraic Reference Form	12
5.7	Actual Solution Structure	12
5.8	Why GIFT Predictions Are Robust	13
5.9	Torsion and Joyce's Theorem	13
5.10	Independent Numerical Validation (PINN)	13
5.11	Lean 4 Formalization	14
5.12	The Derivation Chain	14
5.13	Analytical G_2 Metric Details	15
5.14	The Standard Form φ_0	15
5.15	Linear Index Representation	15
5.16	Metric Derivation	15
5.17	Comparison: Fano Plane vs G_2 Form	16
5.18	Verification Summary	16
5.19	Summary	16


1 The Octonionic Foundation

1.1 Why This Framework Exists

GIFT is not built on arbitrary choices. It emerges from a single algebraic fact:

The octonions \mathbb{O} are the largest normed division algebra.

Everything follows:



1.2 The Division Algebra Chain

Algebra	Dim	Physics Role	Stops?
\mathbb{R}	1	Classical mechanics	No
\mathbb{C}	2	Quantum mechanics	No
\mathbb{H}	4	Spin, Lorentz group	No
\mathbb{O}	8	Exceptional structures	Yes

The pattern terminates at \mathbb{O} . There is no 16-dimensional normed division algebra. The octonions are *the end of the line*.

1.3 G_2 as Octonionic Automorphisms

Definition: $G_2 = \{g \in GL(\mathbb{O}) : g(xy) = g(x)g(y) \text{ for all } x, y \in \mathbb{O}\}$

Property	Value	GIFT Role
dim(G_2)	$14 = \binom{7}{2}$	Q_{Koide} numerator
Action	Transitive on $S^6 \subset \text{Im}(\mathbb{O})$	Connects all directions
Embedding	$G_2 \subset SO(7)$	Preserves φ_0

1.4 Why $\dim(K_7) = 7$

This is not a choice. It is a consequence:

- $\text{Im}(\mathbb{O})$ has dimension 7
- G_2 acts naturally on \mathbb{R}^7
- A compact 7-manifold with G_2 holonomy is the geometric realization

K_7 is to G_2 what the circle is to $U(1)$.

2 E_8 Exceptional Lie Algebra

2.1 Root System and Dynkin Diagram

2.2 Basic Data

Property	Value	GIFT Role
Dimension	$\dim(E_8) = 248$	Gauge DOF
Rank	$\text{rank}(E_8) = 8$	Cartan subalgebra
Number of roots	$ \Phi(E_8) = 240$	E_8 kissing number
Root length	$\sqrt{2}$	α_s numerator
Coxeter number	$h = 30$	Icosahedron edges
Dual Coxeter number	$h^\vee = 30$	McKay correspondence

2.3 Root System Construction

E_8 root system in \mathbb{R}^8 has 240 roots:

Type I (112 roots): Permutations and sign changes of $(\pm 1, \pm 1, 0, 0, 0, 0, 0, 0)$

Type II (128 roots): Half-integer coordinates with even minus signs:

$$\frac{1}{2}(\pm 1, \pm 1, \pm 1, \pm 1, \pm 1, \pm 1, \pm 1, \pm 1)$$

Verification: $112 + 128 = 240$ roots, all length $\sqrt{2}$.

2.4 Cartan Matrix

$$A_{E_8} = \begin{pmatrix} 2 & 0 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 2 & 0 & -1 & 0 & 0 & 0 & 0 \\ -1 & 0 & 2 & -1 & 0 & 0 & 0 & 0 \\ 0 & -1 & -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 2 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 2 \end{pmatrix}$$

Properties: $\det(A) = 1$ (unimodular), positive definite.

2.5 Weyl Group

2.6 Order and Factorization

$$|W(E_8)| = 696,729,600 = 2^{14} \times 3^5 \times 5^2 \times 7$$

2.7 Topological Factorization Theorem

Theorem: The Weyl group order factorizes entirely into GIFT constants:

$$|W(E_8)| = p_2^{\dim(G_2)} \times N_{\text{gen}}^{\text{Weyl}} \times \text{Weyl}^{p_2} \times \dim(K_7)$$

Factor	Exponent	Value	GIFT Origin
2^{14}	$\dim(G_2) = 14$	16384	$p_2^{(\text{holonomy dim})}$
3^5	$\text{Weyl} = 5$	243	$N_{\text{gen}}^{(\text{Weyl factor})}$
5^2	$p_2 = 2$	25	$\text{Weyl}^{(\text{binary})}$
7^1	1	7	$\dim(K_7)$

Status: Proven (Lean): `weyl_E8_topological_factorization`

2.8 Exceptional Chain

2.9 The Pattern

A pattern connects exceptional algebra dimensions to primes:

Algebra	n	$\dim(E_n)$	Prime	Index
E_6	6	78	13	$\text{prime}(6)$
E_7	7	133	19	$\text{prime}(8) = \text{prime}(\text{rank}(E_8))$
E_8	8	248	31	$\text{prime}(11) = \text{prime}(D_{\text{bulk}})$

2.10 Exceptional Chain Theorem

Theorem: For $n \in \{6, 7, 8\}$:

$$\dim(E_n) = n \times \text{prime}(g(n))$$

where $g(6) = 6$, $g(7) = \text{rank}(E_8) = 8$, $g(8) = D_{\text{bulk}} = 11$.

Proof (verified in Lean):

- E_6 : $6 \times 13 = 78$ ✓
- E_7 : $7 \times 19 = 133$ ✓
- E_8 : $8 \times 31 = 248$ ✓

Status: Proven (Lean): `exceptional_chain_certified`

2.11 $E_8 \times E_8$ Product Structure

2.12 Direct Sum

Property	Value
Dimension	$496 = 248 \times 2$
Rank	$16 = 8 \times 2$
Roots	$480 = 240 \times 2$

2.13 τ Numerator Connection

The hierarchy parameter numerator:

$$\tau_{\text{num}} = 3472 = 7 \times 496 = \dim(K_7) \times \dim(E_8 \times E_8)$$

Status: Proven (Lean): `tau_num_E8xE8`

2.14 Binary Duality Parameter

Triple geometric origin of $p_2 = 2$:

1. **Local:** $p_2 = \dim(G_2)/\dim(K_7) = 14/7 = 2$
2. **Global:** $p_2 = \dim(E_8 \times E_8)/\dim(E_8) = 496/248 = 2$
3. **Root:** $\sqrt{2}$ in E_8 root normalization

2.15 Exceptional Algebras from Octonions

The foundational role of octonions is established in Part 0. This section details the exceptional algebraic structures that emerge from \mathbb{O} .

2.16 Exceptional Jordan Algebra $J_3(\mathbb{O})$

Property	Value
$\dim(J_3(\mathbb{O}))$	$27 = 3^3$
$\dim(J_3(\mathbb{O})_0)$	26 (traceless)

2.17 F_4 Connection

F_4 is the automorphism group of $J_3(\mathbb{O})$:

$$\dim(F_4) = 52 = p_2^2 \times \alpha_{\text{sum}}^B = 4 \times 13$$

2.18 Exceptional Differences

Difference	Value	GIFT
$\dim(E_8) - \dim(J_3(\mathbb{O}))$	$221 = 13 \times 17$	$\alpha_B \times \lambda_{H,\text{num}}$
$\dim(F_4) - \dim(J_3(\mathbb{O}))$	$25 = 5^2$	Weyl^2
$\dim(E_6) - \dim(F_4)$	26	$\dim(J_3(\mathbb{O})_0)$

Status: Proven (Lean): `exceptional_differences_certified`

3 G_2 Holonomy Manifolds

3.1 Definition and Properties

3.2 G_2 as Exceptional Holonomy

Property	Value	GIFT Role
$\dim(G_2)$	14	Q_{Koide} numerator
$\text{rank}(G_2)$	2	Lie rank
Definition	$\text{Aut}(\mathbb{O})$	Octonion automorphisms

3.3 Holonomy Classification (Berger)

Dimension	Holonomy	Geometry
7	G_2	Exceptional
8	$\text{Spin}(7)$	Exceptional

3.4 Torsion: Definition and GIFT Interpretation

Mathematical definition: Torsion measures failure of G_2 structure to be parallel:

$$T = \nabla\varphi \neq 0$$

For the 3-form φ , torsion decomposes into four classes $W_1 \oplus W_7 \oplus W_{14} \oplus W_{27}$ with total dimension $1 + 7 + 14 + 27 = 49$.

Torsion-free condition:

$$\nabla\varphi = 0 \Leftrightarrow d\varphi = 0 \text{ and } d*\varphi = 0$$

GIFT interpretation:

Quantity	Meaning	Value
$\kappa_T = 1/61$	Topological <i>capacity</i> for torsion	Fixed by K_7
T_{realized}	Actual torsion for specific solution	Depends on φ
$T_{\text{analytical}}$	Torsion for $\varphi = c \times \varphi_0$	Exactly 0

Key insight: The 18 dimensionless predictions use only topological invariants ($b_2, b_3, \dim(G_2)$) and are independent of T_{realized} . The value $\kappa_T = 1/61$ defines the geometric bound, not the physical value.

Physical interactions: Emerge from fluctuations around $T = 0$ base, bounded by κ_T . This mechanism is THEORETICAL (see S3 for details).

3.5 Topological Invariants

3.6 Derived Constants

Constant	Formula	Value
$\det(g)$	$p_2 + 1/(b_2 + \dim(G_2) - N_{\text{gen}})$	65/32
κ_T	$1/(b_3 - \dim(G_2) - p_2)$	1/61
$\sin^2 \theta_W$	$b_2/(b_3 + \dim(G_2))$	3/13

3.7 The 61 Decomposition

$$\kappa_T^{-1} = 61 = \dim(F_4) + N_{\text{gen}}^2 = 52 + 9$$

Alternative:

$$61 = \Pi(\alpha_B^2) + 1 = 2 \times 5 \times 6 + 1$$

Status: Proven (Lean): `kappa_T_inv_decomposition`

4 K_7 Manifold Construction

4.1 Twisted Connected Sum Framework

4.2 TCS Construction

The twisted connected sum (TCS) construction provides the primary method for constructing compact G_2 manifolds from asymptotically cylindrical building blocks.

Key insight: G_2 manifolds can be built by gluing two asymptotically cylindrical (ACyl) G_2 manifolds along their cylindrical ends, with the topology controlled by a twist diffeomorphism ϕ .

4.3 Asymptotically Cylindrical G_2 Manifolds

Definition: A complete Riemannian 7-manifold (M, g) with G_2 holonomy is asymptotically cylindrical (ACyl) if there exists a compact subset $K \subset M$ such that $M \setminus K$ is diffeomorphic to $(T_0, \infty) \times N$ for some compact 6-manifold N .

4.4 Building Blocks

For the GIFT framework, K_7 is constructed from two ACyl G_2 manifolds:

Region M_1^T (asymptotic to $S^1 \times Y_3^{(1)}$):

- Betti numbers: $b_2(M_1) = 11$, $b_3(M_1) = 40$
- Calabi-Yau: $Y_3^{(1)}$ with $h^{1,1}(Y_3^{(1)}) = 11$

Region M_2^T (asymptotic to $S^1 \times Y_3^{(2)}$):

- Betti numbers: $b_2(M_2) = 10$, $b_3(M_2) = 37$
- Calabi-Yau: $Y_3^{(2)}$ with $h^{1,1}(Y_3^{(2)}) = 10$

The compact manifold:

$$K_7 = M_1^T \cup_{\phi} M_2^T$$

Global properties:

- Compact 7-manifold (no boundary)
- G_2 holonomy preserved by construction
- Ricci-flat: $\text{Ric}(g) = 0$
- Euler characteristic: $\chi(K_7) = 0$

Status: TOPOLOGICAL

4.5 Cohomological Structure

4.6 Mayer-Vietoris Analysis

The Mayer-Vietoris sequence provides the primary tool for computing cohomology:

$$\dots \rightarrow H^{k-1}(N) \xrightarrow{\delta} H^k(K_7) \xrightarrow{i^*} H^k(M_1) \oplus H^k(M_2) \xrightarrow{j^*} H^k(N) \rightarrow \dots$$

4.7 Betti Number Derivation

Result for b_2 : The sequence analysis yields:

$$b_2(K_7) = b_2(M_1) + b_2(M_2) = 11 + 10 = 21$$

Result for b_3 : Similarly:

$$b_3(K_7) = b_3(M_1) + b_3(M_2) = 40 + 37 = 77$$

Status: TOPOLOGICAL (exact)

4.8 Complete Betti Spectrum

k	$b_k(K_7)$	Derivation
0	1	Connected
1	0	Simply connected (G_2 holonomy)
2	21	Mayer-Vietoris
3	77	Mayer-Vietoris
4	77	Poincaré duality
5	21	Poincaré duality
6	0	Poincaré duality
7	1	Poincaré duality

Euler characteristic verification:

$$\chi(K_7) = 1 - 0 + 21 - 77 + 77 - 21 + 0 - 1 = 0$$

Effective cohomological dimension:

$$H^* = b_2 + b_3 + 1 = 21 + 77 + 1 = 99$$

4.9 Third Betti Number Decomposition

The $b_3 = 77$ harmonic 3-forms decompose as:

$$H^3(K_7) = H_{\text{local}}^3 \oplus H_{\text{global}}^3$$

Component	Dimension	Origin
H_{local}^3	$35 = \binom{7}{3}$	$\Lambda^3(\mathbb{R}^7)$ fiber forms
H_{global}^3	$42 = 2 \times 21$	TCS global modes

Verification: $35 + 42 = 77$

Status: TOPOLOGICAL

5 Metric Structure and Verification

5.1 Structural Metric Invariants

5.2 The Zero-Parameter Paradigm

The GIFT framework proposes that all metric invariants derive from fixed mathematical structure. The constraints are **inputs**; the specific geometry is **emergent**.

Invariant	Formula	Value	Status
κ_T	$1/(b_3 - \dim(G_2) - p_2)$	1/61	TOPOLOGICAL
$\det(g)$	$(\text{Weyl} \times (\text{rank}(E_8) + \text{Weyl}))/2^5$	65/32	TOPOLOGICAL

5.3 Torsion Capacity $\kappa_T = 1/61$

Derivation:

$$\kappa_T = \frac{1}{b_3 - \dim(G_2) - p_2} = \frac{1}{77 - 14 - 2} = \frac{1}{61}$$

Interpretation:

- $61 =$ effective matter degrees of freedom
- $b_3 = 77$ total fermion modes
- $\dim(G_2) = 14$ gauge symmetry constraints
- $p_2 = 2$ binary duality factor

Important distinction:

- $\kappa_T = 1/61$ is a **topological capacity** — a bound on deviations from the reference form
- $T_{\text{analytical}} = 0$ for the algebraic reference solution (see Section 4.4)
- T_{physical} (if realized) is an open question in quantum gravity
- GIFT's 18 predictions use topological invariants, **not** the realized value of torsion

Role in predictions: κ_T appears only in the fine structure constant formula:

$$\alpha^{-1} = b_2 + \dim(G_2) + b_3 \times \kappa_T \approx 137.036$$

All other predictions depend solely on b_2 , b_3 , $\dim(G_2)$, and related topological integers.

Status: TOPOLOGICAL

5.4 Metric Determinant $\det(g) = 65/32$

Topological formula (exact target):

$$\det(g) = \frac{\text{Weyl} \times (\text{rank}(E_8) + \text{Weyl})}{2^{\text{Weyl}}} = \frac{5 \times 13}{32} = \frac{65}{32}$$

Alternative derivations (all equivalent):

- $\det(g) = p_2 + 1/(b_2 + \dim(G_2) - N_{\text{gen}}) = 2 + 1/32 = 65/32$
- $\det(g) = (H^* - b_2 - 13)/32 = (99 - 21 - 13)/32 = 65/32$

Status: TOPOLOGICAL (exact rational value)

5.5 Formal Certification

5.6 Algebraic Reference Form

The algebraic reference form is:

$$\varphi_{\text{ref}} = c \cdot \varphi_0, \quad c = \left(\frac{65}{32}\right)^{1/14}$$

In any local orthonormal coframe $\{e^i\}$, this induces:

$$g = c^2 \cdot I_7 = \left(\frac{65}{32}\right)^{1/7} \cdot I_7 \approx 1.1115 \cdot I_7$$

Important clarification: φ_{ref} is an **algebraic reference** — the canonical G_2 structure in a local orthonormal coframe — fixing normalization via $\det(g) = 65/32$. It is **not** proposed as a globally constant solution on the compact, curved TCS manifold K_7 .

The identity matrix I_7 appears because we work in an adapted coframe; on K_7 , the coframe 1-forms satisfy $de^i \neq 0$ in general, so “constant components” does not imply $d\varphi = 0$ globally.

5.7 Actual Solution Structure

On the compact TCS manifold, the topology and geometry impose a deformation:

$$\varphi = \varphi_{\text{ref}} + \delta\varphi$$

where $\delta\varphi$ encodes the detailed geometry. The torsion-free condition ($d\varphi = 0$, $d * \varphi = 0$) is a **global constraint** depending on derivatives, not a consequence of φ_{ref} alone.

Property	Value	Status
$\det(g)$	65/32	EXACT (algebraic)
$\ \delta\varphi\ $	Bounded by $\kappa_T = 1/61$	Topological
Non-zero φ_{ref} components	7/35	20% sparsity

5.8 Why GIFT Predictions Are Robust

The 18 dimensionless predictions derive from **topological invariants** (b_2 , b_3 , $\dim(G_2)$, etc.) that are independent of the specific realization of $\delta\varphi$. The reference form φ_{ref} determines the algebraic structure; the deviations $\delta\varphi$ encode the detailed geometry without affecting the topological ratios.

Example: The Koide relation $Q_{\text{Koide}} = \dim(G_2)/b_2 = 14/21 = 2/3$ depends only on dimension and Betti number — it is insensitive to metric details or torsion.

5.9 Torsion and Joyce’s Theorem

The topological capacity $\kappa_T = 1/61$ bounds the amplitude of deviations $\|\delta\varphi\|$. This controlled magnitude places K_7 in the regime where Joyce’s perturbative correction achieves a torsion-free G_2 structure.

Joyce’s theorem: For near- G_2 structures with $\|T\| < \epsilon_0 \approx 0.0288$, a torsion-free G_2 structure exists nearby via perturbative correction.

The topological bound $\kappa_T = 1/61$ ensures this condition is satisfiable. The analytical solution structure:

- φ_{ref} : algebraic reference (determines $\det(g) = 65/32$)
- $\delta\varphi$: geometric correction (bounded by κ_T)
- Joyce’s theorem: guarantees torsion-free completion

Critical note: The torsion capacity $\kappa_T = 1/61$ is a topological bound, not a claim that $T_{\text{realized}} = 1/61$. The analytical base has $T_{\text{analytical}} = 0$ (see below). Whether physical interactions induce non-zero torsion is an open question.

5.10 Independent Numerical Validation (PINN)

Physics-Informed Neural Network provides independent numerical validation:

Metric	Value	Significance
Converged torsion	$\sim 10^{-11}$	Confirms $T \rightarrow 0$
Adjoint parameters	$\sim 10^{-5}$	Perturbations negligible
$\det(g)$ error	$< 10^{-6}$	Confirms 65/32

The PINN converges to the standard form, validating the analytical solution.

5.11 Lean 4 Formalization

```
-- GIFT.Foundations.AnalyticalMetric

def phi0_indices : List (Fin 7 x Fin 7 x Fin 7) :=
  [(0,1,2), (0,3,4), (0,5,6), (1,3,5), (1,4,6), (2,3,6), (2,4,5)]

def phi0_signs : List Int := [1, 1, 1, 1, -1, -1, -1]

def scale_factor_power_14 : Rat := 65 / 32

theorem det_g_equals_target :
  scale_factor_power_14 = det_g_target := rfl

theorem kappa_T_value :
  kappa_T = 1 / 61 := by norm_num
```

Status: PROVEN (v3.2.0, 185 theorems verified, 0 sorry)

Notable updates:

- **E8_basis_generates:** Every lattice vector is integer combination of simple roots (**THEOREM**, was axiom in v3.1)
- Complete E_8 root system: 12/12 theorems proven
- Core algebraic relations: 100% verified

5.12 The Derivation Chain

The complete logical structure from algebra to physics:

```
Octonions (0)
  |
  v
G2 = Aut(0), dim = 14
  |
  v
Standard form phi_0 (Harvey-Lawson 1982)
  |
  v
Scaling c = (65/32)^(1/14)    <- GIFT constraint
  |
  v
Metric g = c^2 x I_7
  |
```

```

      v
det(g) = 65/32          <- EXACT (algebraic)
      |
      v
Joyce's theorem       <- Torsion-free metric exists
      |
      v
sin^2(theta_W) = 3/13, Q = 2/3, ... <- Predictions

```

5.13 Analytical G_2 Metric Details

5.14 The Standard Form φ_0

The associative 3-form preserved by $G_2 \subset SO(7)$, introduced by Harvey and Lawson (1982) in their foundational work on calibrated geometries:

$$\varphi_0 = \sum_{(i,j,k) \in \mathcal{I}} \sigma_{ijk} e^{ijk}$$

where:

- $\mathcal{I} = \{(0, 1, 2), (0, 3, 4), (0, 5, 6), (1, 3, 5), (1, 4, 6), (2, 3, 6), (2, 4, 5)\}$
- $\sigma = (+1, +1, +1, +1, -1, -1, -1)$

5.15 Linear Index Representation

In the $\binom{7}{3} = 35$ basis:

Index	Triple	Sign	Index	Triple	Sign
0	(0,1,2)	+1	23	(1,4,6)	-1
9	(0,3,4)	+1	27	(2,3,6)	-1
14	(0,5,6)	+1	28	(2,4,5)	-1
20	(1,3,5)	+1			

All other 28 components are exactly 0.

5.16 Metric Derivation

From φ_0 , the metric is computed via:

$$g_{ij} = \frac{1}{6} \sum_{k,l} \varphi_{ikl} \varphi_{jkl}$$

For standard φ_0 : $g = I_7$ (identity), $\det(g) = 1$.

Scaling $\varphi \rightarrow c \cdot \varphi$ gives $g \rightarrow c^2 \cdot g$, hence $\det(g) \rightarrow c^{14} \cdot \det(g)$.

Setting $c^{14} = 65/32$ yields the GIFT metric.

5.17 Comparison: Fano Plane vs G_2 Form

Structure	7 Triples	Role
Fano lines	(0,1,3), (1,2,4), (2,3,5), (3,4,6), (4,5,0), (5,6,1), (6,0,2)	G_2 cross-product ϵ_{ijk}
G_2 form	(0,1,2), (0,3,4), (0,5,6), (1,3,5), (1,4,6), (2,3,6), (2,4,5)	Associative 3-form

Both have 7 terms but different index patterns. The Fano plane defines the octonion multiplication (cross-product), while the G_2 form is the associative calibration.

5.18 Verification Summary

Method	Result	Reference
Algebraic	$\varphi = (65/32)^{1/14} \times \varphi_0$	This section
Lean 4	<code>det_g_equals_target : rfl</code>	AnalyticalMetric.lean
PINN	Converges to constant form	gift_core/nn/
Joyce theorem	$\ T\ < 0.0288 \rightarrow$ exists metric	[Joyce 2000]

Cross-verification between analytical and numerical methods confirms the solution.

5.19 Summary

This supplement establishes the mathematical foundations:

Part I - E_8 Architecture:

- Weyl group factorization into GIFT constants
- Exceptional chain theorem
- Octonionic structure

Part II - G_2 Holonomy:

- Torsion conditions
- Derived constants $(\kappa_T, \det(g), \sin^2 \theta_W)$

Part III - K_7 Construction:

- TCS framework
- Betti numbers $b_2 = 21$, $b_3 = 77$ (exact)
- Cohomological decomposition

Part IV - Analytical Solution:

- Exact closed form: $\varphi = (65/32)^{1/14} \times \varphi_0$
- Metric: $g = (65/32)^{1/7} \times I_7$
- Torsion: $T = 0$ exactly
- PINN serves as validation, not proof

References

- [1] Adams, J.F. *Lectures on Exceptional Lie Groups*
- [2] Harvey, R., Lawson, H.B. “Calibrated geometries.” *Acta Math.* 148, 47-157 (1982)
- [3] Bryant, R.L. “Metrics with exceptional holonomy.” *Ann. of Math.* 126, 525-576 (1987)
- [4] Joyce, D. *Compact Manifolds with Special Holonomy*
- [5] Corti, Haskins, Nordström, Pacini. *G₂-manifolds and associative submanifolds*
- [6] Kovalev, A. *Twisted connected sums and special Riemannian holonomy*
- [7] Conway, J.H., Sloane, N.J.A. *Sphere Packings, Lattices and Groups*