

Development of a myocardial perfusion phantom

Gijs de Vries, s1854526

Revision 0.100

Detailed Design

Development of a myocardial perfusion phantom

G.J. de Vries, s1854526

Thursday 14th February, 2019

5 **Preface**

6 [todo] this

7 G.J. (Gijs) de Vries

8 Enschede, 13th of February 2019

9 **Contents**

10	1 Introduction	1
11	2 Functional system overview	2
12	3 Technical system overview	3
13	A Appendix: Mind map	4

1 Introduction

Myocardial Perfusion Imaging (MPI), or, simply put, the imaging of the blood flow in the heart muscle, plays an important role in diagnosing heart failure or detecting Coronary Artery Disease (CAD). Imaging systems like Computed Tomography (CT), Magnetic Resonance Imaging (MRI), Single-Photon Emission Computed Tomography (SPECT), or Positron Emission Tomography (PET) can visualise a (radioactive) contrast bolus in the supplying arteries and in underlying myocardial tissue, whose flow can give an indication of narrowed or blocked blood vessels.

Many variations in the visualisation process of myocardial perfusion, including variations in hard- and software, can (significantly) influence the outcome and in turn have consequences for patient treatment. These variations need to be validated against a well-known baseline.

A myocardial perfusion phantom will be developed that is able to simulate the blood flow in the heart muscle, i.e. the myocardium, and is able to mimic cardiac defects like (significant) stenosis.

Document overview

[todo] This section

Abbreviations

CAD Coronary Artery Disease

CT Computed Tomography

MPI Myocardial Perfusion Imaging

MRI Magnetic Resonance Imaging

PET Positron Emission Tomography

SPECT Single-Photon Emission Computed
Tomography

38

2 Concept design

3 Detailed design

40

A Appendix: Mind map




Figure A.1: Mind map for concept designs