

Main results - MASA03

Probability spaces

Definition. A *probability space* is a triple $(\Omega, \mathcal{F}, P)$ where:

- Ω : *sample space* (set of outcomes).
- $\mathcal{F} \subseteq 2^\Omega$: σ -algebra of events: $\Omega \in \mathcal{F}$; if $A \in \mathcal{F}$ then $A^c \in \mathcal{F}$; if $A_1, A_2, \dots \in \mathcal{F}$ then $\bigcup_{n \geq 1} A_n \in \mathcal{F}$.
- $P : \mathcal{F} \rightarrow [0, 1]$: *probability measure* satisfying the axioms:
 - (i) Non-negativity: $P(A) \geq 0$ for all $A \in \mathcal{F}$.
 - (ii) Normalization: $P(\Omega) = 1$.
 - (iii) σ -additivity: if $A_i \in \mathcal{F}$ are pairwise disjoint, $P\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} P(A_i)$.

Immediate consequences (useful): $P(\emptyset) = 0$, $P(A^c) = 1 - P(A)$, $A \subseteq B \Rightarrow P(A) \leq P(B)$, if $A \cap B = \emptyset$ then $P(A \cup B) = P(A) + P(B)$.

0.1 Random Variable

Definition. A function $X : \Omega \rightarrow \mathbb{R}$ is a *random variable* on the probability space $(\Omega, \mathcal{F}, P)$ if it satisfies the condition of $\mathcal{F}$ -measurability:

$$\text{For every Borel set } B \in \mathcal{B}(\mathbb{R}), \quad X^{-1}(B) = \{\omega \in \Omega : X(\omega) \in B\} \in \mathcal{F}$$

Required Conditions

The crucial condition is that the pre-image of any set B for which a probability is defined in $\mathbb{R}$ must be an event in the σ -algebra $\mathcal{F}$. The probability measure P on $\mathcal{F}$ itself is not involved in defining X as a random variable, only the structure of $\mathcal{F}$.

Law of Total Probability (LTP)

If $A_1, \dots, A_k$ form a partition of Ω (pairwise disjoint, $\bigcup_i A_i = \Omega$), then for any event B ,

$$P(B) = \sum_{i=1}^k P(B \mid A_i) P(A_i).$$

(Countable version holds similarly.)

Bayes' Rule

For events A_j and B with $P(B) > 0$,

$$P(A_j | B) = \frac{P(B | A_j) P(A_j)}{\sum_i P(B | A_i) P(A_i)}.$$

The denominator is $P(B)$ from the LTP over the partition $\{A_i\}$.

1 Transforms of random variables

1.1 Sums (convolution)

Continuous, independent: If $Z = X + Y$ and $X \perp Y$ with densities f_X, f_Y ,

$$f_Z(z) = (f_X * f_Y)(z) = \int_{-\infty}^{\infty} f_X(x) f_Y(z - x) dx.$$

Support rule: integrate x over the intersection of supports $\{x : f_X(x) > 0\}$ and $\{x : f_Y(z - x) > 0\}$; this often yields piecewise limits.

Discrete, independent: If p_X, p_Y are pmfs and $Z = X + Y$,

$$p_Z(z) = \sum_{k \in \mathbb{Z}} p_X(k) p_Y(z - k).$$

General (possibly dependent): If the joint density $f_{X,Y}$ is known,

$$f_Z(z) = \int_{-\infty}^{\infty} f_{X,Y}(x, z - x) dx.$$

1.2 Monotone scalar transforms

If $Z = g(X)$ and g is strictly monotone with inverse g^{-1} ,

$$f_Z(z) = f_X(g^{-1}(z)) \left| \frac{d}{dz} g^{-1}(z) \right|.$$

(CDF method: $F_Z(z) = P(g(X) \leq z)$ also works when g is not monotone.)

1.3 Two-to-two transforms (Jacobian)

If $(U, V) = T(X, Y)$ is a bijection with inverse $(X, Y) = T^{-1}(U, V)$,

$$f_{U,V}(u, v) = f_{X,Y}(T^{-1}(u, v)) \left| \det \frac{\partial(x, y)}{\partial(u, v)} \right|.$$

Marginals follow by integrating out the other variable.

1.4 Products and quotients

Product $Z = XY$ (general case):

$$f_Z(z) = \int_{-\infty}^{\infty} f_{X,Y}\left(x, \frac{z}{x}\right) \frac{1}{|x|} dx \quad (\text{integrate over } x \text{ where } (x, z/x) \text{ is in support}).$$

If X, Y are independent: replace $f_{X,Y}$ with $f_X f_Y$ inside the integral.

Quotient $Z = X/Y$:

$$f_Z(z) = \int_{-\infty}^{\infty} f_{X,Y}(zy, y) |y| dy.$$

1.5 Min and Max (use CDFs)

Let $M = \max(X, Y)$ and $m = \min(X, Y)$. If X, Y are independent with cdfs F_X, F_Y and densities f_X, f_Y ,

$$F_M(t) = P(M \leq t) = F_X(t)F_Y(t), \quad f_M(t) = f_X(t)F_Y(t) + F_X(t)f_Y(t).$$

$$F_m(t) = 1 - P(X > t \text{ or } Y > t) = 1 - (1 - F_X(t))(1 - F_Y(t)), \quad f_m(t) = f_X(t)(1 - F_Y(t)) + f_Y(t)(1 - F_X(t)).$$

1.6 Quick recipe

1. **Identify the map:** $Z = X + Y$, $Z = g(X)$, or $(U, V) = T(X, Y)$.
2. **Pick the tool:** convolution (sum), monotone transform (one-to-one in 1D), Jacobian (two-to-two), or the CDF method.
3. **Set the limits by geometry:** intersect supports after the change of variables.
4. **Normalize/check:** verify $\int f_Z = 1$ (or $\sum p_Z = 1$).

2 Joint distributions and tools

2.1 Joint density and normalization

A pair (X, Y) has joint density $f_{X,Y}(x, y) \geq 0$ with

$$\iint_{\mathbb{R}^2} f_{X,Y}(x, y) dx dy = 1.$$

2.2 Marginals

$$f_X(x) = \int_{-\infty}^{\infty} f_{X,Y}(x, y) dy, \quad f_Y(y) = \int_{-\infty}^{\infty} f_{X,Y}(x, y) dx.$$

2.3 Independence

X and Y are independent iff

$$f_{X,Y}(x, y) = f_X(x) f_Y(y) \quad (\text{almost everywhere on the support}).$$

2.4 Conditional density

For $f_X(x) > 0$,

$$f_{Y|X}(y | x) = \frac{f_{X,Y}(x, y)}{f_X(x)}.$$

2.5 Cumulative distribution function (cdf)

For any random variable Z ,

$$F_Z(z) = \Pr(Z \leq z), \quad \Pr(a < Z \leq b) = F_Z(b) - F_Z(a).$$

If Z has a density f_Z , then

$$f_Z(z) = \frac{d}{dz} F_Z(z).$$

2.6 Product variable via the cdf method

For $Z = XY$ with (X, Y) supported on $[a, b] \times [c, d] \subseteq [0, \infty)^2$,

$$F_Z(z) = \Pr(XY \leq z) = \iint_{\{(x,y) \in [a,b] \times [c,d] : xy \leq z\}} f_{X,Y}(x, y) dx dy,$$

and $f_Z(z) = \frac{d}{dz} F_Z(z)$ on the interior of the support of Z .

3 Expectation and related results

3.1 Definition

For a random variable X with pdf f_X or pmf p_X ,

$$E[X] = \begin{cases} \sum x p_X(x), & (\text{discrete}), \\ \int_{-\infty}^{\infty} x f_X(x) dx, & (\text{continuous}). \end{cases}$$

More generally (Law of the Unconscious Statistician, LOTUS),

$$E[g(X)] = \begin{cases} \sum g(x) p_X(x), \\ \int g(x) f_X(x) dx. \end{cases}$$

3.2 Conditional expectation and total laws

For densities,

$$E[X | Y = y] = \int x f_{X|Y}(x | y) dx, \quad f_{X|Y}(x | y) = \frac{f_{X,Y}(x, y)}{f_Y(y)}.$$

Law of total expectation (tower property):

$$E[X] = E(E[X | Y]).$$

Law of total probability (continuous form): For any event A ,

$$P(A) = \int P(A | Y = y) f_Y(y) dy \quad \text{or, for discrete } Y, \quad P(A) = \sum_y P(A | Y = y) P(Y = y).$$

This expresses the overall probability $P(A)$ as the average (expectation) of the conditional probabilities $P(A | Y = y)$ weighted by how likely each y is. Equivalently,

$$P(A) = E[P(A | Y)].$$

Hence, probabilities are special cases of expectations:

$$P(X < Y) = E[P(X < Y | X)].$$

3.3 Variance and Covariance

$$\begin{aligned} \text{Var}(X) &= E[X^2] - (E[X])^2 \\ \text{Cov}(X, Y) &= E[XY] - E[X]E[Y] \end{aligned}$$

3.3.1 Law of Total Variance (Conditional Version)

$$\text{Var}(X) = E[\text{Var}(X | Y)] + \text{Var}(E[X | Y])$$

3.3.2 Relationship Between Cov and Var

$$\text{Var}(X) = \text{Cov}(X, X)$$

3.3.3 Correlation Coefficient (ρ)

The correlation coefficient is the standardized covariance.

$$\rho_{X,Y} = \text{Corr}(X, Y) = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)}\sqrt{\text{Var}(Y)}} = \frac{\text{Cov}(X, Y)}{\sigma_X \sigma_Y}$$

Identity: The identity for correlation is its range:

$$-1 \leq \rho_{X,Y} \leq 1$$

Covariance in terms of Correlation:

$$\text{Cov}(X, Y) = \rho_{X,Y} \sigma_X \sigma_Y$$

4 Common Probability Distributions

4.1 Discrete distributions

Binomial distribution

$$\begin{aligned} X &\sim \text{Bin}(n, p) \\ P(X = k) &= \binom{n}{k} p^k (1-p)^{n-k}, \quad k = 0, 1, \dots, n. \\ E[X] &= np, \quad \text{Var}(X) = np(1-p). \end{aligned}$$

Bernoulli distribution

$$\begin{aligned} X &\sim \text{Bern}(p) \\ P(X = 1) &= p, \quad P(X = 0) = 1 - p. \\ E[X] &= p, \quad \text{Var}(X) = p(1-p). \end{aligned}$$

Geometric distribution

Two common parameterizations with success probability $p \in (0, 1)$:

(Trials until first success) $X \sim \text{Geom}_1(p)$ with support $\{1, 2, \dots\}$.

$$\begin{aligned} P(X = k) &= (1-p)^{k-1} p, \quad k = 1, 2, \dots \\ E[X] &= \frac{1}{p}, \quad \text{Var}(X) = \frac{1-p}{p^2}. \end{aligned}$$

Memoryless: $P(X > m + n \mid X > m) = (1-p)^n$.

(Failures before first success) $Y \sim \text{Geom}_0(p)$ with support $\{0, 1, 2, \dots\}$.

$$\begin{aligned} P(Y = k) &= (1-p)^k p, \quad k = 0, 1, 2, \dots \\ E[Y] &= \frac{1-p}{p}, \quad \text{Var}(Y) = \frac{1-p}{p^2}. \end{aligned}$$

Relation: $X = Y + 1$.

CDFs:

$$F_X(k) = 1 - (1-p)^k, \quad k \in \{1, 2, \dots\}; \quad F_Y(k) = 1 - (1-p)^{k+1}, \quad k \in \{0, 1, 2, \dots\}.$$

Notes: $\text{Geom}_1(p)$ is the discrete analog of $\text{Exp}(\lambda)$ and is a special case of the negative binomial (number of trials until the r -th success with $r = 1$).

Poisson distribution

$$X \sim \text{Poisson}(\lambda)$$

$$P(X = k) = \frac{e^{-\lambda} \lambda^k}{k!}, \quad k = 0, 1, 2, \dots$$

$$E[X] = \lambda, \quad \text{Var}(X) = \lambda.$$

4.2 Continuous distributions

Uniform distribution

$$X \sim \text{Unif}(a, b)$$

$$f_X(x) = \begin{cases} \frac{1}{b-a}, & a \leq x \leq b, \\ 0, & \text{otherwise.} \end{cases}$$

$$E[X] = \frac{a+b}{2}, \quad \text{Var}(X) = \frac{(b-a)^2}{12}.$$

Exponential distribution

$$X \sim \text{Exp}(\lambda)$$

$$f_X(x) = \begin{cases} \lambda e^{-\lambda x}, & x \geq 0, \\ 0, & \text{otherwise.} \end{cases}$$

$$E[X] = \frac{1}{\lambda}, \quad \text{Var}(X) = \frac{1}{\lambda^2}.$$

Memoryless property: $P(X > s+t \mid X > s) = P(X > t)$.

Normal (Gaussian) distribution

$$X \sim N(\mu, \sigma^2)$$

$$f_X(x) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right), \quad x \in \mathbb{R}.$$

$$E[X] = \mu, \quad \text{Var}(X) = \sigma^2.$$

Standard Normal distribution

$$Z \sim N(0, 1)$$

$$f_Z(z) = \frac{1}{\sqrt{2\pi}} e^{-z^2/2}, \quad F_Z(z) = \Phi(z) = \int_{-\infty}^z f_Z(t) dt.$$

Gamma distribution

$$X \sim \text{Gamma}(\alpha, \lambda)$$

$$f_X(x) = \begin{cases} \frac{\lambda^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\lambda x}, & x > 0, \\ 0, & \text{otherwise.} \end{cases}$$

$$E[X] = \frac{\alpha}{\lambda}, \quad \text{Var}(X) = \frac{\alpha}{\lambda^2}.$$

Special case: $\text{Exp}(\lambda) = \text{Gamma}(1, \lambda)$.

4.3 Summary Table

Distribution	Support	Mean	Variance
Bernoulli(p)	$\{0, 1\}$	p	$p(1-p)$
Binomial(n, p)	$\{0, \dots, n\}$	np	$np(1-p)$
Poisson(λ)	$\mathbb{N}_0$	λ	λ
Uniform(a, b)	$[a, b]$	$(a+b)/2$	$(b-a)^2/12$
Exponential(λ)	$[0, \infty)$	$1/\lambda$	$1/\lambda^2$
Normal(μ, σ^2)	$\mathbb{R}$	μ	σ^2
Gamma(α, λ)	$[0, \infty)$	α/λ	α/λ^2

5 Limit Theorems: LLN and CLT

5.1 Law of Large Numbers (LLN)

Let $X_1, X_2, \dots$ be i.i.d. random variables with $E[X_i] = \mu$ and $\text{Var}(X_i) = \sigma^2 < \infty$. Define the sample mean

$$\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i.$$

Weak Law of Large Numbers (WLLN)

$$\bar{X}_n \xrightarrow{P} \mu \quad \text{as } n \rightarrow \infty.$$

That is, for every $\varepsilon > 0$,

$$P(|\bar{X}_n - \mu| > \varepsilon) \rightarrow 0.$$

Interpretation: The sample mean converges in probability to the true mean μ .

Strong Law of Large Numbers (SLLN)

$$\bar{X}_n \xrightarrow{\text{a.s.}} \mu \quad \text{as } n \rightarrow \infty.$$

That is,

$$P\left(\lim_{n \rightarrow \infty} \bar{X}_n = \mu\right) = 1.$$

Interpretation: The sample mean converges almost surely (with probability 1) to the true mean μ .

Difference between weak and strong forms: Both state that $\bar{X}_n$ approaches μ as n grows, but:

- **Weak LLN:** convergence in probability (approximate closeness for large n).
- **Strong LLN:** almost sure convergence (the entire sequence converges for almost every outcome).

5.2 Central Limit Theorem (CLT)

Theorem (Lindeberg-Lévy CLT). Let $X_1, X_2, \dots, X_n$ be i.i.d. random variables with

$$E[X_i] = \mu, \quad \text{Var}(X_i) = \sigma^2 < \infty.$$

Then as $n \rightarrow \infty$,

$$\frac{\sqrt{n}(\bar{X}_n - \mu)}{\sigma} \xRightarrow{d} N(0, 1),$$

where $\xRightarrow{d}$ denotes convergence in distribution.

Interpretation: For large n , the sample mean $\bar{X}_n$ is approximately normal:

$$\bar{X}_n \approx N\left(\mu, \frac{\sigma^2}{n}\right).$$

Hence, the CLT justifies using the normal distribution to approximate sums or averages of many independent random variables.

Practical consequence: For large n ,

$$P\left(\frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}} \leq z\right) \approx \Phi(z),$$

where $\Phi(z)$ is the standard normal cdf. This result underpins confidence intervals and hypothesis testing.