

Topology: Exercises 1

Simon Gustafsson

Problem 1

Proof. (\Rightarrow) Assume there exists a bijection $h : A \rightarrow B$. Since A is countable, there exists a set C and a bijection $f : A \rightarrow C$, where $C = \{1, \dots, n\}$ if A is finite and $C = \mathbb{N}$ if A is countably infinite. Define $g := f \circ h^{-1} : B \rightarrow C$. Since compositions of bijections are bijections, g is a bijection. Hence there exists a bijection from both A and B to the same set C , so they have the same cardinality.

(\Leftarrow) Assume A and B have the same cardinality. Then there exists a set $C \subseteq \mathbb{N}$ and bijections $f : A \rightarrow C$ and $g : B \rightarrow C$. Define $h := g^{-1} \circ f : A \rightarrow B$. Since f and g are bijections, h is a bijection. \square

Problem 2

Proof. Let $k \in \mathbb{N}$ and choose distinct primes p_1, \dots, p_k . Define

$$f : \mathbb{N}^k \rightarrow \mathbb{N}, \quad (n_1, \dots, n_k) \mapsto p_1^{n_1} \cdots p_k^{n_k}.$$

Let $x, y \in \mathbb{N}^k$ and suppose $f(x) = f(y)$. Then

$$p_1^{x_1} \cdots p_k^{x_k} = p_1^{y_1} \cdots p_k^{y_k}.$$

By uniqueness of prime factorization, we must have $x_i = y_i$ for each $i \in \{1, \dots, k\}$. Hence $x = y$, and f is injective. Since \mathbb{N} is countable and \mathbb{N}^k injects into \mathbb{N} , it follows by Theorem 7.1 that \mathbb{N}^k is countable. \square