

LUND
UNIVERSITY

The sharp A_p bound for the Hardy–Littlewood maximal operator

SIMON GUSTAFSSON





Weights and weighted L^p spaces

Definition 1 (Weights)

A *weight* is a function $w \in L^1_{\mathrm{loc}}(\mathbb{R}^d)$ with $w \geq 0$ that is not almost everywhere zero.

Definition 2 (Weighted L^p spaces)

Let w be a weight and let $1 \leq p < \infty$. The weighted L^p space is

$$L^p_w(\mathbb{R}^d) := \left\{ f : \mathbb{R}^d \to \mathbb{R} \text{ measurable} : \int_{\mathbb{R}^d} |f(y)|^p \, w(y) \, dy < \infty \right\},$$

with norm $\|f\|_{L^p_w} := \left(\int_{\mathbb{R}^d} |f(y)|^p \, w(y) \, dy\right)^{1/p}$. Moreover, $L^\infty_w(\mathbb{R}^d)$ is the space of measurable functions f with $\|f\|_{L^\infty_w} := \inf\{t \geq 0 : w(\{|f| > t\}) = 0\} < \infty$.

LUND UNIVERSITY

The Hardy–Littlewood maximal operator

Definition 3 (Hardy–Littlewood maximal operator)

$$M_w f(x) := \sup_{Q \ni x} \frac{1}{w(Q)} \int_Q |f(y)|\, w(y)\, dy.$$

M_w^c takes the supremum over centred cubes $Q(x,r)$ only, and $w \equiv 1$ gives M and M^c.

M bounded on $L^p_w(\mathbb{R}^d)$

$$\|Mf\|_{L^p_w} = \left(\int_{\mathbb{R}^d} (Mf)^p \, w \, dy \right)^{1/p} \leq C \left(\int_{\mathbb{R}^d} |f|^p \, w \, dy \right)^{1/p} = C \, \|f\|_{L^p_w}$$

LUND UNIVERSITY

A necessary condition on w

Proposition 1 (Muckenhoupt)

If M is bounded on $L^p_w(\mathbb{R}^d)$ with constant C, then every cube Q satisfies

$$\left(\frac{1}{|Q|}\int_Q w\,dy\right)\left(\frac{1}{|Q|}\int_Q w^{-\frac{1}{p-1}}\,dy\right)^{p-1} \leq C^p.$$

LUND UNIVERSITY

A necessary condition on w

Proposition 1 (Muckenhoupt)

If M is bounded on $L^p_w(\mathbb{R}^d)$ with constant C, then every cube Q satisfies

$$\left(\frac{1}{|Q|}\int_Q w\,dy\right)\left(\frac{1}{|Q|}\int_Q w^{-\frac{1}{p-1}}\,dy\right)^{p-1} \leq C^p.$$

Definition 4 (A_p characteristic)

$$[w]_{A_p} := \sup_Q \left(\frac{1}{|Q|}\int_Q w\,dy\right)\left(\frac{1}{|Q|}\int_Q w^{-\frac{1}{p-1}}\,dy\right)^{p-1},$$

and $w \in A_p$ if $[w]_{A_p} < \infty$.

LUND UNIVERSITY

Dual weights and A_p as a necessary condition

Definition 5 (Dual weight)

The *dual weight* of w is $\sigma := w^{-\frac{1}{p-1}}$.

Corollary 1

If M is bounded on $L^p_w(\mathbb{R}^d)$ with constant C, then $w \in A_p$ with

$$[w]_{A_p} \leq C^p.$$

LUND UNIVERSITY



LUND
UNIVERSITY