

Abstract Algebra - Homework 1

Simon Gustafsson

1 Group Action on the Projective Line

Let $k = \mathbb{F}_7 = \mathbb{Z}/7\mathbb{Z}$. We define $\mathrm{GL}_2(k)$ as the group of all invertible 2×2 matrices with entries in the field k . In other words,

$$\mathrm{GL}_2(k) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in M_2(k) \mid ad - bc \neq 0 \right\}.$$

$\mathrm{GL}_2(k)$ Forms a Group

Let $A, B \in \mathrm{GL}_2(k)$. Then the product AB is defined as:

$$AB = \begin{pmatrix} ae + bg & af + bh \\ ce + dg & cf + dh \end{pmatrix},$$

where all entries involve only addition and multiplication of elements in k . Since k is a field, it is closed under addition and multiplication, so all entries of AB lie in k . Furthermore, $\det(AB) = \det(A)\det(B) \neq 0$, so $AB \in \mathrm{GL}_2(k)$. Thus, $\mathrm{GL}_2(k)$ is closed under matrix multiplication.

Since matrix multiplication is associative, the operation on $\mathrm{GL}_2(k)$ is associative.

The identity element in $\mathrm{GL}_2(k)$ is the identity matrix

$$I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

For any $A \in \mathrm{GL}_2(k)$, we have $AI = IA = A$, and since the entries 1 and 0 are in k , we conclude $I \in \mathrm{GL}_2(k)$.

Since every element of $\mathrm{GL}_2(k)$ is invertible by definition, the inverse of

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{GL}_2(k)$$

is given by:

$$A^{-1} = \frac{1}{\det(A)} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}.$$

Since $a, b, c, d \in k$, and $\det(A) = ad - bc \neq 0$, we have $\frac{1}{\det(A)} \in k$, because k is a field. As k is closed under addition, subtraction, and multiplication, all entries of A^{-1} lie in k . Hence, $A^{-1} \in \text{GL}_2(k)$.

Therefore, $\text{GL}_2(k)$ satisfies the group axioms under matrix multiplication and is a group.

The Center of $\text{GL}_2(k)$

Let $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{GL}_2(k)$. Suppose $A \in Z(\text{GL}_2(k))$. Then for all $B = \begin{pmatrix} e & f \\ g & h \end{pmatrix} \in \text{GL}_2(k)$, we must have

$$AB = BA.$$

Compute both sides:

$$AB = \begin{pmatrix} ae + bg & af + bh \\ ce + dg & cf + dh \end{pmatrix}, \quad BA = \begin{pmatrix} ea + fc & eb + fd \\ ga + hc & gb + hd \end{pmatrix}.$$

For these to be equal for all $e, f, g, h \in k$, compare the corresponding entries:

$$\begin{aligned} ae + bg &= ea + fc & \Rightarrow & b = c = 0 \\ af + bh &= eb + fd & \Rightarrow & a = d, \text{ using } b = 0 \\ ce + dg &= ga + hc & \Rightarrow & c = b = 0 \\ cf + dh &= gb + hd & \Rightarrow & d = a \end{aligned}$$

Hence, A must be of the form:

$$A = \begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix} = \lambda I,$$

where $\lambda \in k^\times = k \setminus \{0\}$ (since A must be invertible). Thus,

$$Z(\text{GL}_2(\mathbb{F}_7)) = \{\lambda I \mid \lambda \in k^\times\}.$$

This center consists of 6 elements and forms an abelian subgroup of $\text{GL}_2(\mathbb{F}_7)$.

Group Action on $\mathbb{P}^1(k)$

We define the projective line over k as $\mathbb{P}^1(k) = k \cup \{\infty\}$. Let $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{GL}_2(k)$. Then the action of $\text{GL}_2(k)$ on $\mathbb{P}^1(k)$ is given by the formula:

$$A \cdot z = \begin{cases} \frac{az + b}{cz + d}, & \text{if } z \in k \text{ and } cz + d \neq 0, \\ \infty, & \text{if } z \in k \text{ and } cz + d = 0, \\ \frac{a}{c}, & \text{if } z = \infty \text{ and } c \neq 0, \\ \infty, & \text{if } z = \infty \text{ and } c = 0. \end{cases}$$

We verify that this defines a group action.

Let $I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$. Then for all $z \in \mathbb{P}^1(k)$,

$$I \cdot z = \frac{1 \cdot z + 0}{0 \cdot z + 1} = \frac{z}{1} = z, \quad \text{and} \quad I \cdot \infty = \frac{1}{0} := \infty.$$

So the identity acts as the identity function.

Let $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ and $B = \begin{pmatrix} e & f \\ g & h \end{pmatrix}$ be in $\text{GL}_2(k)$, and let $z \in \mathbb{P}^1(k)$.

Then:

$$B \cdot z = \frac{ez + f}{gz + h}, \quad \text{and} \quad A \cdot (B \cdot z) = \frac{a \cdot \left(\frac{ez+f}{gz+h} \right) + b}{c \cdot \left(\frac{ez+f}{gz+h} \right) + d}.$$

Simplifying this gives:

$$A \cdot (B \cdot z) = \frac{(aez + af + bgz + bh)}{(cez + cf + dgz + dh)} = \frac{(ae + bg)z + (af + bh)}{(ce + dg)z + (cf + dh)},$$

which is the action of the product matrix AB on z :

$$(AB) \cdot z.$$

Thus, the compatibility condition holds.

Therefore, the formula defines a group action of $\text{GL}_2(k)$ on $\mathbb{P}^1(k)$.

Transitivity and the Stabilizer of 0

Let $z \in \mathbb{P}^1(k)$ be arbitrary. We want to find a matrix $A \in \mathrm{GL}_2(k)$ such that $A \cdot 0 = z$. Let

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{GL}_2(k).$$

Then the action on 0 is given by

$$A \cdot 0 = \frac{b}{d},$$

provided $d \neq 0$. For any $z \in k$, we can take $b = z$ and $d = 1$, with $a \in k^\times$, $c \in k$ arbitrary, such that $\det(A) = ad - bc \neq 0$.

To obtain ∞ , we require $d = 0$, in which case the formula becomes

$$A \cdot 0 = \frac{b}{d} = \infty,$$

provided $b \neq 0$. We can choose $a, c \in k$ arbitrarily so that $\det(A) = -bc \neq 0$, ensuring $A \in \mathrm{GL}_2(k)$.

Hence, for any $z \in \mathbb{P}^1(k)$, there exists a matrix $A \in \mathrm{GL}_2(k)$ such that $A \cdot 0 = z$. Therefore, the action is transitive.

We require $A \cdot 0 = 0$, so $\frac{b}{d} = 0 \Rightarrow b = 0$. Therefore, the stabilizer of 0 is the set of all invertible lower triangular matrices:

$$B = \left\{ \begin{pmatrix} a & 0 \\ c & d \end{pmatrix} \in \mathrm{GL}_2(k) \mid a, d \in k^\times, c \in k \right\}.$$