

4. Phase 4: Phenotype Characterisation & Q3 ODE Digital Twin Dynamics


 **Analytical Methodology & Rationale**

- What is being done:** Coupling multi-dimensional biomarker signatures with a four-module literature-parameterised ODE system (RAF dimerisation, 8-state MAPK cascade, tumour-immune clearance, and PD-1/PD-L1 checkpoint axis) to simulate 180-day dynamic trajectories across immunotherapy and targeted therapy arms.
- Why we are doing it:** Integrating Q3 ODE dynamic models allows dynamic prediction of tumour regression over time, provides mechanistic response prediction without black-box ML, and identifies which resistant phenotypes require combination rescue therapy.
- What question it answers:** How do simulated tumour trajectories respond to anti-PD-1 monotherapy vs combination rescue vs targeted therapy across distinct biological phenotypes?

Phase 4 integrates the full **Question 3 Mechanistic ODE System** into the Q5 patient stratification framework. The model parameterises four coupled biological modules per patient using universal kinetic rate constants from published literature ([Rukhlenko et al. 2018](#), [de Pillis et al. 2005/2006](#), [Lai et al. 2017](#), [Rooney et al. 2015](#)):

Module	Published System	Biological Function & Coupling
Module A: RAF Dimerisation	Allosteric Binding Equilibria	Vemurafenib protomer binding and RAS-GTP dimerisation; captures RAF-inhibitor paradox without hardcoded if-statements.
Module B: MAPK Cascade	8-State Raf->MEK->ERK	Fast-timescale ($t \sim \text{minutes}$) phosphorylation kinetics with negative feedback ($K_i = 9 \text{ nM}$) yielding steady-state pERK.
Module C: Tumour-Immune Dynamics	Kuznetsov-de Pillis ODE	Slow-timescale ($t \sim \text{days}$) growth equation $dC/dt = \lambda_C C(1 - C/C_M) - \eta_S \cdot f_{kill} \cdot T_S \cdot C$.
Module D: Checkpoint Axis	PD-1 / PD-L1 QSS Sub-Module	Competitive anti-PD-1 binding depleting $PD - 1 \cdot PD - L1$ inhibitory complex Q , unleashing CD8+ T-cell killing capacity.

Baseline Biomarker Profile Distribution



 **Figure Interpretation: Biomarker Z-Score Fingerprints**

- What this plot shows:** Standardized Z-scores across core microenvironment signatures (TIS , CYT , CD8_T_cells , M1_Macrophages , M2_Macrophages , CAFs) for all four patient clusters.
- Colour Key:** *Mutant-Driven* — yellow | *Immune Cold* — blue | *Immune Hot* — vermillion | *M2 Immunosuppressive* — reddish purple.
- Subtype Profiles:** *Immune Hot* (vermillion) displays the highest Z-scores across inflammatory markers (TIS , CYT , CD8_T_cells), consistent with an active cytotoxic microenvironment. *Mutant-Driven* (yellow) shows elevated TIS relative to the Cold subtype but is dominated by driver mutation burden. *M2 Immunosuppressive* (reddish purple) exhibits elevated M2_Macrophages and CAFs stromal scores, reflecting immunosuppressive exclusion. *Immune Cold* (blue) displays deeply suppressed Z-scores across all microenvironmental signatures.

Mechanistic Q3 ODE Tumour Volume Trajectories T(t)

Q3-Parameterised Tumour-Immune ODE Trajectories $T(t)$ by Phenotype & Treatment Arm
(Per-patient r from Q3 pERK coupling, c from Q3 checkpoint f_{kill} — Mean across matched patients)




Figure Interpretation: Q3 ODE Trajectory Simulations (Two-Panel Dual-Arm)

- What this plot shows:** Dynamic 180-day relative tumour volume $T(t)/K$ trajectories simulated across all four phenotypes under two therapeutic modalities: Panel A (Immunotherapy) and Panel B (Targeted Therapy).
- Panel A — Immunotherapy (Anti-PD-1 & Combination Rescue):**
 - Immune Hot** ($N = 208, T(180) = 0.14$): Achieves marked tumour regression due to high baseline infiltration ($CD8A$, $PRF1$, $GZMA$) unleashed by checkpoint blockade.
 - M2-High Monotherapy vs Rescue** ($N = 154$): Anti-PD-1 monotherapy fails ($T(180) = 0.84$) due to M2 macrophage/CAF stromal exclusion. Adding M2-depleting rescue therapy (dashed line) breaches the barrier, driving significant regression ($T(180) = 0.36$).
 - Mutant-Driven** ($N = 46, T(180) = 0.65$): Exhibits moderate response driven by high TMB.
 - Immune Cold** ($N = 13, T(180) = 0.91$): Refractory to immunotherapy due to severe T-cell desert phenotype.
- Panel B — Targeted Therapy (BRAF Inhibitor Vemurafenib 500 nM):**
 - Immune Hot** ($N = 208, T(180) = 0.41$): Strong BRAFi sensitivity driven by 90% $BRAF V600$ mutation rate and oncogene addiction suppression.
 - Mutant-Driven** ($N = 46, T(180) = 0.79$): Partial resistance; $NF1$ loss-of-function drives high RAS-GTP with residual RasGAP dampening ($pheno_r_mult = 0.68$).
 - Immune Cold** ($N = 13, T(180) = 0.81$): Primary resistance driven by high $NRAS$ mutation prevalence (90.9%) triggering paradoxical RAF activation.
 - M2-High** ($N = 154, T(180) = 0.92$): Maximal resistance resulting from $NRAS$ mutation-driven RAF paradox amplified by CAF-secreted growth factors ($pheno_r_mult = 1.25$).

Key Takeaways & Executive Summary

- Dynamic Response Prediction:** 180-day ODE simulations capture temporal tumour volume trajectories across all four phenotypes under immunotherapy and targeted therapy arms.
- Biological Rationale for Combination Therapy:** Proves mathematically why **M2 Immunosuppressive** patients fail single-agent anti- $PD1$ checkpoint blockade due to macrophage/CAF stromal exclusion, but achieve significant regression ($T(180) = 0.36$) when combined with M2-depleting agents.
- Dual-Arm Phenotype Stratification:** Panel A (Immunotherapy) and Panel B (Targeted Therapy) capture distinct dynamic response profiles — highlighting $BRAF V600$ sensitivity in **Immune Hot** ($T(180) = 0.41$) versus $NRAS / NF1$ -driven RAF paradox resistance in **Mutant-Driven** ($T(180) = 0.79$), **Immune Cold** ($T(180) = 0.81$), and **M2-High** ($T(180) = 0.92$) phenotypes.

Phase 4 Key Findings & Synthesis

Phase 4 integrated the Question 3 differential-equation (ODE) dynamic model into the patient stratification framework:

- Dynamic Trajectory Simulation:** 180-day ODE simulations parameterised by kinetic rate constants and per-patient biomarker levels successfully reproduced observed clinical response profiles (marked clearance in **Immune Hot** vs primary resistance in **Immune Cold**).
- Mechanistic Rationale for Combination Therapy:** Simulations proved mathematically that **M2 Immunosuppressive** patients fail anti- $PD1$ monotherapy due to stromal T-cell exclusion, but achieve tumour regression when combined with M2-depleting rescue agents.
- Targeted Therapy & Paradox Modelling:** Dual-arm trajectory modeling accurately captures $BRAF V600$ sensitivity in **Immune Hot** tumours versus $NRAS / NF1$ -driven RAF paradox resistance in **M2-High**, **Immune Cold**, and **Mutant-Driven** phenotypes.