# Review of the Greenhouse Effect

EES 3310/5310
Global Climate Change
Jonathan Gilligan

Class #7: Wednesday, Sept. 5 2018



#### Notes on Labs

- Check frequently whether you can knit your document.
  - If you have trouble, ask for help
    - Office hours or Email both me and Kelsea
    - Remember to put "EES 3310" or "EES 5310" in the subject line
- Remember that either PDF or Word documents are acceptable for turning in your work, so if you have trouble knitting to PDF, knit to a Word document.
  - It should only take a one-time set-up to be able to knit smoothly to PDF. Kelsea and I are happy to help you set up your computer to knit to PDF.

# Vertical Structure of the Atmosphere

# Terminology


#### Environmental Lapse

- Measured temperature of actual atmosphere at a single time
- Compares one bit of air at one height with another bit at another height.
- Environmental lapse is different from one time or place to another.


#### Adiabatic Lapse

Change in a single parcel of air as it moves up or down

# Moist Convection



#### Moist Convection




- Latent heat warms air
- Reduces adiabatic cooling
- Moist adiabatic lapse < Dry adiabatic lapse</li>
- Smaller lapse = less stable
- Humid air is less stable than dry air

#### Perspective

- Stable:
  - Environmental lapse ≤ adiabatic lapse
- Unstable:
  - Environmental lapse > adiabatic lapse
- Adiabatic lapse:
  - Dry: 10 K/km
  - Moist: 4-8 K/km (depends on humidity)
- Pure radiative equilibrium:
  - Would produce lapse of 16 K/km: unstable
- Radiative-Convective equilibrium:
  - Convection modifies environmental lapse
  - Normal environmental lapse is roughly 6 K/km
     (typical moist adiabatic lapse rate)


# Greenhouse effect

#### Greenhouse effect



#### Greenhouse effect

- 1.  $T_{\text{skin}} = 254 \text{ K}$
- 2.  $T_{\text{ground}} = T_{\text{skin}} + \text{lapse rate} \times h_{\text{skin}}$
- 3. Increase greenhouse gases
- 4. Skin height rises by  $\Delta h_{skin}$
- 5.  $\tau_{\text{ground}}$  rises by lapse rate  $\times \Delta h_{\text{skin}}$





# Vertical Structure and Saturation


#### Set up MODTRAN:


Go to MODTRAN (http://climatemodels.uchicago.edu/modtran/)

- Set altitude to 70 km and location to "Tropical".
- Set CO<sub>2</sub> to 1 ppm, all other gases to zero.
- Now increase by factors of 10 (10, 100, 1000, ...)











# 10,000 ppm CO<sub>2</sub>



# Question

Why do we see the spike in the middle of the CO<sub>2</sub> absorption feature?

#### Question



- Water vapor absorption is completely saturated.
  - Why does water vapor emit at warmer temperatures than CO<sub>2</sub>?

# Review Perspective

#### Review Perspective

#### 1. Start with bare-rock temperature

This becomes skin temperature

#### 2. Add simple atmosphere:

- Completely absorbs longwave radiation
- Top of atmosphere: skin temperature (same as bare-rock)
- More layers ⇒ bigger greenhouse effect

#### 3. Realistic longwave absorption:


Atmosphere is not a black body

#### 4. Radiative-Convective equilibrium:


- Pure radiative equilibrium would have huge lapse
- Big lapse is unstable ⇒ convection
  - Convection mixes hot & cold air ⇒
     modifies environmental lapse
  - Reduces greenhouse effect









# Around the planet




# Cloud blocks sunlight



# Exercise 3-3



# Temperature drops



# Volcanoes and Temperature

