

C^* -Algebras, and
the Gelfand-Naimark Theorem

Luke Armitage

April 13, 2017

Todo list

write this!	4
state and prove	4
remark on work in real C^* -algebras?	4
cauchy-schwarz, C^* subalgebra generated by a set	4
maybe move this to as early as we can?	5
example of $C(X)$?	5
tidy up proof	5
normal?	6
state in 'spectrum' - 4.1.5	7
is unital necessary?	7
show that $\ \rho\ = \rho(\mathbb{1})$. explain extreme point.	7
justify. 4.3.13.	7
show	8
show a state.	8
define weak* topology on $P(S)$	9
example of $\mathcal{B}(\mathcal{H})$ at end of section to retro-motivate notation. discuss nomenclature (state etc) coming from QM	11
to include all representation theory, including GNS, CGN and GN . . .	11
remark about nonunital commutative.	11
why?	12
sentence	12
start sentence properly	13
cont of what?	14
cont of what?	14
example of this construction on $C(X)$? may just be a short explana- tion of how $B(H)$ and $C(X)$ link together. can then talk about noncommutative topology!	15
further topics: K-theory, group C^* algebras, amenable algebras, von neumann algebras,	15
references!!!!	15

Contents

0.1	Preliminaries	2
0.1.1	History of C^* -Algebras	2
0.1.2	Background Mathematics and Resources.	3
0.1.3	Aims	3
0.2	Definitions	4
0.2.1	$C(X)$ - an example.	5
0.2.2	Unitization	5
0.2.3	the spectrum	6
0.2.4	more definitions	6
0.2.5	$\mathcal{B}(\mathcal{H})$ - an example.	9
0.3	Representations of C^* -algebras	11

0.1 Preliminaries

0.1.1 History of C^* -Algebras

The noncommutative nature of Werner Heisenberg's work in 1925 on a new quantum mechanics [10] led to Born and Jordan [2], together with Heisenberg [3], developing the matrix mechanics required to concisely summarise the new quantum mechanical model. From 1935-1943, John von Neumann, together with F.J. Murray, developed the theory of *rings of operators* acting on a Hilbert space [17, 18, 19, 25], in an attempt to establish a general framework for this matrix mechanics. These rings of operators are now considered part of the theory of *von Neumann algebras*, a subsection of C^* -algebra theory. Discussion of the seminal quantum mechanical works of Heisenberg can be found in [16], and similarly [23] gives a summary of the works of Jordan expanding on this.

In 1943 [9], Gelfand and Naimark established an abstract characterisation of C^* -algebras, free from dependence on the operators acting on a Hilbert space. The Gelfand-Naimark theorem, which we will be considering here at length, gives the link between these abstract C^* -algebras and the rings

of operators previously studied. Used in the proof of the GN theorem is the Gelfand-Naimark-Segal construction, a pair of results relating cyclic $*$ -representations of C^* -algebras to certain linear functionals on that algebra.

0.1.2 Background Mathematics and Resources.

The following is some mathematics which may prove useful throughout the project, with relevant resources; we will of course be making definitions as needed, this is for further background and related theory.

We will be assuming some familiarity with the following theory, giving some explanation as necessary:

- Rings, algebras and linear spaces.
- Normed spaces, inner product spaces, Banach and Hilbert spaces.
- Point-set topology.

A good broad background on all of these can be found in [24].

Some texts which cover C^* -algebras: Dixmier [4] presents a summary of the general theory up to that time (1977), with [5] focussing on reworking and developing the theory of von Neumann algebras. Sakai [22] gives a treatment of C^* - and von Neumann algebras from a more topological point of view. In [11, 12], the authors aim to make accessible the “vast recent research literature” in this subsection of functional analysis. Blackadar [1] gives a much faster, more encyclopaedic coverage of the theory of operator algebras, and covering more specialised material and applications.

0.1.3 Aims

The aims for this project are:

- Give a good background understanding on C^* -algebras, including topological and geometric interpretation of results where possible.
- Consider the representation theory of C^* -algebras, using the Gelfand-Naimark-Segal construction as a starting point.
- Consider the commutative and general versions of the Gelfand-Naimark theorem, and understand their contents and proof.

List of stuff we’re gonna go right ahead and assume:

- Familiarity with algebras, Banach spaces, Hilbert spaces and other guff.

we will assume knowledge on... (algebra homos map 1 to 1, hausdorff/compact spaces)

brief(er than asst 3) history

most texts start from $\mathcal{B}(\mathcal{H})$ to justify the whole thing. we're algebraists, who don't need no justification. we jump right in at the deep (abstract) end.

write this!

0.2 Definitions

We begin this section with some definitions and an example, then give some results we will need later and finish with a fundamental example.

Definition 1. A *Banach algebra* is a complex Banach space $(A, \|\cdot\|)$ which forms an algebra, such that

$$\|ab\| \leq \|a\|\|b\| \text{ for all } a, b \in A.$$

A **-algebra* is an algebra A with an *involution* map $a \mapsto a^*$ on A such that, for all $a, b \in A$ and for $\alpha \in \mathbb{C}$,

- (i) $a^{**} = (a^*)^* = a$,
- (ii) $(\alpha a + b)^* = \bar{\alpha}a^* + b^*$,
- (iii) $(ab)^* = b^*a^*$.

The element a^* is referred to as the *adjoint* of a .

A *C*-algebra* is a Banach algebra $(A, \|\cdot\|)$ with involution map $a \mapsto a^*$ making it a *-algebra, with the condition that

$$\|a^*a\| = \|a\|^2 \text{ for all } a \in A.$$

This condition is known as the *C* axiom*. There is a weaker, but ultimately equivalent, axiom called the *B* axiom*.

Here we consider complex C*-algebras. The theory of real C*-algebras has advanced....

remark on work in real C*-algebras?

state
and
prove

Unless specified otherwise, by an *ideal* of a Banach algebra, we mean a two-sided ideal. Given a subset S of a C*-algebra A , let $C^*(S)$ denote the *C*-subalgebra of A generated by S* , which is the smallest S^* -subalgebra of A containing S .

cauchy-schwarz, C* subalgebra generated by a set

0.2.1 $C(X)$ - an example.

Given a locally compact Hausdorff space X , let $C(X)$ be the algebra of continuous functions $f : X \rightarrow \mathbb{C}$, with addition and multiplication defined pointwise. Define $\|\cdot\|$ on $C(X)$ by

$$\|f\| := \sup_{x \in X} |f(x)|,$$

that is the norm inherited from the Banach space $\ell^2(X, \mathbb{C})$.

example of $C(X)$?

maybe
move
this
to as
early
as
we
can?

0.2.2 Unitization

If a C^* -algebra A contains an identity element $\mathbb{1}$ such that $a \cdot \mathbb{1} = a = \mathbb{1} \cdot a$ for all $a \in A$, call $\mathbb{1}$ the *unit* in A , and A is then a *unital* C^* -algebra.

Proposition 1. *Any non-unital C^* -algebra A can be isometrically embedded in a unital C^* -algebra \tilde{A} as a maximal ideal.*

tidy up proof

Proof. Let $\tilde{A} = A \oplus \mathbb{C}$ with pointwise addition, and define

$$\begin{aligned} (a, \lambda)(b, \mu) &:= (ab + \lambda b + \mu a, \lambda\mu), \\ (a, \lambda)^* &:= (a^*, \bar{\lambda}), \\ \|(a, \lambda)\| &:= \sup_{\|b\|=1} \|ab + \lambda b\|. \end{aligned}$$

Then \tilde{A} is a $*$ -algebra. The norm $\|(a, \lambda)\|$ is the norm in $\mathcal{B}(A)$ of left-multiplication by a on something iunno??? Thus \tilde{A} is a Banach $*$ -algebra with unit $(0, 1)$. By design, A is a maximal ideal of codimension 1. The embedding $a \mapsto (a, 0)$ is isometric as

$$\|a\| = \|a \cdot \frac{a}{\|a\|}\| \leq \|(a, 0)\| \leq \sup_{\|b\|=1} \|ab\| \leq \|a\|.$$

It remains to verify the C^* -axiom:

$$\begin{aligned}
\|(a, \lambda)\|^2 &= \sup_{\|b\|=1} \|ab + \lambda b\|^2 \\
&= \sup_{\|b\|=1} \|b^*a^*ab + \lambda b^*a^*b + \bar{\lambda}b^*ab + |\lambda|^2b^*b\| \\
&\leq \sup_{\|b\|=1} \|a^*ab + \lambda a^*b + \bar{\lambda}ab + |\lambda|^2b\| \\
&= \|(a^*a + \lambda a^* + \bar{\lambda}a, |\lambda|^2)\| \\
&= \|(a, \lambda)^*(a, \lambda)\| \\
&\leq \|(a, \lambda)^*\| \|(a, \lambda)\|.
\end{aligned}$$

By symmetry of $*$, $\|(a, \lambda)^*\| = \|(a, \lambda)\|$. Hence, the above inequality becomes equality and we have that

$$\|(a, \lambda)^*(a, \lambda)\| = \|(a, \lambda)\|^2.$$

□

In light of this result, we take all C^* -algebras from here to be unital unless specified otherwise. For the results we will consider, we can simply consider the unital case. However, there are important circumstances in advanced theory in which one needs to relax the unital condition.

0.2.3 the spectrum

Given an element $a \in A$ of a C^* -algebra, define its spectrum $\text{sp}(a)$:

$$\text{sp}(a) := \{\lambda \in \mathbb{C} \mid a - \lambda \mathbb{1} \text{ is not invertible in } A\}.$$

state without proof results? AAAAAAAAAAAH I NEED SO MUCH SPECTRAL THEORY AND I DON'T KNOW ANY

0.2.4 more definitions

An element $a \in A$ of a C^* -algebra is called normal?

- *self-adjoint* if $a^* = a$;
- *unitary* if $aa^* = a^*a = \mathbb{1}$;
- *positive* if it is self-adjoint and $\text{sp}(a) \subseteq \mathbb{R}^+$.

Denote the set of self-adjoint elements in A by A_{sa} , and the subset of positive elements in A_{sa} by A^+ . The set of positive elements A_{sa} forms a *partially ordered (real) vector space*, with *positive cone* A^+ . That is to say, all $f, g \in A^+$ satisfy

- (i) $f, -f \in A^+$ implies $f = 0$,
- (ii) $\alpha f \in A^+$ for all $\alpha \in \mathbb{R}^+$,
- (iii) $f + g \in A^+$.

The unit $\mathbb{1}$ is positive, and for any $a \in A_{sa}$ we have $-\|a\|\mathbb{1} \leq a \leq \|a\|\mathbb{1}$. With commuting elements $a, b \in A_{sa}$, we have $(ab)^* = b^*a^* = ba = ab$, so ab is self-adjoint. Since a, b, ab have the same spectrum in A as in the Abelian C^* -subalgebra $C^*(\mathbb{1}, a, b)$, by our spectral theory we have

$$\text{sp}(ab) \subseteq \text{sp}(a)\text{sp}(b).$$

Definition 2. Given Banach $*$ -algebras A and B , a map $\varphi : A \rightarrow B$ is a *$*$ -homomorphism* if it is an algebra homomorphism for which $\varphi(a^*) = \varphi(a)^*$ for all $a \in A$. If A and B are both unital algebras and a homomorphism φ maps $\mathbb{1}_A$ to $\mathbb{1}_B$, say φ is a *unital* homomorphism. If a $*$ -homomorphism φ is one-to-one, call it a *$*$ -isomorphism*.

Proposition 2. Suppose A and B are C^* -algebras and $\varphi : A \rightarrow B$ is a $*$ -homomorphism. Then $\|\varphi(a)\| \leq \|a\|$ for all $a \in A$. If φ is a $*$ -isomorphism, then $\|\varphi(a)\| = \|a\|$ for all $a \in A$.

Proof. □

Definition 3. A *linear functional* on a C^* -algebra A is a linear operator $\rho : A \rightarrow \mathbb{C}$. A linear functional ρ is *positive* if $\rho(a) \geq 0$ for all $a \in A^+$. A *multiplicative* linear functional ρ satisfies $\rho(ab) = \rho(a)\rho(b)$ for all $a, b \in A$. A *state* on A is a positive linear functional ρ such that $\|\rho\| = 1$ and $\rho(a) \geq 0$ for all positive elements $a \in A^+$. Denote by $\mathcal{S}(A)$ the set of all states on A . An extreme point of $\mathcal{S}(A)$ is called a *pure state* on A , and the set of pure states on A is denoted by $\mathcal{P}(A)$.

It is a simple exercise, using the fact that $A^+ \subseteq A_{sa}$, to verify that a linear functional ρ is a pure state on A if and only if the restriction $\rho|_{A_{sa}}$ is a pure state on A_{sa} . Every pure state on A_{sa} extends to a pure state on A . We will need the following few results on pure states later.

state
in
'spec-
trum',
-
4.1.5

is
uni-
tal
nec-
es-
sary?

show
that
 $\|\rho\| = \rho(\mathbb{1})$.
ex-
plain
ex-
treme
point.

justify.
4.3.13.

Proposition 3. *A state ρ on A_{sa} is pure if and only if, for all positive linear functionals τ on A_{sa} such that $0 \leq \tau \leq \rho$, we have $\tau = \lambda\rho$ for some $\lambda \in \mathbb{R}$.*

Proof. (Adapted from 3.4.6). Suppose that $\tau = \lambda\rho$ for all $0 \leq \tau \leq \rho$, and suppose we can write $\rho = \alpha\rho_1 + (1 - \alpha)\rho_2$ for some $0 \leq \alpha \leq 1$ and some $\rho_1, \rho_2 \in \mathcal{S}(A_{sa})$. Then $0 \leq \alpha\rho_1 \leq \rho$, so $\alpha\rho_1 = \lambda\rho$. Then $\rho_1(\mathbb{1}) = 1 = \rho(\mathbb{1})$, so $\alpha = \lambda$ so $\rho_1 = \rho$. Similarly, we can show that $\rho_2 = \rho$, and so we conclude that ρ is pure.

Conversely, suppose that ρ is a pure state and $0 \leq \tau \leq \rho$. Applying this to $\mathbb{1}$, we get $0 \leq \tau(\mathbb{1}) \leq \rho(\mathbb{1}) = 1$. Let $\lambda = \tau(\mathbb{1})$. If $\lambda = 0$, then for any $a \in A_{sa}$, applying τ to $-\|a\|\mathbb{1} \leq a \leq \|a\|\mathbb{1}$ gives

$$0 = -\|a\|\lambda = \tau(-\|a\|\mathbb{1}) \leq \tau(a) \leq \tau(\|a\|\mathbb{1}) = \|a\|\lambda = 0,$$

so $\tau = 0 = \lambda\rho$. A similar argument shows that $\lambda = 1$ implies $\tau - \rho = 0$ so that $\tau = \rho = \lambda\rho$. If $0 < \lambda < 1$, we can write $\rho = \lambda\rho_1 + (1 - \lambda)\rho_2$ for $\rho_1 = \lambda^{-1}\tau$ and $\rho_2 = (1 - \lambda)^{-1}(\rho - \tau)$. ρ is pure so $\tau = \lambda\rho_1 = \lambda\rho$. □

Proposition 4. *The set of pure states on an Abelian C^* -algebra A is precisely the set of multiplicative linear functionals on A .*

Proof. (Adapted from $K\mathcal{E}R$, 4.4.1). Suppose ρ is a pure state on A . To show that $\rho(ab) = \rho(a)\rho(b)$ for $a, b \in A$, we restrict attention to the case where $0 \leq b \leq \mathbb{1}$. Linearity gives us the general case. In this case, for $h \in A^+$ we show have that $0 \leq hb \leq h$, so $0 \leq \rho(hb) \leq \rho(h)$. Hence $\rho_b(a) := \rho(ab)$ for $a \in A$ defines a positive linear functional on A with $\rho_b \leq \rho$. The restriction $\rho|_{A_{sa}}$ is a pure state on A_{sa} and $\rho_b|_{A_{sa}} \leq \rho|_{A_{sa}}$, and it follows from Proposition 3 that $\rho_b|_{A_{sa}} = \alpha\rho|_{A_{sa}}$ for some $\alpha \in \mathbb{R}^+$. Hence $\rho_b = \alpha\rho$ and so for $a \in A$:

$$\rho(ab) = \rho_b(a) = \alpha\rho(a) = \alpha\rho(\mathbb{1})\rho(a) = \rho_b(\mathbb{1})\rho(a) = \rho(b)\rho(a)$$

Conversely, suppose ρ is a multiplicative linear functional. Suppose we show a state. can write $\rho = \alpha\rho_1 + \beta\rho_2$ for states ρ_1, ρ_2 on A and $\alpha, \beta > 0$ such that $\alpha + \beta = 1$. For $c \in A_{sa}$, by the Cauchy-Schwarz inequality we have for $j = 1, 2$:

$$(\rho_j(c))^2 = (\rho_j(\mathbb{1}c))^2 \leq \rho_j(\mathbb{1})\rho_j(c^2) = \rho_j(c^2).$$

Then:

$$\begin{aligned} 0 &= \rho(c^2) - \rho(c)^2 \\ &= \alpha\rho_1(c^2) + \beta\rho_2(c^2) - (\alpha\rho_1(c) + \beta\rho_2(c))^2 \\ &\geq \alpha(\alpha + \beta)\rho_1(c)^2 + \beta(\alpha + \beta)\rho_2(c)^2 - (\alpha\rho_1(c) + \beta\rho_2(c))^2 \\ &= \alpha\beta(\rho_1(c) - \rho_2(c))^2. \end{aligned}$$

Hence $\rho_1(c) = \rho_2(c)$, for all $c \in A_{sa}$, so $\rho_1 = \rho_2$ and we conclude that ρ is a pure state. \square

define weak* topology on $P(S)$

0.2.5 $\mathcal{B}(\mathcal{H})$ - an example.

This section concerns the fundamental example of a C^* -algebra - the set $\mathcal{B}(\mathcal{H})$ of bounded linear operators on a Hilbert space \mathcal{H} . Here we will demonstrate that $\mathcal{B}(\mathcal{H})$ is a C^* -algebra and give some basic results.

Claim. $\mathcal{B}(\mathcal{H})$ is a C^* -algebra with the operator norm

$$\|T\| := \sup_{\|x\|=1} \|Tx\|$$

and involution taking T to its adjoint map T^* . The identity map $I : x \mapsto x$ is a unit for $\mathcal{B}(\mathcal{H})$

Proof. $\|\cdot\|$ is a norm on $\mathcal{B}(\mathcal{H})$. Let $\{T_n\}_{n \in \mathbb{N}}$ be a Cauchy sequence in $\mathcal{B}(\mathcal{H})$. Then for any positive ϵ , there is a positive integer N such that

$$\|T_m - T_n\| < \epsilon \text{ for all } m, n \geq N.$$

Applying $T_m - T_n$ to $x \in \mathcal{H}$, we have

$$\|T_m x - T_n x\| \leq \|T_m - T_n\| \|x\| < \epsilon \|x\|, \quad (1)$$

so $\{T_n x\}_{n \in \mathbb{N}}$ is a Cauchy sequence in \mathcal{H} , converging to an element in \mathcal{H} . Define a linear operator $T : \mathcal{H} \rightarrow \mathcal{H}$ by

$$Tx := \lim_{n \rightarrow \infty} T_n x \text{ for } x \in \mathcal{H}.$$

Taking limits as m tends to infinity in equation (1), we obtain

$$\|Tx - T_n x\| < \epsilon \|x\| \text{ for all } n \geq N,$$

and so we have that $T - T_n$ (and hence $T = (T - T_n) + T_n$) is a bounded operator and

$$\|T - T_n\| < \epsilon \text{ for all } n \geq N.$$

We conclude that $T_n \rightarrow T$, and so $\mathcal{B}(\mathcal{H})$ is complete.

Since boundedness is equivalent to continuity on \mathcal{H} , given $S, T \in \mathcal{B}(\mathcal{H})$, the operator $ST : \mathcal{H} \rightarrow \mathcal{H}; x \mapsto (S \circ T)(x)$ is bounded on \mathcal{H} . Given $x \in \mathcal{H}$ and $\lambda \in \mathbb{C}$,

$$\begin{aligned} ((\lambda S)T)(x) &= ((\lambda S) \circ T)(x) \\ &= \lambda S(Tx) \\ &= \lambda(S \circ T)(x) \\ &= \lambda ST(x), \end{aligned}$$

so that $(\lambda S)T = \lambda ST$ in $\mathcal{B}(\mathcal{H})$, whence $\mathcal{B}(\mathcal{H})$ is an algebra. We have

$$\begin{aligned} \|ST\| &= \sup_{\|x\|=1} \|STx\| \\ &= \sup_{\|x\|=1} \|S(Tx)\| \\ &\leq \|S\| \sup_{\|x\|=1} \|Tx\| \\ &= \|S\| \|T\|. \end{aligned}$$

To see that $*$ is an involution, use the fact that the adjoint operator is unique for each operator and the following equalities.

$$\begin{aligned} \text{(i)} \quad \langle (\alpha T + S)^* x, y \rangle &= \langle x, \alpha T + Sy \rangle \\ &= \overline{\alpha} \langle x, Ty \rangle + \langle x, Sy \rangle \\ &= \overline{\alpha} \langle T^* x, y \rangle + \langle S^* x, y \rangle \\ &= \langle (\overline{\alpha} T^* + S^*) x, y \rangle. \end{aligned}$$

$$\begin{aligned} \text{(ii)} \quad \langle (T^*)^* x, y \rangle &= \langle x, T^* y \rangle \\ &= \overline{\langle T^* y, x \rangle} \\ &= \overline{\langle y, Tx \rangle} \\ &= \langle Tx, y \rangle. \end{aligned}$$

$$\begin{aligned} \text{(iii)} \quad \langle (ST)^* x, y \rangle &= \langle x, STy \rangle \\ &= \langle S^* x, Ty \rangle \\ &= \langle T^* S^* x, y \rangle. \end{aligned}$$

It remains to demonstrate the C^* -axiom on $\mathcal{B}(\mathcal{H})$. For all $x \in \mathcal{H}$, we have

$$\|Tx\|^2 = \langle Tx, Tx \rangle = \langle T^* Tx, x \rangle \leq \|T^* T\| \|x\|^2,$$

so that

$$\|T\|^2 \leq \|T^*T\| \leq \|T^*\|\|T\| = \|T\|^2.$$

It is clear that I is a unit. Hence, the claim. \square

example of $\mathcal{B}(\mathcal{H})$ at end of section to retro-motivate notation. discuss nomenclature (state etc) coming from QM

0.3 Representations of C*-algebras

to include all representation theory, including GNS, CGN and GN

Theorem 1 (Gelfand-Naimark, commutative). *Every commutative C*-algebra A is *-isomorphic to $C(X)$, the algebra of continuous functions on a compact Hausdorff space X .*

Proof. Our compact topological space will be the set $\mathcal{P}(A)$ of pure states, endowed with the weak* topology as defined above. \square

Definition 4. Given a C*-algebra A , a *representation of A on a Hilbert space \mathcal{H}* is a *-homomorphism $\varphi : A \rightarrow \mathcal{B}(\mathcal{H})$. An isomorphic representation is called *faithful*. If there exists an element $x \in \mathcal{H}$ such that the set $\{\varphi(a)x \mid a \in A\}$ is everywhere-dense in \mathcal{H} , say that φ is a *cyclic* representation, with *cyclic vector* x .

remark about nonunital commutative.

Theorem 2 (Gelfand-Naimark-Segal construction). *If ρ is a state on a C*-algebra A , then there exists a cyclic representation π_ρ of A on a Hilbert space \mathcal{H}_ρ , with unit cyclic vector x_ρ , such that*

$$\rho(a) = \langle \pi_\rho(a)x_\rho, x_\rho \rangle, \quad \forall a \in A.$$

Proof. We will construct from ρ the space \mathcal{H}_ρ , representation π_ρ , and vector x_ρ , and demonstrate the required properties.

Consider the *left kernel* of ρ :

$$L_\rho := \{t \in A \mid \rho(t^*t) = 0\}.$$

For $a, b \in A$, define $\langle a, b \rangle_0 := \rho(b^*a)$. Then $L_\rho = \{t \in A \mid \langle t, t \rangle_0 = 0\}$, and $\langle \cdot, \cdot \rangle_0$ satisfies

(i) Linearity in 1st argument: for $a, b \in A$, $\alpha, \beta \in \mathbb{C}$:

$$\begin{aligned}\langle \alpha a + \beta b, c \rangle_0 &= \rho(c^*(\alpha a + \beta b)) \\ &= \rho(\alpha c^*a + \beta c^*b) \\ &= \alpha \rho(c^*a) + \beta \rho(c^*b) \\ &= \alpha \langle a, c \rangle_0 + \beta \langle b, c \rangle_0.\end{aligned}$$

(ii) Conjugate symmetric: for $a, b \in A$:

$$\begin{aligned}\langle b, a \rangle_0 &= \rho(a^*b) \\ &= \rho((b^*a)^*) \\ &= \overline{\rho(b^*a)} \\ &= \overline{\langle a, b \rangle_0}.\end{aligned}$$

(iii) Positive semi-definite.

why?

Note that $\langle \cdot, \cdot \rangle$ is not necessarily positive definite on $A - L_\rho$ is exactly where this fails.

L_ρ is a linear subspace of A : Consider

sentence

$$L := \{t \in A \mid \langle t, a \rangle_0 = 0, \forall a \in A\} \subseteq L_\rho.$$

For $t \in L_\rho$, by Cauchy-Schwarz we have

$$|\langle t, a \rangle_0|^2 \leq \langle t, t \rangle_0 \langle a, a \rangle_0, \quad \forall a \in A;$$

that is,

$$\langle t, a \rangle_0 = 0, \quad \forall a \in A,$$

so $t \in L$ and $L_\rho = L$. Now, for $a, b \in L$, $\alpha \in \mathbb{C}$ and $c \in A$:

$$\langle \alpha a + b, c \rangle_0 = \alpha \langle a, c \rangle_0 + \langle b, c \rangle_0 = 0,$$

so $\alpha a + b \in L$; also, $\langle 0, c \rangle_0 = 0$ so $0 \in L$. Hence, $L(= L_\rho)$ is a linear subspace of A .

For $s \in A$, $t \in L_\rho$, by the Cauchy Schwarz inequality [ref] we have

$$\begin{aligned}|\rho(s^*t)|^2 &= |\langle t, s \rangle_0|^2 \\ &\leq \langle t, t \rangle_0 \cdot \langle s, s \rangle_0 \\ &= \rho(t^*t) \cdot \rho(s^*s) \\ &= 0,\end{aligned}$$

so $\rho(s^*t) = 0$. Letting $s = a^*at$ for $a \in A$, then

$$\begin{aligned}\rho((at)^*at) &= \rho(at^*a^*at) \\ &= \rho((a^*at)^*t) \\ &= \rho(s^*t) \\ &= 0,\end{aligned}$$

so that $at \in L_\rho$, for all $a \in A$ and $t \in L_\rho$; we conclude that L_ρ is a left ideal in A . L_ρ is the preimage in A of $\{0\}$ under the continuous map $t \mapsto \rho(t^*t)$, so is closed.

Consider now $V_\rho := A/L_\rho$, with $\langle \cdot, \cdot \rangle$ defined by

$$\langle a + L_\rho, b + L_\rho \rangle := \langle a, b \rangle_0, \quad \text{for } a + L_\rho, b + L_\rho \in V_\rho.$$

It follows from properties *i*), *ii*) and *iii*) of $\langle \cdot, \cdot \rangle_0$ that $\langle \cdot, \cdot \rangle$ is an inner product on V_ρ - with

$$\begin{aligned}\langle a + L_\rho, a + L_\rho \rangle = 0 &\iff \langle a, a \rangle = 0 \\ &\iff a \in L_\rho \\ &\iff a + L_\rho = 0 + L_\rho\end{aligned}$$

giving positive definiteness. The completion of V_ρ with respect to $\langle \cdot, \cdot \rangle$ is a Hilbert space - this is the Hilbert space \mathcal{H}_ρ we're looking for.

Now we fix $a \in A$, and consider the map

$$\pi_a : V_\rho \rightarrow V_\rho; b + L_\rho \mapsto ab + L_\rho.$$

Let $b_1, b_2 \in A$ be such that $b_1 + L_\rho = b_2 + L_\rho$. Then:

$$\begin{aligned}\implies b_1 - b_2 &\in L_\rho \\ \implies a(b_1 - b_2) &\in L_\rho \\ \implies ab_1 - ab_2 &\in L_\rho \\ \implies ab_1 + L_\rho &= ab_2 + L_\rho \\ \implies \pi_a(b_1 + L_\rho) &= \pi_a(b_2 + L_\rho).\end{aligned}$$

Hence π_a defines a linear operator on V_ρ .

For $b + L_\rho \in V_\rho$:

$$\begin{aligned}\|a\|^2 \cdot \|b + L_\rho\| - \|\pi_a(b + L_\rho)\| &= \|a\|^2 \cdot \|b + L_\rho\| - \|ab + L_\rho\| \\ &= \|a\|^2 \cdot \langle b + L_\rho, b + L_\rho \rangle - \langle ab + L_\rho, ab + L_\rho \rangle \\ &= \|a\|^2 \cdot \rho(b^*b) - \rho((ab)^*ab) \\ &= \rho(\|a\|^2 b^*b - b^*a^*ab) \\ &= \rho(b^*(\|a\|^2 \mathbb{1} - a^*a)b) \\ &\geq 0.\end{aligned}$$

start
sen-
tence
prop-
erly

Thus π_a is a bounded operator, with $\|\pi_a\| \leq \|a\|$. By continuity, π_a extends to a bounded operator on \mathcal{H}_ρ – say $\pi_\rho(a) : \mathcal{H}_\rho \rightarrow \mathcal{H}_\rho$ such that

$$\pi_\rho(a)(v) = \pi_a(v)$$

for $v \in V_\rho$. Then $\pi_\rho(a) \in \mathcal{B}(\mathcal{H}_\rho)$ for each $a \in A$, so π_ρ defines a map $A \rightarrow \mathcal{B}(\mathcal{H}_\rho)$ such that $a \mapsto \pi_\rho(a)$. This will be our representation.

Now, for $a, b \in A$, $c + L_\rho \in V_\rho$ and $\alpha \in \mathbb{C}$:

$$\begin{aligned} \pi_{\alpha a + b}(c + L_\rho) &= (\alpha a + b)(c + L_\rho) \\ &= (\alpha ac + L_\rho) + (bc + L_\rho) \\ &= \alpha \pi_a(c + L_\rho) + \pi_b(c + L_\rho), \end{aligned}$$

so that $\pi_{\alpha a + b} = \alpha \pi_a + \pi_b$ on V_ρ .

For $a, b \in A$ and $c + L_\rho \in V_\rho$:

$$\begin{aligned} \pi_{ab}(c + L_\rho) &= abc + L_\rho \\ &= \pi_a(bc + L_\rho) \\ &= \pi_a(\pi_b(c + L_\rho)) \\ &= (\pi_a \cdot \pi_b)(c + L_\rho), \end{aligned}$$

so that $\pi_{ab} = \pi_a \cdot \pi_b$ on V_ρ .

For $a \in A$ and $b + L_\rho, c + L_\rho \in V_\rho$:

$$\begin{aligned} \langle b + L_\rho, \pi_a^*(c + L_\rho) \rangle &= \langle \pi_a(b + L_\rho), c + L_\rho \rangle \\ &= \langle ab + L_\rho, c + L_\rho \rangle \\ &= \rho(c^*ab) \\ &= \rho((a^*c)^*b) \\ &= \langle b + L_\rho, a^*c + L_\rho \rangle \\ &= \langle b + L_\rho, \pi_{a^*}(c + L_\rho) \rangle, \end{aligned}$$

so that $\pi_a^* = \pi_{a^*}$ on V_ρ .

$V_\rho \subset \mathcal{H}_\rho$ is a dense subset, so the three properties above hold on \mathcal{H}_ρ by continuity. Hence, $\pi_\rho : A \rightarrow \mathcal{B}(\mathcal{H}_\rho)$ is a representation of A . As to the unit vector, consider $x_\rho := \mathbb{1} + L_\rho \in V_\rho$. Then for $a \in A$,

$$\begin{aligned} \langle \pi_\rho(a)x_\rho, x_\rho \rangle &= \langle \pi_a(\mathbb{1} + L_\rho), \mathbb{1} + L_\rho \rangle \\ &= \langle a + L_\rho \mathbb{1} + L_\rho \rangle \\ &= \rho(a); \end{aligned}$$

in particular, $\langle x_\rho, x_\rho \rangle = \rho(\mathbb{1}) = 1$, so x_ρ is a unit vector in \mathcal{H}_ρ . \square

cont
of
what?

cont
of
what?

example of this construction on $C(X)$? may just be a short explanation of how $B(H)$ and $C(X)$ link together. can then talk about noncommutative topology!

Theorem 3 (Gelfand-Naimark). *Every C^* -algebra has a faithful representation.*

Proof. for this we just take the direct sum representation of the representations given from GNS by some set of states containing all pure states. \square

further topics: K-theory, group C^* algebras, amenable algebras, von neumann algebras,

references!!!!

Bibliography

- [1] Blackadar, B., *Operator Algebras: Theory of C^* -Algebras and von Neumann Algebras*. Encyclopaedia of Mathematical Sciences, 122. Operator Algebras and Non-commutative Geometry, III. Springer-Verlag, Berlin (2006).
- [2] Born, M. & Jordan, P., *Zur Quantenmechanik*. Z. Physik (1925) 34: 858.
- [3] Born, M.; Heisenberg, W. & Jordan, P., *Zur Quantenmechanik. II*. Z. Physik (1926) 35: 557.
- [4] Dixmier, J., *C^* -algebras*. Holland Mathematical Library, Vol. 15. North-Holland Publishing Co., Amsterdam – New York – Oxford (1977).
- [5] Dixmier, J., *von Neumann algebras*. North-Holland Publishing Co., Amsterdam – New York (1981).
- [6] Elliott, G. A., *The classification problem for amenable C^* -algebras*. Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Zürich, 1994), pp. 922-932, Birkhuser, Basel (1995).
- [7] Fell, J. M. G., *The structure of algebras of operator fields*. Acta Math. 106, 1961, 233-280.
- [8] Gardella, E. E., *Compact group actions on C^* -algebras: classification, non-classifiability, and crossed products and rigidity results for L^p -operator algebras*. Thesis (Ph.D.) - University of Oregon (2015).
- [9] Gelfand, I. & Neumark, M., *On the imbedding of normed rings into the ring of operators in Hilbert space*. Rec. Math. [Mat. Sbornik] N.S. 12(54) (1943), pp. 197-213.
- [10] Heisenberg, W., *Über quantentheoretische Umdeutung kinematischer und mechanischer Beziehungen*. Z. Physik (1925) 33: 879.

- [11] Kadison, R. V. & Ringrose, J. R., *Fundamentals of the theory of operator algebras: Vol. I. Elementary theory*. Pure and Applied Mathematics, 100. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York (1983).
- [12] Kadison, R. V. & Ringrose, J. R., *Fundamentals of the theory of operator algebras: Vol. II. Advanced theory*. Pure and Applied Mathematics, 100. Academic Press, Inc., Orlando, FL (1986).
- [13] Lin, H., *An introduction to the classification of amenable C^* -algebras*. World Scientific Publishing Co., Inc., River Edge, NJ (2001).
- [14] Lin, H. & Niu, Z., *Lifting KK -elements, asymptotic unitary equivalence and classification of simple C^* -algebras*. Adv. Math. 219 (2008), no. 5, pp. 1729-1769.
- [15] Lin, H., *Asymptotic unitary equivalence and classification of simple amenable C^* -algebras*. Invent. Math. 183 (2011), no. 2, pp. 385-450.
- [16] MacKinnon, E., *Heisenberg, Models, and the Rise of Matrix Mechanics*. Hist. Stud. Phys. Sci., Vol. 8 (1977), pp. 137–188
- [17] Murray, F. J. & von Neumann, J., *On rings of operators*. Ann. of Math. (2) 37 (1936), no. 1, pp. 116-229.
- [18] Murray, F. J. & von Neumann, J., *On rings of operators. II*. Trans. Amer. Math. Soc. 41 (1937), no. 2, pp. 208-248.
- [19] Murray, F. J. & von Neumann, J., *On rings of operators. IV*. Ann. of Math. (2) 44, (1943), pp. 716-808.
- [20] Niemiec, P., *Elementary Approach to Homogeneous C^* -algebras*. Rocky Mountain J. Math. 45 (2015), no. 5, pp 1591–1630.
- [21] Pedersen, G. K., *C^* -algebras and their automorphism groups*. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], London-New York (1979).
- [22] Sakai, S., *C^* -algebras and W^* -algebras*. Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 60. Springer-Verlag, New York-Heidelberg (1971).
- [23] Schroer, B., *Pascual Jordan, Glory and Demise and his legacy in contemporary local quantum physics*. Unpublished manuscript (2003).

- [24] Simmons, G., *Introduction to Topology and Modern Analysis*. Robert E. Krieger Publishing Co., Inc., Melbourne, Fl. (1983).
- [25] von Neumann, J., *On rings of operators. III*. Ann. of Math. (2) 41, (1940), pp. 94-161.