

C*-ALGEBRAS, THE GELFAND-NAIMARK THEOREM, AND [OTHER THING]

LUKE ARMITAGE

1. INTRODUCTION

1.1. History of C*-Algebras. The noncommutative nature of W. Heisenberg's work on (finish sentence) [4] lead to Born and Jordan [1], together with Heisenberg [2], developing the matrix mechanics required to concisely convey the new quantum mechanical model. From 1930-1943, J. von Neumann, together with F. J. Murray, developed the theory of *rings of operators* acting on a Hilbert space [7], [8], [10], [9], in an attempt to establish a general framework for this matrix mechanics. These rings of operators are now considered part of the theory of *von Neumann algebras*, a subsection of C*-algebra theory. Discussion of the seminal quantum mechanical works of Heisenberg can be found in [5], and likewise in [6], for the works of Jordan expanding on this.

In [3], Gelfand and Naimark established an abstract characterisation of C*-algebras, free from dependence on the operators acting on a Hilbert space. The Gelfand-Naimark theorem, which we will be considering here at length, gives the link between these abstract C*-algebras and the rings of operators previously studied. Used in the proof of the G-N theorem is the Gelfand-Naimark-Segal construction, a pair of results relating cyclic *-representations of C*-algebras to certain linear functionals on that algebra.

1.2. C*-Algebras. A brief overview of what a C*-algebra is.

- banach space
- banach algebra
- *-algebra
- C* algebra
- representations

Definition 1. A *C*-algebra* is a Banach algebra $(A, \|\cdot\|)$ with involution $*$: $A \rightarrow A$ with the condition that

$$\|x^*x\| = \|x\|^2 \text{ for all } x \in A.$$

This condition is known as the *C* axiom*. The study of C*-algebras started with the consideration of matrix mechanics by H

Theorem (Gelfand-Naimark, Commutative). *Every commutative, unital C^* -algebra A is isometrically $*$ -isomorphic to the algebra of continuous functions on the algebra of characters on A .*

The theorem first appeared in [gelfand-naimark] in a form relating ‘normed rings’ and a closed subrings of the set of bounded operators on a Hilbert space.

1.3. **Aims.** The aims for my project are, provisionally:

- Take the Gelfand-Naimark theorem, and understand its contents and proof.
- Consider the representation theory of C^* -algebras, using the Gelfand-Naimark-Segal construction as a starting point.
- (as ‘further reading’) Give an overview of areas in which operator algebra theory can be taken (for example, Cuntz algebras and operator K-theory, von Neumann algebras and Factors, abstract harmonic analysis).

REFERENCES

- [1] Born, M. & Jordan, P., *Zur Quantenmechanik*. Z. Physik (1925) 34: 858.
- [2] Born, M.; Heisenberg, W. & Jordan, P., *Zur Quantenmechanik. II*. Z. Physik (1926) 35: 557.
- [3] Gelfand, I. & Neumark, M., *On the imbedding of normed rings into the ring of operators in Hilbert space*. Rec. Math. [Mat. Sbornik] N.S. 12(54), (1943), pp. 197-213.
- [4] Heisenberg, W., *Über quantentheoretische Umdeutung kinematischer und mechanischer Beziehungen*. Z. Physik (1925) 33: 879.
- [5] MacKinnon, E., *Heisenberg, Models, and the Rise of Matrix Mechanics*. Hist. Stud. Phys. Sci., Vol. 8 (1977), pp. 137–188
- [6] Schroer, B., *Pascual Jordan, Glory and Demise and his legacy in contemporary local quantum physics*. (no idea where published)
- [7] Murray, F. J. & von Neumann, J., *On rings of operators*. Ann. of Math. (2) 37 (1936), no. 1, pp. 116-229.
- [8] Murray, F. J. & von Neumann, J., *On rings of operators. II*. Trans. Amer. Math. Soc. 41 (1937), no. 2, pp. 208-248.
- [9] Murray, F. J. & von Neumann, J., *On rings of operators. IV*. Ann. of Math. (2) 44, (1943), pp. 716-808.
- [10] von Neumann, J., *On rings of operators. III*. Ann. of Math. (2) 41, (1940), pp. 94-161.