CSci 423 Homework 1

Due: 1:00 pm, Wednesday, 9/12 Eric Shih

1. (5 points) Prove by induction that for all integers $n \ge 4$ the inequality $2^n < n!$ holds.

Basis Step: $2^n < n!$ when n = 4 $\implies 2^4 < 4!$ $\implies 2^4 < 4 \cdot 3 \cdot 2 \cdot 1$ $\implies 16 < 24$

The basis step is confirmed

Inductive Hypothesis:

 $\forall k \text{ assume } 2^k < k! \text{ for some } k \le n \text{ where } k > 4$

Inductive Step:

n = k + 1 such that $2^{k+1} < (k+1)!$ $\implies 2^k \cdot 2 < k \cdot k!$

Since k > 2 as shown before, we can assume that by the inductive hypothesis that $2^{k+1} < (k+1)!$

2. (5 points) Prove by contradiction that $2 - \sqrt{2}$ is not a rational number.

We know that any any sum of a rational and irrational number is irrational, so we will first focus on $\sqrt{2}$. To prove by contradiction we first assume $\sqrt{2}$ is a rational number.

 \exists int a, b with gcd(a,b) = 1, such that $\sqrt{2} = \frac{a}{b}$ $\implies 2 = \frac{a^2}{b^2}$

We then find the greatest common denominator (gcd) of a and b.

 $\exists k \ a = 2k \implies (2k)^2 = 2b^2$ $\exists l \ b = 2l$

From this, we deduce that the $gcd(a,b) \ge 2$, which contradicts the fact that gcd(a,b) = 1, and proves $\sqrt{2}$ irrational. Therefore if $\sqrt{2}$ is irrational, $2 - \sqrt{2}$ is irrational by the properties of irrational numbers.


3. (5 points) Find the error in the following proof that all horses are the same color.

There is an incorrect claim when H_1 is presented as having k horses where h = k. If a horse is removed from the base case of h = 1, then there are no horses left, therefore no way to know what color of the set of horses is. The entire proof relies on this property, and since it cannot be verified, the rest of the proof is false.


4. (5 points for each question) Give state diagrams of DFAs recognizing the following languages. In all parts the alphabet is $\{0,1\}$.

1

a. $\{w|w \text{ begins with a 1 and ends with a 0}\}$




e. $\{w|w \text{ starts with } 0 \text{ and has odd length, or starts with } 1 \text{ and has even length}\}$




d = odd, e = even

i. $\{w | \text{ every odd position of } w \text{ is a } 1\}$



d = odd, e = even

n. All strings except the empty string



5. (5 points) Give the state diagram of a DFA that recognizes (accepts) the set of strings such that in each string the number of 0s is divisible by 5 and the number of 1s is divisible by 3.

