CSCI3230 / ESTR3108 2026-27 First Term Assignment 1

I declare that the assignment here submitted is original except for source material explicitly acknowledged, and that the same or closely related material has not been previously submitted for another course. I also acknowledge that I am aware of University policy and regulations on honesty in academic work, and of the disciplinary guidelines and procedures applicable to breaches of such policy and regulations, as contained in the following websites.

University Guideline on Academic Honesty:
`http://www.cuhk.edu.hk/policy/academichonesty/`

Faculty of Engineering Guidelines to Academic Honesty:
`http://www.erg.cuhk.edu.hk/erg-intra/upload/documents/ENGG_Discipline.pdf`

Student Name: Jiang Huanyun **Student ID:** 1155233165

Question 1

The RSS objective is $J(\Theta) = \sum_{i=1}^{n} \big(f_\Theta(x^{(i)}) - y^{(i)}\big)^2$. Stacking the samples into a design matrix $\mathbf{X}$ (rows $1, x^{(i)}, \dots$) and the labels into $\mathbf{y}$, the minimizer solves the *normal equations* $\mathbf{X}^\top \mathbf{X}\Theta = \mathbf{X}^\top \mathbf{y}$.

(a) Analytic solution of $f_\Theta(x) = \theta_0 + \theta_1 x$

Here $\mathbf{X}$ has rows $\big(1,\, x^{(i)}\big)$, $i = 1, \dots, 10$. The needed sums are

$$\sum_i x^{(i)} = 56.1, \qquad \sum_i \big(x^{(i)}\big)^2 = 353.09, \qquad \sum_i y^{(i)} = 8.5, \qquad \sum_i x^{(i)} y^{(i)} = 52.88,$$

so the normal equations are

$$\begin{pmatrix} 10 & 56.1 \\ 56.1 & 353.09 \end{pmatrix} \begin{pmatrix} \theta_0 \\ \theta_1 \end{pmatrix} = \begin{pmatrix} 8.5 \\ 52.88 \end{pmatrix}.$$

The determinant is $\Delta = 10 \times 353.09 - 56.1^2 = 383.69 > 0$, hence the unique solution

$$\theta_1 = \frac{10 \times 52.88 - 56.1 \times 8.5}{383.69} = \frac{51.95}{383.69} = 0.1354, \quad \theta_0 = \bar{y} - \theta_1 \bar{x} = 0.85 - 0.1354 \times 5.61 = 0.0904.$$

Therefore

$$\boxed{f_\Theta(x) = 0.0904 + 0.1354\, x}, \qquad \text{training RSS} = 4.7816.$$

Figure 1 shows the fitted line together with the training points.




Figure 1: Training points and the fitted linear model $f_\Theta(x) = 0.0904 + 0.1354\,x$.

(b) Analytic solution of $f_\Theta(x) = \theta_0 + \theta_1 x + \theta_2 x^2$

Now the rows of $\mathbf{X}$ are $\big(1,\, x^{(i)},\, (x^{(i)})^2\big)$ and the normal equations form the 3×3 system

$$\begin{pmatrix} 10 & 56.1 & 353.09 \\ 56.1 & 353.09 & 2367.981 \\ 353.09 & 2367.981 & 16523.1077 \end{pmatrix} \begin{pmatrix} \theta_0 \\ \theta_1 \\ \theta_2 \end{pmatrix} = \begin{pmatrix} 8.5 \\ 52.88 \\ 373.386 \end{pmatrix},$$

where $\sum_i (x^{(i)})^3 = 2367.981$, $\sum_i (x^{(i)})^4 = 16523.1077$ and $\sum_i (x^{(i)})^2 y^{(i)} = 373.386$. Solving this system,

$$\boxed{f_\Theta(x) = 3.0589 - 1.2711\,x + 0.1394\,x^2}, \qquad \text{training RSS} = 1.8765.$$

Figure 2 shows the fitted parabola.




Figure 2: Training points and the fitted quadratic model $f_\Theta(x) = 3.0589 - 1.2711\,x + 0.1394\,x^2$.

(c) Analytic solution of the quartic model $f_\Theta(x) = \theta_0 + \theta_1 x + \theta_2 x^2 + \theta_3 x^3 + \theta_4 x^4$

The rows of $\mathbf{X}$ are now $\big(1,\ x^{(i)},\ (x^{(i)})^2,\ (x^{(i)})^3,\ (x^{(i)})^4\big)$ and the normal equations form a 5×5 system

$$\begin{pmatrix} 10 & 56.1 & 353.09 & 2367.981 & 16523.1077 \\ 56.1 & 353.09 & 2367.981 & 16523.1077 & 118449.431 \\ 353.09 & 2367.981 & 16523.1077 & 118449.431 & 865703.6178 \\ 2367.981 & 16523.1077 & 118449.431 & 865703.6178 & 6418500.4415 \\ 16523.1077 & 118449.431 & 865703.6178 & 6418500.4415 & 48113659.523 \end{pmatrix} \begin{pmatrix} \theta_0 \\ \theta_1 \\ \theta_2 \\ \theta_3 \\ \theta_4 \end{pmatrix} = \begin{pmatrix} 8.5 \\ 52.88 \\ 373.386 \\ 2745.6248 \\ 20503.1486 \end{pmatrix},$$

where $\sum_i (x^{(i)})^k y^{(i)}$ for $k = 3, 4$ are 2745.6248 and 20503.1486. Solving,

$$\boxed{f_\Theta(x) = -12.1769 + 16.4458\,x - 6.3777\,x^2 + 0.9547\,x^3 - 0.0485\,x^4}, \quad \text{training RSS} = 0.7712.$$

Figure 3 shows the fitted curve together with the training points.




Figure 3: Training points and the fitted quartic model. The curve wiggles through the training points and takes extreme values outside the data range.

(d) Underfitting / overfitting and test-set verification

Qualitative comparison.

- **Degree 1 — underfitting.** The line cannot capture the curvature of the data (the dip around $x \approx 4.5$ is ignored). The residuals are systematically positive at the ends and negative in the middle; the model has high *bias* and its errors are large on *both* training and test sets.
- **Degree 4 — overfitting.** The quartic is flexible enough to nearly interpolate the 10 training points (smallest training RSS, 0.7712), but it bends aggressively between and beyond the points: e.g. $f_\Theta(0) = -12.18$ and $f_\Theta(9) = -2.89$, far outside the label range $[-0.3, 2.1]$. This high *variance* is not supported by the data, and indeed its test error is the *largest* of the three.
- **Degree 2 — relatively good.** The parabola captures the main curvature while keeping the model simple (three parameters). It attains the *smallest test RSS* (2.4768) with a still-moderate training RSS (1.8765): the best bias–variance trade-off of the three models.

Quantitative verification. Table 1 collects the training/test errors and Table 2 lists the per-point test predictions.

Model	Train RSS	Train MSE	Test RSS	Test MSE
Degree 1	4.7816	0.4782	4.1244	0.4124
Degree 2	1.8765	0.1877	**2.4768**	**0.2477**
Degree 4	**0.7712**	**0.0771**	5.1835	0.5184

Table 1: Training/test errors of the three models (4 d.p.). The smallest training error belongs to the quartic (overfitting), while the smallest test error belongs to the quadratic.

x	y	$f_1(x)$	$f_2(x)$	$f_4(x)$	e_1^2	e_2^2	e_4^2
7.2	0.6	1.0653	1.1333	1.6514	0.2165	0.2844	1.1055
1.1	1.0	0.2394	1.8294	−0.6040	0.5786	0.6879	2.5727
3.6	0.3	0.5779	0.2896	0.7699	0.0772	0.0001	0.2208
4.8	−0.4	0.7403	0.1694	−0.3381	1.3004	0.3242	0.0038
4.3	0.0	0.6726	0.1707	−0.0570	0.4524	0.0291	0.0033
6.6	0.2	0.9840	0.7418	1.0222	0.6147	0.2935	0.6760
7.8	1.8	1.1465	1.6251	1.6661	0.4270	0.0306	0.0179
5.6	0.6	0.8486	0.3123	−0.1109	0.0618	0.0828	0.5054
7.3	1.7	1.0788	1.2083	1.7122	0.3859	0.2418	0.0001
8.2	1.3	1.2007	2.0088	1.0209	0.0099	0.5024	0.0779
				Test RSS:	4.1244	2.4768	5.1835

Table 2: Test-set predictions and squared errors (f_1, f_2, f_4: degree 1/2/4; $e_k^2 = (f_k(x) - y)^2$; 4 d.p.). The degree-2 model wins on most intermediate points; the quartic's large errors occur on the two left-most test points, where its oscillation hurts.

Comparing the residual magnitudes in Table 2: the degree-1 model is uniformly mediocre (large errors everywhere), the quartic achieves very small errors on most points but blows up on points $x = 1.1$, 7.2 (squared errors 2.57, 1.11), where its oscillation hurts, while the quadratic keeps all squared errors below 0.69. The numbers confirm the qualitative picture: **degree 1 underfits, degree 4 overfits, and degree 2 is the relatively good model**.

Question 2

Write $\mathbf{X}^{(i)}$ for the i-th sample (row vector), $\bar{\mathbf{X}} = \frac{1}{m}\sum_{i=1}^{m} \mathbf{X}^{(i)}$ for the mean vector and consider

$$J(\Theta, \theta_0) = \sum_{i=1}^{m} \Big(\mathbf{X}^{(i)}\Theta + \theta_0 - Y^{(i)}\Big)^2 + \lambda\|\Theta\|^2, \qquad \lambda > 0.$$

(a) Proof that $\theta_0 = \bar{Y} - \bar{\mathbf{X}}\Theta$ minimizes J

Rewrite each squared term as

$$\mathbf{X}^{(i)}\Theta + \theta_0 - Y^{(i)} = \underbrace{\big(\mathbf{X}^{(i)} - \bar{\mathbf{X}}\big)\Theta}_{=:\, a^{(i)}} + \underbrace{\theta_0 + \bar{\mathbf{X}}\Theta}_{=:\, b} - Y^{(i)}.$$

For fixed Θ, differentiating w.r.t. θ_0 (note $\partial a^{(i)}/\partial\theta_0 = 0$ and $\partial b/\partial\theta_0 = 1$):

$$\frac{\partial J}{\partial \theta_0} = 2\sum_{i=1}^{m}\Big(a^{(i)} + b - Y^{(i)}\Big) = 2\Big(\sum_{i=1}^{m} a^{(i)} + mb - \sum_{i=1}^{m} Y^{(i)}\Big).$$

Since $\sum_{i=1}^{m}\big(\mathbf{X}^{(i)} - \bar{\mathbf{X}}\big) = \mathbf{0}$, the first sum vanishes:

$$\sum_{i=1}^{m} a^{(i)} = \Big(\sum_{i=1}^{m}\big(\mathbf{X}^{(i)} - \bar{\mathbf{X}}\big)\Big)\Theta = 0, \qquad \text{and} \qquad \sum_{i=1}^{m} Y^{(i)} = m\bar{Y},$$

hence

$$\frac{\partial J}{\partial \theta_0} = 2m\Big(\theta_0 + \bar{\mathbf{X}}\Theta - \bar{Y}\Big).$$

Setting $\partial J/\partial\theta_0 = 0$ gives

$$\theta_0 = \bar{Y} - \bar{\mathbf{X}}\Theta.$$

Moreover $\dfrac{\partial^2 J}{\partial \theta_0^2} = 2m > 0$ (the term $\lambda\|\Theta\|^2$ does not depend on θ_0), so J is a strictly convex quadratic in θ_0: this stationary point is the **unique global minimizer**. ■

(b) Substituting θ_0: $J(\Theta, \theta_0) = \tilde{J}(\Theta)$

Define the centered input $\tilde{\mathbf{X}}^{(i)} = \mathbf{X}^{(i)} - \bar{\mathbf{X}}$ and the centered label $\tilde{Y}^{(i)} = Y^{(i)} - \bar{Y}$. Substituting $\theta_0 = \bar{Y} - \bar{\mathbf{X}}\Theta$,

$$a^{(i)} + b - Y^{(i)} = \big(\mathbf{X}^{(i)} - \bar{\mathbf{X}}\big)\Theta + \bar{Y} - Y^{(i)} = \tilde{\mathbf{X}}^{(i)}\Theta - \tilde{Y}^{(i)}.$$

Therefore

$$J(\Theta, \theta_0) = \sum_{i=1}^{m}\Big(\tilde{\mathbf{X}}^{(i)}\Theta - \tilde{Y}^{(i)}\Big)^2 + \lambda\|\Theta\|^2 =: \tilde{J}(\Theta). \qquad \blacksquare$$

(c) Analytic solution of ridge regression

In matrix form, with $\tilde{\mathbf{X}}$ the matrix whose i-th row is $\tilde{\mathbf{X}}^{(i)}$ and $\tilde{\mathbf{Y}} = \big(\tilde{Y}^{(1)}, \ldots, \tilde{Y}^{(m)}\big)^\top$,

$$\tilde{J}(\Theta) = \big\|\tilde{\mathbf{X}}\Theta - \tilde{\mathbf{Y}}\big\|^2 + \lambda\|\Theta\|^2.$$

Using $\nabla_\Theta\big\|\tilde{\mathbf{X}}\Theta - \tilde{\mathbf{Y}}\big\|^2 = 2\tilde{\mathbf{X}}^\top\big(\tilde{\mathbf{X}}\Theta - \tilde{\mathbf{Y}}\big)$ and $\nabla_\Theta\|\Theta\|^2 = 2\Theta$,

$$\frac{\partial \tilde{J}}{\partial \Theta} = 2\tilde{\mathbf{X}}^\top\big(\tilde{\mathbf{X}}\Theta - \tilde{\mathbf{Y}}\big) + 2\lambda\Theta = 2\Big[\big(\tilde{\mathbf{X}}^\top\tilde{\mathbf{X}} + \lambda I\big)\Theta - \tilde{\mathbf{X}}^\top\tilde{\mathbf{Y}}\Big].$$

Setting the gradient to zero gives the *ridge normal equations*

$$\big(\tilde{\mathbf{X}}^\top\tilde{\mathbf{X}} + \lambda I\big)\Theta = \tilde{\mathbf{X}}^\top\tilde{\mathbf{Y}}.$$

Because $\lambda > 0$, for every $\mathbf{v} \neq \mathbf{0}$,

$$\mathbf{v}^\top\big(\tilde{\mathbf{X}}^\top\tilde{\mathbf{X}} + \lambda I\big)\mathbf{v} = \big\|\tilde{\mathbf{X}}\mathbf{v}\big\|^2 + \lambda\|\mathbf{v}\|^2 > 0,$$

so $\tilde{\mathbf{X}}^\top\tilde{\mathbf{X}} + \lambda I$ is positive definite and thus invertible. The analytic solution is

$$\boxed{\Theta^* = \big(\tilde{\mathbf{X}}^\top\tilde{\mathbf{X}} + \lambda I\big)^{-1}\tilde{\mathbf{X}}^\top\tilde{\mathbf{Y}}} \qquad \text{with} \qquad \theta_0^* = \bar{Y} - \bar{\mathbf{X}}\Theta^*. \qquad \blacksquare$$

Question 3

(a) Logistic model and cross-entropy error

With the linear model $f_\Theta(x_1, x_2) = \theta_0 + \theta_1 x_1 + \theta_2 x_2$ and $z = \theta_0 + \theta_1 x_1 + \theta_2 x_2$, the logistic regression model is

$$P(\hat{y} = 1 \mid x_1, x_2) = \sigma(z) = \frac{1}{1 + e^{-z}} = \frac{1}{1 + e^{-(\theta_0 + \theta_1 x_1 + \theta_2 x_2)}}, \qquad P(\hat{y} = 0 \mid x_1, x_2) = 1 - \sigma(z).$$

For the 6 training samples the cross-entropy (negative log-likelihood) error function is

$$E(\Theta) = -\sum_{i=1}^{6} \Big[y^{(i)} \ln \sigma\big(z^{(i)}\big) + \big(1 - y^{(i)}\big) \ln\big(1 - \sigma\big(z^{(i)}\big)\big) \Big], \qquad z^{(i)} = \theta_0 + \theta_1 x_1^{(i)} + \theta_2 x_2^{(i)}.$$

(b) One gradient-descent iteration

Since $\sigma'(z) = \sigma(z)\big(1 - \sigma(z)\big)$, differentiating E gives

$$\frac{\partial E}{\partial \theta_j} = \sum_{i=1}^{6} \Big(\sigma\big(z^{(i)}\big) - y^{(i)} \Big) x_j^{(i)}, \qquad x_0^{(i)} := 1.$$

With the initial parameters $(\theta_0, \theta_1, \theta_2) = (-1,\, 1.5,\, 0.5)$, Table 3 lists $z^{(i)}$ and $\sigma\big(z^{(i)}\big)$ for all training samples.

i	$x_1^{(i)}$	$x_2^{(i)}$	$y^{(i)}$	$z^{(i)}$	$\sigma\big(z^{(i)}\big)$
1	0.15	0.35	0	-0.6000	0.3543
2	0.30	0.45	0	-0.3250	0.4195
3	0.55	0.25	0	-0.0500	0.4875
4	0.30	0.80	1	-0.1500	0.4626
5	0.55	0.60	1	0.1250	0.5312
6	0.70	0.50	1	0.3000	0.5744

Table 3: Training samples with pre-activations and model outputs under the initial parameters (4 d.p.).

Writing $\sigma_i := \sigma(z^{(i)})$ and using the values of Table 3 (full precision carried in the computation):

$$\frac{\partial E}{\partial \theta_0} = \sum_i \big(\sigma_i - y^{(i)}\big) = \big(0.3543 + 0.4195 + 0.4875\big) - \big(0.5374 + 0.4688 + 0.4256\big) \approx -0.1705,$$

$$\begin{aligned} \frac{\partial E}{\partial \theta_1} &= \sum_i \big(\sigma_i - y^{(i)}\big) x_1^{(i)} = 0.3543(0.15) + 0.4195(0.30) + 0.4875(0.55) \\ &\qquad - 0.5374(0.30) - 0.4688(0.55) - 0.4256(0.70) \approx -0.2698, \end{aligned}$$

$$\begin{aligned} \frac{\partial E}{\partial \theta_2} &= \sum_i \big(\sigma_i - y^{(i)}\big) x_2^{(i)} = 0.3543(0.35) + 0.4195(0.45) + 0.4875(0.25) \\ &\qquad - 0.5374(0.80) - 0.4688(0.60) - 0.4256(0.50) \approx -0.4893. \end{aligned}$$

(The analytic gradient was verified against a central finite-difference estimate; the two agree to

within 4×10^{-10}.) The gradient-descent update $\Theta \leftarrow \Theta - \eta\,\nabla E$ with $\eta = 0.1$ gives

$$\begin{aligned}
\theta_0 &= -1 - 0.1(-0.170474) = -0.98295,\\
\theta_1 &= 1.5 - 0.1(-0.269839) = 1.52698,\\
\theta_2 &= 0.5 - 0.1(-0.489345) = 0.54893.
\end{aligned}$$

The updated logistic regression model is therefore

$$\boxed{P(\hat{y} = 1 \mid x_1, x_2) = \frac{1}{1 + e^{-(-0.98295 + 1.52698\,x_1 + 0.54893\,x_2)}}}$$

and the cross-entropy error decreases from $E = 3.6077$ to $E = 3.5756$.

(c) Test-set predictions, accuracy, precision, recall

Applying the updated model to the test set (prediction rule $\hat{y} = 1$ iff $\sigma(z) \geq 0.5$, i.e. $z \geq 0$) gives Table 4.

i	$x_1^{(i)}$	$x_2^{(i)}$	$y^{(i)}$	$z^{(i)}$	$\sigma\big(z^{(i)}\big)$	$\hat{y}^{(i)}$
1	0.28	0.82	1	−0.1053	0.4737	0
2	0.45	0.75	1	0.1159	0.5289	1
3	0.58	0.62	1	0.2430	0.5605	1
4	0.72	0.48	1	0.3800	0.5939	1
5	0.22	0.40	0	−0.4274	0.3947	0
6	0.50	0.26	0	−0.0767	0.4808	0

Table 4: Test-set predictions of the updated model.

The confusion counts are: TP = 3 (samples 2–4), FP = 0, FN = 1 (sample 1), TN = 2 (samples 5–6). Hence

$$\text{Accuracy} = \frac{TP + TN}{6} = \frac{3+2}{6} = \frac{5}{6} \approx 0.8333, \qquad \text{Precision} = \frac{TP}{TP + FP} = \frac{3}{3+0} = 1.0000,$$

$$\text{Recall} = \frac{TP}{TP + FN} = \frac{3}{3+1} = 0.7500.$$

Only sample 1 (true label 1, predicted 0 with probability $0.4737 < 0.5$) is misclassified. After a single gradient step the model already makes no false-positive errors (precision = 1), at the cost of one missed positive (recall = 0.75).

Question 4

Let $z_k = \mathbf{X}^\top \Theta_k$ for class k and write

$$P_i := P(\hat{y}_i = 1 \mid \mathbf{X}) = \frac{e^{z_i}}{\sum_{k=1}^{K} e^{z_k}}, \qquad i = 1, \ldots, K.$$

(a) Derivative of $P(\hat{y}_i = 1 \mid \mathbf{X})$ w.r.t. $\mathbf{X}^\top \Theta_j$

Case $j = i$: the numerator depends on z_i,

$$\frac{\partial P_i}{\partial z_i} = \frac{e^{z_i} \sum_k e^{z_k} - e^{z_i} e^{z_i}}{\left(\sum_k e^{z_k}\right)^2} = P_i - P_i^2 = P_i\big(1 - P_i\big).$$

Case $j \neq i$: the numerator does not contain e^{z_j},

$$\frac{\partial P_i}{\partial z_j} = \frac{0 \cdot \sum_k e^{z_k} - e^{z_i} e^{z_j}}{\left(\sum_k e^{z_k}\right)^2} = -P_i P_j.$$

In summary,

$$\frac{\partial P(\hat{y}_i = 1 \mid \mathbf{X})}{\partial\big(\mathbf{X}^\top \Theta_j\big)} = \begin{cases} P(\hat{y}_i = 1 \mid \mathbf{X})\big[1 - P(\hat{y}_i = 1 \mid \mathbf{X})\big], & j = i, \\ -P(\hat{y}_i = 1 \mid \mathbf{X})\, P(\hat{y}_j = 1 \mid \mathbf{X}), & j \neq i. \end{cases}$$

(Compactly: $P_i(\delta_{ij} - P_j)$, where δ_{ij} is the Kronecker delta.)

(b) Derivative of $P(\hat{y}_i = 1 \mid \mathbf{X})$ w.r.t. Θ_j

Since $z_j = \mathbf{X}^\top \Theta_j$, we have $\partial z_j / \partial \Theta_j = \mathbf{X}$, so by the chain rule

$$\frac{\partial P_i}{\partial \Theta_j} = \frac{\partial P_i}{\partial z_j} \cdot \frac{\partial z_j}{\partial \Theta_j} = \begin{cases} P_i\big(1 - P_i\big)\, \mathbf{X}, & j = i, \\ -P_i P_j\, \mathbf{X}, & j \neq i. \end{cases}$$

(c) Derivative of the negative log-likelihood loss

The loss is

$$E(\Theta) = -\sum_{i=1}^{K} y_i \ln P_i, \qquad P_i = P(\hat{y}_i = 1 \mid \mathbf{X}),$$

where exactly one y_i equals 1 (one-hot labels). Fix j and differentiate w.r.t. z_j:

$$\frac{\partial E}{\partial z_j} = -\sum_{i=1}^{K} \frac{y_i}{P_i} \cdot \frac{\partial P_i}{\partial z_j} = -\frac{y_j}{P_j} P_j\big(1 - P_j\big) - \sum_{i \neq j} \frac{y_i}{P_i}\big(-P_i P_j\big) = -y_j\big(1 - P_j\big) + P_j \sum_{i \neq j} y_i.$$

Because $\sum_{i=1}^{K} y_i = 1$, we have $\sum_{i \neq j} y_i = 1 - y_j$, so

$$\frac{\partial E}{\partial z_j} = -y_j + y_j P_j + P_j\big(1 - y_j\big) = P_j - y_j.$$

Finally, by the chain rule (as in part (b), with $\partial z_j / \partial \Theta_j = \mathbf{X}$):

$$\frac{\partial E(\Theta)}{\partial \Theta_j} = \frac{\partial E}{\partial z_j} \cdot \frac{\partial z_j}{\partial \Theta_j} = \big(P_j - y_j\big)\mathbf{X} = \mathbf{X}\, P(\hat{y}_j = 1 \mid \mathbf{X}) - \mathbf{X}\, y_j. \qquad \blacksquare$$