Amoxicillin Trihydrate (500mg) — Antimicrobial Therapy Guide

PulseAI Clinical Reference Library — Evidence-Based Patient Information

1. THERAPEUTIC INDICATIONS

Amoxicillin is a moderate-spectrum penicillin-class antibiotic indicated for bacterial infections including streptococcal pharyngitis, acute otitis media, sinusitis, bronchitis, and uncomplicated urinary tract infections. It is completely ineffective against viral illnesses such as the common cold or influenza.

2. IMPORTANT: COURSE COMPLETION INSTRUCTIONS

Standard dosage: 500mg orally every 8 hours for 5 to 7 days as prescribed. Swallow whole with water. CRITICAL: Complete the full prescribed course even if symptoms resolve after 48 hours. Stopping early increases the risk of bacterial relapse and antimicrobial resistance.

3. DRUG INTERACTIONS & PRECAUTIONS

Inform your doctor if you have known penicillin allergies. Amoxicillin may reduce the efficacy of oral hormonal contraceptives; barrier protection is advised during therapy. Stagger probiotic supplements at least 2 hours apart from amoxicillin.

4. ADVERSE REACTIONS & WHEN TO CALL THE CLINIC

Common side effects: Mild diarrhea, nausea, and stomach upset. Contact the clinic immediately if you develop severe watery diarrhea (>5 times daily), unexplained fever, or an itchy spreading skin rash.