Ibuprofen Oral Tablets (400mg) — Patient Information Leaflet

PulseAI Clinical Reference Library — Evidence-Based Patient Information

1. WHAT IS IBUPROFEN AND WHAT IS IT USED FOR?

Ibuprofen is a Non-Steroidal Anti-Inflammatory Drug (NSAID). It works by inhibiting cyclooxygenase (COX-1 and COX-2) enzymes, reducing prostaglandins that cause pain, fever, and inflammation. It is indicated for mild to moderate musculoskeletal pain, headache, dental pain, reduction of fever, and inflammatory joint conditions.

2. DOSAGE AND ADMINISTRATION

Adults and children over 12 years: Take one 400mg tablet every 4 to 6 hours as needed with water. Do not exceed 1200mg (3 tablets) in 24 hours. Take immediately after meals or with a full glass of milk to prevent gastrointestinal irritation. Use the lowest effective dose for the shortest necessary duration.

3. CONTRAINDICATIONS & SAFETY WARNINGS

Do not take if you have an active peptic ulcer, gastrointestinal bleeding, severe heart failure, or severe renal impairment. Avoid concurrent alcohol consumption, which elevates gastric bleeding risk. Inform your physician if you have a history of aspirin-sensitive asthma.

4. POSSIBLE ADVERSE EFFECTS & EMERGENCY SIGNS

Common mild effects include nausea, dyspepsia, and mild dizziness. Discontinue immediately and seek emergency medical care if you observe black or tarry stools, sudden chest pain, difficulty breathing, or facial/throat swelling.