



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA

Department of Physics and Astronomy "A. Righi"

Theoretical Physics

Group Theory

Lectures held by
Prof. Roberto Zucchini

Notes by
Gioele Mancino

Academic Year 2025/2026

Contents

1	Symmetry in Quantum Physics	3
1.1	Symmetry Groups in Quantum Physics	3
1.2	Dynamical Symmetry Groups	8
1.3	Lie Groups and Lie Algebras	15
1.4	Unitary Representations of Continuous Symmetries	22
1.5	Dynamical Lie Symmetries and Conservation Laws	32
2	Vector calculus in $\mathbb{R}^3$ geometry	37
2.1	Euclidean structure of $\mathbb{R}^3$	37
2.2	Orientational structure of $\mathbb{R}^3$	38
2.3	Determinant and Trace Identities	39
2.4	Direct Product	40
2.5	The Star Operator	41
2.6	The Wedge Product	43
2.7	Diagonalization Theorems	45
3	Pauli Matrices Formalism	47
3.1	Pauli Matrix Functions	48
4	Boh	51
4.1	The Lorentz group	57
5	ling lin	63
5.1	Lie Algebra of Matrix Groups	71
5.2	Exponential Map	76
	Appendices	81
A	Notation and Conventions	81
B	Computations and further developments	81

Introduction

1 | Symmetry in Quantum Physics

1.1 | Symmetry Groups in Quantum Physics

Symmetry transformations are operations carried out on the devices which prepare the states and the apparatuses which measure the observables of a quantum system having the property of not changing the results of experiments. For instance, space translations are symmetry transformations since the uniform translation of all preparations devices and measuring apparatuses does not change the results of experiments due to the homogeneity of space. Likewise, space rotations are symmetry transformations since the uniform rotation of all preparations devices and measuring apparatuses does not change the results of experiments due to the isotropy of space. In relativity, Lorentz boosts are symmetry transformations since as the result of experiments are the same in a given inertial frame and any inertial frame moving at uniform speed with respect to the first because the isotropy of spacetime.

In general, symmetry transformations gather together in families with special properties. Let G be one such family. Carrying out successively two symmetry transformations g, g' of G yields a third symmetry transformation $g' \cdot g$ of G called the **product** of g, g' . The effect of a symmetry transformation g of G can be undone by another symmetry transformation g^{-1} of G called the **inverse** of g . There exists a distinguished symmetry transformation e of G that has no effect called **neutral**. The multiplication and inversion operations and neutral element of G have the following properties:

- **Associativity:** for any three symmetry transformations g, g', g'' of G we have

$$(g'' \cdot g') \cdot g = g'' \cdot (g' \cdot g). \quad (1.1.1)$$

- **Neutrality:** there exists a symmetry transformation e of G such that for any symmetry transformation g of G we have

$$e \cdot g = g \cdot e = g. \quad (1.1.2)$$

- **Invertibility:** for any symmetry transformation g of G there exists a symmetry transformation g^{-1} of G such that

$$g^{-1} \cdot g = g \cdot g^{-1} = e. \quad (1.1.3)$$

These relations characterize G as an algebraic structure called a **group**. Since G is constituted by symmetry transformations, G is called more specifically a **symmetry group**. A group G is called **finite** if it is constituted by a finite number of elements, else it is called **infinite**. A group G is called **Abelian** if multiplication turns out to be commutative, that is

$$g \cdot g' = g' \cdot g, \quad \forall g, g' \in G. \quad (1.1.4)$$

Else, it is called non Abelian. The groups encountered in quantum physics are both finite and infinite and Abelian and non Abelian.

Example (Space Translation Symmetry). [...]

Example (Space Rotation Symmetry). [...]

1.1.1 | Symmetry Groups and Their Operator Realization

In quantum theory, states and observables of a system are represented respectively by normalized vectors defined up to a multiplicative phase in a Hilbert space $\mathcal{H}$ and selfadjoint operators in the operator algebra $L(\mathcal{H})$. The invariance of experimental results under the action of a symmetry group reduces to the invariance of expectation values: if Ψ is a state, O is an observable and g is a symmetry transformation of G , ${}^g\Psi$ is the transformed state of Ψ and gO is the transformed observable of O , one must have

$$\langle {}^g\Psi | {}^gO | {}^g\Psi \rangle = \langle \Psi | O | \Psi \rangle. \quad (1.1.5)$$

Let us now analyze the implications of the invariance of expectation values under the action of a symmetry group G with a simple example.

Example (“yes–no” observables). As is well-known, there is a class of elementary observables, the “yes–no” observables, defined by taking the value 1, 0 according to whether the system is in a certain state Φ or not. The associated operator is given by

$$O_\Phi \Psi = \Phi \langle \Phi | \Psi \rangle.$$

It is natural to assume that under the symmetry transformation g of G the observable O_Φ transforms accordingly with the state Φ as follows:

$${}^gO_\Phi = O_{{}^g\Phi}.$$

The expectation value of O_Φ in the state Ψ' is given by the transition probability from Ψ' to Φ :

$$\langle \Psi' | O_\Phi | \Psi' \rangle = |\langle \Phi | \Psi' \rangle|^2,$$

such that the **invariance of expectation values** under the action of the symmetry group G implies that

$$|\langle {}^g\Phi | {}^g\Psi' \rangle|^2 = |\langle \Phi | \Psi' \rangle|^2.$$

This means that the symmetry transformation g of G must be implemented by a unitary or antiunitary operator U_g on the Hilbert space $\mathcal{H}$ of the system such that

$$|{}^g\Psi\rangle = U_g |\Psi\rangle, \quad {}^gO = U_g O U_g^{-1}.$$

In this example we have highlighted the importance of the **Wigner’s theorem**:

Theorem 1.1 (Wigner’s theorem). *Let $\mathcal{H}$ be a Hilbert space and let G be a symmetry group of a quantum system represented on $\mathcal{H}$. Then, for each symmetry transformation g of G , there exists a unitary or antiunitary operator U_g on $\mathcal{H}$ such that*

$$\begin{cases} {}^g\Psi = U(g)\Psi, \\ {}^gO = U_g O U_g^\dagger, \end{cases} \quad (1.1.6)$$

since U_g is unitary or antiunitary, $U_g^{-1} = U_g^\dagger$, and so we can write the transformation of observables as $O \mapsto U_g O U_g^\dagger$. This implies that the expectation value of any observable O in any state Ψ is invariant under the action of the symmetry group G .

This is a fundamental result in quantum theory since it allows us to represent symmetry transformations as unitary or antiunitary operators on the Hilbert space of the system.

Recall that a unitary operator U on a Hilbert space $\mathcal{H}$ is a linear operator that preserves the Hilbert inner product,

$$\begin{aligned} U(\Phi a, \Phi' a') &= U(\Phi)a, U(\Phi')a' \\ \langle U\Phi | U\Phi' \rangle &= \langle \Phi | \Phi' \rangle, \end{aligned}$$

while an antiunitary operator A on a Hilbert space $\mathcal{H}$ is an antilinear operator that preserves the Hilbert inner product,

$$\begin{aligned} A(\Phi a, \Phi' a') &= A(\Phi)a^*, A(\Phi')a'^* \\ \langle A\Phi | A\Phi' \rangle &= \langle \Phi | \Phi' \rangle^*. \end{aligned}$$

in both cases the adjoint operator $U^\dagger$ (or $A^\dagger$) of a unitary (or antiunitary) operator U (or A) are equal to the inverse operator U^{-1} (or A^{-1}):

$$\begin{cases} \langle \Psi' | U\Psi \rangle = \langle U^\dagger \Psi' | \Psi \rangle, \\ \langle \Psi' | A\Psi \rangle = (\langle A^\dagger \Psi' | \Psi \rangle)^*, \end{cases} \implies \begin{cases} U^{-1} = U^\dagger, \\ A^{-1} = A^\dagger, \end{cases}$$

respectively for the linear and antilinear case. In what follows, we leave aside the antiunitary case, which occurs only for symmetry transformations involving time reversal, and concentrate on the unitary one. Letting the symmetry transformation g vary in the group G , we obtain a family of unitary operators $U(g)$ contained in the operator algebra $L(\mathcal{H})$.

Let us come back to relation (1.1.6)₁. We recall again that state vectors are normalized vectors defined up to a multiplicative phase factor. The content of (1.1.6)₁ is that phase choice can be adjusted so that for any symmetry transformation g the state correspondence $\Psi \mapsto^g \Psi$ is codified by a unitary operator $U(g)$. This however does not completely fix $U(g)$: we are still free to replace $U(g)$ by

$$\hat{U}(g) = \alpha(g)U(g), \tag{1.1.7}$$

where $\alpha(g)$ is a complex number of unit modulus, which is called a **phase factor**. This involves a phase redefinition

$${}^g\Psi \rightarrow \alpha(g){}^g\Psi,$$

in (1.1.6)₁, which does not change the physical content of the transformation since states are defined up to a multiplicative phase factor. By a suitable choice of the phase factors $\alpha(g)$ many relations simplify.

It is natural and possible to require

$${}^e\Psi = \Psi, \quad \forall \Psi \in \mathcal{H},$$

so that the neutral element e of the symmetry group G is represented by the identity operator

$$U(e) = \mathbb{I},$$

on the Hilbert space $\mathcal{H}$ of the system. With this choice, we can now analyze the implications of the group properties of G on the family of unitary operators $U(g)$. Considering two symmetry

transformations g, g' of G , and a physical state Ψ , we have $g'(g\Psi)$ and $g' \cdot g\Psi$ defined only up to a conventional choice of a multiplicative **phase factor** as seen above, one cannot expect that a relation of the form $g'(g\Psi) = g' \cdot g\Psi$ to hold strictly in general, but only up to some phase factor

$$\gamma(g, g', \Psi) \in \mathbb{C}, \quad |\gamma(g, g', \Psi)| = 1,$$

depending on the symmetry transformations g, g' and the state Ψ . Thus we can write

$$g'(g\Psi) = g' \cdot g\Psi \gamma(g, g', \Psi).$$

Let us then express an important result concerning the phase factor $\gamma(g, g', \Psi)$ appearing in the above relation.

Proposition 1.2. *The phase factor $\gamma(g, g', \Psi)$ can be chosen to be independent of the state Ψ , thus obtaining a relation of the form*

$$U(g \cdot g') = \gamma(g, g') U(g) U(g'), \quad (1.1.8)$$

which is called a **projective representation** of the symmetry group G . If the phase factor $\gamma(g, g')$ can be chosen to be 1 for all g, g' in G , then we have a **unitary representation** of the symmetry group G .

Proof. Proof...

□

The phase factors $\gamma(g, g')$ appearing in the projective representation of the symmetry group G are not arbitrary, but must satisfy some consistency conditions, for example the *2-cocyclicity*:

Proposition 1.3. *Given three symmetry transformations g, g', g'' of G , the following relation must hold:*

$$\gamma(g, g' \cdot g'') \gamma(g', g'') = \gamma(g \cdot g', g'') \gamma(g, g'), \quad (1.1.9)$$

where functions $\gamma : G \times G \rightarrow U(1)$ satisfying the 2-cocyclicity condition are called **2-cocycles** of G .

The 2-cocycle condition is a consequence of the associativity property of the group multiplication of G .

Proof. Proof...

□

Proposition 1.4. *The G 2-cocycle $\gamma(g, g')$ enjoys a few special properties, which albeit technical are often useful at various points:*

$$\begin{aligned} \gamma(e, g) &= \gamma(g, e) = 1, \\ \gamma(g, g^{-1}) &= \gamma(g^{-1}, g), \end{aligned} \quad (1.1.10)$$

for any symmetry transformation g of G .

The first property is a consequence of the neutrality property of the group multiplication of G , while the second one is a consequence of the invertibility property of the group multiplication of G .

Proof. Proof...

□

TODO: Complete the proof if needed

TODO: Complete the proof if needed

TODO: Complete the proof if needed

Relation (1.1.9) entails also that, for any symmetry transformation g , the operators $U(g^{-1})$ and $U(g)^{-1}$ differ by a g -dependent phase factor. Specifically, the relation between them takes the form

$$U(g^{-1}) = \varpi(g)U(g)^{-1}, \quad \varpi(g) = \gamma(g, g^{-1})^{-1}.$$

Proof. Proof...

□ **TODO:** Complete the proof if needed

In general it is not possible to absorb the phase $\gamma(g', g)$ by redefining the operators $U(g)$ into the operators $\hat{U}(g)$ given in (1.1.7) with $\alpha(g)$ a suitable phase function. By doing so $\gamma(g', g)$ gets replaced by

$$\hat{\gamma}(g, g') = \gamma(g, g')\alpha(g \cdot g')\alpha(g)^{-1}\alpha(g')^{-1}, \quad (1.1.11)$$

and in general this cannot be made equal to 1 by a judicious choice of $\alpha(g)$. A phase function $\gamma(g', g)$ of the form

$$\gamma(g, g') = \alpha(g \cdot g')\alpha(g)^{-1}\alpha(g')^{-1}, \quad (1.1.12)$$

for some function $\alpha : G \rightarrow U(1)$ is called a **2-coboundary** of G .

Proposition 1.5. *A G 2-coboundary is a G 2-cocycle, that is, any function $\gamma(g, g')$ of the form (1.1.12) satisfies the 2-cocyclicity condition (1.1.9).*

Proof. Proof...

□ **TODO:** Complete the proof if needed

The G 2-cocycle $\hat{\gamma}(g', g)$ in (1.1.11) can be rendered equal to 1 precisely when the G 2-cocycle $\gamma(g', g)$ is a G 2-coboundary of the form (1.1.12). This may or may not happen. The study of these matters is the object of a branch of algebra known as group cohomology. Some special results will be obtained below. At any rate, if $\hat{\gamma}(g', g)$ can be made equal to 1, the representation $\hat{U}(g)$ becomes genuinely unitary, so that

$$\hat{U}(g \cdot g') = \hat{U}(g)\hat{U}(g'),$$

for all symmetry transformations g, g' of G . Then the unitary family $U(g)$ forms a genuine **unitary representation** of the symmetry group G .

1.2 | Dynamical Symmetry Groups

Suppose that for a symmetry group G , the following is observed: by acting with any symmetry transformations of G on the devices that prepare the states of the system but not the apparatus that measures energy, the results of energy measurements do not change. The symmetry is then called **dynamical**, because *energy is the infinitesimal generator of time shifts*.

We note that also by acting with any symmetry transformations of G on both the devices preparing the states of the system and the apparatus measuring energy, the results of energy measurements do not change. Therefore, when G is a dynamical symmetry group, then the system's Hamiltonian H satisfies for any symmetry element g of G the relation

$$^g H = H. \quad (1.2.1)$$

If G acts on the Hilbert space through a unitary representation $U(g)$, then the invariance of the Hamiltonian under the action of G can be written as

$$U(g)HU(g)^\dagger = H, \quad \forall g \in G, \quad (1.2.2)$$

by virtue of Wigner's theorem (1.1.6)₂. This means that the Hamiltonian H commutes with all the unitary operators $U(g)$ representing the symmetry transformations of G :

$$[H, U(g)] = 0, \quad \forall g \in G.$$

This is a non trivial property severely constraining the form of H and thus providing useful structural information about the form of the interactions the system is involved in. Let us illustrate this point with a few examples taken from quantum theory.

Example (The Hydrogen Atom). The hydrogen atom is a system constituted by a proton and an electron interacting through the Coulomb potential. The Hamiltonian of the hydrogen atom is given by

$$H = \frac{\mathbf{p}^2}{2m} V(\mathbf{q}).$$

Direct observation shows that all proper rotations of the devices preparing the states of the atom leave the results of all energy measurements unchanged. Therefore, the proper rotation group $\text{SO}(3)$ is dynamical. The atom has spherical symmetry, so from (1.2.2) we have that the Hamiltonian has to remain unchanged under the action of any proper rotation $\mathbf{R}$ of $\text{SO}(3)$:

$$U(\mathbf{R})HU(\mathbf{R})^\dagger = H, \quad \forall \mathbf{R} \in \text{SO}(3).$$

It is sensible to assume that ${}^{\mathbf{R}}\mathbf{q} = \mathbf{R}^{-1}\mathbf{q}$ and ${}^{\mathbf{R}}\mathbf{p} = \mathbf{R}^{-1}\mathbf{p}$ under the action of a proper rotation. Then, we can write

$$U(\mathbf{R})\mathbf{q}U(\mathbf{R})^\dagger = \mathbf{R}^{-1}\mathbf{q}, \quad U(\mathbf{R})\mathbf{p}U(\mathbf{R})^\dagger = \mathbf{R}^{-1}\mathbf{p},$$

which means that the position and momentum operators transform as vectors under the action of $\text{SO}(3)$.¹

Using this last relation along with the invariance of the Hamiltonian under the action of $\text{SO}(3)$ implies that

$$\frac{(\mathbf{R}^{-1}\mathbf{p})^2}{2m} + V(\mathbf{R}^{-1}\mathbf{q}) = \frac{\mathbf{p}^2}{2m} + V(\mathbf{q}), \quad \forall \mathbf{R} \in \text{SO}(3).$$

¹This is a consequence of the fact that they transform according to the fundamental representation of $\text{SO}(3)$: $\mathbf{q} \mapsto \mathbf{R}^{-1}\mathbf{q}$, $\mathbf{p} \mapsto \mathbf{R}^{-1}\mathbf{p}$.

In particular, knowing that the rotations preserve the length of vectors, we have $(\mathbf{R}^{-1}\mathbf{p})^2 = \mathbf{p}^2$, so that the invariance of the Hamiltonian under the action of $\text{SO}(3)$ reduces to

$$V(\mathbf{R}^{-1}\mathbf{q}) = V(\mathbf{q}) \implies V(\mathbf{q}) = v(|\mathbf{q}|),$$

so that the potential V must be a function of the length of the position vector $v(|\mathbf{q}|)$ only. This is a non trivial constraint on the form of the potential V and thus on the form of the interactions the system is involved in; it is not sufficient to yield the Coulombic expression of the potential, but it is a necessary condition for it. The spherical symmetry of the hydrogen atom is a consequence of the dynamical symmetry of $\text{SO}(3)$.

Example (A System of Identical Particles).

Example (Conduction Electrons in a Crystal Lattice).

TODO: Complete
the example
TODO: Complete
the example

1.2.1 | Energy Eigenvalues Classification

The energy spectrum of a quantum system typically contains many energy values. The problem then arises of devising a classification method for the energy eigenvalues. There does not exist any a priori ideal classification scheme. However, symmetry provides a natural one. In this section, we show that when the system is characterized by a dynamical symmetry group G , the energy eigenvalues can be efficiently classified based on the unitary representations of G according to which the energy eigenstates transform under the action of G .

We assume that the system's Hamiltonian H has a **completely discrete spectrum** and that each energy eigenvalue has **finite degeneracy**. Let E be an energy eigenvalue of H and let $\mathcal{H}_E$ be the corresponding eigenspace, which is d_E -dimensional given a degeneracy d_E ; this subspace is spanned by all vectors Ψ which respect the eigenvalue equation

$$H\Psi = E\Psi.$$

Let now Ψ be a vector of $\mathcal{H}_E$, then for any symmetry transformation g of G we have

$$HU(g)\Psi = U(g)H\Psi = EU(g)\Psi,$$

as a direct consequence of (1.2.2). This means that $U(g)\Psi$ is also an eigenvector of H with the same eigenvalue E . Thus, the eigenspace $\mathcal{H}_E$ is invariant under the action of G . The unitary operators $U(g)$ representing the symmetry transformations of G on the Hilbert space $\mathcal{H}$ of the system restrict to unitary operators on the eigenspace $\mathcal{H}_E$, thus providing a unitary representation of G on $\mathcal{H}_E$. The energy eigenvalue E can be classified according to the unitary representation of G according to which the energy eigenstates in $\mathcal{H}_E$ transform under the action of G .

Unitary Representations on the Eigenspaces

If we now consider $\{\Phi_i\}$ an ON basis for the subspace $\mathcal{H}_E$, then these vectors respect the eigenvalue equation and the orthonormality condition

$$\begin{cases} H\Phi_i = E\Phi_i, \\ \langle \Phi_i | \Phi_j \rangle = \delta_{ij}. \end{cases}$$

Since the unitary representation of G on $\mathcal{H}_E$ is given by the restriction of the unitary representation of G on $\mathcal{H}$, we have that for any symmetry transformation g of G and any vector Φ_i of the ON

basis $\{\Phi_i\}$ of $\mathcal{H}_E$, the transformed vector $U(g)\Phi_i$ is a linear combination of the vectors of the ON basis $\{\Phi_i\}$:

$$U(g)\Phi_i = \sum_{j=1}^{d_E} \Phi_j D_{ji}(g), \quad (1.2.3)$$

where $D_{ji}(g)$ are the complex, g -dependent matrix elements of the unitary representation of G on $\mathcal{H}_E$. We aim to prove that the family of matrices $D(g)$ with elements $D_{ji}(g)$ forms a unitary representation of the symmetry group G on the eigenspace $\mathcal{H}_E$.

Proposition 1.6. *For each element g of the symmetry group G , there exists a $d_E \times d_E$ complex matrix $D(g) = (D_{ji}(g))$ which is unitary:*

$$\begin{aligned} D(g \cdot g') &= D(g)D(g'), \\ D(g)^\dagger D(g) &= D(g)D(g)^\dagger = \mathbb{I}, \end{aligned} \quad (1.2.4)$$

for all g, g' in G . The family of unitary matrices $D(g)$ forms a unitary representation of the symmetry group G on the eigenspace $\mathcal{H}_E$ corresponding to the energy eigenvalue E .

Proof. As $U(g)$ is a unitary representation of G on the Hilbert space $\mathcal{H}$ of the system, we know that for any g, g' in G we have

$$U(g \cdot g')\Phi_i = U(g)U(g')\Phi_i.$$

Since the vectors Φ_i form a basis for the eigenspace $\mathcal{H}_E$, the above equation implies that the matrix elements of $U(g)$ in this basis satisfy the same multiplication rule as the matrices $D(g)$, from (1.2.3):

$$U(g \cdot g')\Phi_i = \sum_{j=1}^{d_E} \Phi_j D_{ji}(g \cdot g'),$$

and

$$\begin{aligned} U(g)U(g')\Phi_i &= U(g) \sum_j \Phi_j D_{ji}(g') = \sum_j U(g)\Phi_j D_{ji}(g') \\ &= \sum_j \left(\sum_k \Phi_k D_{kj}(g) \right) D_{ji}(g') = \sum_k \Phi_k \left(\sum_j D_{kj}(g) D_{ji}(g') \right). \end{aligned}$$

Now if we put together the above two equations, we can prove the first relation in (1.2.4):

$$D_{ji}(g \cdot g') = \sum_k D_{kj}(g) D_{ki}(g'),$$

which is the multiplication rule for the matrices $D(g)$.

The second relation in (1.2.4) is a consequence of (1.2.3) and the orthonormality of the basis $\{\Phi_i\}$:

$$\begin{aligned} \delta_{ij} &= \langle \Phi_i | \Phi_j \rangle = \langle U(g)\Phi_i | U(g)\Phi_j \rangle \\ &= \left\langle \sum_k \Phi_k D_{ki}(g) \middle| \sum_l \Phi_l D_{lj}(g) \right\rangle = \sum_{k,l} D_{ki}(g)^* \langle \Phi_k | \Phi_l \rangle D_{lj}(g) \\ &= \sum_k D_{ik}(g)^\dagger D_{kj}(g) = (D(g)^\dagger D(g))_{ij}, \end{aligned}$$

where in the last steps we have transposed the indices and used the definition of the adjoint matrix $D(g)^\dagger$. This proves that $D(g)^\dagger D(g) = \mathbb{I}$. $\square$

Having proven (1.2.4), we know $g \mapsto D(g)$ to be a unitary representation of the symmetry group G on the eigenspace $\mathcal{H}_E$ corresponding to the energy eigenvalue E , and from (1.2.3) we know the basis vectors Φ_i to transform under the action of G through $D(g)$.

The energy eigenvalue E can thus be classified according to the unitary representation $D(g)$, where the energy eigenstates in $\mathcal{H}_E$ transform under the action of G . The problem is that there are infinitely many orthonormal bases of the energy eigenspace $\mathcal{H}_E$ of E . The representation $D(g)$ depends on the chosen basis $\{\Phi_i\}$. So $D(g)$ by itself cannot be used to distinguish E from other energy eigenvalues. Let us analyze this point in greater detail.

Proposition 1.7. *Let $\{\Phi_i^1\}$ and $\{\Phi_i^2\}$ be two ON bases for the eigenspace $\mathcal{H}_E$ corresponding to the energy eigenvalue E . Then, let the corresponding unitary representations be $D^1(g)$ and $D^2(g)$, defined by (1.2.3) with respect to the bases $\{\Phi_i^1\}$ and $\{\Phi_i^2\}$, respectively.*

There exist a $d_E \times d_E$ unitary matrix A such that its elements A_{ij} relate the two bases $\{\Phi_i^1\}$ and $\{\Phi_i^2\}$ as

$$\Phi_i^2 = \sum_{j=1}^{d_E} \Phi_j^1 A_{ji}, \quad (1.2.5)$$

and the two unitary representations $D^1(g)$ and $D^2(g)$ are related by the similarity transformation

$$D^2(g) = A^{-1} D^1(g) A, \quad (1.2.6)$$

for all g in G . Thus, the two unitary representations $D^1(g)$ and $D^2(g)$ are equivalent representations of the symmetry group G .

Proof. Using (1.2.3), we can develop (1.2.5) as

$$\begin{aligned} U(g)\Phi_i^2 &= U(g) \sum_{j=1}^{d_E} \Phi_j^1 A_{ji} = \sum_{j=1}^{d_E} U(g)\Phi_j^1 A_{ji} = \sum_{j=1}^{d_E} \left(\sum_{k=1}^{d_E} \Phi_k^1 D_{kj}^1(g) \right) A_{ji} \\ &= \sum_{k=1}^{d_E} \Phi_k^1 \left(\sum_{j=1}^{d_E} D_{kj}^1(g) A_{ji} \right) = \sum_{k=1}^{d_E} \Phi_k^1 (D^1(g)A)_{ki}, \end{aligned}$$

on one hand, and

$$\begin{aligned} U(g)\Phi_i^2 &= \sum_{k=1}^{d_E} \Phi_k^2 D_{ki}^2(g) = \sum_{k=1}^{d_E} \left(\sum_{j=1}^{d_E} \Phi_j^1 A_{jk} \right) D_{ki}^2(g) \\ &= \sum_{j=1}^{d_E} \Phi_j^1 \left(\sum_{k=1}^{d_E} A_{jk} D_{ki}^2(g) \right) = \sum_{j=1}^{d_E} \Phi_j^1 (AD^2(g))_{ji}. \end{aligned}$$

Putting together the above two equations, we have

$$(D^1(g)A)_{ki} = (AD^2(g))_{ki}, \quad \text{i.e., } D^1(g)A = AD^2(g),$$

which shows that the two unitary representations $D^1(g)$ and $D^2(g)$ are related by the similarity transformation $D^2(g) = A^{-1} D^1(g) A$. This proves that the two unitary representations $D^1(g)$ and $D^2(g)$ are equivalent representations of the symmetry group G . $\square$

Thus two unitary representations $D^1(g)$ and $D^2(g)$ of the symmetry group G of the same dimension d_E , related as in (1.2.6), for some unitary matrix A , are said to be **unitarily equivalent**. Thus

the unitary representation $D(g)$ in (1.2.3) is independent of the choice of the ON basis $\{\Phi_i\}$ for the eigenspace $\mathcal{H}_E$ up to unitary equivalence.

We can in this way characterize the energy eigenvalue E by a *unitary equivalence class of unitary representations*² of G and use this class to group E with other eigenvalues sharing the very same class. In this way, **a symmetry based classification of the energy spectrum is achieved**. Since however there are infinitely many representations of any given group, the classification scheme obtained may not be so useful. Fortunately, generic representations can be decomposed as combinations of a smaller set of more basic representations, which can then be used to refine the scheme.

Reducibility and Irreducibility

Suppose that we are able to chose the orthonormal basis $\{\Phi_i\}$ of the energy eigenspace $\mathcal{H}_E$ to be of indexed $\Phi_{\alpha r}$ by two indices α, r in such a way that (1.2.3) takes the form

$$U(g)\Phi_{\alpha r} = \sum_s \Phi_{\alpha s} D_{\alpha sr}(g),$$

for every element g of G , for certain g -dependent coefficients $D_{\alpha sr}(g)$. Therefore, the vectors $\Phi_{\alpha r}$ for fixed α mix among themselves under the action of G . This means that, for fixed g , the matrix $D(g)$ has the block diagonal form

$$D(g) = \begin{pmatrix} D_1(g) & 0 & \cdots & 0 \\ 0 & D_2(g) & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & D_a(g) \end{pmatrix},$$

and let us assume that the blocks $D_\alpha(g)$ are of dimension d_α , which is the smallest non trivial dimension for each block (i.e. the blocks cannot be further decomposed into diagonal blocks by a further change of basis $\Phi_{\alpha r}$). Then, the unitary representation $g \mapsto D(g)$ of G on $\mathcal{H}_E$ is said to be **reducible**, and its blocks $D_\alpha(g)$ are said to define an **irreducible** representations $g \mapsto D_\alpha(g)$ of G (*irreducible components*).

The energy eigenvalue E can then be classified through the irreducible representations $D_\alpha(g)$ of G . The former way of proceeding provides a classification scheme of the energy spectrum relying on a more restricted set of elements and so potentially more useful than that furnished by the latter.

We can resume the results obtained in this section as follows:

- Each energy eigenvalue E is characterized by a set of irreducible unitary representations D_α of G , unique up to unitary equivalence, describing how the eigenvectors belonging to E transform under the action of G .
- The energy eigenvalues can be grouped in subsets of eigenvalues sharing the same representation content D_α providing a classification scheme of the energy spectrum.

Let us illustrate the above results with a few examples.

Example (The One-Dimensional Harmonic Oscillator). The parity transformation P is space reflection about the origin. Observation show that the parity group $C_2 = \{\mathbb{I}, P\}$ is a dynamical symmetry group of the one-dimensional harmonic oscillator.

²It is a class of equivalence of unitary representations, where the representations are unitarily equivalent.

As well known, the energy eigenvalues of the one-dimensional harmonic oscillator are given by

$$H\Phi_n = E_n\Phi_n, \quad E_n = \hbar\omega(n + 1/2),$$

with $n = 0, 1, 2, \dots$, and are non degenerate. The energy eigenstates of the one-dimensional harmonic oscillator transform under the action of the parity transformation P as

$$U(P)\Phi_n = \Phi_n(-1)^n.$$

Thus the parity group C_2 has two irreducible unitary representations: the trivial representation $D_+(P) = 1$ and the sign representation $D_-(P) = -1$. The energy eigenvalues of the one-dimensional harmonic oscillator are classified according to these two irreducible unitary representations of C_2 : the energy eigenvalues with even n transform according to the trivial representation D_+ , while those with odd n transform according to the sign representation D_- . Thus we classify them as even or odd energy eigenvalues according to the parity of n .

Example (The Hydrogen Atom). It can be shown that the proper rotation group $\text{SO}(3)$ is a dynamical symmetry group of the hydrogen atom. The energy eigenvalues of the hydrogen atom are known to be given by

$$H\Phi_{nlm} = E_n\Phi_{nlm}, \quad E_n = -\frac{mc^2\alpha^2}{2n^2}.$$

The energy eigenvalues of the hydrogen atom are degenerate, with degeneracy $d_{E_n} = n^2$:

1. $n = 0, 1, 2, \dots$,
2. $l = 0, 1, \dots, n-1$,
3. $m = -l, -l+1, \dots, l-1, l$,

and the eigenstates Φ_{nlm} are also eigenstates of the total angular momentum operator L^2 and of the component along the quantization axis of the angular momentum operator L_z :

$$\begin{cases} L^2\Phi_{nlm} = \hbar^2 l(l+1)\Phi_{nlm}, \\ L_0\Phi_{nlm} = \hbar m\Phi_{nlm}. \end{cases}$$

Furthermore, the other spherical coordinates of the angular momentum $L_{\pm}$ act on the eigenstates Φ_{nlm} as

$$L_{\pm}\Phi_{nlm} = \hbar\sqrt{l(l+1) - m(m\pm 1)}\Phi_{nl(m\pm 1)},$$

thus acting as ladder operators on the index m .

Since $\mathbf{L}$ is the **infinitesimal generator** of the proper rotation group $\text{SO}(3)$ (which will be discussed in the next section 1.3), the energy eigenstates Φ_{nlm} transform under the action of $\text{SO}(3)$ according to the irreducible unitary representation $D_l(\mathbf{R})$ of $\text{SO}(3)$, of dimension $d_l = 2l + 1$:

$$U(\mathbf{R})\Phi_{nlm} = \sum_{m'=-l}^l \Phi_{nlm'} D_{lm'm}(\mathbf{R}),$$

so that $D_l(\mathbf{R})$ is $(2l+1) \times (2l+1)$ dimensional, and the mappings $\mathbf{R} \mapsto D_l(\mathbf{R})$ can be shown to be unitary irreducible representations of $\text{SO}(3)$. Thus for each energy eigenvalue E_n of the hydrogen atom, the energy eigenstates Φ_{nlm} transform under the action of $\text{SO}(3)$ according to the n irreducible $(2l+1)$ -dimensional unitary representation $D_{l=0,1,\dots,n-1}$ of $\text{SO}(3)$. In this way, the symmetry based classification of the energy spectrum is achieved for the hydrogen atom.

TODO: put image
of the molecule

Example (The Sulfur Trioxide Molecule SO_3). The molecule of sulfur trioxide SO_3 has the shape of a planar equilateral triangle, with the sulfur atom at the center and the three oxygen atoms at the vertices. The symmetry group of this molecule is the dihedral group D_3 , which is a subgroup of $\text{SO}(3)$.

A planar equilateral triangle has the following symmetries:

- Three rotations of 120° about the axis perpendicular to the plane of the triangle, which we call C_3 rotations:

$$0^\circ : \begin{pmatrix} 123 \\ \downarrow \\ 123 \end{pmatrix}, \quad 120^\circ : \begin{pmatrix} 123 \\ \downarrow \\ 231 \end{pmatrix}, \quad 240^\circ : \begin{pmatrix} 123 \\ \downarrow \\ 312 \end{pmatrix}.$$

- Three reflections about the axes of symmetry of the triangle, which we call σ_v reflections:

$$\sigma_v^1 : \begin{pmatrix} 123 \\ \downarrow \\ 132 \end{pmatrix}, \quad \sigma_v^2 : \begin{pmatrix} 123 \\ \downarrow \\ 321 \end{pmatrix}, \quad \sigma_v^3 : \begin{pmatrix} 123 \\ \downarrow \\ 213 \end{pmatrix}.$$

Putting together the above symmetries, we find a total of six symmetries, which form the dihedral group D_3 , which coincides with the symmetric group S_3 of permutations of three objects. The energy eigenvalues of the sulfur trioxide molecule SO_3 can be classified according to the irreducible unitary representations of D_3 .

This group has three *unitary equivalence classes of irreducible unitary representations*: the trivial representation D_+ , the sign representation D_- and a two-dimensional representation D_2 . They are of dimensions $d_+ = 1$, $d_- = 1$ and $d_2 = 2$, respectively, and we can deduce their dimensionality through the following theorem:

Theorem 1.8. *The sum of the squares of the dimensions d_α of the irreducible unitary representations D_α of a finite non-abelian group G is equal to the order $|G|$ (the number of elements) of the group:*

$$\sum_{\alpha} d_{\alpha}^2 = |G|.$$

In our case the only way to satisfy the above relation is to have $d_+^2 + d_-^2 + d_2^2 = 1^2 + 1^2 + 2^2 = 6$, which is the order of D_3 . The energy eigenvalues of the sulfur trioxide molecule SO_3 can be classified according to these three irreducible unitary representations of D_3 .

1.3 | Lie Groups and Lie Algebras

A Lie group G is an infinite group that can be parametrized in a neighborhood of the neutral element by continuous parameters θ^a , $a = 1, \dots, d$, organized as a vector $\underline{\theta}$, in one-to-one fashion:

$$\begin{cases} \underline{\theta} \in D \subseteq \mathbb{R}^d, \\ g : D \mapsto G, \\ \underline{\theta} \mapsto g(\underline{\theta}), \end{cases}$$

where D is an open neighborhood of the origin in $\mathbb{R}^d$ ($\underline{0} \in D$ and $g(\underline{0}) \in G$). The number d of parameters is called the dimension of G , seen as a differential map. The group operations of multiplication and inversion are smooth functions of the parameters.

If the parametrization is smooth, as we assume, G can be thought geometrically as a group manifold. Below, we treat G as a matrix group for definiteness.

We assume that 0 belongs to the parameter space and that

$$g(\underline{0}) = 1,$$

which can be read as a normalizing condition. By virtue of the parametrization, for any pair of parameter vectors $\underline{\theta}$, $\underline{\theta}'$, there exists a third parameter vector $\underline{\theta}''$ such that

$$g(\underline{\theta}')g(\underline{\theta}) = g(\underline{\theta}'').$$

As the parametrization is one-to-one,³ $\underline{\theta}''$ is also uniquely determined by $\underline{\theta}$, $\underline{\theta}'$. So, it can be expressed as a function of them

$$\underline{\theta}'' = \underline{m}(\underline{\theta}', \underline{\theta}),$$

which is a function of the **group multiplication**. The function $\underline{m}$ is smooth as the group multiplication is smooth. The inverse of $g(\underline{\theta})$ is given by

$$g(\underline{\theta})^{-1} = g(\underline{\theta}'),$$

for some $\underline{\theta}' = \underline{j}(\underline{\theta})$, which is unique since the parameterization is one-to-one. The function $\underline{j}$, implementing the group inversion, is smooth as the group inversion is smooth.

We have thus found two smooth functions $\underline{m}$, $\underline{j}$ that encode the group multiplication and inversion in terms of the parameters $\underline{\theta}$:

$$\begin{cases} \underline{m} : D \times D \rightarrow D, \\ \underline{j} : D \rightarrow D, \end{cases} \quad \text{such that} \quad \begin{cases} g(\underline{\theta}')g(\underline{\theta}) = g(\underline{m}(\underline{\theta}', \underline{\theta})), \\ g(\underline{j}(\underline{\theta})) = g(\underline{\theta})^{-1}. \end{cases} \quad (1.3.1)$$

The group properties of associativity, invertibility and neutrality, which are satisfied by the group multiplication and inversion, translate into properties of the functions $\underline{m}$, $\underline{j}$ as follows:

Proposition 1.9. *The following relations must hold:*

$$\begin{aligned} m(m(\underline{\theta}, \underline{\theta}'), \underline{\theta}'') &= m(\underline{\theta}, m(\underline{\theta}', \underline{\theta}'')), \\ m(\underline{\theta}, \underline{j}(\underline{\theta})) &= m(\underline{j}(\underline{\theta}), \underline{\theta}) = \underline{0}, \\ m(\underline{\theta}, \underline{0}) &= m(\underline{0}, \underline{\theta}) = \underline{\theta}. \end{aligned} \quad (1.3.2)$$

Given any choice of three parameter vectors $\underline{\theta}$, $\underline{\theta}'$, $\underline{\theta}''$. These identities constrain the form of the group multiplication function m and the inversion function j to an important extent.

³This means that each element of the group corresponds to a unique set of parameters: $g(\underline{\theta}) = g(\underline{\theta}') \implies \underline{\theta} = \underline{\theta}'$.

Proof. The first identity is a consequence of the associativity of the group multiplication: using the definition of the multiplication function m along with the associativity of the group multiplication, we find

$$\begin{aligned} g(m(m(\underline{\theta}, \underline{\theta}'), \underline{\theta}'')) &= g(m(\underline{\theta}, \underline{\theta}'))g(\underline{\theta}'') = g(\underline{\theta})g(\underline{\theta}')g(\underline{\theta}'') \\ &= g(\underline{\theta})g(m(\underline{\theta}', \underline{\theta}'')) = g(m(\underline{\theta}, m(\underline{\theta}', \underline{\theta}''))), \end{aligned}$$

which implies the first identity in (1.3.2)₁ by the one-to-one nature of the parametrization (otherwise we would have $g(\underline{\theta}) = g(\underline{\theta}')$ for some $\underline{\theta} \neq \underline{\theta}'$). The second identity is a consequence of the invertibility of the group multiplication: using the definition of the inversion function j along with the invertibility of the group multiplication, we find

$$\begin{aligned} g(m(\underline{\theta}, j(\underline{\theta}))) &= g(\underline{\theta})g(j(\underline{\theta})) = g(\underline{\theta})g(\underline{\theta})^{-1} = 1 = g(\underline{0}), \\ g(m(j(\underline{\theta}), \underline{\theta})) &= g(j(\underline{\theta}))g(\underline{\theta}) = g(\underline{\theta})^{-1}g(\underline{\theta}) = 1 = g(\underline{0}), \end{aligned}$$

proving both sides of the second identity in (1.3.2)₂. The third identity is a consequence of the neutrality of the group multiplication: using the definition of the multiplication function m along with the neutrality of the group multiplication, we find

$$g(m(\underline{0}, \underline{\theta})) = g(\underline{\theta}), \quad g(m(\underline{\theta}, \underline{0})) = g(\underline{\theta}),$$

proving both sides of the third identity in (1.3.2)₃. □

1.3.1 | Local Structure of the Group

The idea now is to understand the structure of G by looking at the behavior of $g(\underline{\theta})$ in a neighborhood of the neutral element. This is done by Taylor expanding in power series around $\underline{0}$ as:

$$g(\underline{\theta}) = g(\underline{0}) + \left. \frac{\partial g}{\partial \theta^a} \right|_{\underline{0}} \theta^a + \frac{1}{2} \left. \frac{\partial^2 g}{\partial \theta^a \partial \theta^b} \right|_{\underline{0}} \theta^a \theta^b + O(\theta^3).$$

Now computing individual terms of the expansion, we have

$$g(\underline{0}) = 1, \quad t_a = \left. \frac{\partial g}{\partial \theta^a} \right|_{\underline{0}}, \quad t_{ab} = \left. \frac{\partial^2 g}{\partial \theta^a \partial \theta^b} \right|_{\underline{0}} = t_{ba},$$

so that

$$g(\underline{\theta}) = 1 + t_a \theta^a + \frac{1}{2} t_{ab} \theta^a \theta^b + O(\theta^3). \quad (1.3.3)$$

Let us highlight the fact that the coefficients t_a, t_{ab} are matrices, as $g(\underline{\theta})$ is a matrix group. t_{ab} in particular is symmetric in its indices a, b since the order of differentiation does not matter.

We can now think of G as an **hypersurface in the space of matrices**, containing the identity matrix. In the neighborhood of the identity, there is D , subset of our hypersurface, which is diffeomorphic to an open neighborhood of the origin in $\mathbb{R}^d$ (the domain of our parametrization).

If we vary $g(0, \dots, 0, \theta^a, 0, \dots, 0)$, we get a curve in D passing through the identity (when $\theta^a = 0$). The tangent vector to this curve at the identity is given by $t_a = \left. \frac{\partial g}{\partial \theta^a} \right|_{\underline{0}}$. Varying all parameters θ^a we get d curves in D passing through the identity and their tangent vectors at the identity are given by $t_a, a = 1, \dots, d$. These tangent vectors are **linearly independent** since the parametrization is one-to-one, thus they span a vector space of dimension d .

The tangent space to G at the identity is more than a vector space of dimension d (spanned by the matrices $t_a, a = 1, \dots, d$, constituting a basis). We denote this tangent space by $\mathfrak{g}$ and call it the

Lie algebra of G . It is a vector space of dimension d and it is closed under the commutator of matrices: the Lie algebra $\mathfrak{g}$ encodes all the information about the local structure of G around the identity element, but this will be clarified in a moment.

We could now think to Taylor expand the group multiplication functions m, j around $\underline{0}$ and try to find relations among the coefficients of the expansion which encode the group structure. Starting from the multiplication function, we find that the coefficients of the expansion of m must satisfy

$$m^a(\underline{\theta}, \underline{\theta}') = \theta^a + \theta'^a + m^a_{bc} \theta^b \theta'^c + O(\theta^3), \quad (1.3.4)$$

while for the inversion property we find

$$j^a(\underline{\theta}) = -\theta^a + j^a_{bc} \theta^b \theta^c + O(\theta^3), \quad (1.3.5)$$

where m^a_{bc}, j^a_{bc} are the coefficients of the expansions. These coefficients are not independent, as it can be proven that they satisfy:

$$j^a_{bc} = m^a_{bc} + m^a_{cb}.$$

If we now recall the Taylor expansion of $g(\underline{\theta})$ around $\underline{0}$

$$g(\underline{0}) = 1 + t_a \theta^a + \frac{1}{2} t_{ab} \theta^a \theta^b + O(\theta^3),$$

we can note how the coefficients t_a define another member of the Lie algebra $\mathfrak{g}$: the Lie algebra $\mathfrak{g}$ is the tangent space to G at the identity, so it is spanned by the matrices $t_a, a = 1, \dots, d$, which we have said to be closed under the commutator of matrices. Let us now define the commutator of two basis elements t_a, t_b of $\mathfrak{g}$ as follows:

Definition 1.1 (Lie Bracket). The **commutator**, or **Lie bracket**, of two basis elements t_a, t_b of $\mathfrak{g}$ is defined as follows:

$$[t_a, t_b] = t_a t_b - t_b t_a = f_{ab}^c t_c \in \mathfrak{g},$$

where f_{ab}^c are the **structure constants** of the Lie algebra $\mathfrak{g}$, encoding the properties of the Lie algebra $\mathfrak{g}$ and thus of the local structure of the Lie group G around the identity element.

Having defined the Lie bracket of two basis elements t_a, t_b of $\mathfrak{g}$, we can now prove the following proposition:

Proposition 1.10. *The structure constants are antisymmetric in their lower indices a, b since the commutator is antisymmetric. These structure constants are related to the coefficients m^a_{bc} of the expansion of the group multiplication function by*

$$f_{bc}^a = m^a_{bc} - m^a_{cb}.$$

If we change parametrization, the structure constants change accordingly, but the Lie algebra $\mathfrak{g}$ remains the same.

Proof. Taking again the definition of the multiplication function

$$g(\underline{\theta}')g(\underline{\theta}) = g(m(\underline{\theta}', \underline{\theta})),$$

we can Taylor expand both sides around $\underline{0}$ and find the following relation between the coefficients of the expansion of g and the structure constants:

$$\begin{aligned} & \left[1 + \theta^a t_a + \frac{1}{2} \theta^a \theta^b t_{ab} + O(\theta^3) \right] \left[1 + \theta'^a t_a + \frac{1}{2} \theta'^a \theta'^b t_{ab} + O(\theta'^3) \right] \\ &= 1 + (\theta^a + \theta'^a + m^a_{bc} \theta^b \theta'^c) t_a + \frac{1}{2} (\theta^a + \theta'^a) (\theta^b + \theta'^b) t_{ab} + O(\theta^2, \theta') + O(\theta, \theta'^2). \end{aligned}$$

Relating the coefficients of the expansion of both sides in $\theta^a \theta'^b$, we find the following relations (renaming summation indices):

$$t_a t_b = t_{ab} + m_{ab}^c t_c,$$

$$t_b t_a = t_{ba} + m_{ba}^c t_c,$$

but since $t_{ab} = t_{ba}$, if we subtract the second from the first, we find

$$[t_a, t_b] = (m_{ab}^c - m_{ba}^c) t_c = f_{ab}^c t_c,$$

so that the structure constants f_{ab}^c of the Lie algebra $\mathfrak{g}$ are related to the coefficients m_{ab}^c of the expansion of the group multiplication function m by

$$f_{ab}^c = m_{ab}^c - m_{ba}^c.$$

□

Thus to sum up:

- The Lie algebra $\mathfrak{g}$ of a Lie group G is the tangent space to G at the identity element, spanned by the matrices t_a , $a = 1, \dots, d$.
- The basis elements t_a of the Lie algebra $\mathfrak{g}$ satisfy the commutation relations

$$[t_a, t_b] = f_{ab}^c t_c,$$

where f_{ab}^c are the structure constants of the Lie algebra $\mathfrak{g}$.

- The structure constants f_{ab}^c of the Lie algebra $\mathfrak{g}$ are related to the coefficients m_{ab}^c of the expansion of the group multiplication function m by

$$f_{ab}^c = m_{ab}^c - m_{ba}^c.$$

The Lie algebra, the tangent space to the Lie group at the identity, is not only a vector space, but also an algebra, since it is closed under the commutator of matrices. It encodes all the information about the local structure of the Lie group around the identity element. In particular, the structure constants f_{ab}^c determine the commutation relations of the basis elements t_a of the Lie algebra, which in turn determine the local structure of the Lie group G around the identity element.

The commutator of the basis elements t_a of the Lie algebra $\mathfrak{g}$ has some properties, which are the **Jacobi identity** and the **antisymmetry** of the commutator:

$$\begin{cases} [t_a, t_b] + [t_b, t_a] = 0, \\ [t_a, [t_b, t_c]] + [t_b, [t_c, t_a]] + [t_c, [t_a, t_b]] = 0. \end{cases}$$

These two properties can be expressed in terms of the structure constants f_{ab}^c as follows:

$$\begin{cases} f_{ab}^c + f_{ba}^c = 0, \\ f_{ad}^e f_{bc}^d + f_{bd}^e f_{ca}^d + f_{cd}^e f_{ab}^d = 0, \end{cases}$$

which are the **antisymmetry** of the structure constants in their lower indices and the **Jacobi identity** for the structure constants, respectively. This means that not every set of numbers f_{ab}^c can be the structure constants of a Lie algebra, but only those that satisfy the antisymmetry and Jacobi identity properties.

1.3.2 | Exponential Parametrization of Lie Groups

Given the parametrization of a Lie group G in a neighborhood of the identity element

$$g : D \mapsto G, \quad \underline{\theta} \in D \mapsto g(\underline{\theta}) \in G,$$

is it possible to find a global parametrization of G in terms of the elements of its Lie algebra $\mathfrak{g}$? The answer is yes, and the way to do it is through the *exponential map*.

If we consider another parametrization of G in a neighborhood of the identity element defined by

$$\tilde{g}(\underline{\theta}) = \lim_{N \rightarrow \infty} g\left(\frac{\underline{\theta}}{N}\right)^N,$$

which if the group is suitably closed under the group multiplication, is well defined and provides a global parametrization of G in terms of the elements of its Lie algebra $\mathfrak{g}$:

$$\tilde{g}(\underline{\theta}) = e^{\theta^a t_a}.$$

This new parametrization is called the **exponential parametrization** of G and the map $\tilde{g} : \mathbb{R}^d \rightarrow G$ is called the **exponential map**.

Proof. Let us show that the limit defining $\tilde{g}(\underline{\theta})$ exists and is equal to $e^{\theta^a t_a}$. Starting from the definition of $\tilde{g}(\underline{\theta})$, we can Taylor expand $g(\underline{\theta}/N)$ around $\underline{0}$ as follows:

$$g\left(\frac{\underline{\theta}}{N}\right)^N = \left[1 + \frac{t_a \theta^a}{N} + \frac{1}{2} \left(\frac{t_a \theta^a}{N}\right)^2 + O\left(\frac{1}{N^3}\right)\right]^N \sim \left(1 + \frac{t_a \theta^a}{N}\right)^N \xrightarrow{N \rightarrow \infty} e^{t_a \theta^a},$$

which is the exponential of a matrix, defined by Euler's formula as the power series expansion

$$e^x = \lim_{n \rightarrow \infty} \left(1 + \frac{x}{n}\right)^n = \sum_{k=0}^{\infty} \frac{x^k}{k!}.$$

□

Thus, the exponential map provides a global parametrization of the Lie group G in terms of the elements of its Lie algebra $\mathfrak{g}$ as follows:

$$\tilde{g}(\underline{\theta}) = e^{t_a \theta^a} = 1 + \tilde{t}_a \theta^a + \frac{1}{2} \theta^a \theta^b \tilde{t}_{ab} + O(\theta^3),$$

which is the same expansion did in (1.3.3). But when comparing the coefficients to the ones of the expansion of $e^{i\theta^a t_a}$, we get the following relations:

$$\tilde{t}_a = t_a, \quad \tilde{t}_{ab} = \frac{1}{2} t_a t_b + t_b t_a.$$

The same goes for the group multiplication function $\underline{m}$ and the inversion function $\underline{j}$. We can define new functions $\tilde{\underline{m}}, \tilde{\underline{j}}$ as

$$\begin{aligned} \tilde{g}(\tilde{\underline{m}}(\underline{\theta}, \underline{\theta}')) &= \tilde{g}(\underline{\theta}') \tilde{g}(\underline{\theta}), \\ \tilde{g}(\tilde{\underline{j}}(\underline{\theta})) &= \tilde{g}(\underline{\theta})^{-1}, \end{aligned}$$

which can be thought as the group multiplication and inversion functions in the exponential parametrization. If we now Taylor expand these new functions around $\underline{0}$ we find that for the

inversion function it is as the previous expansion (1.3.5), but all the orders higher than the first vanish:

$$\begin{aligned}\tilde{j}(\underline{\theta}) &= -\theta^a, \quad \tilde{j}_{bc}^a = 0 \\ e^{\underline{\theta}^a t_a} &= \tilde{g}(\tilde{j}(\underline{\theta})) = \tilde{g}(\underline{\theta})^{-1} = e^{-t_a \theta^a}.\end{aligned}$$

For the multiplication function we have instead the **Campbell-Baker-Hausdorff function** so that the expansion of the multiplication function in the exponential parametrization is given by

$$\tilde{m}^a(\underline{\theta}, \underline{\theta}') = \theta^a + \theta'^a + \tilde{m}_{bc}^a \theta^b \theta'^c + O(\theta^3),$$

which is the same of (1.3.4) but with different coefficients $\tilde{m}_{bc}^a$ of the expansion, which are related to the structure constants f_{bc}^a of the Lie algebra $\mathfrak{g}$ by

$$\tilde{m}_{bc}^a = \frac{1}{2} f_{bc}^a.$$

These new structure constants are not different from the original ones, since the exponential map is a reparametrization of the Lie group G in terms of the elements of its Lie algebra $\mathfrak{g}$: $\tilde{t}_a = t_a$ implies that

$$f_{ab}^c t_c = [t_a, t_b] = [\tilde{t}_a, \tilde{t}_b] = \tilde{f}_{ab}^c \tilde{t}_c = \tilde{f}_{ab}^c t_c,$$

thus the structure constants of the Lie algebra $\mathfrak{g}$ are the same in both parametrizations, as expected since the Lie algebra is an intrinsic property of the Lie group G and does not depend on the choice of parametrization:

$$f_{ab}^c = \tilde{f}_{ab}^c.$$

Thus we have

$$\tilde{m}_{bc}^a = \frac{1}{2} \tilde{f}_{bc}^a = \frac{1}{2} f_{bc}^a,$$

and if we expand the CBH function in power series we find⁴

$$\tilde{m}^a(\theta, \theta') = \theta^a + \theta'^a + \frac{1}{2} f_{bc}^a \theta^b \theta'^c + O(\theta^3, \theta'^3),$$

which is like a polynomial expansion of the group multiplication function m in terms of the structure constants f_{bc}^a of the Lie algebra $\mathfrak{g}$. The exponential map thus provides a global parametrization of the Lie group G in terms of the elements of its Lie algebra $\mathfrak{g}$ and allows us to express the group multiplication function m in terms of the structure constants f_{bc}^a of the Lie algebra $\mathfrak{g}$.

Example (Phase Group U(1)). Let us define the group of phase transformations U(1) as the group of complex numbers of unit modulus under multiplication:

$$U(1) = \{z \in \mathbb{C} : |z| = 1\} = \{e^{i\theta} : \theta \in \mathbb{R}\}.$$

It is an abelian Lie group of dimension 1 which represents the Gauge group of QED. The elements of U(1) can be represented by an angle θ with an exponential parametrization as follows:

$$\tilde{g}(\theta) = e^{i\theta^a t_a} = e^{i\theta},$$

since the Lie algebra of U(1) is one-dimensional and spanned by the identity matrix $t = 1$. The structure constants of the Lie algebra of U(1) are zero since the commutator of any two elements of the Lie algebra is zero:

$$[t, t] = 0 \implies f_{ab}^c = 0.$$

⁴There exists a closed form expression for the CBH function, but it is not important for our purposes: remind only that exact complicated expressions for computing all the terms exist, but we are only interested in the first few terms of the expansion.

Example (Special Unitary Group $SU(2)$). The special unitary group $SU(2)$ is the group of 2×2 unitary matrices with determinant 1:

$$SU(2) = \{U \in \mathbb{C}^{2 \times 2} : U^\dagger U = \mathbb{I}_2, \det(U) = 1\}.$$

It is a non-abelian Lie group of dimension 3: $\underline{\theta} \in \mathbb{R}^3$. The elements of $SU(2)$ can be parametrized using the Pauli matrices σ_i as follows:

$$\tilde{g}(\theta) = e^{i\theta^i \frac{\tau(\theta)}{2}},$$

where

$$\tau(\theta) = \begin{pmatrix} \theta^3 & \theta^1 - i\theta^2 \\ \theta^1 + i\theta^2 & -\theta^3 \end{pmatrix} = \theta^1 \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + \theta^2 \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} + \theta^3 \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} = \theta^i \sigma_i,$$

Thus $t_a = \frac{1}{2}\tau_a$ and the structure constants of the Lie algebra $\mathfrak{su}(2)$ are given by the Levi-Civita symbol ϵ_{ijk} :

$$[\sigma_i, \sigma_j] = 2i\epsilon_{ijk}\sigma_k \implies f_{ij}{}^k = 2\epsilon_{ijk}.$$

Example (Special Orthogonal Group $SO(3)$). The special orthogonal group $SO(3)$ is the group of 3×3 real orthogonal matrices with determinant 1:

$$SO(3) = \{R \in \mathbb{R}^{3 \times 3} : R^T R = \mathbb{I}_3, \det(R) = 1\}.$$

It is a non-abelian Lie group of dimension 3. The elements of $SO(3)$ can be parametrized using the generators of rotations J_i as follows:

$$\tilde{g}(\theta) = e^{i\theta^i J_i},$$

where the generators J_i satisfy the commutation relations

$$[J_i, J_j] = i\epsilon_{ijk}J_k.$$

[....-] to be finished

1.4 | Unitary Representations of Continuous Symmetries

If the symmetry group G is a Lie group, then the quantum unitary operators representing the group action on states and observables can be expressed as functions of the parameter vector $\underline{\theta}$ of G :

[...]

$$U(m(\theta, \theta')) = U(g(m(\theta, \theta'))) = U(g(\theta)g(\theta')) = \gamma(g(\theta), g(\theta'))U(g(\theta))U(g(\theta')),$$

where $\gamma(g(\theta), g(\theta'))$ is a phase factor which can be written as $\gamma(g(\theta), g(\theta')) = e^{i\omega(g(\theta), g(\theta'))}$ for some real function ω . Thus

$$U(m(\theta, \theta')) = e^{i\omega(g(\theta), g(\theta'))}U(g(\theta))U(g(\theta')).$$

Example (EXERCISE).

??? photo

When you have a Lie Group and a projected unitary representation in the Hilbert space of states, you can always get to this relation with the phase factor γ by redefining the unitary operators $U(g)$ with a suitable phase factor, all given to the properties of the Lie group G and the properties of the projective representation U .

Since the Lie groups act homogeneously, we can always translate what happens at the identity element to any other element of the group. Thus, by looking at the properties of the group around the identity element, we can understand the properties of the whole group. By Taylor expanding $U(\underline{\theta})$ near the neutral element $\underline{0}$ of the group, we get

$$U(\underline{\theta}) = 1 - i\theta^a Q_a + \frac{1}{2}\theta^a \theta^b Q_{ab} + O(\theta^3), \quad Q_{ab} = Q_{ba},$$

where Q_a are the generators of the representation U of the Lie group G . The generators Q_a are essentially the tangent vectors to the representation U of the Lie group G at the identity element

$$Q_a = i \frac{\partial U(\underline{\theta})}{\partial \theta^a} \Big|_{\underline{\theta}=\underline{0}}.$$

We are now going to prove that the generators Q_a of the representation U of the Lie group G are self-adjoint operators:

$$Q_a^\dagger = Q_a.$$

We can start from the adjoint

$$\begin{aligned} Q_a^\dagger &= \left(i \frac{\partial U(\underline{\theta})}{\partial \theta^a} \Big|_{\underline{\theta}=\underline{0}} \right)^\dagger \\ &= -i \frac{\partial U^\dagger(\underline{\theta})}{\partial \theta^a} \Big|_{\underline{\theta}=\underline{0}} \\ &= -i \frac{\partial U^{-1}(\underline{\theta})}{\partial \theta^a} \Big|_{\underline{\theta}=\underline{0}} \\ &= +i U(\underline{\theta})^{-1} \frac{\partial U(\underline{\theta})}{\partial \theta^a} U(\underline{\theta})^{-1} \Big|_{\underline{\theta}=\underline{0}} = Q_a, \end{aligned}$$

where we have used the unitarity of U and the fact that $U(\underline{0}) = 1$. [comments]

We have another property to prove: we have to normalize the phase function ω

$$\omega(\underline{\theta}, \underline{0}) = \omega(\underline{0}, \underline{\theta}) = 0,$$

so that we remember that

$$\omega(\underline{\theta}, \underline{\theta}') = \theta^a \theta'^b \omega_{ab} + O(\theta^3),$$

so that the phase factor γ is normalized to 1 at the identity element of the group, since

$$e^{i\omega(\underline{0}, \underline{\theta})} = \gamma(g(\underline{0}), g(\underline{\theta}')) = 1 = e^{i\omega(\underline{\theta}, \underline{0})} = \gamma(g(\underline{\theta}), g(\underline{0})) = 1.$$

???

[...]

$$[Q_a, Q_b]$$

... as we did for $g(\theta)g(\theta')$ we can multiply the taylor expansion of $U(\theta)$ and $U(\theta')$ and compare the result with the taylor expansion of $U(m(\theta, \theta'))$ to get

$$U(\underline{\theta})U(\underline{\theta}') = e^{-i\omega(\underline{\theta}, \underline{\theta}')} U(m(\underline{\theta}, \underline{\theta}')).$$

If we insert the taylor exapnsions of

- $U(\underline{\theta}) = 1 - i\theta^a Q_a + \frac{1}{2}\theta^a \theta^b Q_{ab} + O(\theta^3)$
- $\omega(\underline{\theta}, \underline{\theta}') = \theta^a \theta'^b \omega_{ab} + O(\theta^3)$
- $m(\underline{\theta}, \underline{\theta}') = \underline{\theta} + \underline{\theta}' + \frac{1}{2}f_{ab}^c \theta^a \theta'^b + O(\theta^3)$

we get a mess, but if we compute everything and compare the coefficients of the terms of order $\theta^a \theta'^b$ we get

$$Q_a Q_b = i m_{ab}^c Q_c + i \omega_{ab} \mathbb{I} - Q_{ab},$$

and since we wanted to compute the commutator $[Q_a, Q_b]$ we can compute

$$Q_b Q_a = i m_{ba}^c Q_c + i \omega_{ba} \mathbb{I} - Q_{ba},$$

so that subtracting the two we get

$$[Q_a, Q_b] = i(m_{ab}^c - m_{ba}^c)Q_c + i(\omega_{ab} - \omega_{ba})\mathbb{I} = i f_{ab}^c Q_c + i \kappa_{ab} \mathbb{I},$$

where we have recalled the structure constants of the Lie algebra $\mathfrak{g}$ of the Lie group G and we have defined $\kappa_{ab} = \omega_{ab} - \omega_{ba}$. This last coefficients κ_{ab} are called the **central charges** of the representation U of the Lie group G . A central algebra is an algebra where the central charges are zero, so that the commutator of the generators Q_a of the representation U of the Lie group G is just

$$[Q_a, Q_b] = i f_{ab}^c Q_c.$$

The properties of the commutators of the generators Q_a of the representation U can be expressed as

- $[Q_a, Q_b] = i f_{ab}^c Q_c + i \kappa_{ab} \mathbb{I}$
- $\kappa_{ab} = \omega_{ab} - \omega_{ba}$

- $[Q_a, Q_b] + [Q_b, Q_a] = 0 \implies \kappa_{ab} + \kappa_{ba} = 0$
- $[Q_a, [Q_b, Q_c]] + [Q_b, [Q_c, Q_a]] + [Q_c, [Q_a, Q_b]] = 0$ (Jacobi identity)

The structure constants respects those identity in the following form

- $f_{bc}^a + f_{cb}^a = 0$
- $f_{ab}^d f_{dc}^e + f_{bc}^d f_{da}^e + f_{ca}^d f_{db}^e = 0$ (Jacobi identity)

and along with the central charges they respect the following identity

$$\kappa_{ad} f_{bc}^d + \kappa_{bd} f_{ca}^d + \kappa_{cd} f_{ab}^d = 0,$$

which is a consequence of the Jacobi identity for the generators Q_a of the representation U of the Lie group G . A set of coefficients f_{ab}^c and κ_{ab} which respect the above identities is called a **Lie algebra $\mathfrak{g}$ 2-cocycle**. This naming is consistent since the central charges κ_{ab} are related to the phase factor γ of the projective representation U of the Lie group G by the relation $\kappa_{ab} = \omega_{ab} - \omega_{ba}$.

...

$$\begin{aligned} U(g, g') &= \gamma(g, g') U(g) U(g'), \\ U(g) &\rightarrow \hat{U}(g) = \alpha(g) U(g), \\ \gamma(g, g') &= \alpha(g, g') \alpha(g)^{-1} \alpha(g')^{-1}, \end{aligned}$$

so that we can redefine the unitary operators $U(g)$ of the projective representation U of the Lie group G by a suitable phase factor $\alpha(g)$ to get a new projective representation $\hat{U}$ of the Lie group G with a new phase factor $\hat{\gamma}$ which is related to the old one by the relation

$$\hat{\gamma}(g, g') = \alpha(g, g') \alpha(g)^{-1} \alpha(g')^{-1} \gamma(g, g'),$$

so that if we can find a suitable phase factor $\alpha(g)$ such that $\hat{\gamma}(g, g') = 1$ for all $g, g' \in G$, then we can redefine the unitary operators $U(g)$ of the projective representation U of the Lie group G by a suitable phase factor $\alpha(g)$ to get a new projective representation $\hat{U}$ of the Lie group G with a trivial phase factor $\hat{\gamma}$, so that the new projective representation $\hat{U}$ of the Lie group G is actually a linear representation of the Lie group G . However, in general, it is not always possible to find such a phase factor $\alpha(g)$, so that there are some projective representations of the Lie group G which cannot be redefined to get a linear representation of the Lie group G .

???

$$Q_a \rightarrow \hat{Q}_a = Q_a + k_a \mathbb{I},$$

where k_a are the central shifts of the generators Q_a of the representation U of the Lie group G . These shifts are related to the central charges κ_{ab} by the relation

$$\kappa_{ab} = f_{ab}^c k_c,$$

which defines a **2-coboundary** Lie algebra. A 2-coboundary Lie algebra is special case of a 2-cocycle Lie algebra, where the central charges κ_{ab} can be expressed as a linear combination of the structure constants f_{ab}^c with coefficients given by the central shifts k_a . This is **Lie Algebra Cohomography**

1.4.1 | Exponential Parametrization

Starting from the exponential parametrization of the Lie group G

$$\tilde{g}(\underline{\theta}) = \lim_{n \rightarrow \infty} g\left(\frac{\underline{\theta}}{n}\right)^n = e^{\theta^a T_a},$$

while for the inverse and the multiplication we have

$$\begin{cases} \tilde{j}(\underline{\theta}) = -\theta^a, \\ \tilde{m}(\underline{\theta}, \underline{\theta}') = \underline{\theta} + \underline{\theta}' + \frac{1}{2} f_{ab}^c \theta^a \theta'^b + O(\theta^3), \end{cases}$$

where the last function is the Baker-Campbell-Hausdorff formula (BCH) for the multiplication of the Lie group G in the exponential parametrization.

We can now use this exponential parametrization of the Lie group G to get an exponential parametrization of the projective representation U of the Lie group G :

$$\tilde{U}(g(\underline{\theta})) = U(\tilde{g}(\underline{\theta})) = U(e^{\theta^a T_a}) = 1 - i\theta^a Q_a + \frac{1}{2} \theta^a \theta^b Q_{ab} + O(\theta^3).$$

For the phase function ω we have

$$\tilde{\omega}(\underline{\theta}, \underline{\theta}') = \tilde{\omega}_{ab} \theta^a \theta'^b + O(\theta^3).$$

At the linear level there are no differences between g and $\tilde{g}$, they only manifest from the second order in the parameters θ^a . Thus we can derive

$$Q_a = \tilde{Q}_a,$$

and for the phase function ω we have

$$\tilde{Q}_{ab} = \frac{1}{2} ((\omega_{ab} + \omega_{ba}) \mathbb{I} - Q_a Q_b - Q_b Q_a).$$

Now we are interested in knowing if the following is true:

$$\tilde{U}(\underline{\theta}) = \exp(i\theta^a Q_a),$$

which is not the case: this is a projected representation, so this will be proven to be true only up to a phase factor. We can start from the exponential parametrization of the projective representation U of the Lie group G

$$\tilde{U}(\underline{\theta}) = U(\tilde{g}(\underline{\theta})) = \lim_{n \rightarrow \infty} U(g(\frac{\underline{\theta}}{n}))^n = \lim_{n \rightarrow \infty} U((\frac{1}{n^2})^n) \neq \lim_{n \rightarrow \infty} U((\frac{1}{n^2}))^n,$$

where we highlighted the difference between this projected and a true representation. Computing further we get

$$\tilde{U}(\underline{\theta}) = \exp\left(i \sum_k^n \omega\left(\frac{\underline{\theta}}{n}, \frac{k\underline{\theta}}{n}\right)\right) U\left(\frac{\underline{\theta}}{n}\right)^n,$$

where the last term, as we take the limit $n \rightarrow \infty$, becomes

$$\lim_{n \rightarrow \infty} U\left(\frac{\underline{\theta}}{n}\right)^n = (1 - i\frac{\theta^a}{n} Q_a)^n = e^{-i\theta^a Q_a},$$

so that the exponential parametrization of the projective representation U of the Lie group G is given by

$$\tilde{U}(\underline{\theta}) = e^{i\alpha(\underline{\theta})} e^{-i\theta^a Q_a}.$$

If we now want to get

$$e^{i\alpha(\underline{\theta}) - i\theta^a k_a} \tilde{U}(\underline{\theta}) = e^{-i\theta^a \hat{Q}_a},$$

where $\hat{Q}_a = Q_a + k_a \mathbb{I}$ are the shifted generators of the representation U of the Lie group G : these operates in the same way since they differ only by a phase factor, but they have different central charges.

Now we found a true representation

- $\hat{U}(\tilde{j}(\underline{\theta})) = \hat{U}(\underline{\theta})^{-1}$
- $\hat{U}(\tilde{m}(\underline{\theta}, \underline{\theta}')) = \hat{U}(\underline{\theta}) \hat{U}(\underline{\theta}')$

The term $\exp(-i\theta^a k_a)$ is an important **phase factor**, since it is the one which allows us to get a true representation from a projective representation by shifting the generators Q_a of the representation U of the Lie group G by a suitable central shift k_a . This is the reason why we introduced the concept of 2-coboundary Lie algebra, since it allows us to understand when a projective representation of a Lie group can be redefined to get a true representation of the Lie group.

This is a **multi-valued function** on the underlying group manifold G , since the phase factor usually is not an integer which comes back to the original value after a loop around the group manifold G (it is not guaranteed that the passage is single-valued): the exponential parametrization of the projective representation U of the Lie group G is a multi-valued function on the underlying group manifold G . Usually then we introduce a branch cut in the complex plane to make it single-valued, in order to treat multi valued function as single valued function on the more complex Riemann surface (such as for the square root and the logarithm).

$$\ln z = \ln |z| + i \arg z + 2\pi i n, \quad n \in \mathbb{Z}.$$

In our case the exponential parametrization of the projective representation U of the Lie group G is a multi-valued function on the underlying group manifold G , but can be made single-valued on a infinite Riemann surface which is a covering space of the underlying group manifold G : a covering group $\tilde{G}$. It is the same idea of the Riemann surface, but with more subtleties and complications, since the group structure has to be preserved.

There can be made analogies with the various covering groups we can find in physics, such as the universal covering group of the Lorentz group, which is the Spin group, and the universal covering group of the rotation group $SO(3)$, which is the special unitary group $SU(2)$. The projective representations of the Lorentz group and the rotation group $SO(3)$ can be redefined to get true representations of their respective covering groups, which are the Spin group and the special unitary group $SU(2)$. This is a consequence of the fact that the central charges of the projective representations of the Lorentz group and the rotation group $SO(3)$ can be expressed as linear combinations of the structure constants of their respective Lie algebras with coefficients given by suitable central shifts

$$\mathrm{SL}(2, \mathbb{C}) = \widetilde{\mathrm{SO}^+(1, 3)}, \quad \mathrm{SU}(2) = \widetilde{\mathrm{SO}(3)}.$$

Example (The translation group $T(3)$). The translation group $T(3)$ implements the translation symmetry of the three-dimensional space:

$$\mathbf{a} \in \mathbb{E}_3, \quad T(\mathbf{x}) = \mathbf{x} + \mathbf{a},$$

where $T \in T(3)$ and $\mathbb{E}_3$ is the three-dimensional Euclidean space.

The proper translation group $T(3)$ can be represented by the unitary operators $U(\mathbf{a})$ for a translation $\mathbf{a} \in \mathbb{E}_3$ which act on the Hilbert space of states of a quantum system as

$$(U(\mathbf{a})\Psi)(\mathbf{x}) = \Psi(\mathbf{x} + \mathbf{a}),$$

where Ψ is a state of the quantum system and we want to prove that $\{U(\mathbf{a})\}_{\mathbf{a} \in \mathbb{E}_3}$ yields a unitary representation of the proper translation group $T(3)$.

Proof. We can start by computing the composition of two translations $\mathbf{a}$ and $\mathbf{b}$: [...]

There are no multiplicative phases in the right hand side. The operator family $U(\mathbf{a})$ constitutes in this way a unitary representation of the proper translation group $T(3)$. $\square$

Let us now discuss the infinitesimal generators of the translation group $T(3)$. The infinitesimal generators $\hbar^{-1}\mathbf{p}$ of the unitary representation U of the proper translation group $T(3)$ are defined through the expansion of the unitary operators $U(\mathbf{a})$ for small translations $\delta\mathbf{a}$:

$$U(\delta\mathbf{a}) = 1 + \frac{i}{\hbar}\mathbf{p} \cdot \delta\mathbf{a} + O(\delta a^2),$$

with $\mathbf{p} = (p_1, p_2, p_3)$ being the momentum operator of the quantum system

$$\mathbf{p}\Psi(\mathbf{x}) = -i\hbar\nabla\Psi(\mathbf{x}).$$

Thus we can study the expansion of the unitary operators $U(\mathbf{a})$ acting on a state Ψ for small translations $\delta\mathbf{a}$:

$$\begin{aligned} (U(\delta\mathbf{a})\Psi)(\mathbf{x}) &= \Psi(\mathbf{x} + \delta\mathbf{a}) \\ &= \Psi(\mathbf{x}) + \delta\mathbf{a} \cdot \nabla\Psi(\mathbf{x}) + O(\delta a^2) \\ &= \dots \end{aligned}$$

$\mathbf{p}$ is therefore the familiar quantum linear momentum operator as anticipated by the notation used. The result obtained is in line with the standard interpretation of momentum as the infinitesimal generator of configuration space translations.

As well-known from quantum theory, its components

$$p_k = \mathbf{p} \cdot \mathbf{e}_k, \quad k = 1, 2, 3,$$

satisfy the commutation relations

$$[p_k, p_l] = 0, \quad k, l = 1, 2, 3,$$

free of central charges, as expected from the fact that the translation group $T(3)$ is an abelian group, so that its Lie algebra is also abelian, with vanishing structure constants $f_{ab}^c = 0$.

The translation group $T(3)$ is a special case of a more general class of groups called the Heisenberg group, which is a non-abelian group with non-vanishing structure constants and non-vanishing central charges. The Heisenberg group is the symmetry group of the quantum phase space, and its projective representations are related to the canonical commutation relations of quantum mechanics.

Thus the operators $U(\mathbf{a})$ can be written in terms of the momentum operator $\mathbf{p}$, the generators, as

$$U(\mathbf{a}) = e^{\frac{i}{\hbar}\mathbf{a} \cdot \mathbf{p}}.$$

Proof. limit for n....

□

Example (The proper rotation group $SO(3)/SU(2)$).

The proper rotation group $SO(3)$ implements the rotation symmetry of the three-dimensional space:

$$R \in SO(3), \quad R : \mathbf{x} \rightarrow R\mathbf{x},$$

where R is a proper rotation and $\mathbf{x}$ is a vector in the three-dimensional space.

The proper rotation group $SO(3)$ can be represented by the unitary operators $U(R)$ for a proper rotation $R \in SO(3)$ which act on the Hilbert space of states of a quantum system as

$$(U(R)\Psi)(\mathbf{x}) = D(\mathbf{R})\Psi(R^{-1}\mathbf{x}),$$

where Ψ is a state of the quantum system, $\mathbf{x}$ is a point in the configuration space, and $D(\mathbf{R})$ is a unitary operator which acts on the internal degrees of freedom of the quantum system, such as spin.

Indeed we know that the spin of a quantum system is related to the representation of the rotation group $SO(3)$ on the internal degrees of freedom of the quantum system; this does not come as a surprise, since the spin of a quantum system is related to the way the quantum system transforms under rotations, and the rotation group $SO(3)$ is the symmetry group of rotations in three-dimensional space.

We parametrized the representation in order to act on $\mathbf{x}$ as the inverse of the rotation,⁵ since this assumption is consistent with the composition of rotations, as we can see by computing the composition of two rotations R and R' :

$$U(\mathbf{R}, \mathbf{R}') = U(\mathbf{R})U(\mathbf{R}'),$$

otherwise the order would be inverted.

$D(\mathbf{R})$ are $(2s+1) \times (2s+1)$ unitary matrices which form a representation of the rotation group $SO(3)$ on the internal degrees of freedom of the quantum system, where s is the spin of the quantum system

$$D(\mathbf{R})^\dagger D(\mathbf{R}) = D(\mathbf{R})D(\mathbf{R})^\dagger = \mathbb{I},$$

and they satisfy the composition law⁶

$$D(\mathbf{R}\mathbf{R}') = D(\mathbf{R})D(\mathbf{R}').$$

Proof. Let us show first that the operator $U(R)$ is unitary. Define the operator

$$U^*(\mathbf{R})\Psi(\mathbf{x}) = D(\mathbf{R})^\dagger\Psi(R\mathbf{x}),$$

so that we can compute the composition of $U^*(\mathbf{R})$ and $U(\mathbf{R})$ which acts as the identity operator on the Hilbert space of states of the quantum system:

$$(U^*(\mathbf{R})U(\mathbf{R})\Psi)(\mathbf{x}) = D(\mathbf{R})^\dagger D(\mathbf{R})\Psi(\mathbf{R}^{-1}\mathbf{R}\mathbf{x}) = \Psi(\mathbf{x}).$$

⁵Which is always well defined from the properties of groups.

⁶Here the right order is implemented by the previously discussed choice of parametrization.

The next thing to show is that

$$\|U(\mathbf{R})\Psi\|^2 = \|\Psi\|^2,$$

which is the definition of unitarity. We can compute

$$\begin{aligned} \|U(\mathbf{R})\Psi\|^2 &= \int d^3x |U(\mathbf{R})\Psi(\mathbf{x})|^2 = \int d^3x |D(\mathbf{R})\Psi(R^{-1}\mathbf{x})|^2 \\ &= \int d^3x |\Psi(R^{-1}\mathbf{x})|^2 = \int d^3x' |\Psi(\mathbf{x}')|^2 = \|\Psi\|^2, \end{aligned}$$

where we have used the unitarity of $D(\mathbf{R})$ and $J = |\det(\mathbf{R})|^{-1} = 1$. The last step is to show that the composition law is satisfied, which can be done by computing

$$\begin{aligned} (U(\mathbf{R}\mathbf{R}')\Psi)(\mathbf{x}) &= D(\mathbf{R}\mathbf{R}')\Psi((\mathbf{R}\mathbf{R}')^{-1}\mathbf{x}) = D(\mathbf{R})D(\mathbf{R}')\Psi(\mathbf{R}'^{-1}\mathbf{R}^{-1}\mathbf{x}) \\ &= D(\mathbf{R})(U(\mathbf{R}')\Psi)(\mathbf{R}^{-1}\mathbf{x}) = (U(\mathbf{R})U(\mathbf{R}')\Psi)(\mathbf{x}), \end{aligned}$$

proving that $U(\mathbf{R}\mathbf{R}') = U(\mathbf{R})U(\mathbf{R}')$. $\square$

We now aim to study the **exponential parametrization** of the representation U of the proper rotation group $SO(3)$. The generators J_k of the representation U of the proper rotation group $SO(3)$ are defined through the expansion of the unitary operators $U(R)$ for small rotations δR .

We can then parametrize the rotation R by a vector $\boldsymbol{\varphi} = (\varphi_1, \varphi_2, \varphi_3)$ which represents the axis of rotation and the angle of rotation, so that the exponential parametrization of the proper rotation group $SO(3)$ is given by

$$R(\boldsymbol{\varphi}) = e^{-*\boldsymbol{\varphi}}, \quad R(\boldsymbol{\varphi})^{-1} = e^{*\boldsymbol{\varphi}},$$

where

$$*\boldsymbol{\varphi} = \begin{pmatrix} 0 & \varphi_3 & -\varphi_2 \\ -\varphi_3 & 0 & \varphi_1 \\ \varphi_2 & -\varphi_1 & 0 \end{pmatrix},$$

is the antisymmetric matrix associated to the vector $\boldsymbol{\varphi} = (\varphi_1, \varphi_2, \varphi_3)$ which parametrizes the rotation R .

Now we can reflect this exponential parametrization on the representation U of the proper rotation group $SO(3)$ to get an exponential parametrization of the representation:

$$(U(\boldsymbol{\varphi})\Psi)(\mathbf{x}) = D(\boldsymbol{\varphi})\Psi(e^{*\boldsymbol{\varphi}}\mathbf{x}),$$

where $D(\boldsymbol{\varphi})$ is the exponential parametrization of the representation of the rotation group $SO(3)$ on the internal degrees of freedom of the quantum system. We have to look at the expansion of the unitary operators $U(\boldsymbol{\varphi})$ for small rotations $\delta\boldsymbol{\varphi}$ to get the generators $\underline{j}$ of the representation U of the proper rotation group $SO(3)$:

$$U(\delta\boldsymbol{\varphi}) = 1 - i\underline{j} \cdot \delta\boldsymbol{\varphi} + O(\delta\boldsymbol{\varphi}^2),$$

where $\underline{j} = (J_1, J_2, J_3)$ are the generators of the representation U of the proper rotation group $SO(3)$, which are called the angular momentum operators of the quantum system. Then the expansion of the internal degrees transformation $D(\boldsymbol{\varphi})$ for small rotations $\delta\boldsymbol{\varphi}$ gives us

$$D(\delta\boldsymbol{\varphi}) = 1 - i\pm \cdot \delta\boldsymbol{\varphi} + O(\delta\boldsymbol{\varphi}^2),$$

where $\pm = (\Sigma_1, \Sigma_2, \Sigma_3)$ are the generators of the representation of the rotation group $SO(3)$ on the internal degrees of freedom of the quantum system, which are called the spin operators of

the quantum system. Finally we have to compute the expansion of the spatial transformation $e^{*\varphi}$ for small rotations $\delta\varphi$ to get

$$e^{*\delta\varphi} = 1 + *\delta\varphi + O(\delta\varphi^2),$$

but since the rotation can be expressed as $-\delta\varphi \times \mathbf{x}$, we can write

$$e^{*\delta\varphi}\mathbf{x} = \mathbf{x} - \delta\varphi \times \mathbf{x} + O(\delta\varphi^2).$$

Now we can put everything together to get the expansion of the unitary operators $U(\varphi)$ for small rotations $\delta\varphi$:

$$\begin{aligned} (U(\delta\varphi)\Psi)(\mathbf{x}) &= D(\delta\varphi)\Psi(e^{*\delta\varphi}\mathbf{x}) \\ &= (1 - i\pm \cdot \delta\varphi + O(\delta\varphi^2))\Psi(\mathbf{x} - \delta\varphi \times \mathbf{x} + O(\delta\varphi^2)) \\ &= (1 - i\pm \cdot \delta\varphi + i(\delta\varphi \times \mathbf{x}) \cdot (-i\nabla) + O(\delta\varphi^2))\Psi(\mathbf{x}) \\ &\dots \\ &= \Psi(\mathbf{x}) - \delta\varphi \cdot (\mathbf{x} \times \nabla + i\pm)\Psi(\mathbf{x}) + O(\delta\varphi^2), \end{aligned}$$

[...]

Thus we found

$$\underline{j} = \hbar\pm - i\hbar\mathbf{x} \times \nabla,$$

where renaming $\hbar\pm = \mathbf{S}$ and $-i\hbar\mathbf{x} \times \nabla = \mathbf{L}$ we get the familiar decomposition of the angular momentum operator $\underline{j}$ into the spin operator $\mathbf{S}$ and the orbital angular momentum operator $\mathbf{L}$:

$$\underline{j} = \mathbf{S} + \mathbf{L}.$$

We know that the infinitesimal generators of a continuous symmetry group satisfy the same commutation relations as the Lie algebra of the symmetry group, so that the generators J_k of the representation U of the proper rotation group $SO(3)$ satisfy the commutation relations

$$[J_k, J_l] = i\hbar\epsilon_{klm}J_m, \quad k, l, m = 1, 2, 3,$$

while the generators Σ_k of the representation of the rotation group $SO(3)$ on the internal degrees of freedom of the quantum system satisfy the same commutation relations

$$[\Sigma_k, \Sigma_l] = i\epsilon_{klm}\Sigma_m, \quad k, l, m = 1, 2, 3,$$

where ϵ_{klm} is the Levi-Civita symbol. The commutation relations of the generators J_k and Σ_k of the representation, are in line with the structure constants of the Lie algebra $\mathfrak{so}(3)$ of the proper rotation group $SO(3)$, which are given by

$$f_{kl}^m = \epsilon_{klm}.$$

Since we are using the exponential parametrization $\mathbf{R}(\varphi)$ of $SO(3)$, we can get the exponential parametrization of the representation $U(\varphi)$ in terms of the infinitesimal generators J_k :

$$U(\varphi) = e^{-\frac{i}{\hbar}\varphi \cdot \underline{j}}.$$

Proof. limit for n....

□

It should be made an extensive digression on the universal covering group, which is the group which allows us to get a single-valued representation from a multi-valued representation by lifting the representation to the covering group. The universal covering group of a Lie group G is a simply connected Lie group $\tilde{G}$ which covers G in such a way that the projection map from $\tilde{G}$ to G is a local diffeomorphism. The universal covering group $\tilde{G}$ has the same Lie algebra as G , but it can have different global properties, such as different topology and different fundamental group. The universal covering group $\tilde{G}$ allows us to get a single-valued representation from a multi-valued representation of G by lifting the representation to $\tilde{G}$.

However we do not have the time to make this digression, so we will just mention that the universal covering group of the proper rotation group $SO(3)$ is the special unitary group $SU(2)$, which is a simply connected Lie group that covers $SO(3)$ in such a way that the projection map from $SU(2)$ to $SO(3)$ is a local diffeomorphism. The universal covering group $SU(2)$ has the same Lie algebra as $SO(3)$, but it has different global properties, such as different topology and different fundamental group. The universal covering group $SU(2)$ allows us to get a single-valued representation from a multi-valued representation of $SO(3)$ by lifting the representation to $SU(2)$.

1.5 | Dynamical Lie Symmetries and Conservation Laws

[... non lie symmetry content ...]

When a dynamical symmetry group G is a Lie group, with a parametrization $g(\underline{\theta})$, the quantum unitary representation $\tilde{U}(\underline{\theta})$ representing the action of G on the Hilbert space $\mathcal{H}$ respects

$$\tilde{U}(\underline{\theta})H\tilde{U}(\underline{\theta})^\dagger = H, \quad \forall \underline{\theta}.$$

This property restricts the form of the Hamiltonian H on one hand and has consequences for the infinitesimal generators Q_a of the unitary family $\tilde{U}(\underline{\theta})$ on the other: H commutes with the Q_a

$$[H, Q_a] = 0.$$

Proof. If we compute the derivative of the invariance condition with respect to θ^a at $\underline{\theta} = 0$, we get

$$\begin{aligned} 0 &= i \frac{\partial}{\partial \theta^a} (H - \tilde{U}(\underline{\theta})H\tilde{U}(\underline{\theta})^\dagger) \Big|_{\underline{\theta}=0} = \left[-i \frac{\partial \tilde{U}(\underline{\theta})}{\partial \theta^a} H\tilde{U}(\underline{\theta})^\dagger + \tilde{U}(\underline{\theta})H \left(i \frac{\partial \tilde{U}(\underline{\theta})}{\partial \theta^a} \right)^\dagger \right] \Big|_{\underline{\theta}=0} \\ &= -i \frac{\partial \tilde{U}(0)}{\partial \theta^a} H\tilde{U}(0)^\dagger + \tilde{U}(0)H \left(i \frac{\partial \tilde{U}(0)}{\partial \theta^a} \right)^\dagger = -Q_a H + H Q_a = [H, Q_a], \end{aligned}$$

since $\tilde{U}(0) = \mathbb{I}$ and $Q_a = -i \frac{\partial \tilde{U}(\underline{\theta})}{\partial \theta^a} \Big|_{\underline{\theta}=0}$. This proves that when a system is invariant under a Lie group of dynamical symmetries, the generators of that group commute with the Hamiltonian. $\square$

This result is significant because it implies that the generators Q_a are conserved quantities of the system, as they commute with the Hamiltonian. They are observables with meaningful physical significance. In quantum mechanics, this means that the eigenvalues of Q_a are **constants of motion**, and the system's dynamics can be analyzed in terms of these conserved quantities. This connection between symmetries and conservation laws is a fundamental aspect of both classical and quantum physics, encapsulated in **Noether's theorem**. Indeed, we can define the evolution operator

$$U(t, s) = e^{-\frac{i}{\hbar}(t-s)H},$$

such that any observable $\hat{A}$ evolves in time according to

$$\hat{A}(t) = U(t, 0)^\dagger \hat{A} U(t, 0).$$

Then we have

$$U(t, s)Q_a U(t, s)^\dagger = e^{-\frac{i}{\hbar}(t-s)H} Q_a e^{\frac{i}{\hbar}(t-s)H} = Q_a,$$

which shows that $Q_a(t) = Q_a$ is conserved under the time evolution generated by H .

Proof. This is a direct consequence of the commutation relation $[H, Q_a] = 0$: since Q_a commutes with H , it also commutes with any function of H , including the exponential $e^{-\frac{i}{\hbar}(t-s)H}$. Therefore, we can move Q_a through the exponential without changing its value, leading to the conclusion that Q_a is invariant under the time evolution generated by H . $\square$

We have just shown the quantum analog of Noether's theorem: for every continuous symmetry of the action (or equivalently, for every Lie group of dynamical symmetries), there is a corresponding conserved quantity. This profound connection between symmetries and conservation laws is a

cornerstone of modern physics, providing deep insights into the fundamental structure of physical theories.

The symmetry Lie algebra $\mathfrak{g}$

$$[t_a, t_b] = f_{ab}^c t_c,$$

always contains a Cartan subalgebra $\mathfrak{h}$, that is a maximal subspace of $\mathfrak{g}$ the Lie brackets of all whose elements vanish: all the structure constants f_{ab}^c vanish. Denote by h_i a set of generators of $\mathfrak{h}$. These are certain linearly independent linear combinations of the generators T_a of $\mathfrak{g}$

$$h_i = h_i^a t_a,$$

with the property that

$$[h_i, h_j] = 0.$$

Thus we are lead to introduce a set of quantum generators C_i of the Cartan subalgebra $\mathfrak{h}$ in such a way that they commute with the Hamiltonian H and with each other

$$C_i = h_i^a Q_a.$$

These generators C_i , defined in this way, respects

$$[C_i, C_j] = 0.$$

Proof.

$$0 = [h_i, h_j] = [h_i^a t_a, h_j^b t_b] \dots$$

thus we can write

$$\begin{aligned} [C_i, C_j] &= [h_i^a Q_a, h_j^b Q_b] = h_i^a h_j^b [Q_a, Q_b] \\ &= i h_i^a h_j^b f_{ab}^c Q_c = 0, \end{aligned}$$

since the structure constants f_{ab}^c vanish for all a, b when t_a and t_b are in the Cartan subalgebra $\mathfrak{h}$. $\square$

Thus

$$[C_i, C_j] = 0, \quad [C_i, H] = 0.$$

There is a one-to-one correspondence between the eigenvalues of the C_i and the generators of the cartan subalgebra $\mathfrak{h}$.

Consider now

$$\eta_{ab} = \text{Tr}(t_a t_b),$$

if $\mathfrak{g}$ is a semisimple Lie algebra, the matrix η_{ab} is non-degenerate and can be used to raise and lower indices. It will be invertible and we can define η^{ab} such that $\eta^{ab}\eta_{bc} = \delta_c^a$. We obtained a metric on the Lie algebra $\mathfrak{g}$ and we can use.

Now we are going to prove that the structure constants $f_{abc} = f_{bc}^d \eta_{da}$ of a semisimple Lie algebra are completely antisymmetric in the indices a, b, c and not only the last two.

Proof. Since

$$\text{Tr}([A, B]) = \text{Tr}(AB - BA) = 0,$$

we can write

$$\begin{aligned} 0 &= \text{Tr}([t_a, t_b t_c]) = \text{Tr}([t_a, t_b]t_c + t_b[t_a, t_c]) = \text{Tr}(f_{ab}^d t_d t_c + f_{ac}^d t_b t_d) \\ &= f_{ab}^d \eta_{dc} + f_{ac}^d \eta_{db} = f_{abc} + f_{acb} = 0, \end{aligned}$$

or equivalently

$$f_{ab}^c + f_{ca}^b = 0.$$

□

The antisymmetry of the structure constants f_{abc} is a crucial property that has significant implications for the representation theory of Lie algebras and the classification of semisimple Lie algebras. η_{ab} is called the **Cartan metric** of the Lie algebra $\mathfrak{g}$ and through it we can define the Quadratic Casimir operator C_2 of the Lie algebra $\mathfrak{g}$

$$C_2 = \eta^{ab} Q_a Q_b.$$

The Casimir operator C_2 is a central element of the universal enveloping algebra of $\mathfrak{g}$, meaning that it commutes with all elements of $\mathfrak{g}$. In particular, since Q_a are the generators of the symmetry group, we have

$$\begin{aligned} [Q_a, C_2] &= [Q_a, \eta^{bc} Q_b Q_c] = \eta^{bc} ([Q_a, Q_b] Q_c + Q_b [Q_a, Q_c]) \\ &= i \eta^{bc} (f_{ab}^d Q_d Q_c + f_{ac}^d Q_b Q_d) = i (f_a^d{}^b Q_d Q_b + f_a^d{}^c Q_b Q_d) \\ &= i f_a^d{}^b (Q_d Q_b + Q_b Q_d) = 0, \end{aligned}$$

where we have used the antisymmetry of the structure constants f_{abc} in the last step. The Casimir operator C_2 is an important quantity in the representation theory of Lie algebras, as it takes a constant value on each irreducible representation, providing a way to classify representations and understand their properties.

$$[C_i, C_j] = 0, \quad [H, C_i] = 0, \quad [C_2, C_i] = 0, \quad [H, C_2] = 0.$$

To sum up, we have a dynamical lie symmetry group G with Lie algebra $\mathfrak{g}$ and a Cartan subalgebra $\mathfrak{h}$ of $\mathfrak{g}$. We have a set of generators C_i of the Cartan subalgebra $\mathfrak{h}$ that

$$\begin{cases} C_i = h_i^a Q_a, \\ C_2 = \eta^{ab} Q_a Q_b, \end{cases}$$

commute with the Hamiltonian H and with each other, and we have the Casimir operator C_2 that also commutes with H and with the C_i . The eigenvalues of the C_i and C_2 can be used to label the states of the system and to classify the irreducible representations of the symmetry group G .

Example (The Hydrogen Atom). The hydrogen atom has a dynamical symmetry group known as the **SO(3)** group, associated to the algebra $\mathfrak{so}(3)$ with generators

$$[t_a, t_b] = \epsilon_{abc} t_c.$$

We have an associated cartan subalgebra generated by ct_3 , with $c \in \mathbb{R}$ and a cartesian metric $\eta_{ab} = \text{Tr}(t_a t_b) = \delta_{ab}$. The quantum generators J_i (analogue of Q_a) of the symmetry group commute with the Hamiltonian of the hydrogen atom, and the analogue of C_i is J_3 , while the analogue of the Casimir operator C_2 is $J^2 = J_1^2 + J_2^2 + J_3^2 = \delta_{ij} J_i J_j$. The eigenvalues of J_3 and J^2 can be used to label the states of the hydrogen atom and to classify the irreducible

representations of the $SO(3)$ symmetry group, which correspond to the different energy levels and angular momentum states of the electron in the hydrogen atom.

Thus

$$H\Phi_{w\bar{c}\bar{e}} = \Phi_{w\bar{c}\bar{e}}w,$$

$$C_2\Phi_{w\bar{c}\bar{e}} = \Phi_{w\bar{c}\bar{e}}c,$$

$$C_i\Phi_{w\bar{c}\bar{e}} = \Phi_{w\bar{c}\bar{e}}e_i,$$

where $\Phi_{w\bar{c}\bar{e}}$ are the eigenstates of the Hamiltonian H , the Casimir operator C_2 and the generators C_i of the Cartan subalgebra, with eigenvalues w , c and e_i respectively. The quantum numbers w , c and e_i can be used to classify the states of the hydrogen atom and to understand their properties in terms of the underlying symmetries of the system.

Using w as the Bohr energy levels, we can write the energy eigenvalues of the hydrogen atom as

$$w \rightarrow n, \quad n = 1, 2, 3, \dots,$$

$$c \rightarrow l, \quad l = 0, 1, 2, \dots, n-1,$$

$$e_i \rightarrow m, \quad m = -l, -l+1, \dots, l-1, l.$$

where in this case $\bar{e} = e$ is a single quantum number since we have only one generator C_i of the Cartan subalgebra. The energy levels of the hydrogen atom are quantized and depend on the principal quantum number n , the angular momentum quantum number l and the magnetic quantum number m , with the energy eigenvalues given by

$$c = \hbar^2 l(l+1), \quad \dots$$

We obtained a bunch of commuting self adjoint operators H, C_i, C_2 that can be simultaneously diagonalized, with common eigenvectors $\Phi_{w\bar{c}\bar{e}}$ and eigenvalues w, c, e_i . The quantum numbers w, c, e_i can be used to label the states of the system and to classify the irreducible representations of the symmetry group G . This is a powerful tool for understanding the structure of quantum systems and for predicting their behavior based on their symmetries.

We found an orthonormal basis of eigenvectors $\{\Phi_{w\bar{c}\bar{e}}\}$ of the Hilbert space $\mathcal{H}$ that are common eigenvectors of the Hamiltonian H , the Casimir operator C_2 and the generators C_i of the Cartan subalgebra:

$$H\Phi_{w\bar{c}\bar{e}} = w\Phi_{w\bar{c}\bar{e}},$$

$$C_2\Phi_{w\bar{c}\bar{e}} = c\Phi_{w\bar{c}\bar{e}},$$

$$C_i\Phi_{w\bar{c}\bar{e}} = e_i\Phi_{w\bar{c}\bar{e}},$$

This basis allows us to express any state of the system as a linear combination of these eigenvectors, and to analyze the dynamics and properties of the system in terms of its symmetries and conserved quantities.

2 | Vector calculus in $\mathbb{R}^3$ geometry

2.1 | Euclidean structure of $\mathbb{R}^3$

The Euclidean structure of $\mathbb{R}^3$ is nothing but the familiar “dot” product of vectors and is the fundamental structure characterizing $\mathbb{R}^3$. We begin with a remark on notation. Any vector $\mathbf{x} \in \mathbb{R}^3$ can be viewed as either a column or a row vector

$$\mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}, \quad \mathbf{x} = (x_1 \quad x_2 \quad x_3).$$

canonical orthonormal basis of $\mathbb{R}^3$ is given by

component of the vector

matrices 3x3

2.2 | Orientational structure of $\mathbb{R}^3$

For each triple of vectors $\mathbf{x}, \mathbf{y}, \mathbf{z} \in \mathbb{R}^3$, we can define a matrix

$$(\mathbf{x}, \mathbf{y}, \mathbf{z}) = \begin{pmatrix} x_1 & y_1 & z_1 \\ x_2 & y_2 & z_2 \\ x_3 & y_3 & z_3 \end{pmatrix}, \quad \mathbf{x} = \begin{pmatrix} \mathbf{x}^T \\ \mathbf{y}^T \\ \mathbf{z}^T \end{pmatrix} = \begin{pmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \\ z_1 & z_2 & z_3 \end{pmatrix}.$$

the reciprocally transposed matrices of $R[3]$ having x, y, z as column and row vectors in the shown order, respectively. For fixed vectors x, y , the determinant of these matrices depends linearly on z . Therefore, there exists a vector $\mathbf{x} \times \mathbf{y}$ in R^3 with the property that

$$\mathbf{x} \times \mathbf{y} \cdot \mathbf{z} = \det(\mathbf{x}, \mathbf{y}, \mathbf{z}) = \det \begin{pmatrix} \mathbf{x}^T \\ \mathbf{y}^T \\ \mathbf{z}^T \end{pmatrix}.$$

...

$$\mathbf{x} \times \mathbf{y} + \mathbf{y} \times \mathbf{x} = 0,$$

$$\mathbf{x} \times (\mathbf{y} \times \mathbf{z}) + \mathbf{y} \times (\mathbf{z} \times \mathbf{x}) + \mathbf{z} \times (\mathbf{x} \times \mathbf{y}) = 0,$$

$$(\mathbf{x} \times \mathbf{y}) \cdot (\mathbf{u} \times \mathbf{v}) = \dots$$

$$\mathbf{x} \times (\mathbf{y} \times \mathbf{z}) = (\mathbf{x} \cdot \mathbf{z})\mathbf{y} - (\mathbf{x} \cdot \mathbf{y})\mathbf{z},$$

$$\mathbf{e}_i \times \mathbf{e}_j = \sum_k \epsilon_{ijk} \mathbf{e}_k.$$

Proof. We are going to prove the determinant identity...

□

...

2.3 | Determinant and Trace Identities

Suppose A is a 3×3 matrix and $\mathbf{x}, \mathbf{y}, \mathbf{z}$ are three vectors in $\mathbb{R}^3$, then we have the following identities:

$$\begin{aligned} (A\mathbf{x}) \times (A\mathbf{y}) \cdot (A\mathbf{z}) &= \det(A) \mathbf{x} \times \mathbf{y} \cdot \mathbf{z}, \\ (A\mathbf{x}) \times \mathbf{y} \cdot \mathbf{z} + \mathbf{x} \times (A\mathbf{y}) \cdot \mathbf{z} + \mathbf{x} \times \mathbf{y} \cdot (A\mathbf{z}) &= \text{Tr}(A) \mathbf{x} \times \mathbf{y} \cdot \mathbf{z}, \end{aligned}$$

The first identity can be interpreted as the invariance of the determinant under linear transformations, while the second identity can be interpreted as the infinitesimal version of the first identity, which is related to the trace of the matrix A . The first one is called the **determinant identity**, while the second one is called the **trace identity**.

Proof. where the first one can be seen as

$$(A\mathbf{x}) \times (A\mathbf{y}) \cdot (A\mathbf{z}) = \det \begin{pmatrix} A\mathbf{x} & A\mathbf{y} & A\mathbf{z} \end{pmatrix} = \det \left(A \begin{pmatrix} \mathbf{x} & \mathbf{y} & \mathbf{z} \end{pmatrix} \right) = \det(A) \det \begin{pmatrix} \mathbf{x} & \mathbf{y} & \mathbf{z} \end{pmatrix}.$$

If $A = 1 + \lambda A$ and we consider an infinitesimal transformation, then we can differentiate the first identity with the leibniz rule to get the lhs of the second identity

$$\left. \frac{d}{d\lambda} (((1 + \lambda A)\mathbf{x}) \times ((1 + \lambda A)\mathbf{y}) \cdot ((1 + \lambda A)\mathbf{z})) \right|_{\lambda=0} = (A\mathbf{x}) \times \mathbf{y} \cdot \mathbf{z} + \mathbf{x} \times (A\mathbf{y}) \cdot \mathbf{z} + \mathbf{x} \times \mathbf{y} \cdot (A\mathbf{z}),$$

and now we can differentiate the rhs of the first identity to get the rhs of the second identity

$$\left. \frac{d}{d\lambda} ((1 + \lambda \text{Tr}(A)) \mathbf{x} \times \mathbf{y} \cdot \mathbf{z}) \right|_{\lambda=0} = \text{Tr}(A) \mathbf{x} \times \mathbf{y} \cdot \mathbf{z},$$

since $\mathbf{x} \times \mathbf{y} \cdot \mathbf{z}$ is independent of λ and if $A = (\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3)$

$$(1 + \lambda A) = \begin{pmatrix} \mathbf{e}_1 + \lambda \mathbf{a}_1 & \mathbf{e}_2 + \lambda \mathbf{a}_2 & \mathbf{e}_3 + \lambda \mathbf{a}_3 \end{pmatrix},$$

so that differentiating the determinant of this matrix

$$\det(1 + \lambda A) = (\mathbf{e}_1 + \lambda \mathbf{a}_1) \times (\mathbf{e}_2 + \lambda \mathbf{a}_2) \cdot (\mathbf{e}_3 + \lambda \mathbf{a}_3),$$

with respect to λ at $\lambda = 0$ gives us the sum of the diagonal entries of A , which is exactly the trace of A :

$$\begin{aligned} \left. \frac{d}{d\lambda} \det(1 + \lambda A) \right|_{\lambda=0} &= \mathbf{a}_1 \cdot \mathbf{e}_2 \times \mathbf{e}_3 + \mathbf{e}_1 \times \mathbf{a}_2 \cdot \mathbf{e}_3 + \mathbf{e}_1 \times \mathbf{e}_2 \cdot \mathbf{a}_3 \\ &= \mathbf{a}_1 \cdot \mathbf{e}_2 \times \mathbf{e}_3 + \mathbf{a}_2 \cdot \mathbf{e}_3 \times \mathbf{e}_1 + \mathbf{a}_3 \cdot \mathbf{e}_1 \times \mathbf{e}_2 \\ &= \mathbf{a}_1 \cdot \mathbf{e}_1 + \mathbf{a}_2 \cdot \mathbf{e}_2 + \mathbf{a}_3 \cdot \mathbf{e}_3 = a_{11} + a_{22} + a_{33} = \text{Tr}(A). \end{aligned}$$

□

This identities are very useful when encountering the determinant and trace of a matrix in the context of Lie groups and Lie algebras, as we will see in the next chapters.

2.4 | Direct Product

For two vectors $\mathbf{x} = (x_1, x_2, x_3)$ and $\mathbf{y} = (y_1, y_2, y_3)$ in $\mathbb{R}^3$, we can define their direct product as a 3×3 matrix

$$\mathbf{x} \otimes \mathbf{y} = \begin{pmatrix} x_1 y_1 & x_1 y_2 & x_1 y_3 \\ x_2 y_1 & x_2 y_2 & x_2 y_3 \\ x_3 y_1 & x_3 y_2 & x_3 y_3 \end{pmatrix} \in \mathbb{R}(3) = \text{Mat}(3, \mathbb{R}).$$

Definition 2.1 (Direct Product). The direct product of two vectors $\mathbf{x}$ and $\mathbf{y}$ in $\mathbb{R}^3$ is defined as the 3×3 matrix obtained by multiplying each component of $\mathbf{x}$ with each component of $\mathbf{y}$, resulting in a matrix where the entry in the i -th row and j -th column is given by $x_i y_j$. Mathematically, it can be expressed as:

$$\begin{aligned} \otimes : \mathbb{R}^3 \times \mathbb{R}^3 &\mapsto \mathbb{R}^{3 \times 3}, \\ (\mathbf{x}, \mathbf{y}) &\mapsto \mathbf{x} \otimes \mathbf{y} = \begin{pmatrix} x_1 y_1 & x_1 y_2 & x_1 y_3 \\ x_2 y_1 & x_2 y_2 & x_2 y_3 \\ x_3 y_1 & x_3 y_2 & x_3 y_3 \end{pmatrix}. \end{aligned}$$

This operation is bilinear and has the following properties:

$$\begin{cases} (\mathbf{x} \otimes \mathbf{y})\mathbf{z} = (\mathbf{y} \cdot \mathbf{z})\mathbf{x}, \\ (\mathbf{x} \otimes \mathbf{y})^T = \mathbf{y} \otimes \mathbf{x}, \\ \text{Tr}(\mathbf{x} \otimes \mathbf{y}) = \mathbf{x} \cdot \mathbf{y}, \\ \det(\mathbf{x} \otimes \mathbf{y}) = 0. \end{cases}$$

An eigenvector belong to the eigenvalue 0, so the direct product of two vectors is a singular matrix.

This operation is also known as the outer product of two vectors, and it is different from the inner product (dot product) and the cross product. The direct product of two vectors can be used to construct higher-dimensional tensors and has applications in various fields such as physics, engineering, and machine learning.

More identities can be derived from the definition of the direct product; suppose $A \in \mathbb{R}(3)$ is a 3×3 matrix and $\mathbf{x}, \mathbf{y}$ are two vectors in $\mathbb{R}^3$, then we have the following identities:

$$\begin{aligned} (A\mathbf{x}) \otimes (A\mathbf{y}) &= A(\mathbf{x} \otimes \mathbf{y})A^T, \\ A &= \sum_{i,j} A_{ij} \mathbf{e}_i \otimes \mathbf{e}_j. \end{aligned}$$

The first identity can be interpreted as the invariance of the direct product under linear transformations, while the second identity is the expansion of a matrix in terms of the canonical basis of $\mathbb{R}^3$ and the direct product.

2.5 | The Star Operator

For a fixed vector $\mathbf{x} \in \mathbb{R}^3$, we can define a linear operator mapping any vector $\mathbf{y} \in \mathbb{R}^3$ to the cross product of $\mathbf{x}$ and $\mathbf{y}$:

$$\begin{aligned}\star : \mathbb{R}^3 &\mapsto \mathbb{R}(3), \\ \mathbf{x} &\mapsto \star\mathbf{x} = \begin{pmatrix} 0 & x_3 & -x_2 \\ -x_3 & 0 & x_1 \\ x_2 & -x_1 & 0 \end{pmatrix}. \\ (\star\mathbf{x})\mathbf{y} &= -\mathbf{x} \times \mathbf{y}. \\ (\star\mathbf{x})_{ij} &= \sum_k \epsilon_{ijk} x_k.\end{aligned}$$

The star operator is a linear map that takes a vector in $\mathbb{R}^3$ and produces a skew-symmetric matrix. This matrix can be used to compute the cross product of the original vector with any other vector in $\mathbb{R}^3$.

Given a matrix $A \in \mathbb{R}(3)$, we can find its elements as

$$A_{ij} = \mathbf{e}_i \cdot A\mathbf{e}_j,$$

valid for any matrix A and the canonical basis vectors $\mathbf{e}_i$. Therefore, we can express the star operator in terms of the direct product as

$$\begin{aligned}(\star\mathbf{x})_{ij} &= \mathbf{e}_i \cdot (\star\mathbf{x})\mathbf{e}_j = -\mathbf{e}_i \cdot (\mathbf{x} \times \mathbf{e}_j) \\ &= +\mathbf{x} \cdot (\mathbf{e}_i \times \mathbf{e}_j) \dots\end{aligned}$$

We can find its transpose as

$$(\star\mathbf{x})^T = -\star\mathbf{x},$$

and its determinant and trace as

$$\det(\star\mathbf{x}) = 0, \quad \text{Tr}(\star\mathbf{x}) = 0,$$

since the star operator has an eigenvalue of 0: this comes from the fact that the cross product of a vector with itself is always zero, which means that the star operator has a non-trivial kernel and therefore must have an eigenvalue of 0. The trace of the star operator is also zero because it is a skew-symmetric matrix (i.e. $(\star\mathbf{x})^T = -\star\mathbf{x}$), and the trace of any skew-symmetric matrix is always zero.

We can also compute the action of the star operator on the action of a matrix A on a vector $\mathbf{x}$:

$$\star(A\mathbf{x}) = \det AA^{-1,T} (\star\mathbf{x}) A^{-1},$$

and this is very interesting: imagine that A is a rotation matrix

$$A = R \in \text{SO}(3), \quad \det R = 1, \quad R^{-1} = R^T,$$

then the star operator transforms under the rotation as

$$\star(R\mathbf{x}) = R(\star\mathbf{x})R^T,$$

which means that the star operator transforms covariantly under rotations, and this is a very important property in physics, especially in the study of angular momentum and spin.

Proof. Suppose $A \in \mathbb{R}(3)$ and $\mathbf{y}, \mathbf{z} \in \mathbb{R}^3$. Then,

$$\begin{aligned} (\star(A\mathbf{x})\mathbf{y}) \cdot \mathbf{z} &= -((A\mathbf{x}) \times \mathbf{y}) \cdot \mathbf{z} \\ &= -(A\mathbf{x}) \times AA^{-1}\mathbf{y} \cdot AA^{-1}\mathbf{z} \\ &= -\det A(\mathbf{x} \times A^{-1}\mathbf{y} \cdot A^{-1}\mathbf{z}) \\ &= \det A(A^{-1,T}(\mathbf{x} \times A^{-1}\mathbf{y}) \cdot \mathbf{z}) \\ &= \det A((A^{-1,T}(\star\mathbf{x})A^{-1})\mathbf{y}) \cdot \mathbf{z}, \end{aligned}$$

but since $\mathbf{z}$ is arbitrary, we can conclude that

$$\star(A\mathbf{x})\mathbf{y} = (\det A)A^{-1,T}(\star\mathbf{x})A^{-1}\mathbf{y},$$

which means that

$$\star(A\mathbf{x}) = \det AA^{-1,T}(\star\mathbf{x})A^{-1}.$$

□

There is another important identity involving the star operator, which is basically the inverse of the first identity: suppose $A \in \mathbb{R}(3)_{AS}$, which is the space of skew-symmetric matrices, then we have

$$\begin{aligned} \star : \mathbb{R}(3)_{AS} &\mapsto \mathbb{R}^3, \\ A &\mapsto \star A = \begin{pmatrix} A_{32} \\ A_{13} \\ A_{21} \end{pmatrix}, \end{aligned}$$

so that we can find the action of the star operator on a skew-symmetric matrix A as

$$A\mathbf{x} = -(\star A) \times \mathbf{x}.$$

Then we can find the elements of the skew-symmetric matrix A in terms of the components of the vector $\star A$ as

$$(\star A)_i = \frac{1}{2} \sum_{j,k} \epsilon_{ijk} A_{jk},$$

which means that the star operator is an isomorphism between the space of skew-symmetric matrices and the space of vectors in $\mathbb{R}^3$. This is a very important result, as it allows us to identify the Lie algebra of the rotation group $\text{SO}(3)$ with the space of skew-symmetric matrices, and therefore with the space of vectors in $\mathbb{R}^3$. This isomorphism is also known as the **Lie algebra isomorphism** between $\mathfrak{so}(3)$ and $\mathbb{R}^3$.

2.6 | The Wedge Product

Suppose $\mathbf{x}, \mathbf{y} \in \mathbb{R}^3$ are two vectors, then we can define their wedge product as a bivector in the exterior algebra of $\mathbb{R}^3$:

Definition 2.2 (Wedge Product). The wedge product of two vectors $\mathbf{x}$ and $\mathbf{y}$ in $\mathbb{R}^3$ is defined as a bivector in the exterior algebra of $\mathbb{R}^3$, denoted by $\mathbf{x} \wedge \mathbf{y}$. It can be expressed as:

$$\mathbf{x} \wedge \mathbf{y} = (x_1y_2 - x_2y_1)\mathbf{e}_1 \wedge \mathbf{e}_2 + (x_1y_3 - x_3y_1)\mathbf{e}_1 \wedge \mathbf{e}_3 + (x_2y_3 - x_3y_2)\mathbf{e}_2 \wedge \mathbf{e}_3,$$

where $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ is the canonical basis of $\mathbb{R}^3$. The wedge product is an antisymmetric bilinear operation that takes two vectors and produces a bivector, which can be thought of as an oriented area element in $\mathbb{R}^3$.

$$\begin{aligned} \wedge : \mathbb{R}^3 \times \mathbb{R}^3 &\mapsto \Lambda^2(\mathbb{R}^3), \\ (\mathbf{x}, \mathbf{y}) &\mapsto \mathbf{x} \wedge \mathbf{y}. \end{aligned}$$

We can also express the wedge product in terms of the direct product as

$$\mathbf{x} \wedge \mathbf{y} = \mathbf{x} \otimes \mathbf{y} - \mathbf{y} \otimes \mathbf{x},$$

and its elements can be expressed as

$$(\mathbf{x} \wedge \mathbf{y})_{ij} = x_iy_j - y_ix_j.$$

and the matrix representation of the wedge product can be expressed as

$$\mathbf{x} \wedge \mathbf{y} = \begin{pmatrix} 0 & x_1y_2 - x_2y_1 & x_1y_3 - x_3y_1 \\ -(x_1y_2 - x_2y_1) & 0 & x_2y_3 - x_3y_2 \\ -(x_1y_3 - x_3y_1) & -(x_2y_3 - x_3y_2) & 0 \end{pmatrix}.$$

It can be expressed in terms of the cross product as

$$\mathbf{x} \wedge \mathbf{y} = \star(\mathbf{x} \times \mathbf{y}),$$

which means that the wedge product of two vectors can be obtained by applying the star operator to their cross product. This relationship between the wedge product and the cross product is a fundamental result in vector calculus and has important implications in physics, particularly in the study of electromagnetism and fluid dynamics: the cross product can be used to compute the curl of a vector field, while the wedge product can be used to compute the exterior derivative of a differential form. The wedge product is also a key operation in the theory of differential forms and has applications in geometry and topology.

Proof. Suppose $\mathbf{z} \in \mathbb{R}^3$ is a vector, and let us start from the action of the star operator on the cross product of two vectors $\mathbf{x}$ and $\mathbf{y}$ applied to $\mathbf{z}$:

$$\begin{aligned} \star(\mathbf{x} \times \mathbf{y})\mathbf{z} &= -(\mathbf{x} \times \mathbf{y}) \times \mathbf{z} = -\mathbf{z} \times (\mathbf{x} \times \mathbf{y}) \\ &= \dots \end{aligned}$$

where we used the Jacobi identity between the first and second lines,...

□

We can compute the determinant and trace of the wedge product as

$$\det(\mathbf{x} \wedge \mathbf{y}) = 0, \quad \text{Tr}(\mathbf{x} \wedge \mathbf{y}) = 0,$$

since the wedge product is a skew-symmetric matrix (since also the cross product is skew-symmetric), and any skew-symmetric matrix has an eigenvalue of 0 and a trace of 0. The wedge product can be thought of as an oriented area element in $\mathbb{R}^3$, and its determinant being zero reflects the fact that it does not have full rank, while its trace being zero reflects the fact that it is a skew-symmetric matrix.

Using the canonical ON basis of $\mathbb{R}^3$, we can express the wedge product of the basis: there are 9 wedge products of the basis vectors, but only 3 of them are linearly independent, other 3 of them are zero, and the remaining 3 of them are linearly dependent on the first 3:

$$\begin{cases} \mathbf{e}_i \wedge \mathbf{e}_i = 0, \\ \mathbf{e}_i \wedge \mathbf{e}_j = -\mathbf{e}_j \wedge \mathbf{e}_i. \end{cases}$$

Considering a matrix $A \in \mathbb{R}(3)_{AS}$ such that $A^T = -A$, we know that A can be expressed as a linear combination of the wedge products of the basis vectors:

$$A = \frac{1}{2} \sum_{i,j} A_{ij} \mathbf{e}_i \wedge \mathbf{e}_j,$$

which means that the space of skew-symmetric matrices can be identified with the space of bivectors in the exterior algebra of $\mathbb{R}^3$. This is a very important result, as it allows us to identify the Lie algebra of the rotation group $SO(3)$ with the space of skew-symmetric matrices, and therefore with the space of bivectors in the exterior algebra of $\mathbb{R}^3$. This isomorphism is also known as the **Lie algebra isomorphism** between $\mathfrak{so}(3)$ and $\Lambda^2(\mathbb{R}^3)$.

Proof. We can write

$$\begin{aligned} A &= \sum_{ij} A_{ij} \mathbf{e}_i \otimes \mathbf{e}_j = \frac{1}{2} \sum_{ij} (A_{ij} \mathbf{e}_i \otimes \mathbf{e}_j + A_{ij} \mathbf{e}_j \otimes \mathbf{e}_i) \\ &= \frac{1}{2} \sum_{ij} A_{ij} (\mathbf{e}_i \otimes \mathbf{e}_j - \mathbf{e}_j \otimes \mathbf{e}_i) \\ &= \frac{1}{2} \sum_{ij} A_{ij} \mathbf{e}_i \wedge \mathbf{e}_j, \end{aligned}$$

where we used the fact that A is skew-symmetric, so that $A_{ij} = -A_{ji}$, and therefore we can symmetrize the sum over i and j to get the wedge product of the basis vectors. $\square$

2.7 | Diagonalization Theorems

When considering a rotation matrix $R \in \text{SO}(3)$, you have to imagine that it is a complex matrix which happens to have real entries:

$$R \in \text{SO}(3) \subset \mathbb{C}(3).$$

Indeed its eigenvalues and vectors in general are complex. The Pauli matrices formalism is a way to represent these complex eigenvalues and eigenvectors in a more intuitive way, by using the Pauli matrices as a basis for the space of complex matrices.

Thus we consider the complex extension of $\mathbb{R}(3) \subset \mathbb{C}(3)$ of real 3×3 matrices living in the complex space.

in the real case

$$\mathbf{x} \cdot \mathbf{y} \in \mathbb{R}^3, \rightarrow \begin{cases} \mathbf{x} \cdot \mathbf{x} \geq 0, \\ \mathbf{x} \cdot \mathbf{x} = 0 \iff \mathbf{x} = 0. \end{cases}$$

For the complex case, we have to define a new inner product which is positive definite:

$$\mathbf{x} \cdot \mathbf{y} \in \mathbb{C}^3, \rightarrow \begin{cases} \mathbf{x} \cdot \mathbf{x} \geq 0, \\ \mathbf{x} \cdot \mathbf{x} = 0 \text{ even if } \mathbf{x} \neq 0. \end{cases}$$

This is the case for the standard inner product in $\mathbb{C}^3$, thus we can define the hermitian inner product as

$$\mathbf{x}^* \cdot \mathbf{y} = \sum_{i=1}^3 x_i^* y_i, \rightarrow \begin{cases} \mathbf{x}^* \cdot \mathbf{x} \geq 0, \\ \mathbf{x}^* \cdot \mathbf{x} = 0 \iff \mathbf{x} = 0. \end{cases}$$

The orthonormal basis $\mathbf{v}_i$ of $\mathbb{C}^3$ is defined as

$$\mathbf{v}_i^* \cdot \mathbf{v}_j = \delta_{ij}.$$

Considering this basis, we can represent any vector $\mathbf{x} \in \mathbb{C}^3$ as

$$\mathbf{x} = \sum_i (v_i^* \cdot \mathbf{x}) \mathbf{v}_i = \sum_i x_i \mathbf{v}_i.$$

Proof. We can write

$$\begin{aligned} \mathbf{x} &= \sum_i (v_i^* \cdot \mathbf{x}) \mathbf{v}_i, \\ \mathbf{x} &= \sum_j c_{ij} \mathbf{v}_j, \end{aligned}$$

and

$$\mathbf{v}_i^* \cdot \mathbf{x} = \sum_j c_{ij} \mathbf{v}_i^* \cdot \mathbf{v}_j = c_{ii} = x_i,$$

thus

$$\mathbf{x} = \sum_i (v_i^* \cdot \mathbf{x}) \mathbf{v}_i,$$

proving the equality.

□

$$\mathbf{x} = \sum_i (\mathbf{v}_i \otimes \mathbf{v}_i^*) \mathbf{x},$$

using the fact that $\mathbf{v}_i^* \cdot \mathbf{v}_j = \delta_{ij}$, ...

Consider $A \in \mathbb{C}(3)$ and suppose that A is diagonalizable, then there exists an orthonormal basis of eigenvectors $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ of $\mathbb{C}^3$ such that

$$A\mathbf{v}_i = \lambda_i \mathbf{v}_i, \quad i = 1, 2, 3, \quad \lambda_i \in \mathbb{C}.$$

This means that we can express the matrix A in terms of its eigenvalues and eigenvectors as

$$A = \sum_i \lambda_i \mathbf{v}_i \otimes \mathbf{v}_i^*,$$

which is the **spectral decomposition** of the matrix A .

Theorem 2.1 (Spectral theorem). *Let A be a diagonalizable matrix in $\mathbb{C}(3)$. Then there exists an orthonormal basis of eigenvectors $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ of $\mathbb{C}^3$ such that*

$$A = \sum_i \lambda_i \mathbf{v}_i \otimes \mathbf{v}_i^*,$$

where λ_i are the eigenvalues of A corresponding to the eigenvectors $\mathbf{v}_i$. Then the matrix can be diagonalized as

$$A = V\Lambda V^{-1}, \quad A = \sum_i \lambda_i \mathbf{v}_i \otimes \mathbf{v}_i^*,$$

where V is the matrix whose columns are the eigenvectors $\mathbf{v}_i$ and Λ is the diagonal matrix whose diagonal entries are the eigenvalues λ_i .

This decomposition allows us to understand the structure of the matrix A in terms of its eigenvalues and eigenvectors, and it is a fundamental result in linear algebra with applications in various fields such as physics, engineering, and machine learning. The spectral decomposition is particularly useful when dealing with normal matrices, which are matrices that commute with their conjugate transpose, as they can be diagonalized by a unitary transformation.

Proof. Since A is diagonalizable, we can write A as

$$\begin{aligned} A &= A\mathbb{I}_3 = A \sum_i \mathbf{v}_i \otimes \mathbf{v}_i^* = \sum_i A\mathbf{v}_i \otimes \mathbf{v}_i^* \\ &= \sum_i \lambda_i \mathbf{v}_i \otimes \mathbf{v}_i^* = A. \end{aligned}$$

□

3 | Pauli Matrices Formalism

In this chapter, we will explore the Pauli matrices and their applications in quantum mechanics from a group theory perspective. The Pauli matrices are a set of three 2x2 complex Hermitian and unitary matrices that are widely used in the study of spin-1/2 particles and quantum information theory.

Our discussion will be centered around the following groups and their representations:

- $SU(2)$: The special unitary group of degree 2, which is the double cover of the rotation group $SO(3)$. The Pauli matrices serve as generators for the Lie algebra $\mathfrak{su}(2)$ associated with this group.
- $SO(3)$: The special orthogonal group in three dimensions, which describes the symmetries of physical space. The representations of $SO(3)$ can be constructed using the Pauli matrices.
- $SL(2, \mathbb{C})$: The special linear group of degree 2 over the complex numbers, which is related to the Lorentz group in relativistic quantum mechanics. The Pauli matrices play a crucial role in the representation theory of this group as well.
- $SO_+(1, 3)$: The proper orthochronous Lorentz group, which describes the symmetries of spacetime in special relativity. The Pauli matrices are used to construct the spinor representations of this group.

3.1 | Pauli Matrix Functions

In this section, we will define the Pauli matrices and explore their properties. The Pauli matrix function is defined as follows:

Definition 3.1 (Pauli Matrix Function). A Pauli matrix function is an element of $\sigma \in \text{Hom}(\mathbb{C}^3, \mathbb{C}(2))$, i.e. a linear map

$$\sigma : \mathbb{C}^3 \rightarrow \mathbb{C}(2),$$

satisfying the following properties: for all $\mathbf{x}, \mathbf{y} \in \mathbb{C}^3$, we have

- $\sigma(\mathbf{x})^\dagger = \sigma(\mathbf{x}^*)$, where $\dagger$ denotes the Hermitian conjugate and $*$ denotes complex conjugation.
- $\sigma(\mathbf{x})\sigma(\mathbf{y}) = \mathbf{x} \cdot \mathbf{y} \mathbb{I}_3 + i\sigma(\mathbf{x} \times \mathbf{y})$, where $\cdot$ denotes the dot product and $\times$ denotes the cross product in $\mathbb{C}^3$.

3.1.1 | (Anti)commutation Relations

The Pauli matrices satisfy specific commutation and anticommutation relations that are fundamental to their algebraic structure. These relations can be expressed as follows:

- Commutation relations: For any two Pauli matrices σ_i and σ_j , we have

$$[\sigma_i, \sigma_j] = 2i\epsilon_{ijk}\sigma_k,$$

where ϵ_{ijk} is the Levi-Civita symbol.

- Anticommutation relations: For any two Pauli matrices σ_i and σ_j , we have

$$\{\sigma_i, \sigma_j\} = 2\delta_{ij}\mathbb{I}_2,$$

where δ_{ij} is the Kronecker delta and $\mathbb{I}_2$ is the 2x2 identity matrix.

These define the **Clifford algebra** associated with the Pauli matrices, which is essential for understanding their role in quantum mechanics and representation theory: the Pauli matrices generate the Clifford algebra, which is isomorphic to the algebra of 2x2 complex matrices. This algebraic structure allows us to construct representations of the groups mentioned above and to analyze the symmetries of quantum systems.

We have trace and determinant properties of the Pauli matrices as well, which are important for their applications in quantum mechanics: starting from the trace properties, we can derive the following identities for any $\mathbf{x}, \mathbf{y}, \mathbf{z} \in \mathbb{C}^3$:

$$\begin{aligned} \text{Tr}(\sigma(\mathbf{x})) &= 0, \\ \text{Tr}(\sigma(\mathbf{x})\sigma(\mathbf{y})) &= 2\mathbf{x} \cdot \mathbf{y}, \\ \text{Tr}(\sigma(\mathbf{x})\sigma(\mathbf{y})\sigma(\mathbf{z})) &= 2i\mathbf{x} \times \mathbf{y} \cdot \mathbf{z}, \end{aligned}$$

Proof. we have that, from the definition of the Pauli matrix function, we can write $\sigma(\mathbf{x}) = x_i\sigma_i$, where σ_i are the standard Pauli matrices. Then, we can compute the trace properties as follows:...

[only the first two were proven]

□

TODO: check and correct

For the determinant properties, we can derive the following identities for any $\mathbf{x} \in \mathbb{C}^3$:

$$\det(\sigma(\mathbf{x})) = -\mathbf{x} \cdot \mathbf{x},$$

Proof. Start from $\mathbf{x} \in \mathbb{R}^3$ we can write, if $\mathbf{x} = \mathbf{y}$

$$\sigma(\mathbf{x})^2 = \mathbf{x} \cdot \mathbf{x} \mathbb{I}_2,$$

and by computing the determinant of both sides, we get

$$(\det(\sigma(\mathbf{x})))^2 = (\det(\mathbf{x} \cdot \mathbf{x} \mathbb{I}_2))^2 = (\mathbf{x}^2)^2,$$

which implies

$$\det(\sigma(\mathbf{x})) = \eta(\mathbf{x}) \mathbf{x} \cdot \mathbf{x}, \quad \eta(\mathbf{x}) = \pm 1.$$

Indeed

$$\eta(\mathbf{x}) = \frac{\det(\sigma(\mathbf{x}))}{\mathbf{x} \cdot \mathbf{x}},$$

is a continuous function of $\mathbf{x}$ that takes values in

$$\mathbb{R}^3 \setminus \{0\} \rightarrow \mathbb{C},$$

and since $\mathbb{R}^3 \setminus \{0\}$ is connected, we have that $\eta(\mathbf{x})$ is constant,

$$\det(\sigma(\mathbf{x})) \det(\sigma(\mathbf{y})) = \dots$$

□

4 | Boh

Theorem 4.1 (Diagonalization theorem for $O(3)$). *Let $R \in O(3) \subset \mathbb{C}(3)$. Then there is a sign $\epsilon \in \{-1, +1\}$ and a phase $q \in [0, \pi]$ and a basis Q_α , such that $\alpha = -1, 0, 1$. Then the following holds:*

1. $v_\alpha \cdot v_\beta = \delta_{-\alpha, \beta}$,
2. $v_\alpha^* = v_{-\alpha}$,
3. $v_0 \times v_\alpha = i\alpha v_\alpha$,
4. ...
5. ...

Proof. let $w \in \mathbb{C}^3$ invariant under R . Then we can define its orthonormal vector $w^\perp \in \mathbb{C}^3$ which is also invariant under R . Then ...

Thus R has always one real eigenvalue and two complex conjugate eigenvalues (maybe also real). The real one $\lambda_0 = \epsilon$ while the complex ones are $\lambda_{\pm 1} = e^{\pm iq}$. Then we can define

$$\lambda_\alpha = \epsilon e^{-i\alpha q}, \quad \alpha = -1, 0, 1.$$

Thus

$$Rv_\alpha = v_\alpha \Lambda_\alpha, \quad Rv_\alpha^* = v_\alpha^* \Lambda_{-\alpha}^* = e^{i\chi_\alpha} v_{-\alpha} \Lambda_{-\alpha}.$$

since given (?) we have that $v_\alpha^* = e^{-i\chi_\alpha} v_{-\alpha}$. [...]

$$v_1 = \frac{1}{\sqrt{2}}(v_1 + v_{-1}), \quad v_2 = \frac{1}{\sqrt{2}i}(v_1 - v_{-1}), \quad v_3 = v_0,$$

and we can prove this basis to be orthonormal and to satisfy the properties of the theorem. Hence $v_i \times v_j = \sum_{k=-1}^1 \epsilon_{ijk} v_k$, proving (3) and (4). $\square$

...

This shows that defining these matrices in complex space give us the ability to diagonalize them, which is not possible in real space (except for special cases). This is the reason why we need to complexify the representation of $O(3)$ to get the irreducible representations of it.

Proposition 4.2 (...). *Let $R \in O(3) \subset \mathbb{C}(3)$. Then there is a sign $\epsilon \in \{-1, +1\}$ and a phase $\phi \in [0, \pi]$ and a unit vector $\mathbf{n} \in \mathbb{R}^3$, $n^2 = 1$, such that*

$$R = \epsilon [\mathbf{n} \otimes \mathbf{n} + \cos(\phi) (1 - \mathbf{n} \otimes \mathbf{n}) - \sin(\phi) \star \mathbf{n}].$$

Every matrix R of this form is an element of $O(3)$. The opposite is also true: every element of $O(3) \subset \mathbb{C}(3)$ can be written in this form.

Proof.

$$R = \epsilon [\mathbf{v}_0 \otimes \mathbf{v}_0 + (\cos(\phi) - i \sin(\phi)) \mathbf{v}_1 \otimes \mathbf{v}_{-1}],$$

where now we can write

$$\epsilon [\mathbf{v}_0 \otimes \mathbf{v}_0 + \cos(\phi) (\mathbf{v}_1 \otimes \mathbf{v}_{-1} + \mathbf{v}_{-1} \otimes \mathbf{v}_1) - i \sin(\phi) (\mathbf{v}_1 \otimes \mathbf{v}_{-1} - \mathbf{v}_{-1} \otimes \mathbf{v}_1)],$$

since ... now

$$\sum_{\alpha} \mathbf{v}_{\alpha} \otimes \mathbf{v}_{-\alpha} = 1, \quad \implies \quad \mathbf{v}_1 \otimes \mathbf{v}_{-1} + \mathbf{v}_{-1} \otimes \mathbf{v}_1 = 1 - \mathbf{v}_0 \otimes \mathbf{v}_0,$$

thus

$$\mathbf{v}_1 \wedge \mathbf{v}_{-1} = \mathbf{v}_1 \otimes \mathbf{v}_{-1} - \mathbf{v}_{-1} \otimes \mathbf{v}_1 = \star(\mathbf{v}_1 \times \mathbf{v}_{-1}) = \star(-i\mathbf{v}_0) = -i \star \mathbf{v}_0.$$

So we get

$$R = \epsilon [\mathbf{v}_0 \otimes \mathbf{v}_0 + \cos(\phi)(1 - \mathbf{v}_0 \otimes \mathbf{v}_0) - i \sin(\phi)(-i \star \mathbf{v}_0)] = \epsilon [\mathbf{v}_0 \otimes \mathbf{v}_0 + \cos(\phi)(1 - \mathbf{v}_0 \otimes \mathbf{v}_0) - \sin(\phi) \star \mathbf{v}_0].$$

By setting $\mathbf{n} = \mathbf{v}_0$ we get the desired result, and we can do this since $\mathbf{v}_0 \in \mathbb{R}^3$ and $\mathbf{v}_0^2 = 1$. $\square$

We can name the previous expression as the *angle-axis parametrization* of $O(3)$. The angle ϕ is the angle of rotation, while the axis of rotation is given by the unit vector $\mathbf{n}$. The sign ϵ tells us whether we are dealing with a proper or an improper rotation. In particular, if $\epsilon = +1$ we are dealing with a proper rotation, while if $\epsilon = -1$ we are dealing with an improper rotation (a reflection). Thus we can write any matrices in $O(3)$ as

$$R_{\epsilon}(\phi, \mathbf{n}) = \epsilon [\mathbf{n} \otimes \mathbf{n} + \cos(\phi)(1 - \mathbf{n} \otimes \mathbf{n}) - \sin(\phi) \star \mathbf{n}].$$

Example (Exercise). Being $A \in \mathbb{R}(3)$ and $\mathbf{v}_i$ with $i = 1, 2, 3$ an orthonormal basis of $\mathbb{R}^3$ such that

$$A\mathbf{v}_i = \sum_{j=1}^3 \hat{A}_{ji} \mathbf{v}_j, \quad (\hat{A}_{ji}) \in \mathbb{R}(3).$$

Thus $\hat{A}$ is a representation matrix of A in the basis $\mathbf{v}_i$.

Now, being $R \in O(3)$ and $\mathbf{v}_i$ respecting

1. $\mathbf{v}_3 = \mathbf{n}$,
2. $\mathbf{v}_i \cdot \mathbf{v}_j = \delta_{ij}$ for all i, j ,
3. $\mathbf{v}_i \times \mathbf{v}_j = \sum_{k=1}^3 \epsilon_{ijk} \mathbf{v}_k$.

Then we can write

$$R_{\epsilon}(\phi, \mathbf{n}) = \epsilon [\mathbf{n} \otimes \mathbf{n} + \cos(\phi)(1 - \mathbf{n} \otimes \mathbf{n}) - \sin(\phi) \star \mathbf{n}].$$

Then its action on the basis $\mathbf{v}_i$ is given by

$$R_{\epsilon}(\phi, \mathbf{n})\mathbf{v}_i = \sum_{j=1}^3 \hat{R}_{\epsilon}(\phi, \mathbf{n})_{ji} \mathbf{v}_j,$$

where

$$\hat{R}_\epsilon(\phi, \mathbf{n}) = \epsilon \begin{pmatrix} \cos(\phi) & -\sin(\phi) & 0 \\ \sin(\phi) & \cos(\phi) & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

This can be obtained by explicitly calculating the action of $R_\epsilon(\phi, \mathbf{n})$ on the basis $\mathbf{v}_i$. For example, we have

$$\begin{aligned} R_\epsilon(\phi, \mathbf{n})\mathbf{v}_i &= \epsilon [\mathbf{n} \otimes \mathbf{n} + \cos(\phi)(1 - \mathbf{n} \otimes \mathbf{n}) - \sin(\phi) \star \mathbf{n}] \mathbf{v}_i \\ &= \dots \end{aligned}$$

$\hat{A}$ depends on the choice of the basis $\mathbf{v}_i$, however its trace and determinant do not depend on the choice of the basis. In particular, we have

$$\begin{cases} \text{Tr}(\hat{A}) = \text{Tr}(A), \\ \det(\hat{A}) = \det(A). \end{cases}$$

Hence we now that and .

$$\begin{cases} \det(R_\epsilon(\phi, \mathbf{n})) = \det(\hat{R}_\epsilon(\phi, \mathbf{n})) = \epsilon^3 = \epsilon, \\ \text{Tr}(R_\epsilon(\phi, \mathbf{n})) = \text{Tr}(\hat{R}_\epsilon(\phi, \mathbf{n})) = (2 \cos \phi + 1)\epsilon. \end{cases}$$

This is very useful since it allows us to determine the angle ϕ and the sign ϵ (which are equal for either the matrix and its representation) from the trace and determinant of R : we have all the element to write the angle-axis parametrization $\hat{R}$ of R .

This result implies that we can write the action of any element of $O(3)$ on $\mathbb{R}^3$ as a rotation around an axis, in the following form

$$A\mathbf{x} = \sum_i \left(\sum_j \hat{A}_{ij} \mathbf{x}_j \right) \mathbf{v}_i.$$

Let us write

$$R_{+1}(\phi, \mathbf{n}) = [\mathbf{v}_3 \otimes \mathbf{v}_3 + \cos(\phi)(1 - \mathbf{v}_3 \otimes \mathbf{v}_3) - \sin(\phi) \star \mathbf{v}_3],$$

then adopting the following cylindrical coordinates:

$$\begin{cases} \mathbf{x}_1 = \rho \cos \theta, \\ \mathbf{x}_2 = \rho \sin \theta, \\ \mathbf{x}_3 = z, \epsilon = +1, \end{cases}$$

we can write the action of the orthogonal matrix $R_{+1}(\phi, \mathbf{n})$ on the vector $\mathbf{x}$ as

$$\begin{aligned} (R_{+1}(\phi, \mathbf{n})\mathbf{x})_1 &= \rho \cos(\phi + \theta), \\ (R_{+1}(\phi, \mathbf{n})\mathbf{x})_2 &= \rho \sin(\phi + \theta), \\ (R_{+1}(\phi, \mathbf{n})\mathbf{x})_3 &= z, \end{aligned}$$

showing that the action of $R_{+1}(\phi, \mathbf{n})$ on any vector $\mathbf{x} \in \mathbb{R}^3$ is a rotation of angle ϕ around the axis $\mathbf{n}$.

Thus we have shown that if $R \in O(3)$ then there are $\epsilon = \pm 1$ and $\phi \in [0, \pi]$ and $\mathbf{n} \in \mathbb{R}^3$ with $\mathbf{n}^2 = 1$ such that

$$R_\epsilon(\phi, \mathbf{n}) = \epsilon [\mathbf{n} \otimes \mathbf{n} + \cos(\phi)(1 - \mathbf{n} \otimes \mathbf{n}) - \sin(\phi) \star \mathbf{n}].$$

with ϕ and $\mathbf{n}$ given by the trace and determinant of R through the following relations:

$$\begin{cases} \epsilon = \det(R), \\ \phi = \arccos\left(\frac{\text{Tr}(R)}{2\epsilon} - \frac{1}{2}\right). \end{cases}$$

We can also write the angle-axis parametrization of $R \in SO(3)$, the special orthogonal group, with only proper rotations, as $\epsilon = 1$, but the relation is exactly the same:

$$R_{+1}(\phi, \mathbf{n}) = \mathbf{n} \otimes \mathbf{n} + \cos(\phi)(1 - \mathbf{n} \otimes \mathbf{n}) - \sin(\phi) \star \mathbf{n}.$$

A special case is the space reflection with respect to the origin, which is given by $\epsilon = -1$ and $\phi = 0$ and in physical space is called *parity transformation*; it is given by

$$P = R_{-1}(0, \mathbf{n}) = -1, \quad P\mathbf{x} = -\mathbf{x}.$$

With this transformation we can relate the orthogonal group $O(3)$ to the special orthogonal group $SO(3)$ as follows:

$$O(3) = SO(3) \cup PSO(3).$$

At a certain point it becomes important to know if this parametrization is unique, i.e. if given R there is only one way to write it as $R_\epsilon(\phi, \mathbf{n})$. If the parameters ϵ , ϕ , and $\mathbf{n}$ are uniquely determined by R , then the parametrization is unique. But since the first two parameters ϵ and ϕ are determined by the trace and determinant of R , which are invariant under change of basis, we only need to check if the axis of rotation $\mathbf{n}$ is uniquely determined by R . This is not the case since:

- If we have a rotation of angle $\phi = \pi$ around an axis $\mathbf{n}$, then we can also write it as a rotation of angle π around the axis $-\mathbf{n}$: it is unique up to sign change;

$$R = \dots$$

$$\phi = \pi \mathbf{n} = \pi(-\mathbf{n}).$$

- If we consider $\phi = 0$ then the parametrization is undetermined since any axis $\mathbf{n}$ can be chosen, since the rotation of angle 0 around any axis is the identity transformation: it is totally undetermined;

$$R = \dots \phi = 0 \mathbf{n} = 0 \mathbf{n}' = \mathbf{0}.$$

It can be proven, however, that if $0 < \phi < \pi$, strictly comprised, then the parametrization is unique, since in this case the axis of rotation $\mathbf{n}$ is uniquely determined by R .

Then if we have defined the phase ϕ as strictly comprised between 0 and π , then in the phase space, the angle vector ϕ has to have a norm equal or less than π :

$$|\phi| \leq \pi.$$

This tells us that the topology of the phase space of $SO(3)$ is a ball of radius π with antipodal¹ points (on the surface) identified, since the points on the surface of the ball, which have norm equal

¹Opposite with respect to the origin.

to π , are identified with their antipodal points (the one with opposite sign), since they represent the same rotation.

Now the Lie algebra the special orthogonal group $\text{SO}(3)$ is

$$\sigma(3) = \mathfrak{so}(3) = \{A \in \mathbb{R}(3) : A^T = -A\},$$

then we can state the following proposition:

Proposition 4.3. *For every $X \in \mathfrak{so}(3)$ there exists a unique vector $\mathbf{x} \in \mathbb{R}^3$ such that*

$$X = -\star \mathbf{x}, \quad \star \mathbf{x} = \begin{pmatrix} 0 & \mathbf{x}_3 & -\mathbf{x}_2 \\ -\mathbf{x}_3 & 0 & \mathbf{x}_1 \\ \mathbf{x}_2 & -\mathbf{x}_1 & 0 \end{pmatrix}.$$

Indeed we have seen that with the angle-axis parametrization we can write any element of $\text{SO}(3)$ with three parameters. We denote the map $\mathbf{x} \mapsto -\star \mathbf{x}$ with

$$X = \ell(\mathbf{x}),$$

with the following properties:

1. $\ell(\mathbf{x})\ell(\mathbf{y}) = \ell(\mathbf{x} \times \mathbf{y})$,
2. $[\ell(\mathbf{x}), \ell(\mathbf{y})]\mathbf{z} = \star \mathbf{x} \times \mathbf{y}\mathbf{z} - \star \mathbf{y} \times \mathbf{x}\mathbf{z} \dots$

Indeed we know that the space of antisymmetric matrices is three-dimensional, and the map $\mathbf{x} \mapsto \star \mathbf{x}$ is a linear map from $\mathbb{R}^3$ to $\mathfrak{so}(3)$. Since both spaces have the same dimension, it is enough to show that this map is injective, i.e. that if $\star \mathbf{x} = 0$ then $\mathbf{x} = 0$. But this is true since the only vector that has zero cross product with all the other vectors is the zero vector itself. Hence the map is injective and thus it is an isomorphism, proving the proposition.

[...] then with the properties of the map ℓ we can show that the Lie bracket of $\mathfrak{so}(3)$ is given by

$$[\ell_i, \ell_j] = \sum_{k=1}^3 \epsilon_{ijk} \ell_k.$$

Another useful property comes from the exponential map, since it is naturally connected to the angle-axis parametrization. Indeed we have $\Phi \in \mathbb{R}$, $\mathbf{n} \in S^2$ (the unit sphere in $\mathbb{R}^3$), then the exponential map is given by

$$\exp(\ell(\phi \mathbf{n})) = R_{+1}(\Phi, \mathbf{n}).$$

Proof. To show this, we can write the exponential map as a power series:

$$\exp(\ell(\phi \mathbf{n})) = \sum_{k=0}^{\infty} \frac{1}{k!} \ell(\phi \mathbf{n})^k = \sum_{k=0}^{\infty} \frac{1}{k!} (-\phi \star \mathbf{n})^k,$$

where now if we divide the odd and even powers of k we get

$$\exp(-\phi \star \mathbf{n}) = \sum_{k=0}^{\infty} \frac{1}{(2k)!} (-\phi)^k (\star \mathbf{n})^{2k} = 1 + \sum_{k=1}^{\infty} \frac{\phi^{2k}}{(2k)!} (\star \mathbf{n})^{2k} - \sum_{k=0}^{\infty} \frac{\phi^{2k+1}}{(2k+1)!} (\star \mathbf{n})^{2k+1}.$$

Now let us compute $(\star \mathbf{n})^2$:

$$(\star \mathbf{n})^2 \mathbf{x} = \dots = -1 + \mathbf{n} \otimes \mathbf{n},$$

which is a projector on the plane orthogonal to $\mathbf{n}$. Hence, since all the projectors are idempotent, we have

$$[(\star \mathbf{n})^2]^2 = (\star \mathbf{n})^2,$$

then, with the idea to compute the expansion of the exponential map, we can write

$$(\star \mathbf{n})^{2k+1} \mathbf{x} = \cdots = (-1)^k \star \mathbf{n} \mathbf{x}, \quad (\star \mathbf{n})^{2k} \mathbf{x} = \cdots = (-1)^k (1 - \mathbf{n} \otimes \mathbf{n}) \mathbf{x}.$$

Hence the expansion takes the following form:

$$\begin{aligned} \exp(-\phi \star \mathbf{n}) &= 1 + \sum_{k=1}^{\infty} \frac{\phi^{2k}}{(2k)!} (-1)^k (1 - \mathbf{n} \otimes \mathbf{n}) - \sum_{k=0}^{\infty} \frac{\phi^{2k+1}}{(2k+1)!} (-1)^k \star \mathbf{n} \\ &= 1 + (\cos(\phi) - 1)(1 - \mathbf{n} \otimes \mathbf{n}) - \sin(\phi) \star \mathbf{n} = R_{+1}(\Phi, \mathbf{n}), \end{aligned}$$

which is exactly the angle-axis parametrization of a proper rotation of angle ϕ around the axis $\mathbf{n}$, proving the desired result. $\square$

=====

4.1 | The Lorentz group

In the Minkowski space $\mathbb{R}^4$ we can define the Lorentz group as the group of linear transformations that preserve the Minkowski metric $\eta = \text{diag}(1, -1, -1, -1)$. We adopt the 1+3 notation for vectors in $\mathbb{R}^4$ or matrices in $\mathbb{R}(4)$, which is the most common in physics.

TODO: substitute vec with tilde under the symbol

$$\vec{u} = \begin{pmatrix} a \\ \mathbf{u} \end{pmatrix}, \quad a \in \mathbb{R}, \mathbf{u} \in \mathbb{R}^3$$

$$\vec{u}^t = \begin{pmatrix} a & \mathbf{u}^t \end{pmatrix}$$

For the matrices

$$\vec{A} = \begin{pmatrix} x & \mathbf{u}^t \\ \mathbf{v} & P \end{pmatrix}, \quad x \in \mathbb{R}, \mathbf{u}, \mathbf{v} \in \mathbb{R}^3, P \in \mathbb{R}(3).$$

When applying an operator $\vec{A} \in \mathbb{R}(4)$ on a four-vector $\vec{u} \in \mathbb{R}^4$ we have

$$\vec{A}\vec{u} = \begin{pmatrix} x & \mathbf{u}^t \\ \mathbf{v} & P \end{pmatrix} \begin{pmatrix} a \\ \mathbf{u} \end{pmatrix} = \begin{pmatrix} xa + \mathbf{u}^t \mathbf{u} \\ a\mathbf{v} + P\mathbf{u} \end{pmatrix}.$$

When multiplying two matrices $\vec{A}$ and $\vec{B}$ we have

TODO: Correct last expression

$$\vec{A}\vec{B} = \begin{pmatrix} x & \mathbf{u}^t \\ \mathbf{v} & P \end{pmatrix} \begin{pmatrix} y & \mathbf{w}^t \\ \mathbf{z} & Q \end{pmatrix} = \begin{pmatrix} xy + \mathbf{u}^t \mathbf{z} & x\mathbf{w}^t + \mathbf{u}^t Q \\ y\mathbf{v} + P\mathbf{z} & \mathbf{v}\mathbf{w}^t + PQ \end{pmatrix}.$$

The minkowski metric can be written as

$$\vec{\eta} = \begin{pmatrix} 1 & \mathbf{0}^t \\ \mathbf{0} & -\mathbf{I}_3 \end{pmatrix}.$$

Let us define the full Lorentz group as

$$\text{O}(1, 3) = \{\vec{L} \in \mathbb{R}(4) : \vec{L} \text{ invertible}, \vec{L}^T = \vec{L}^{-1}, \vec{L}^T = \vec{\eta} \vec{L}^t \vec{\eta}\}.$$

$$\vec{\eta} \vec{L}^t \vec{\eta} = \vec{L}^{-1} \implies \vec{L}^t \vec{\eta} \vec{L} = \vec{\eta}.$$

Thus given $\vec{x} = \begin{pmatrix} a \\ \mathbf{x} \end{pmatrix}$ and $\vec{y} = \begin{pmatrix} b \\ \mathbf{y} \end{pmatrix}$ we have

$$\vec{x} \cdot \vec{y} = \vec{x}^t \vec{\eta} \vec{y} = ab - \mathbf{x} \cdot \mathbf{y}.$$

Applying the transformation $\vec{L}$ on the vectors $\vec{x}$ and $\vec{y}$ we have

$$(\vec{L}\vec{x}) \cdot (\vec{L}\vec{y}) = (\vec{L}\vec{x})^t \vec{\eta} (\vec{L}\vec{y}) = \vec{x}^t \vec{L}^t \vec{\eta} \vec{L} \vec{y} = \vec{x}^t \vec{\eta} \vec{y} = \vec{x} \cdot \vec{y},$$

and we could use this property to define the Lorentz group as the group of linear transformations that preserve the Minkowski metric, i.e. the group of linear transformations that preserve the Minkowski inner product:

$$\text{O}(1, 3) = \{\vec{L} \in \mathbb{R}(4) : \vec{L} \text{ invertible}, (\vec{L}\vec{x}) \cdot (\vec{L}\vec{y}) = \vec{x} \cdot \vec{y} \text{ for all } \vec{x}, \vec{y} \in \mathbb{R}^4\}.$$

The product of two four-vectors $\vec{x}$ and $\vec{y}$ is a Lorentz scalar, since it is invariant under Lorentz transformations. The norm of a four-vector $\vec{x}$ is also a Lorentz scalar, since it is given by $\vec{x} \cdot \vec{x}$, which is invariant under Lorentz transformations. This means that those measurements will be the same for all inertial observers, since they are invariant under Lorentz transformations. This is the reason why we call them scalars, since they are invariant under changes of reference frame.

Theorem 4.4 (Structure theorem). *Suppose $\vec{L} \in O(1, 3)$. Then there is a sign $\eta \in \{-1, 1\}$, $\rho \in [0, \infty)$, $\mathbf{u} \in \mathbb{R}^3$ with unit norm $|\mathbf{u}| = 1$ and finally a transformation $\mathbf{R} \in O(3)$ such that*

$$\vec{L} = \begin{pmatrix} \eta \cosh(\rho) & -\eta \sinh(\rho)(\mathbf{u}^t \mathbf{R}) \\ -\sinh(\rho)\mathbf{u} & [\cosh(\rho)\mathbf{u} \otimes \mathbf{u} + (1 - \mathbf{u} \otimes \mathbf{u})] \mathbf{R} \end{pmatrix}$$

[...]

Proof. Let us take an element $\vec{L} \in O(1, 3)$ such that $\vec{L}^t \vec{\eta} \vec{L} = \vec{\eta}$. Then we can write this expression in $(1+3)$ notation as

$$\vec{L} = \begin{pmatrix} a & \mathbf{b}^t \\ \mathbf{c} & \mathbf{D} \end{pmatrix}, \quad a \in \mathbb{R}, \mathbf{b}, \mathbf{c} \in \mathbb{R}^3, \mathbf{D} \in \mathbb{R}(3).$$

The metric in this notation is given by

$$\vec{\eta} = \begin{pmatrix} 1 & \mathbf{0}^t \\ \mathbf{0} & -\mathbf{I}_3 \end{pmatrix}.$$

Then we can write the condition $\vec{L}^t \vec{\eta} \vec{L} = \vec{\eta}$ as

$$\begin{pmatrix} a & \mathbf{c}^t \\ \mathbf{b} & \mathbf{D}^t \end{pmatrix} \begin{pmatrix} 1 & \mathbf{0}^t \\ \mathbf{0} & -\mathbf{I}_3 \end{pmatrix} \begin{pmatrix} a & \mathbf{b}^t \\ \mathbf{c} & \mathbf{D} \end{pmatrix} = \begin{pmatrix} 1 & \mathbf{0}^t \\ \mathbf{0} & -\mathbf{I}_3 \end{pmatrix}.$$

By performing the matrix multiplication we get

$$\begin{pmatrix} a & \mathbf{c}^t \\ \mathbf{b} & \mathbf{D}^t \end{pmatrix} \begin{pmatrix} a & \mathbf{b}^t \\ -\mathbf{c} & -\mathbf{D} \end{pmatrix} = \begin{pmatrix} a^2 - \mathbf{c}^2 & a\mathbf{b}^t - \mathbf{c}^t \mathbf{D} \\ a\mathbf{b} - \mathbf{D}^t \mathbf{c} & \mathbf{D}^t \mathbf{D} - \mathbf{b} \otimes \mathbf{b} \end{pmatrix} = \begin{pmatrix} 1 & \mathbf{0}^t \\ \mathbf{0} & -\mathbf{I}_3 \end{pmatrix},$$

which is equivalent to the following system of equations: the two diagonal blocks are just the transposition of each other, so we have only three independent equations give by

$$\begin{cases} a^2 - \mathbf{c}^2 = 1, \\ a\mathbf{b} - \mathbf{D}^t \mathbf{c} = \mathbf{0}, \\ \mathbf{D}^t \mathbf{D} - \mathbf{b} \otimes \mathbf{b} = \mathbf{I}_3. \end{cases}$$

From this system of equations we can derive the following properties:

- From the first equation we have $a^2 - \mathbf{c}^2 = 1$, which implies that $a^2 \geq 1$ and thus $a \geq 1$ or $a \leq -1$. We can write $a = \eta \cosh(\rho)$ with $\eta = \pm 1 = \text{sgn}(a)$ and $\rho \geq 0$ (since $a = \text{sgn}(a)|a|$).
- From the same equation we get constraints on $\mathbf{c}$ since we have $\mathbf{c}^2 = a^2 - 1$, which implies

$$\mathbf{c}^2 = a^2 - 1 = \eta^2 \cosh^2(\rho) - 1 = \cosh^2(\rho) - 1 = \sinh^2(\rho),$$

thus we can write $\mathbf{c} = -\sinh(\rho)\mathbf{u}$ with $\mathbf{u} \in \mathbb{R}^3$ and $|\mathbf{u}| = 1$ ($\mathbf{u}$ is a unit vector expressing the direction of $\mathbf{c}$: $-\mathbf{u} = \text{vers}(\mathbf{c})$). Thus we have

$$\mathbf{c} = -\sinh(\rho)\mathbf{u}.$$

TODO: change notation: matrix in 3D is bold as vector but with capital letter.

TODO: Check the computation.

- From the second equation we have $a\mathbf{b} = \mathbf{D}^t \mathbf{c}$, which implies that

$$\mathbf{b} = \frac{1}{a} \mathbf{D}^t \mathbf{c} = -\frac{\sinh(\rho)}{\eta \cosh(\rho)} \mathbf{D}^t \mathbf{u}.$$

Now since η is a sign, we can move it up and recognize the hyperbolic tangent function, thus we can write

$$\mathbf{b} = -\eta \tanh(\rho) \mathbf{D}^t \mathbf{u}.$$

We will understand the form of $\mathbf{D}$ and then we will be able to write $\mathbf{b}$ in a more explicit form.

- From the third equation we have $\mathbf{D}^t \mathbf{D} = \mathbf{I}_3 + \mathbf{b} \otimes \mathbf{b}$, which implies that

$$\mathbf{D}^t \mathbf{D} - \tanh^2(\rho) (\mathbf{D}^t \mathbf{u}) \otimes (\mathbf{D}^t \mathbf{u}) = \mathbf{I}_3,$$

which since $(\mathbf{A}\mathbf{x}) \otimes (\mathbf{B}\mathbf{y}) = \mathbf{A}(\mathbf{x} \otimes \mathbf{y})\mathbf{B}$ we can write as

$$\mathbf{D}^t [1 - \tanh^2(\rho) \mathbf{u} \otimes \mathbf{u}] \mathbf{D} = \mathbf{I}_3.$$

Now since the identity is invertible, the other side has to be invertible as well, with a determinant given by the product of the three determinants of the terms, and since the determinant of the identity is 1, we have that each of the three determinants has to be different from zero, in particular the determinant of the middle term has to be different from zero, which implies that $\tanh^2(\rho) \neq 1$, which is true since $\rho \geq 0$ and thus $\tanh^2(\rho) < 1$. Since we have

$$\mathbf{D}^{-1} = \mathbf{D}^t [1 - \tanh^2(\rho) \mathbf{u} \otimes \mathbf{u}],$$

we have that

$$\mathbf{D}\mathbf{D}^t = [1 - \tanh^2(\rho) \mathbf{u} \otimes \mathbf{u}]^{-1}.$$

Now we have to compute the inverse of the middle term, which can be time consuming, but we can do it by knowing that $\mathbf{u}$ is a unit vector: $\mathbf{u} \otimes \mathbf{u}$ is a projector on the direction of $\mathbf{u}$, and $1 - \mathbf{u} \otimes \mathbf{u}$ is a projector on the plane orthogonal to $\mathbf{u}$. Hence we can write

$$[1 - \tanh^2(\rho) \mathbf{u} \otimes \mathbf{u}] = [(1 - \mathbf{u} \otimes \mathbf{u}) + (1 - \tanh^2(\rho)) \mathbf{u} \otimes \mathbf{u}],$$

so that we have written the term as a linear combination of two projectors, which are idempotent, and thus we can easily compute the inverse as

$$\mathbf{A} = \alpha(1 - \mathbf{u} \otimes \mathbf{u}) + \beta \mathbf{u} \otimes \mathbf{u} \implies \mathbf{A}^{-1} = \frac{1}{\alpha}(1 - \mathbf{u} \otimes \mathbf{u}) + \frac{1}{\beta} \mathbf{u} \otimes \mathbf{u},$$

since we know $\alpha, \beta \neq 0$. Thus the inverse of the middle term is given by

$$[1 - \tanh^2(\rho) \mathbf{u} \otimes \mathbf{u}]^{-1} = (1 - \mathbf{u} \otimes \mathbf{u}) + \frac{1}{1 - \tanh^2(\rho)} \mathbf{u} \otimes \mathbf{u} = (1 - \mathbf{u} \otimes \mathbf{u}) + \cosh^2(\rho) \mathbf{u} \otimes \mathbf{u},$$

where we have used the identity $1 - \tanh^2(\rho) = \frac{1}{\cosh^2(\rho)}$. Hence we have

$$\mathbf{D}\mathbf{D}^t = (1 - \mathbf{u} \otimes \mathbf{u}) + \cosh^2(\rho) \mathbf{u} \otimes \mathbf{u}.$$

It is not difficult to guess a solution for $\mathbf{D}$ that satisfies this equation: imagine a matrix which has the same structure but different coefficients, which since the projectors are idempotent, when multiplied by its transposition will give the same structure but with the coefficients squared, thus we can write

$$\mathbf{D}_o = [\cosh(\rho) \mathbf{u} \otimes \mathbf{u} + (1 - \mathbf{u} \otimes \mathbf{u})] = \mathbf{D}_o^t,$$

and if we now imagine a rotation matrix $\mathbf{R} \in \text{O}(3)$ and we write $\mathbf{D} = \mathbf{D}_o \mathbf{R}$, then we have $\mathbf{D}_o \mathbf{R} (\mathbf{D}_o \mathbf{R})^t = \mathbf{D}_o \mathbf{R} \mathbf{R}^t \mathbf{D}_o^t = \mathbf{D}_o \mathbf{D}_o^t$. Hence we have found a solution for $\mathbf{D}$ that satisfies the third equation, which is given by

$$\mathbf{D} = [\cosh(\rho) \mathbf{u} \otimes \mathbf{u} + (1 - \mathbf{u} \otimes \mathbf{u})] \mathbf{R}.$$

- Now we can write $\mathbf{b}$ in a more explicit form by substituting the expression for $\mathbf{D}$ that we have just found, thus we have

$$\mathbf{b} = -\eta \tanh(\rho) \mathbf{D}^t \mathbf{u} = -\eta \tanh(\rho) \mathbf{R}^t [\cosh(\rho) \mathbf{u} \otimes \mathbf{u} + (1 - \mathbf{u} \otimes \mathbf{u})] \mathbf{u} = -\eta \sinh(\rho) (\mathbf{R}^t \mathbf{u}),$$

where we have used the fact that the term in square brackets is a projector on the direction of $\mathbf{u}$, thus when applied to the vector $\mathbf{u}$ gives back the same vector.

The only thing we are left to do is to check that the found parameters are uniquely determined by the matrix $\vec{L}$. The parameters η and ρ are uniquely determined since the first is the sign of a while the second is given by the relation $\rho = \text{arcosh}(|a|)$. The parameter $\mathbf{u}$ is uniquely determined since it is given by the relation $\mathbf{u} = -\frac{1}{\sinh(\rho)} \mathbf{c}$, and finally the parameter $\mathbf{R}$ is uniquely determined. ...

Now if we substitute the found parameters in the expressions found before, they will be all verified and thus we can write the matrix $\vec{L}$ as

$$\vec{L} = \begin{pmatrix} \eta \cosh(\rho) & -\eta \sinh(\rho) (\mathbf{u}^t \mathbf{R}) \\ -\sinh(\rho) \mathbf{u} & [\cosh(\rho) \mathbf{u} \otimes \mathbf{u} + (1 - \mathbf{u} \otimes \mathbf{u})] \mathbf{R} \end{pmatrix},$$

which is the desired result, proving the structure theorem for the Lorentz group. $\square$

4.1.1 | Subsets of the Lorentz Group

Definition 4.1 (Lorentz boost). $\rho \in [0, \infty)$, $\mathbf{u} \in \mathbb{R}^3$ with $|\mathbf{u}| = 1$, we can define an element of the Lorentz group called *Lorentz boost* (SR) as

$$\vec{B}(\rho, \mathbf{u}) = \begin{pmatrix} \cosh(\rho) & -\sinh(\rho) \mathbf{u}^t \\ -\sinh(\rho) \mathbf{u} & [\cosh(\rho) \mathbf{u} \otimes \mathbf{u} + (1 - \mathbf{u} \otimes \mathbf{u})] \end{pmatrix}.$$

Definition 4.2 (Spatial rotation). $\mathbf{R} = \mathbf{R}_{+1}(\phi, \mathbf{n})$, such that $\phi \in [0, \pi]$ and $\mathbf{n} \in S^2$, we can define an element of the Lorentz group called *spatial rotation* as

$$\vec{R}(\phi, \mathbf{n}) = \begin{pmatrix} 1 & \mathbf{0}^t \\ \mathbf{0} & \mathbf{R}_{+1}(\phi, \mathbf{n}) \end{pmatrix}.$$

Definition 4.3 (Time reversal). Given a sign $\eta = \{-1, +1\}$, we can define an element of the Lorentz group called *time reversal* as

$$\vec{T}(\eta) = \begin{pmatrix} \eta & \mathbf{0}^t \\ \mathbf{0} & \mathbf{I}_3 \end{pmatrix},$$

for a sign $\eta = -1$ we get the time reversal transformation, while for $\eta = +1$ we get the identity transformation.

Definition 4.4 (Parity transformation). Given a sign $\epsilon = \{-1, +1\}$, we can define an element of the Lorentz group called *parity transformation* as

$$\vec{P}(\epsilon) = \begin{pmatrix} 1 & \mathbf{0}^t \\ \mathbf{0} & \epsilon \mathbf{I}_3 \end{pmatrix},$$

where $\epsilon = -1$ gives the parity transformation, while $\epsilon = +1$ gives the identity transformation.

Thus we have found four subsets of the Lorentz group, which are given by the Lorentz boosts, the spatial rotations, the time reversal and the parity transformation. There is a theorem, not difficult to prove, that states that any element of the Lorentz group can be written as a product of an element of each of these subsets:

Proposition 4.5 (Factorization theorem for $O(1,3)$). *For every $\vec{L} \in O(1,3)$ there exist two signs $\eta, \epsilon \in \{-1, +1\}$, a rapidity $\rho \in [0, \infty)$, two unit vectors $\mathbf{u}, \mathbf{n} \in S^2$ and an angle $\phi \in [0, \pi]$ such that*

$$\vec{L} = \vec{T}(\eta)\vec{B}(\rho, \mathbf{u})\vec{P}(\epsilon)\vec{R}(\phi, \mathbf{n}).$$

This is a very useful decomposition of the Lorentz group, since it allows us to understand the structure of the group and to find its representations.

Furthermore the parameters $\eta, \epsilon, \rho, \mathbf{u}, \mathbf{n}, \phi$ are uniquely determined by the matrix $\vec{L}$, with exceptions on the unit vectors: $\mathbf{u}$ asks for $\rho \neq 0$ and $\mathbf{n}$ asks for $0 < \phi < \pi$ strictly, to be uniquely determined.

Proof. future work

□

5 | ling lin

[...]

From the Heine Borel theorem we know that a subset of $\mathbb{R}$ is compact if and only if it is closed (there are no limiting points that belong to the set) and bounded. We can use this to show that the set of all subgroups of $\mathbb{R}$ is uncountable.

Example (Compactness of the unitary group). The unitary group $G = U(n)$ satisfies the following properties:

$$\sum_i u_{ij}^* u_{ik} = \delta_{jk},$$

and expanding the sum we understand that for $i = j$ we have no entries greater than 1:

$$|u_{ij}| \leq 1.$$

Furthermore if we have some matrix group defined by a set of polynomial equations, then the set of solutions is closed. Therefore $U(n)$ is a closed and bounded subset of $\mathbb{R}^{n^2}$ and thus compact.

Example (Non compact Lie Groups). $GL(n, \mathbb{R})$ or $GL(n, \mathbb{C})$, $SL(n, \mathbb{R})$, $SO(1, n-1)$ are not compact, since they are not bounded.

Definition 5.1 (Connectedness). A topological space G is *connected* if for any two points $g_1, g_2 \in G$ there exists a continuous path $\gamma : [0, 1] \rightarrow G$ such that $\gamma(0) = g_1$ and $\gamma(1) = g_2$ (it is not necessary to impose the limits on the domain of γ , but it is convenient to do so).

This curve is a continuous map, thus any function cannot vary discontinuously along it. In particular, if we have a continuous homomorphism $f : G \rightarrow H$ and G is connected, then the image of f must be connected as well. This is because the image of a continuous map is always connected.

If we start from a point where $\det(f(g)) = 1$ for some $g \in G$, then we can infer that $\det(f(g)) = 1$ for all $g \in \gamma \subset G$. This is because the determinant is a continuous function, and if it is unitary at one point, it must be unitary along the entire path connecting that point to any other point in G .

[... unitary and orthogonal groups ...]

Definition 5.2 (Simply connected Lie groups). If a group G is connected and any closed curve in G can be continuously contracted (smoothly deformed) to a point, then G is called *simply connected*.

It is easier to think about topological spaces as being made of rubber, and the continuous deformation as being a process of stretching and bending the space without tearing it. In this way, a

simply connected space is one that has no holes or loops that cannot be contracted to a point. As a visual example, the case of a torus, a can connect any two points with a continuous path, but there are loops around the hole of the torus that cannot be contracted to a point. On the other hand, a sphere is simply connected, because any loop on the sphere can be contracted to a point.

Example (Simply connectedness in Lie groups). The unitary group $U(n)$ is not simply connected, since it contains $U(1) \subset U(n)$ which is topologically a circle, and thus has loops that cannot be contracted to a point. On the other hand, the special unitary group $SU(n)$ is simply connected, because it does not contain any loops that cannot be contracted to a point.

Another example is the special orthogonal group $SO(3)$ in three dimensional space: For $n \geq 3$, $SO(n)$ is not simply connected. However, for $n = 2$, $SO(2)$ is simply connected. If we think about $SU(2)$, which is the double cover of $SO(3)$, we can see that it is simply connected: the inherited loops from $SO(3)$ can be contracted to a point in $SU(2)$ because of the double cover structure. In fact, $SU(2)$ is topologically equivalent to the 3-sphere S^3 , which is simply connected. Thus spin- $\frac{1}{2}$ objects, which are described by representations of $SU(2)$, can be thought of as living on a simply connected space, while the rotations in three-dimensional space, described by $SO(3)$, have a non-simply connected structure due to the presence of loops that cannot be contracted to a point.

[Talking about fundamental groups, we are implying that we are working with a connected manifold, since the fundamental group is defined for connected spaces. If the space is not connected, we can consider the fundamental group of each connected component separately.]

5.0.1 | Lie Algebra from Tangent Vectors

Definition 5.3 (Tangent vectors). Let $(\mathcal{M}, \{u_\alpha\})$ and a point $p \in \mathcal{M}$ in the manifold. Let us imagine a smooth curve $\gamma : (a, b) \rightarrow \mathcal{M}$ such that $\gamma(t = 0) = p$. We can define the tangent vector to the curve at the point p , $\dot{\gamma}_p$ as a differential operator acting on smooth functions $\phi \in \mathbb{C}^\infty(\mathcal{M})$ as follows:

$$\dot{\gamma}_p(\phi) = \left. \frac{d}{dt}(\phi \circ \gamma)(t) \right|_{t=t_0}.$$

A set of tangent vectors at a point p spans a m dimensional vector space ($m = \dim \mathcal{M}$), called the tangent space at p , denoted by $T_p \mathcal{M}$. The dimension of the tangent space is thus equal to the dimension of the manifold.

Let us now consider a chart $p \in U$ with coordinates $x \equiv x^i : U \rightarrow \mathbb{R}^m$. We can define a set of tangent vectors at p as follows:

$$\dot{\gamma}_p^i(\phi) = \left. \frac{d}{dt}(\phi \circ x^{-1} \circ x \circ \gamma)(t) \right|_{t=t_0} = \left. \frac{d}{dt}(\phi \circ x^{-1})(x(\gamma(t))) \right|_{t=t_0}.$$

Applying the chain rule, we can write this as:

$$\dot{\gamma}_p^i(\phi) = \left. \frac{dx^j}{dt} \frac{\partial}{\partial x^j}(\phi \circ x^{-1})(x(\gamma(t))) \right|_{t=t_0} = \left. \frac{dx^j}{dt} \right|_{t=t_0} \left. \frac{\partial}{\partial x^j}(\phi \circ x^{-1})(x) \right|_{x=x(p)}.$$

Thus we find that

$$\dot{\gamma}_p(\phi) = \sum_i \gamma^i(t_0) \frac{\partial}{\partial x^i}(\phi \circ x^{-1})(x(p)) = \sum_i \gamma^i(t_0) \frac{\partial}{\partial x^i}(\phi(p)).$$

TODO: correct the expression below, it is not correct

In a local coordinate patch thus the tangent vector can be expressed as a linear combination of the partial derivatives with respect to the coordinates, evaluated at the point p . The coefficients of this linear combination are given by the components of the curve γ at $t = t_0$. Thus the tangent space has dimension equal to the number of coordinates, which is equal to the dimension of the manifold.

Corollary 5.1. *If we have two manifolds $\mathcal{M}_1$ and $\mathcal{M}_2$ of dimensions m_1 and m_2 respectively, then the tangent space of their product is the direct sum of their tangent spaces:*

$$T_p(\mathcal{M}_1 \times \mathcal{M}_2) \cong T_p\mathcal{M}_1 \oplus T_p\mathcal{M}_2 = T_p\mathcal{M},$$

where $p = (p_1, p_2)$ is a point in the product manifold $\mathcal{M}_1 \times \mathcal{M}_2$.

Definition 5.4 (Tangent bundle). If one take the union of all the tangent spaces at all points of the manifold, we obtain the **tangent bundle** $T\mathcal{M} = \bigcup_{p \in \mathcal{M}} T_p\mathcal{M}$. The tangent bundle is itself a manifold of dimension $2m$, where m is the dimension of the original manifold $\mathcal{M}$ (we have to keep the original manifold in the definition of the tangent bundle, since the tangent spaces are defined at each point of the manifold):

$$T\mathcal{M} = \{(p, \mathbf{v}) | p \in \mathcal{M}, \mathbf{v} \in T_p\mathcal{M}\},$$

and furthermore we have a natural projection map $\pi : T\mathcal{M} \rightarrow \mathcal{M}$ defined by $\pi(p, \mathbf{v}) = p$. The tangent bundle is a vector bundle, since each fiber (the tangent space at each point) is a vector space.

Definition 5.5 (Vector fields). A **vector field** on a manifold $\mathcal{M}$ is a smooth section of the tangent bundle, that is, a smooth map $V : \mathcal{M} \rightarrow T\mathcal{M}$ such that $\pi \circ V = \text{id}_{\mathcal{M}}$. In other words, for each point $p \in \mathcal{M}$, the vector field assigns a tangent vector $V(p) \in T_p\mathcal{M}$. It is also called a *section* of the tangent bundle $T\mathcal{M}$.

Thus we associate to each point of the manifold a tangent space, and a vector field is a smooth assignment of a tangent vector to each point of the manifold. We are cutting the fibers of the bundle, and we are left with a single vector at each point of the manifold.

Since locally this vector field is a map from the manifold to the tangent space, if and only if the map is smooth, then the vector field is smooth and can be written as $\Gamma(TM)$.

$$\dot{\gamma}_p(\phi) \in T_p\mathcal{M},$$

$$\dot{\gamma}_p(\phi) = \frac{d}{dt}(\phi \circ \gamma)(t_0) = \sum_i \dot{\gamma}^i(t_0) \frac{\partial}{\partial x^i}(\phi(p)),$$

where $\partial_i = \frac{\partial}{\partial x^i}$ is the basis for the tangent space at the point p . Thus we can express the tangent vector as a linear combination of the basis vectors of the tangent space, with coefficients given by the components of the curve γ at $t = t_0$. This shows that the tangent space at each point of the manifold is a vector space, and that the dimension of the tangent space is equal to the dimension of the manifold.

The vector field is a smooth assignment of a tangent vector to each point of the manifold

$$V : \mathcal{M} \rightarrow T\mathcal{M}, \quad p \mapsto (p, V_p) \in T_p\mathcal{M},$$

and it can be thought of as a smooth map from the manifold to the tangent bundle. The vector field can be expressed in local coordinates as a linear combination of the basis vectors of the tangent space, with coefficients that are smooth functions on the manifold:

$$x^i(p) \in \mathbb{R}^n, \quad x^i(p) \mapsto (x^i(p), V^i(p)) \in T_p\mathcal{M} \equiv \mathbb{R}^n \times \mathbb{R}^n,$$

This allows us to perform calculus on the manifold using the vector fields, and to define concepts such as divergence, curl, and gradient in a coordinate-independent way. If this mapping is smooth for every choice of chart, then the vector field is smooth and can be written as $\Gamma(T\mathcal{M})$, and it forms a Lie algebra under the Lie bracket operation defined by the commutator of vector fields. Thus we managed to generalize the notion of vector fields from Euclidean space to arbitrary manifolds, and we can use this to define the Lie algebra of a Lie group as the tangent space at the identity element of the group.

We can also apply a vector field $V \in \Gamma(T\mathcal{M})$ on a function $\phi \in \mathbb{C}^\infty(\mathcal{M})$ to obtain a new function $V(\phi) : \mathcal{M} \rightarrow \mathbb{R}$ defined by

$$p \mapsto V(\phi)(p) := V_p(\phi) \in \mathbb{R}, \quad V(\phi) \in \mathbb{C}^\infty(\mathcal{M}),$$

where V_p is the tangent vector assigned to the point p by the vector field V . This allows us to define the action of a vector field on functions, and to perform calculus on the manifold using the vector fields.

If we now take two vector fields $V, W \in \Gamma(T\mathcal{M})$, we can apply the previous construction to the function $V(\phi)$ to obtain a new function $W(V(\phi)) : \mathcal{M} \rightarrow \mathbb{R}$ defined by

$$W(V(\phi))(p) := W_p(V(\phi)) \in \mathbb{R}, \quad W(V(\phi)) \in \mathbb{C}^\infty(\mathcal{M}),$$

where W_p is the tangent vector assigned to the point p by the vector field W . This can be computed as follows:

$$W_p(V(\phi)) = \sum_i W_p^i \partial_i (V(\phi))(p) = \sum_{ij} W_p^i \partial_i (V^j(p) \partial_j (\phi \circ x^{-1}))(x(p)),$$

where we have used the local coordinate expression for the vector fields; we can continue to compute this expression by applying the product rule for differentiation:

$$W_p(V(\phi)) = \sum_{ij} W_p^i \partial_i (V^j(p)) \partial_j (\phi \circ x^{-1})(x(p)) + \sum_{ij} W_p^i V_p^j \partial_i \partial_j (\phi \circ x^{-1})(x(p)).$$

Note that a second derivative term has appeared in the expression, which we do not now how to interpret in terms of the original vector fields V and W : this new term is not a vector anymore, it is not an object behaving as a vector field $X(\phi)$ with $X \in \Gamma(T\mathcal{M})$. We cannot interpret this result as another vector field operating on the same function ϕ , since the second derivative term does not have the same structure as the original vector fields.

Definition 5.6 (Lie bracket of vector fields). The Lie bracket of two vector fields (or commutator) $V, W \in \Gamma(T\mathcal{M})$ is defined as the new vector field $[V, W] \in \Gamma(T\mathcal{M})$ such that

$$[V, W](\phi) := V(W(\phi)) - W(V(\phi)),$$

for all smooth functions $\phi \in \mathbb{C}^\infty(\mathcal{M})$. This new vector field is defined by the difference of the two second derivative terms that appeared in the previous expression, and it can be shown that it satisfies the properties of a Lie bracket, as we will see.

With this new object defined, we have an operation that takes two vector fields and produces a new vector field, with in charts can be written as

$$[V, W]^j = \sum_i (V^i \partial_i W^j - W^i \partial_i V^j) \partial_j.$$

Proposition 5.2. *For any three vector fields $X, Y, Z \in \Gamma(TM)$, the Lie bracket satisfies the following properties:*

- **Antisymmetry:** $[X, Y] = -[Y, X]$.
- **Bilinearity:** $[aX + bY, Z] = a[X, Z] + b[Y, Z]$ and $[Z, aX + bY] = a[Z, X] + b[Z, Y]$ for all scalars $a, b \in \mathbb{R}$.
- **Jacobi identity:** $[X, [Y, Z]] + [Y, [Z, X]] + [Z, [X, Y]] = 0$.

Corollary 5.3. *With the previously defined Lie bracket, the set of vector fields $\Gamma(TM)$ can be endowed with the structure of a non-associative algebra over $\mathbb{R}$. We said non-associative because the Lie bracket does not satisfy the associative property, but it satisfies the Jacobi identity instead:*

$$[[X, Y], Z] \neq [X, [Y, Z]].$$

The Jacobi identity is a weaker condition than associativity, and it ensures that the Lie bracket defines a consistent algebraic structure on the space of vector fields.

It is now time to connect this construction to the Lie algebra of a Lie group.

Definition 5.7. Given a smooth map $f : \mathcal{M} \rightarrow \mathcal{N}$ between two manifolds, we can define an induced map (which will be identified with a *vector space homomorphism* for every point in the manifold) on the tangent spaces, called the **pushforward** of f , denoted by $(f_*)_p : T_p \mathcal{M} \rightarrow T_{f(p)} \mathcal{N}$. The pushforward is defined by

$$(f_*)_p(\dot{\gamma}_p) = (f \circ \gamma)_{f(p)} \quad \forall p \in \mathcal{M},$$

then let $h \in \mathbb{C}^\infty(\mathcal{N})$, and $v = \dot{\gamma}_p \in T_p \mathcal{M}$, then we have

$$\implies [(f_*)_p(v)](h) = v(h \circ f).$$

In coordinates x for $\mathcal{M}$ and y for $\mathcal{N}$, we can write the pushforward as

$$[(f_*)_p]^i_k = \partial_k (y^i \circ f \circ x^{-1})(x(p)) = (df)^i_k(x(p)),$$

where df is the Jacobian matrix of the map f at the point p : $f^i(x^k)$. This is just the jacobian matrix of a vector-valued function, which is a linear map that takes a tangent vector at p and maps it to a tangent vector at $f(p)$ by applying the derivative of the function f to the components of the tangent vector.

This concept is known in the field of general relativity as the **Lie dragging** of a vector field along a curve, and it is used to describe how vector fields change as they are transported along curves in a manifold. The pushforward allows us to compare vector fields at different points in the manifold, and to understand how they transform under smooth maps between manifolds.

If the function f is a diffeomorphism (smooth and invertible), then the pushforward is an isomorphism between the tangent spaces at p and $f(p)$, and it allows us to transfer the structure of the tangent space from one point to another. This is particularly useful when we want to define the Lie algebra of a Lie group, as we will see successively.

Proposition 5.4. *If $f : \mathcal{M} \rightarrow \mathcal{N}$ is a diffeomorphism, then the pushforward $(f_*)_p$ is an isomorphism between vector fields on $\mathcal{M}$ and vector fields on $\mathcal{N}$:*

$$f_* : \Gamma(T\mathcal{M}) \rightarrow \Gamma(T\mathcal{N}), \quad V \mapsto f_*(V),$$

$$(f_*(V))_{f(p)} = (f_*)_p(V_p) \in T_{f(p)}\mathcal{N},$$

and since consequently the jacobian matrix of f is invertible, we can also define the inverse pushforward $(f_*^{-1})_{f(p)} : T_{f(p)}\mathcal{N} \rightarrow T_p\mathcal{M}$ as the inverse of the pushforward, and it is also an isomorphism between vector fields on $\mathcal{N}$ and vector fields on $\mathcal{M}$.

We can now imagine to concatenate functions; let $f_1 \in \mathbb{C}^\infty(\mathcal{M}, \mathcal{N})$ and $f_2 \in \mathbb{C}^\infty(\mathcal{N}, \mathcal{Q})$ be two diffeomorphisms, then their composition $f_2 \circ f_1 : \mathcal{M} \rightarrow \mathcal{Q}$ is also a diffeomorphism, and we can define the pushforward of the composition as follows:

$$\forall p \in \mathcal{M}, \quad [(f_2 \circ f_1)_*]_p = [(f_2)_*]_{f_1(p)} \circ [(f_1)_*]_p : T_p\mathcal{M} \rightarrow T_{f_2(f_1(p))}\mathcal{Q},$$

which can be read as the pushforward of the composition is the composition of the pushforwards. This property is known as the **functoriality** of the pushforward, and it ensures that the pushforward behaves well under composition of functions, which is a fundamental property for any mathematical construction that aims to capture the structure of smooth maps between manifolds.

Now we are ready to define the Lie algebra of a Lie group as the tangent space at the identity element of the group, and to show how the Lie bracket of vector fields induces a Lie bracket on this tangent space, which will give us the structure of a Lie algebra on the tangent space at the identity element. This is a crucial step in understanding the relationship between Lie groups and their associated Lie algebras, and it will allow us to study the properties of Lie groups using the tools of Lie algebras.

Definition 5.8. Let G be a Lie group, and let e be the identity element of G . Then any $a \in G$ defines a “left-translation” which maps any element $g \in G$ to ag :

$$L_a : G \rightarrow G, \quad g \mapsto L_ag = ag.$$

This left-translation is a diffeomorphism, since it is smooth and has a smooth inverse given by $(L_a)^{-1} = L_{a^{-1}}$. Left translation are natural diffeomorphisms of the Lie group, and they allow us to transfer the structure of the tangent space at the identity element to any other point in the group. In particular, we can use left translations to define a vector field on the Lie group by translating a tangent vector at the identity element to every other point in the group.

Definition 5.9. Let us consider a class of vector fields $V \in \Gamma(TG)$ on a Lie group G that are invariant under left translations; they are called **left-invariant vector fields**, and they satisfy the following property:

$$\forall a \in G, \quad (L_a)_*V = V,$$

which means that the vector field is unchanged under left translations. In other words, for any point $g \in G$, the value of the vector field

$$\forall a, g \in G, \quad [(L_a)_*V]_{ag} = [(L_a)_*]_g(V_g) = V_{ag},$$

which reads as the value of the vector field at the point ag is equal to the pushforward of the vector field at the point g under the left translation by a . This property ensures that the vector field is consistent with the group structure of G .

Lemma 5.5. *Given a tangent vector $v \in T_e G$ (any point would have been fine, so we chose the identity element for convenience), there exists a unique left-invariant vector field $V^{(v)} \in \Gamma(TG)$ by translating v to every point in the group using left translations, which satisfies the initial conditions*

$$V_e^{(v)} = v.$$

Since it is a left-invariant vector field, for every point $g \in G$ we can use left translation to define the value of the vector field at that point.

On a lie group we have a one to one correspondence between tangent vectors at a point (the identity element for convenience) and left-invariant vector fields, which allows us to transfer the structure of the tangent space at the identity element to the entire group

$$v \in T_e G, \quad v \mapsto V^{(v)} \in \Gamma(TG), \quad (V^{(v)})_g = (L_g)_*(v) \in T_g G.$$

The other ingredient we need to define the Lie algebra of a Lie group is the closeness of the Lie bracket of vector fields, which we have already defined.

Proposition 5.6. *The set of left-invariant vector fields on a Lie group G is closed under the Lie bracket operation $[\cdot, \cdot]$, which means that if $V, W \in \Gamma(TG)$ are left-invariant vector fields, then*

$$[V, W] \in \Gamma(TG) \text{ is left-invariant.}$$

In order to prove this proposition, we need to show another final lemma:

Lemma 5.7. *Let $f : \mathcal{M} \rightarrow \mathcal{N}$ surjective and $\phi \in \mathbb{C}^\infty(\mathcal{N})$ (so that $(\phi \circ f) \in \mathbb{C}^\infty(\mathcal{M})$) and $V \in \Gamma(T\mathcal{M})$ be a vector field on $\mathcal{M}$ such that $(f_*)(V)$ is a well defined vector field on $\mathcal{N}$, then we have*

$$V(\phi \circ f) = (f_*(V))(\phi) \circ f \iff V(\phi \circ f) = (f_*(V))_{f(p)}(\phi) = ((f_*)(V)(\phi) \circ f)(p).$$

This means that ...

Proof. Let $\phi \in \mathbb{C}^\infty(G)$. Then, for any $a, g \in G$ we have

$$\begin{aligned} [(L_a)_*[V, W]]_{ag}(\phi) &= \dots \\ &= [V, W]_g(\phi \circ L_a) = V_g(W(\phi \circ L_a)) - W_g(V(\phi \circ L_a)), \end{aligned}$$

and since we have the previous lemma, we can write for a generic vector field U on G

$$[(L_a)_*U]_{ag} = U_{ag} \implies U(\phi \circ L_a)(g) = [(L_a)_*U]_{ag}(\phi) = U_{ag}(\phi) = U(\phi)(L_a(g)) = (U(\phi) \circ L_a)(g).$$

Thus we can write

$$\begin{aligned} ((L_a)_*[V, W])_{ag}(\phi) &= V_g(W(\phi \circ L_a)) - W_g(V(\phi \circ L_a)) \\ &= (((L_a)_*)_g V)_g(W(\phi)) - (((L_a)_*)_g W)_g(V(\phi)) \\ &= [(L_a)_*V]_{ag}(W(\phi)) - [(L_a)_*W]_{ag}(V(\phi)) \\ &= V_{ag}(W(\phi)) - W_{ag}(V(\phi)) = [V, W]_{ag}(\phi), \end{aligned}$$

since both V and W are left-invariant vector fields, and thus we have shown that $[(L_a)_*[V, W]]_{ag}(\phi) = [V, W]_{ag}(\phi)$ for all $a, g \in G$, which means that $[V, W]$ is left-invariant. $\square$

Thus we have shown that the set of left-invariant vector fields on a Lie group G is closed under the Lie bracket operation, which means that we can define a Lie algebra structure on the tangent space at the identity element of the group by using the Lie bracket of left-invariant vector fields. This is a crucial step in understanding the relationship between Lie groups and their associated Lie algebras.

Definition 5.10. Let G be a Lie group, and let e be the identity element of G . The **Lie algebra** of G (or *associated to* G), denoted by $\mathfrak{g}$, is defined as the $\mathbb{R}$ -vector space $\mathfrak{g} = T_e G$ with the bilinear product given by the Lie bracket of left-invariant vector fields:

$$[\cdot, \cdot]_{\mathfrak{g}} : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}, \quad (v, w) \mapsto [V^{(v)}, W^{(w)}]_{\mathfrak{g}} := [V^{(v)}, W^{(w)}]_e,$$

where $(\mathfrak{g}, [\cdot, \cdot]_{\mathfrak{g}})$ satisfies antisymmetry, bilinearity, and the Jacobi identity, and thus it is abstractly defining a Lie algebra.

Definition 5.11. A vector space basis $\{T_i\}_{i=1, \dots, n}$ with $n = \dim \mathfrak{g}$ of the Lie algebra $\mathfrak{g}$ is called a **basis of generators** of the Lie algebra. Since the Lie brackets of elements of the Lie algebra can be expressed as linear combinations of the basis elements (since the algebra is closed under the Lie bracket), we can write

$$[T_i, T_j] = \sum_{k=1}^n c_{ij}^k T_k,$$

where c_{ij}^k are the **structure constants** of the Lie algebra $\mathfrak{g}$ with the basis of generators $\{T_i\}$.

We can highlight some properties of the structure constants c_{ij}^k that follow from the properties of the Lie bracket:

- They are antisymmetric in the first two (lower) indices: $c_{ij}^k = -c_{ji}^k$, which follows from the antisymmetry of the Lie bracket.
- They satisfy the Jacobi identity: $\sum_{l=1}^n (c_{ij}^l c_{lk}^m + c_{jk}^l c_{li}^m + c_{ki}^l c_{lj}^m) = 0$ for all i, j, k, m , which follows from the Jacobi identity of the Lie bracket.

These comes directly from the properties of the Lie bracket and thus their application onto the generators of the Lie algebra. Furthermore an abstract Lie algebra can be defined by specifying a choice of basis $\{T_i\}$ along with a set of structure constants c_{ij}^k that satisfy the above properties, without any reference to a specific Lie group.

As an easy example, we can consider the tensor ϵ_{ij}^k and check that it respects the above properties: along with a good choice of basis, it can be used to define the structure constants of the Lie algebra $\mathfrak{su}(2)$ associated to the Lie group $SU(2)$.

We used real coefficients for the Lie algebra, but we could have used complex coefficients as well, and thus we would have defined a complex Lie algebra. The choice of real or complex coefficients depends on the context and the applications we have in mind, and it does not affect the general properties of the Lie algebra. The extension and generalization is straightforward, and it is just a matter of changing the field over which the vector space is defined.

5.1 | Lie Algebra of Matrix Groups

If we consider matrices $G \subseteq \text{GL}(V)$, with V ($n = \dim V$) a vector space over $\mathbb{R}$ or $\mathbb{C}$ (with a proper basis chosen), then $\text{GL}(V) \simeq \text{GL}(n, \mathbb{R})$ or $\text{GL}(n, \mathbb{C})$, respectively (for simplicity let us stick to the real case, the complex case is just a straightforward generalization).

Taking than any element $g \in G \subseteq \text{GL}(n, \mathbb{R})$ can be thought of as an $n \times n$ matrix

$$g = (g^{AB})_{AB}, \quad g^{AB} : G \rightarrow \mathbb{R}, \quad g \rightarrow g^{AB},$$

for any fixed pair of indices A, B . Thus we can think of the elements of the group as being labeled by a set of real numbers, which are the entries of the matrix.

Definition 5.12. Let $G \subset \text{GL}(n, \mathbb{R})$ be a matrix group. Let $v \in T_k G$ and $(U, x \equiv x^k)$, such that a test function $\phi \in \mathbb{C}^\infty(G)$ with $k = 1, \dots, \dim G \neq n^2$ and a point $h \in U$, then

$$\begin{aligned} v(\phi) &= \sum_k v^k \partial_k \phi \Big|_{x(h)} \\ &= \sum_{k,A,B} v^k \frac{\partial g^{AB}}{\partial x^k} \Big|_{x(h)} \frac{\partial \phi}{\partial g^{AB}} \\ &= \sum_{A,B} v^{AB} \frac{\partial \phi}{\partial g^{AB}}, \end{aligned}$$

where we can see how the summation over k defines a sort of *tangent matrix*

$$v^{AB} := \sum_k v^k \frac{\partial g^{AB}}{\partial x^k} \Big|_{x(h)}, \quad \frac{d}{dt}(\gamma^{AB}(t)) \Big|_{t=t_0} = v^{AB},$$

which is the tangent vector to the curve $\gamma^{AB}(t)$ in the space of matrices, and it can be thought of as an element of the Lie algebra associated to the matrix group G . Thus we can identify the tangent space at the identity element of the group with a subspace of the space of matrices, and this subspace is precisely the Lie algebra of the group.

Definition 5.13. Let $G \subset \text{GL}(n, \mathbb{R}) \subset \mathbb{R}^{n \times n}$ be a matrix group, where the groups from left to right are included injectively. Then there exists a $g \in G$ such that $g \mapsto (g^{AB}) \in \mathbb{R}^{n^2}$ is a smooth and injective map, and thus we can identify the elements of the group with points in $\mathbb{R}^{n^2}$.

Let now $v \in T_h G$ then $(g^{AB})_*(v) \in T_{h^{AB}} \mathbb{R}^{n \times n} \simeq \mathbb{R}^{n^2}$ is injective. We will denote $(g^{AB})_*(v) = v^{AB}$ is the tangent matrix:

$$\{v^{AB}\} \subseteq T_{h^{AB}} \mathbb{R}^{n \times n} \simeq \mathbb{R}^{n^2}.$$

For $v = \dot{\gamma}(t_0)$ with $\gamma : \mathbb{R} \rightarrow G$ a curve with $\gamma(t_0) = h$, the tangent vector v corresponds to a tangent matrix v^{AB} in $\mathbb{R}^{n^2}$. We have $\forall \phi \in \mathbb{C}^\infty(G)$,

$$[(g^{AB})_*(v)](\phi) = \frac{d}{dt}(\phi \circ g^{AB} \circ \gamma)(t_0) = \frac{d}{dt}(g^{AB} \circ \gamma)(t_0) \frac{\partial \phi}{\partial g^{AB}},$$

which implies

$$v^{AB} = \frac{d}{dt}(g^{AB} \circ \gamma)(t_0).$$

[...]

Example (General Linear group). Let us consider $G = \mathrm{GL}(n, \mathbb{R}) = \{M \in \mathbb{R}^{n \times n} : \det(M) \neq 0\}$, we are interested in finding the Lie algebra of the group as the tangent space at the identity element $e = \mathbb{I}_n$ of the group. We can identify the elements of the group with points in $\mathbb{R}^{n^2}$ by mapping each matrix M to its entries M^{AB} . Locally, around the identity element, the tangent space can be identified with the space of all $n \times n$ matrices:

$$T_e G \simeq \mathfrak{g} = T_e \mathbb{R}^{n \times n} \simeq \mathbb{R}^{n^2},$$

and since the dimension of the algebra, the group, the manifold etc. coincides, we know it to be of dimension n^2 . When passing to the complex counterpart, we have $G = \mathrm{GL}(n, \mathbb{C})$ and the Lie algebra is $\mathfrak{g} = T_e G \simeq \mathbb{C}^{n^2}$, which is of dimension $2n^2$ as a real vector space.

Example (Special Linear group). If we consider the special linear group $G = \mathrm{SL}(n, \mathbb{R}) = \{M \in \mathrm{GL}(n, \mathbb{R}) : \det(M) = 1\}$, we can identify the elements of the group with points in $\mathbb{R}^{n^2}$ by mapping each matrix M to its entries M^{AB} . Let $M(t) : \mathbb{R} \rightarrow G$ be a curve in the group with $M(0) = e = \mathbb{I}_n$. Then the tangent vector at the identity is given by

$$v = \left. \frac{d}{dt} M(t) \right|_{t=0}.$$

Since $M(t)$ is a matrix in $\mathrm{SL}(n, \mathbb{R})$, we have $\det(M(t)) = 1$ for all t . Taking the derivative of this condition with respect to t and evaluating at $t = 0$ gives

$$\left. \frac{d}{dt} \det(M(t)) \right|_{t=0} = 0.$$

Using the formula for the derivative of the determinant

$$\left. \frac{d}{dt} \det(M(t)) \right|_{t=0} = \mathrm{Tr} [\det(M(t))(M^{-1}(t))^T \dot{M}(t)]_{t=0} = \mathrm{Tr}(\dot{M}(0)) = \mathrm{Tr}(v),$$

this implies

$$\mathrm{tr}(v) = 0,$$

which means that the tangent vector v (derivative of an element) at the identity element must be a traceless matrix. Therefore, the Lie algebra of the special linear group is the set of all traceless $n \times n$ matrices, which can be identified with a subspace of $\mathbb{R}^{n^2}$:

$$\mathfrak{sl}(n, \mathbb{R}) = T_e G \simeq \{X \in \mathbb{R}^{n \times n} : \mathrm{tr}(X) = 0\}.$$

The dimension of this Lie algebra is $n^2 - 1$, since the condition of being traceless reduces the number of independent parameters by one (but is the same as the original group, since there we had the condition on the determinant). In the complex case, we have $G = \mathrm{SL}(n, \mathbb{C})$ and the Lie algebra is $\mathfrak{sl}(n, \mathbb{C}) = T_e G \simeq \{X \in \mathbb{C}^{n \times n} : \mathrm{tr}(X) = 0\}$, which is of dimension $2(n^2 - 1)$ as a real vector space.

Example (Orthogonal group). Consider the orthogonal group $G = \mathrm{O}(n, \mathbb{R}) = \{M \in \mathbb{R}^{n \times n} : M^T M = \mathbb{I}_n\}$. To find its Lie algebra, we look at the tangent space at the identity element $e = \mathbb{I}_n$. A curve $M(t) : \mathbb{R} \rightarrow G$ with $M(0) = \mathbb{I}_n$ satisfies $M(t)^T M(t) = \mathbb{I}_n$ for all t . Differentiating this condition with respect to t and evaluating at $t = 0$ gives

$$\left. \frac{d}{dt} (M(t)^T M(t)) \right|_{t=0} = 0.$$

Using the product rule,

$$\dot{M}(0)^T \mathbb{I}_n + \mathbb{I}_n^T \dot{M}(0) = 0,$$

which simplifies to

$$\dot{M}(0)^T + \dot{M}(0) = 0.$$

This means that the tangent vector $v = \dot{M}(0)$ must be skew-symmetric. Therefore, the Lie algebra of the orthogonal group is the set of all skew-symmetric $n \times n$ matrices:

$$\mathfrak{o}(n, \mathbb{R}) = T_e G \simeq \{X \in \mathbb{R}^{n \times n} : X^T + X = 0\}.$$

The dimension of this Lie algebra is $n(n-1)/2$, as there are $n(n-1)/2$ independent parameters for a skew-symmetric matrix.

Example (Special Orthogonal group). For the special orthogonal group $G = \mathrm{SO}(n, \mathbb{R}) = \{M \in \mathrm{O}(n, \mathbb{R}) : \det(M) = 1\}$, the Lie algebra is the same as that of the orthogonal group, since the condition of having determinant equal to one does not impose any additional constraints on the tangent vectors at the identity element. Therefore, we have

$$\mathfrak{so}(n, \mathbb{R}) = T_e G \simeq \{X \in \mathbb{R}^{n \times n} : X^T + X = 0\},$$

which is of dimension $n(n-1)/2$.

We can study this group as the intersection of the special linear group and the orthogonal group, since it is defined by the conditions of being both orthogonal and having determinant equal to one. Thus we can write

$$\mathrm{SO}(n, \mathbb{R}) = \mathrm{SL}(n, \mathbb{R}) \cap \mathrm{O}(n, \mathbb{R}),$$

and consequently we can write the Lie algebra as the intersection of the Lie algebras of the special linear group and the orthogonal group:

$$\mathfrak{so}(n, \mathbb{R}) = \mathfrak{sl}(n, \mathbb{R}) \cap \mathfrak{o}(n, \mathbb{R}) = \{X \in \mathbb{R}^{n \times n} : X^T + X = 0, \mathrm{tr}(X) = 0\}.$$

But since the condition of being skew-symmetric already implies that the trace is zero, we can simplify this to

$$\mathfrak{so}(n, \mathbb{R}) = \{X \in \mathbb{R}^{n \times n} : X^T + X = 0\} = \mathfrak{o}(n, \mathbb{R}),$$

which is the same as the Lie algebra of the orthogonal group.

Example (Unitary group). For the unitary group $G = \mathrm{U}(n, \mathbb{C}) = \{U \in \mathbb{C}^{n \times n} : U^\dagger U = \mathbb{I}_n\}$, we can find its Lie algebra by looking at the tangent space at the identity element $e = \mathbb{I}_n$. A curve $U(t) : \mathbb{R} \rightarrow G$ with $U(0) = \mathbb{I}_n$ satisfies $U(t)^\dagger U(t) = \mathbb{I}_n$ for all t . Differentiating this condition with respect to t and evaluating at $t = 0$ gives

$$\left. \frac{d}{dt} (U(t)^\dagger U(t)) \right|_{t=0} = 0.$$

Using the product rule,

$$\dot{U}(0)^\dagger \mathbb{I}_n + \mathbb{I}_n^\dagger \dot{U}(0) = 0,$$

which simplifies to

$$\dot{U}(0)^\dagger + \dot{U}(0) = 0.$$

This means that the tangent vector $v = \dot{U}(0)$ must be anti-Hermitian. Therefore, the Lie algebra of the unitary group is the set of all anti-Hermitian $n \times n$ matrices:

$$\mathfrak{u}(n, \mathbb{C}) = T_e G \simeq \{X \in \mathbb{C}^{n \times n} : X^\dagger + X = 0\}.$$

The real dimension of this Lie algebra is n^2 , as there are n^2 independent parameters for an anti-Hermitian matrix: the upper half of the matrix has $\frac{n(n-1)}{2}$ independent parameters, and the diagonal elements (all purely imaginary since they have to be changed in sign when daggered) have n independent parameters, for a total of $2\frac{n(n-1)}{2} + n = n^2$ independent parameters.

Example (Special Unitary group). For the special unitary group $G = \mathrm{SU}(n, \mathbb{C}) = \{U \in \mathrm{U}(n, \mathbb{C}) : \det(U) = 1\}$, the Lie algebra is the set of all anti-Hermitian $n \times n$ matrices with trace zero:

$$\mathfrak{su}(n, \mathbb{C}) = T_e G \simeq \{X \in \mathbb{C}^{n \times n} : X^\dagger + X = 0, \mathrm{tr}(X) = 0\}.$$

The real dimension of this Lie algebra is $n^2 - 1$, as the condition of being traceless reduces the number of independent parameters by one.

We are computing the **real dimension** of the lie algebra even in the complex case, since we are considering the Lie algebra as a real vector space. If we were to consider it as a complex vector space, we would have to consider that when the coefficients are complex, the dimension would be halved, since each complex parameter corresponds to two real parameters (the real and imaginary parts). Thus, when we have an odd number of independent parameters, the complex dimension would be a half-integer, which is not possible for a vector space. Therefore, we consider the Lie algebra as a real vector space, even in the complex case, to ensure that the dimension is always an integer.

We want to impose only **holomorphic functions** on the complex Lie groups, since we want to preserve the complex structure of the group and its Lie algebra. If we were to allow non-holomorphic functions, we would lose the complex structure and we would not be able to define a complex Lie algebra. The holomorphic functions (for example the determinant condition, which is just a polynomial equation in complex variables) ensure that the Lie algebra is closed under the Lie bracket and that it has a well-defined complex structure, which is essential for many applications in mathematics and physics.

Lemma 5.8. *Let G be a matrix Lie group, and let $v \in T_h G$ be a tangent vector in h and consider $a \in G$; then the pushforward of v under the left translation by a is given by*

$$[(L_a)_*]_h(v)^{AB} = \sum_{C=1}^n a^{AC} v^{CB} = (a \cdot v)^{AB},$$

where the last equality makes sense only in a matrix group, since we are using the matrix multiplication to define the action of the group on the tangent vector. This shows how the pushforward of a tangent vector under left translation can be expressed in terms of the matrix multiplication of the group element and the tangent vector, which is a key property that allows us to understand how the Lie algebra is related to the structure of the Lie group.

Proof. We can use a test function $\phi \in \mathbb{C}^\infty(G)$ to compute the pushforward of v under left translation by a :

$$[(L_a)_*]_h(v)(\phi) = v(\phi \circ L_a) = \sum_k v^k \partial_k (\phi \circ L_a)|_{x(h)} \equiv v(\tilde{\phi}),$$

where we defined $\tilde{\phi} := \phi(ag)$. Now we can express the function $\tilde{\phi}$ in terms of the matrix entries g^{AB} if we compute

$$[(L_a)_*]_h(v)^{AB} = v^{AB} \frac{\partial \phi}{\partial g^{PQ}} \bigg|_{g=ah} \frac{\partial}{\partial g^{AB}} (ag)^{PQ} \bigg|_{g=h},$$

which can be rewritten as

$$= v^{AB} \frac{\partial \phi}{\partial g^{PQ}} \Big|_{g=ah} a^{PR} \delta^{RA} \delta^{QB} = a^{PR} v^{RQ} \frac{\partial \phi}{\partial g^{PQ}} = (a \cdot v)^{AB} \frac{\partial \phi}{\partial g^{AB}},$$

[...]

□

Proposition 5.9. *Let G be the matrix group and $v, w \in T_e G \implies \mathfrak{g}$, then we can associate to v and w the left-invariant vector fields $V^{(v)}$ and $W^{(w)}$, and we can compute the Lie bracket of these vector fields at the identity element:*

$$[V^{(v)}, W^{(w)}]_e = [v, w]_{\mathfrak{g}},$$

where the Lie bracket on the right-hand side is defined by the commutator of the corresponding tangent matrices:

$$[v, w]_{\mathfrak{g}} = (vw)^{AB} - (wv)^{AB} = v^{AC} w^{CB} - w^{AC} v^{CB},$$

which is the standard Lie bracket for matrix Lie algebras and makes sense in the context of matrix groups, since the tangent vectors can be identified with matrices and the Lie bracket can be defined as the commutator of these matrices.

We are not going to prove this proposition, since it is a straightforward computation that follows from the definition of the Lie bracket of vector fields and the properties of the pushforward under left translations. The key point is that the left-invariant vector fields are defined by translating the tangent vectors at the identity element to every other point in the group using left translations, and thus their Lie bracket can be computed by using the properties of the pushforward under left translations and the definition of the Lie bracket of vector fields.

5.2 | Exponential Map

We are now going to define the exponential map for a Lie group, which is a crucial tool for understanding the relationship between the Lie algebra and the Lie group. The exponential map allows us to relate elements of the Lie algebra (tangent vectors at the identity element) to elements of the Lie group, and it provides a way to construct curves in the Lie group from tangent vectors in the Lie algebra. It is the tool which let us take a finite transformation from an infinitesimal one, and thus it is fundamental for applications in physics, where we often deal with infinitesimal transformations and we want to understand how they relate to finite transformations.

Definition 5.14 (Integral curve). Let $V \in \Gamma(TM)$ be a vector field and let $\gamma : (a, b) \rightarrow \mathcal{M}$ be an integral curve for V , i.e., a smooth curve whose tangent vector at each point $\gamma(t)$ is equal to the value of the vector field at that point $V_{\gamma(t)}$, i.e.

$$\dot{\gamma}(t) = V_{\gamma(t)} \quad \forall t \in (a, b).$$

If we use a chart (U, x) around a point $p \in \mathcal{M}$ such that $\gamma(t_0) = p$ for some $t_0 \in (a, b)$, then we can write the integral curve in coordinates as

$$\dot{\gamma}^i(t) = \frac{d}{dt}\gamma^i(t) = V^i(\gamma(t)) \quad \forall t \in (a, b), \quad (\text{ODE}),$$

which is a system of ordinary differential equations (ODEs) that can be solved to find the integral curve $\gamma(t)$ given the vector field V and the initial condition $\gamma(t_0) = p$.

The existence and uniqueness of solutions to this ODE is guaranteed by the standard theory of ODEs, and thus we can always find an integral curve for any given vector field and initial condition.

Definition 5.15 (One-parameter subgroup). Let a smooth curve $\sigma : \mathbb{R} \rightarrow G$ in a Lie group G is called a **one-parameter subgroup** of G if it satisfies the group homomorphism property

$$\sigma(s + t) = \sigma(s)\sigma(t) \quad \forall s, t \in \mathbb{R},$$

with $\sigma(0) = e$ is the identity element of the Lie group G .

This last definition means that a one-parameter subgroup is a curve in the Lie group that respects the group structure, i.e., the value of the curve at the sum of two parameters is equal to the product (group operation) of the values of the curve at each parameter. This property is crucial for understanding how one-parameter subgroups relate to the Lie algebra and how they can be used to construct finite transformations from infinitesimal ones.

We will adopt the notation in which

$$\sigma \equiv \{\sigma(t) \mid t \in \mathbb{R}\},$$

as a subgroup of G generated by the one-parameter subgroup σ . This notation emphasizes the fact that the one-parameter subgroup generates a subgroup of G that is isomorphic to $\mathbb{R}$ under addition, and it allows us to think of the one-parameter subgroup as a continuous family of group elements parameterized by a real variable t .

Proposition 5.10. *Let G be a Lie group and let $V \in \Gamma(TG)$ be a left-invariant vector field on G . Then the integral curve $\sigma : \mathbb{R} \rightarrow G$ of V with initial condition $\sigma(0) = e$ is a one-parameter subgroup of G .*

The idea here is that by taking the integral curve of a left-invariant vector field starting at the identity element, we obtain a curve that respects the group structure and thus defines a one-parameter subgroup. The left-invariance of the vector field ensures that the flow generated by the vector field is compatible with the group operation, which is essential for the resulting curve to be a one-parameter subgroup.

Proposition 5.11. *Let σ be a one-parameter subgroup of a Lie group G . Then the tangent vector $V_{\sigma(t)} = \frac{d}{dt}\sigma(t)$ to the curve $\sigma(t)$ at any point t can be computed as the pushforward of the tangent vector at the identity element V_e under the left translation by $\sigma(t)$:*

$$V_{\sigma(t)} = [(L_{\sigma(t)})_*]_e(\dot{\sigma}(0)),$$

which implies that $V_{\sigma(t)}$ is a left-invariant vector field along the curve $\sigma(t)$, and thus the one-parameter subgroup σ is generated by the left-invariant vector field V that corresponds to the tangent vector at the identity element.

Thus every one-parameter subgroup of a Lie group is generated by a left-invariant vector field, and conversely, every left-invariant vector field generates a one-parameter subgroup. This establishes a deep connection between the Lie algebra (the space of left-invariant vector fields) and the structure of the Lie group (the one-parameter subgroups), which is fundamental for understanding how the Lie algebra encodes information about the local structure of the Lie group and how it can be used to construct finite transformations from infinitesimal ones.

It is a one-to-one correspondence between the elements of the Lie algebra and the one-parameter subgroups of the Lie group, which is a key result in the theory of Lie groups and Lie algebras.

Corollary 5.12. *Suppose to have two different one-parameter subgroups $\sigma(t)$ and $\mu(t)$ of a Lie group G with $\dot{\sigma}(0) = \dot{\mu}(0) = v \in T_e G$. Then $\sigma(t) = \mu(t)$ for all $t \in \mathbb{R}$.*

This corollary can be read as saying that the one-parameter subgroup generated by a given tangent vector at the identity element is unique. This emphasizes the one-to-one correspondence between the elements of the Lie algebra and the one-parameter subgroups of the Lie group, and it ensures that each element of the Lie algebra generates a unique one-parameter subgroup.

Definition 5.16. Let $X \in \mathfrak{g}$ be an element of the Lie algebra of a Lie group G . We denote as σ_X the unique one-parameter subgroup of G constructed as follows:

1. $X \in \mathfrak{g} \iff V^{(X)} \in \Gamma(TG)$ is left-invariant vector field on G such that $V_e^{(X)} = X$.
2. $\sigma_X : \mathbb{R} \rightarrow G$ is the unique integral curve of $V^{(X)}$ with initial condition $\sigma_X(0) = e$ (passing through the identity element) with $\dot{\sigma}_X(0) = V_e^{(X)} = X$; it is a one-parameter subgroup of G .

Since along the curve $\sigma_X(t)$ we have a possible mapping from the Lie algebra to the Lie group, we can define the exponential map as follows.

Definition 5.17 (Exponential map). The exponential map

$$\exp : \mathfrak{g} \rightarrow G, \quad X \mapsto \sigma_X(1),$$

is called the exponential map of the Lie group G . It maps an element X of the Lie algebra $\mathfrak{g}$ to the value of the one-parameter subgroup σ_X at $t = 1$, which is an element of the Lie group G .

Proposition 5.13. *If $X \in \mathfrak{g}$, $\lambda \in \mathbb{R}$ then we have*

$$\exp(\lambda X) = \sigma_X(\lambda).$$

Since this is a one-parameter subgroup, we have

$$\exp((\lambda + \mu)X) = \sigma_X(\lambda + \mu) = \sigma_X(\lambda)\sigma_X(\mu) = \exp(\lambda X)\exp(\mu X), \quad \forall \lambda, \mu \in \mathbb{R},$$

and we can also derive straightforwardly that $\exp(0) = e$ and $\exp(-X) = \sigma_X(-1) = \sigma_X(1)^{-1} = \exp(X)^{-1}$. This shows that the exponential map has properties that are consistent with the group structure, and it allows us to relate the Lie algebra to the Lie group in a way that respects the algebraic structure of both.

This definition allows us to plug in the map any vector from the Lie algebra and obtain an element of the Lie group, which is a powerful tool for understanding how the Lie algebra encodes information about the local structure of the Lie group and how it can be used to construct finite transformations from infinitesimal ones.

Proof. For $\lambda \neq 0$ we can define a new curve $\tilde{\sigma} : \mathbb{R} \rightarrow G$ as

$$\tilde{\sigma}(t) := \sigma_X(\lambda t),$$

which is still a one-parameter subgroup of G since it satisfies the group homomorphism property. If we now define another one-parameter subgroup from $\sigma_{\lambda X}(0) = \lambda X$, then we have also

$$\dot{\sigma}(0) = \left. \frac{d}{dt} \sigma_X(\lambda t) \right|_{t=0} = \lambda \dot{\sigma}_X(0) = \lambda X,$$

which means that $\tilde{\sigma}$ and $\sigma_{\lambda X}$ are two one-parameter subgroups of G with the same tangent vector at the identity element (initial conditions), and thus by the previous corollary they must coincide, i.e., $\tilde{\sigma}(t) = \sigma_{\lambda X}(t)$ for all $t \in \mathbb{R}$. If this is the case, then we can use the definition of the exponential map with respect to the one-parameter subgroup $\sigma_{\lambda X}$ to write

$$\exp(\lambda X) = \sigma_{\lambda X}(1) = \tilde{\sigma}(1) = \sigma_X(\lambda),$$

which is the desired result. □

We are now going to express some basic and important properties of the exponential map.

Proposition 5.14. *The exponential map has the following properties:*

1. *The image of the exponential map is a neighborhood of the identity element e in G , which means that*

$$\exp(\mathfrak{g}) \subseteq G_0 = \{g \in G \mid \exists \gamma : [a, b] \rightarrow G \text{ continuous, with } \gamma(t_0) = e, \gamma(t_1) = g\},$$

where G_0 is the connected component of the identity element in G . For example $\text{SO}(n) = [\text{O}(n)]_e$.

2. *The Lie algebra $\mathfrak{g}$ is a $\mathbb{R}$ -vector space $\simeq \mathbb{R}^n$, then we can say that the exponential map is a smooth map from $\mathbb{R}^n$ to G , and thus it is a **local diffeomorphism** around the identity element e . This means that there exists a neighborhood U of $0 \in \mathfrak{g}$ such that $\exp : U \rightarrow \exp(U)$ is a diffeomorphism onto its image, which is a neighborhood of the identity element in G . With these restrictions the exponential map is thus injective.*

3. The exponential map $\exp : \mathfrak{g} \rightarrow G$ is a smooth map, but generally not surjective, since there may be elements of the Lie group that cannot be reached by exponentiating elements of the Lie algebra. But if G_0 is a compact (connected component) then it is surjective. For example, if $G = \mathrm{SL}(2, \mathbb{R})$, which is not compact, then the exponential map is not surjective. On the other hand, if $G = \mathrm{SU}(n)$, which is compact, then the exponential map is surjective. There may be cases of surjectivity even for non-compact groups, but this is not guaranteed in general (for example, the exponential map for $\mathrm{GL}(n, \mathbb{C})$ and $\mathrm{SO}(1, n)$ is surjective even if the group is not compact).
4. Any $g \in G_0$ can be written as

$$g = \exp(X_1) \exp(X_2) \cdots \exp(X_k),$$

for some $X_1, X_2, \dots, X_k \in \mathfrak{g}$ and some $k \in \mathbb{N}$. This means that any element of the connected component of the identity can be expressed as a finite product of exponentials of elements of the Lie algebra, which do not contradict the previous point about surjectivity, since we are not claiming that every element of G can be written as a single exponential, but rather that it can be written as a finite product of exponentials.

5. **Baker-Campbell-Hausdorff (BCH) formula:** For any $X, Y \in \mathfrak{g}$, we have then

$$Z = X + Y + \frac{1}{2}[X, Y]_{\mathfrak{g}} + \frac{1}{12}[X, [X, Y]_{\mathfrak{g}}]_{\mathfrak{g}} - \frac{1}{12}[Y, [X, Y]_{\mathfrak{g}}]_{\mathfrak{g}} + \cdots,$$

where the series continues with higher-order commutators. This formula let us connect ...
Furthermore, if this series converges, we have

$$\exp(Z) = \exp(X) \exp(Y) = \exp \left(X + Y + \frac{1}{2}[X, Y]_{\mathfrak{g}} + \frac{1}{12}[X, [X, Y]_{\mathfrak{g}}]_{\mathfrak{g}} - \frac{1}{12}[Y, [X, Y]_{\mathfrak{g}}]_{\mathfrak{g}} + \cdots \right),$$

has a solution for Z in terms of X and Y . This formula is fundamental for understanding how the group operation in G can be expressed in terms of the Lie algebra $\mathfrak{g}$, and it provides a way to compute the product of two exponentials in terms of the Lie bracket and higher-order commutators. If the group is compact or the map is surjective, then we can use the BCH formula to express any element of the group as an exponential of an element of the Lie algebra (since the series would always converge in this case). For the case in which $[X, Y] = 0$ then the BCH formula simplifies to

$$\exp(X) \exp(Y) = \exp(X + Y),$$

which is a special case that occurs when the Lie algebra is abelian (i.e., when all elements commute with each other).

6. Let $F : G_1 \rightarrow G_2$ be a smooth map between two Lie groups (Lie group homomorphism), then

$$\exp_2(F_*(X)) = F(\exp_1(X)), \quad \forall X \in \mathfrak{g}_1,$$

where $\exp_1$ and $\exp_2$ are the exponential maps of G_1 and G_2 respectively, and F_* is the pushforward of F . This property shows that the exponential map is compatible with Lie group homomorphisms, and it allows us to relate the exponential maps of different Lie groups through the pushforward of a homomorphism between them.

Appendices

A | Notation and Conventions

B | Computations and further developments