

Arcara 2024 - Regression Method for normative data - Tutorial

2025-07-16

Introduction

This tutorial explains how to use the code for applying the method in *Arcara G.(2024) Improving Equivalent scores: a new method for regression model selection*.

Set up R

To work with the method you need to load some functions that will be used. Here I included all the required functions in a folder called `R_functions` (these will be wrapped in an R package in the future).

```
source("R_functions/adjscores_A2024_v3.R")
source("R_functions/formula_transf_text.R")
source("R_functions/model_transf_text.R")
source("R_functions/transf_functions.R")
source("R_functions/tolLimits.adjscores.R")
source("R_functions/tolLimits.obs.R")
source("R_functions/ES.R")
source("R_functions/adj_scores_transf_text.R")
```

I then load some required packages.

```
require(effects)
require(car)
require(performance)
```

Import data

In the following lines I import the data and make some check everything is ok

```
Test.dat = read.csv("Original_Data/MOCA_Dataset.csv", sep=",", dec=".")
```

```
dim(Test.dat)
```

```
## [1] 440 5
```

```
head(Test.dat)
```

```
##      ID Age Education Sex Score
## 1 482  58      27.0   M    25
## 2 428  70      26.0   F    24
## 3 236  59      23.0   F    26
## 4 217  58      23.0   F    28
## 5  34  89      22.0   M    23
## 6 414  51      21.5   F    26
```

```
str(Test.dat)
```

```
## 'data.frame': 440 obs. of 5 variables:
## $ ID : int 482 428 236 217 34 414 116 336 104 85 ...
```

```
## $ Age      : int  58 70 59 58 89 51 65 54 61 63 ...
## $ Education: num  27 26 23 23 22 21.5 21 21 20 20 ...
## $ Sex      : chr   "M" "F" "F" "F" ...
## $ Score    : int  25 24 26 28 23 26 24 27 26 26 ...

# fix values for participants with zero Education otherwise some transformations (e.g. 1/x, log) could
Test.dat[Test.dat$Education==0, "Education"] = 1

Test.dat = na.omit(Test.dat)
```

Use the method to select regression model

```
Test.ARC.res = adjscores_A2024(df = Test.dat, dep="Score",
                               age="Age", edu="Education", sex="Sex",
                               dep.range = c(0, 100))
```

Sex converted to numeric

Inspect model results. The results is a list of objects (see `adjscores_A2024.R` file for details):

- the first is a dataset in which a new column with adjusted scores is added

```
head(Test.ARC.res$new.df)
```

```
##      ID Age Education Sex Score dep age  edu sex sex.or age_tr  edu_tr
## 1 482  58      27.0   M    25  25  58 27.0   0     M   3364 0.03703704
## 2 428  70      26.0   F    24  24  70 26.0   1     F   4900 0.03846154
## 3 236  59      23.0   F    26  26  59 23.0   1     F   3481 0.04347826
## 4 217  58      23.0   F    28  28  58 23.0   1     F   3364 0.04347826
## 5  34  89      22.0   M    23  23  89 22.0   0     M   7921 0.04545455
## 6 414  51      21.5   F    26  26  51 21.5   1     F   2601 0.04651163
##      ADJ_SCORES RESIDUALS
## 1    23.00168 -2.466498
## 2    22.90333 -2.564851
## 3    24.18488 -1.283304
## 4    26.11816  0.649978
## 5    23.75248 -1.715703
## 6    23.73792 -1.730263
```


- the second is the linear model estimated (here I use `summary` to better inspect the model) and I plot the partial effects

```
summary(Test.ARC.res$lm.model)
```

```
##
## Call:
## lm(formula = Score ~ quadr(Age) + inv(Education), data = df)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -8.1796 -1.4153  0.2001  1.5930  6.3890
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   3.005e+01  4.407e-01  68.201  < 2e-16 ***
## quadr(Age)    -5.702e-04  1.027e-04  -5.553  4.89e-08 ***
## inv(Education) -1.808e+01  2.272e+00  -7.959  1.50e-14 ***
```

```
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 2.401 on 437 degrees of freedom
## Multiple R-squared:  0.2729, Adjusted R-squared:  0.2696
## F-statistic: 82.02 on 2 and 437 DF,  p-value: < 2.2e-16
```

```
plot(allEffects(Test.ARC.res[[2]], partial.residuals=TRUE))
```



- I can also print the formula for adjusted scores (according to Capitani 1987 formulation)

```
print(Test.ARC.res$adj_text)
```

```
## [1] "Adjusted score = Observed score - [ -0.00057 * quadr(Age - 4583.205) -18.1 * inv(Education - 0.)
```


Model diagnosis

An appropriate model selection should also include some diagnostics

```
plot(check_normality(Test.ARC.res$lm.model))
```

Normality of Residuals


Distribution should be close to the normal curve



```
plot(check_heteroscedasticity(Test.ARC.res$lm.model))
```

Homogeneity of Variance

Reference line should be flat and horizontal



some disomogeneity in variance, but overall a good fit

Calculate Equivalent Scores

To calculate thresholds for Equivalent Scores I can use some some few additional code.

```
Test.ES = ES(adjscores=Test.ARC.res$new.df$ADJ_SCORES)
```

```
print(Test.ES)
```

```
## $Observations
## ES0(oTL)-ES1      ES1-ES2      ES2-ES3      ES3-ES4
##           15           49           119           219
##
## $Adjusted_Scores
## ES0(oTL)-ES1      ES1-ES2      ES2-ES3      ES3-ES4
##    20.67977    22.46747    24.09836    25.63548
```

Citing the Method

If you use the regression method script please cite: Arcara G. (2024) *Improving Equivalent Scores: A new method for regression model selection*

If you use the ES, please cite: Aiello, E. N., & Depaoli, E. G. (2022). Norms and standardizations in neuropsychology via equivalent scores: software solutions and practical guides. *Neurological Sciences*, 43(2), 961-966.