

SDS 383D: Homework 1

Giorgio Paulon

January 22, 2017

Problem 1. Bayesian inference in simple conjugate families

We start with a few of the simplest building blocks for complex multivariate statistical models: the beta/binomial, normal, and inverse-gamma conjugate families.

- (A) Suppose that we take independent observations X_1, \dots, X_N from a Bernoulli sampling model with unknown probability w . That is, the X_i are the results of flipping a coin with unknown bias. Suppose that w is given a $\text{Beta}(a, b)$ prior distribution:

$$p(w) = \Gamma(a + b)w^{a-1}(1 - w)^{b-1},$$

where $\Gamma(\cdot)$ denotes the Gamma function. Derive the posterior distribution $p(w|x_1, \dots, x_N)$.

The following model

$$\begin{aligned} X_1, \dots, X_n | w &\stackrel{\text{iid}}{\sim} \text{Bernoulli}(w) \\ w &\sim \text{Beta}(a, b) \end{aligned}$$

leads to the following posterior distribution:

$$\begin{aligned} p(w|\mathbf{x}) &\propto p(\mathbf{X}|w)p(w) \\ &= \prod_{i=1}^N p(X_i|w)p(w) \\ &\propto \prod_{i=1}^N \{w^{x_i}(1 - w)^{1-x_i}\} w^{a-1}(1 - w)^{b-1} \mathcal{I}_{[0,1]}(w) \\ &= w^{\sum_{i=1}^N x_i} (1 - w)^{n - \sum_{i=1}^N x_i} w^{a-1}(1 - w)^{b-1} \mathcal{I}_{[0,1]}(w) \\ &= w^{a + \sum_{i=1}^N x_i - 1} (1 - w)^{b + (n - \sum_{i=1}^N x_i) - 1} \mathcal{I}_{[0,1]}(w). \end{aligned}$$

Therefore,

$$p(w|\mathbf{x}) = \frac{\Gamma(a + b + n)}{\Gamma(a + \sum_{i=1}^N x_i) \Gamma(b + (n - \sum_{i=1}^N x_i))} w^{a + \sum_{i=1}^N x_i - 1} (1 - w)^{b + (n - \sum_{i=1}^N x_i) - 1} \mathcal{I}_{[0,1]}(w)$$

that is,

$$W|\mathbf{x} \sim \text{Beta}\left(a + \sum_{i=1}^N x_i, b + (n - \sum_{i=1}^N x_i)\right).$$

This is the so-called **Bernoulli-Beta model**.

- (B) The probability density function (PDF) of a gamma random variable, $X \sim \text{Ga}(a, b)$, is

$$p(x) = \frac{b^a}{\Gamma(a)} x^{a-1} e^{-bx}.$$

Suppose that $X_1 \sim \text{Ga}(a_1, 1)$ and $X_2 \sim \text{Ga}(a_2, 1)$. Define two new random variables $Y_1 = X_1/(X_1 + X_2)$ and $Y_2 = X_1 + X_2$. Find the joint density for (Y_1, Y_2) using a direct PDF transformation (and its Jacobian). Use this to characterize the marginals $p(y_1)$ and $p(y_2)$, and propose

a method that exploits this result to simulate beta random variables, assuming you have a source of gamma random variables.

Let $X_1 \sim Ga(a_1, 1)$, $X_2 \sim Ga(a_2, 1)$ and let us define $(Y_1, Y_2) = g(X_1, X_2) = \left(\frac{X_1}{X_1 + X_2}, X_1 + X_2\right)$. The joint density of (Y_1, Y_2) can be found via pdf transformation, i.e.

$$\begin{cases} y_1 = g_1(x_1, x_2) = \frac{x_1}{x_1 + x_2} \\ y_2 = g_2(x_1, x_2) = x_1 + x_2 \end{cases} \Rightarrow \begin{cases} x_1 = g_1^{-1}(y_1, y_2) = y_1 y_2 \\ x_2 = g_2^{-1}(y_1, y_2) = y_2(1 - y_1). \end{cases} \quad (1)$$

The inverse transformation has a unique solution and therefore the mapping is one-to-one. Moreover, the domain $\mathcal{X}_1 \times \mathcal{X}_2 = [0, \infty)^2$ is mapped to $\mathcal{Y}_1 \times \mathcal{Y}_2 = [0, 1] \times [0, \infty)$.

The Jacobian of the transformation is

$$J = \begin{pmatrix} \frac{\partial g_1^{-1}(y_1, y_2)}{\partial y_1} & \frac{\partial g_1^{-1}(y_1, y_2)}{\partial y_2} \\ \frac{\partial g_2^{-1}(y_1, y_2)}{\partial y_1} & \frac{\partial g_2^{-1}(y_1, y_2)}{\partial y_2} \end{pmatrix} = \begin{pmatrix} y_2 & y_1 \\ -y_2 & 1 - y_1 \end{pmatrix}$$

and the absolute value of its determinant is $|J| = |y_2 - y_1 y_2 + y_1 y_2| = y_2$. The joint pdf of (X_1, X_2) is, since $X_1 \perp X_2$,

$$f_{(X_1, X_2)}(x_1, x_2) = f_{X_1}(x_1)f_{X_2}(x_2) = \frac{1}{\Gamma(a_1)\Gamma(a_2)} x_1^{a_1-1} e^{-x_1} x_2^{a_2-1} e^{-x_2} \mathcal{I}_{[0, \infty)}(x_1) \mathcal{I}_{[0, \infty)}(x_2).$$

Therefore

$$\begin{aligned} f_{(Y_1, Y_2)}(y_1, y_2) &= f_{(X_1, X_2)}(g_1^{-1}(y_1, y_2), g_2^{-1}(y_1, y_2)) |J| \\ &= \frac{1}{\Gamma(a_1)\Gamma(a_2)} y_1^{a_1-1} y_2^{a_1-1} e^{-y_1 y_2} y_2^{a_2-1} (1 - y_1)^{a_2-1} e^{-y_2(1-y_1)} \mathcal{I}_{[0, 1]}(y_1) \mathcal{I}_{[0, \infty)}(y_2) y_2 \\ &= \frac{\Gamma(a_1 + a_2)}{\Gamma(a_1)\Gamma(a_2)} y_1^{a_1-1} (1 - y_1)^{a_2-1} \mathcal{I}_{[0, 1]}(y_1) \frac{1}{\Gamma(a_1 + a_2)} \times y_2^{a_1+a_2-1} e^{-y_2} \mathcal{I}_{[0, \infty)}(y_2). \end{aligned}$$

The joint density factors, and therefore we recognize that

$$f_{(Y_1, Y_2)}(y_1, y_2) = \text{Beta}(a_1, a_2) \times \text{Gamma}(a_1 + a_2, 1),$$

that is,

$$Y_1 \sim \text{Beta}(a_1, a_2)$$

$$Y_2 \sim \text{Gamma}(a_1 + a_2, 1)$$

$$Y_1 \perp Y_2$$

Therefore, in order to sample $Y_1 \sim \text{Beta}(a_1, a_2)$ we can sample $X_1 \sim \text{Gamma}(a_1, 1)$, then $X_2 \sim \text{Gamma}(a_2, 1)$ independently. The resulting $Y_1 = \frac{X_1}{X_1 + X_2}$ is a draw from $\text{Beta}(a_1, a_2)$.

- (C) Suppose that we take independent observations X_1, \dots, X_N from a normal sampling model with unknown mean θ and known variance σ^2 : $X_i \stackrel{\text{iid}}{\sim} N(\theta, \sigma^2)$. Suppose that θ is given a normal prior distribution with mean m and variance v . Derive the posterior distribution $p(\theta | x_1, \dots, x_N)$.

The following model

$$\begin{aligned} X_1, \dots, X_n | \theta &\stackrel{\text{iid}}{\sim} N(\theta, \sigma^2) \\ \theta &\sim N(m, v) \end{aligned}$$

leads to the following posterior distribution:

$$\begin{aligned} p(\theta | \mathbf{x}) &\propto p(\mathbf{X} | \theta) p(\theta) \\ &\propto \prod_{i=1}^N \left\{ \left(\frac{1}{2\pi\sigma^2} \right)^{1/2} e^{-\frac{1}{2\sigma^2}(x_i - \theta)^2} \right\} \left(\frac{1}{2\pi v} \right)^{1/2} e^{-\frac{1}{2v}(\theta - m)^2} \\ &\propto e^{-\frac{1}{2\sigma^2} \sum_{i=1}^N (x_i - \theta)^2} e^{-\frac{1}{2v}(\theta - m)^2} \\ &= e^{-\frac{1}{2\sigma^2} (\sum_{i=1}^N x_i^2 + n\theta^2 - 2\theta \sum_{i=1}^N x_i)} e^{-\frac{1}{2v}(\theta^2 + m^2 - 2m\theta)} \\ &\propto e^{-\frac{n}{2\sigma^2}(\theta^2 - 2\bar{x}\theta)} e^{-\frac{1}{2v}(\theta^2 - 2m\theta)} \\ &= e^{-\frac{n}{2\sigma^2 v} (v\theta^2 - 2\bar{x}v\theta + \frac{\sigma^2}{n}\theta^2 - 2\frac{\sigma^2}{n}m\theta)} \\ &= e^{-\frac{n}{2\sigma^2 v} (v\theta^2 - 2\bar{x}v\theta + \frac{\sigma^2}{n}\theta^2 - 2\frac{\sigma^2}{n}m\theta)} \\ &= e^{-\frac{n}{2\sigma^2 v} [(v + \frac{\sigma^2}{n})\theta^2 - 2(\bar{x}v + \frac{\sigma^2}{n}m)\theta]} \\ &= e^{-\frac{nv + \sigma^2}{2\sigma^2 v} (\theta^2 - 2\frac{\bar{x}nv + \sigma^2 m}{nv + \sigma^2} \theta)} \\ &\propto e^{-\frac{nv + \sigma^2}{2\sigma^2 v} (\theta - \frac{\bar{x}nv + \sigma^2 m}{nv + \sigma^2})^2} \\ &\propto N\left(\frac{v}{v + \sigma^2/n} \bar{x} + \frac{\sigma^2/n}{v + \sigma^2/n} m; \left(\frac{n}{\sigma^2} + \frac{1}{v}\right)^{-1}\right) \end{aligned}$$

This is the so-called **Normal-Normal model for unknown mean and known variance**.

- (D) Suppose that we take independent observations X_1, \dots, X_N from a normal sampling model with known mean θ but unknown variance σ^2 (this seems even more artificial than the last, but is conceptually important). To make this easier, we will re-express things in terms of the precision, or inverse variance $\omega = 1/\sigma^2$:

$$p(x_i | \theta, \omega) = \left(\frac{\omega}{2\pi} \right)^{1/2} \exp \left\{ -\frac{\omega}{2} (x_i - \theta)^2 \right\}.$$

Suppose that ω has a gamma prior with parameters a and b , implying that σ^2 has what is called an inverse-gamma prior. Derive the posterior distribution $p(\omega | x_1, \dots, x_N)$. Re-express this as a posterior for σ^2 , the variance.

The following model

$$\begin{aligned} X_1, \dots, X_n | \sigma^2 &\stackrel{\text{iid}}{\sim} N(\theta, \sigma^2) \\ \sigma^2 &\sim IG(a, b) \end{aligned}$$

can be rewritten in terms of the precision parameter $\omega = 1/\sigma^2$ as

$$\begin{aligned} X_1, \dots, X_n | \omega &\stackrel{\text{iid}}{\sim} N(\theta, \omega) \\ \omega &\sim \text{Gamma}(a, b). \end{aligned}$$

This model leads to the following posterior distribution:

$$\begin{aligned}
 p(\omega|\mathbf{x}) &\propto p(\mathbf{X}|\omega)p(\omega) \\
 &\propto \prod_{i=1}^N \left\{ \left(\frac{\omega}{2\pi} \right)^{1/2} e^{-\frac{\omega}{2}(x_i-\theta)^2} \right\} \frac{b^a}{\Gamma(a)} \omega^{a-1} e^{-b\omega} \mathcal{I}_{[0,\infty)}(\omega) \\
 &\propto \omega^{N/2} e^{-\frac{\omega}{2} \sum_{i=1}^N (x_i-\theta)^2} \omega^{a-1} e^{-b\omega} \mathcal{I}_{[0,\infty)}(\omega) \\
 &= \omega^{a+N/2-1} e^{-\omega(b+\frac{1}{2} \sum_{i=1}^N (x_i-\theta)^2)} \mathcal{I}_{[0,\infty)}(\omega) \\
 &\propto \text{Gamma} \left(a + N/2; b + \frac{1}{2} \sum_{i=1}^N (x_i - \theta)^2 \right).
 \end{aligned}$$

and, subsequently, Therefore,

$$\sigma^2|\mathbf{x} \sim IG \left(a + N/2; b + \frac{1}{2} \sum_{i=1}^N (x_i - \theta)^2 \right).$$

In terms of the pdf of σ^2 , let us find the generic pdf of $IG(a, b)$. We know that, if $X \sim \text{Gamma}(a, b)$, then $Y = 1/X \sim IG(a, b)$. The transformation $g(x) = 1/x$ is monotone and its inverse is $g^{-1}(y) = 1/y$. The derivative of the inverse transformation is $\frac{\partial}{\partial y} g^{-1}(y) = -\frac{1}{y^2}$. Thus,

$$\begin{aligned}
 f_Y(y) &= f_X(1/y) \left| \frac{\partial}{\partial y} g^{-1}(y) \right| \\
 &= \frac{b^a}{\Gamma(a)} \left(\frac{1}{y} \right)^{a-1} e^{-\frac{b}{y}} \frac{1}{y^2} \mathcal{I}_{[0,\infty)}(y) \\
 &= \frac{b^a}{\Gamma(a)} y^{-a-1} e^{-\frac{b}{y}} \mathcal{I}_{[0,\infty)}(y).
 \end{aligned}$$

Therefore, if $\sigma^2|\mathbf{x} \sim IG \left(a + N/2; b + \frac{1}{2} \sum_{i=1}^N (x_i - \theta)^2 \right)$, the pdf is

$$f_{\sigma^2|\mathbf{x}}(\sigma^2|\mathbf{x}) = \frac{(b + \frac{1}{2} \sum_{i=1}^N (x_i - \theta)^2)^{a+N/2}}{\Gamma(a + N/2)} \sigma^{2(-a-N/2-1)} e^{-\frac{b+\frac{1}{2} \sum_{i=1}^N (x_i-\theta)^2}{\sigma^2}} \mathcal{I}_{[0,\infty)}(\sigma^2).$$

This is the so-called **Normal-inverse gamma model for known mean and unknown variance**.

- (E) Suppose that, as above, we take independent observations X_1, \dots, X_N from a normal sampling model with unknown, common mean θ . This time, however, each observation has its own idiosyncratic (but known) variance: $X_i \stackrel{\text{iid}}{\sim} N(\theta, \sigma_i^2)$. Suppose that θ is given a normal prior distribution with mean m and variance v . Derive the posterior distribution $p(\theta|x_1, \dots, x_N)$. Express the posterior mean in a form that is clearly interpretable as a weighted average of the observations and the prior mean.

The following model

$$\begin{aligned}
 X_1, \dots, X_N | \theta &\stackrel{\text{iid}}{\sim} N(\theta, \sigma_i^2) \\
 \theta &\sim N(m, v)
 \end{aligned}$$

leads to the following posterior distribution:

$$\begin{aligned} p(\theta|\mathbf{x}) &\propto p(\mathbf{X}|\theta)p(\theta) \\ &\propto \prod_{i=1}^N \left\{ \left(\frac{1}{2\pi\sigma_i^2} \right)^{1/2} e^{-\frac{1}{2\sigma_i^2}(x_i-\theta)^2} \right\} \left(\frac{1}{2\pi v} \right)^{1/2} e^{-\frac{1}{2v}(\theta-m)^2} \\ &\propto e^{-\frac{1}{2} \sum_{i=1}^N \frac{(x_i-\theta)^2}{\sigma_i^2}} e^{-\frac{1}{2v}(\theta-m)^2}. \end{aligned}$$

The exponents can be rewritten as

$$\begin{aligned} &-\frac{1}{2} \left[\sum_{i=1}^N \left(\frac{x_i - \theta}{\sigma_i} \right)^2 + \frac{(\theta - m)^2}{v} \right] \\ &= -\frac{1}{2} \left[\sum_{i=1}^N \left(\frac{x_i^2}{\sigma_i^2} + \frac{\theta^2}{\sigma_i^2} - 2 \frac{x_i \theta}{\sigma_i^2} \right) + \frac{\theta^2 + m^2 - 2m\theta}{v} \right] \\ &\propto -\frac{1}{2} \left[\theta^2 \left(\frac{1}{v} + \sum_{i=1}^N \frac{1}{\sigma_i^2} \right) - 2\theta \left(\frac{m}{v} + \sum_{i=1}^N \frac{x_i}{\sigma_i^2} \right) \right]. \end{aligned}$$

Therefore, we get

$$\theta|\mathbf{x} \sim N \left(\frac{\frac{m}{v} + \sum_{i=1}^N \frac{x_i}{\sigma_i^2}}{\frac{1}{v} + \sum_{i=1}^N \frac{1}{\sigma_i^2}}; \left(\frac{1}{v} + \sum_{i=1}^N \frac{1}{\sigma_i^2} \right)^{-1} \right).$$

This is the so-called **Normal-Normal model for unknown mean and known idiosyncratic variances**.

- (F) Suppose that $(X|\sigma^2) \sim N(0, \sigma^2)$, and that $1/\sigma^2$ has a $\text{Gamma}(a, b)$ prior, defined as above. Show that the marginal distribution of X is Student's t . This is why the t distribution is often referred to as a scale mixture of normals.

Problem 2. The multivariate normal distribution

Basics

We all know the univariate normal distribution, whose long history began with de Moivre's 18th-century work on approximating the (analytically inconvenient) binomial distribution. This led to the probability density function

$$p(x) = \frac{1}{\sqrt{2\pi v}} \exp \left\{ -\frac{(x - m)^2}{2v} \right\}$$

for the normal random variable with mean m and variance v , written $X \sim N(m, v)$.

Here's an alternative characterization of the univariate normal distribution in terms of moment-generating functions: a random variable X has a normal distribution if and only if $E\{\exp(tx)\} = \exp(mt + vt^2/2)$ for some real m and positive real v . Remember that $E(\cdot)$ denotes the expected value of its argument under the given probability distribution. We will generalize this definition to the multivariate normal.

- (A) First, some simple moment identities. The covariance matrix $\text{cov}(\mathbf{X})$ of a vector-valued random variable \mathbf{X} is defined as the matrix whose (i, j) entry is the covariance between X_i and X_j . In matrix notation, $\text{cov}(\mathbf{X}) = E\{(\mathbf{X} - \boldsymbol{\mu})(\mathbf{X} - \boldsymbol{\mu})^T\}$, where $\boldsymbol{\mu}$ is the mean vector whose i th component is $E(X_i)$. Prove the following: (1) $\text{cov}(\mathbf{X}) = E(\mathbf{X}\mathbf{X}^T) - \boldsymbol{\mu}\boldsymbol{\mu}^T$; and (2) $\text{cov}(\mathbf{A}\mathbf{X} + \mathbf{b}) = \mathbf{A}\text{cov}(\mathbf{X})\mathbf{A}^T$ for matrix \mathbf{A} and vector \mathbf{b} .
- (B) Consider the random vector $\mathbf{Z} = (Z_1, \dots, Z_p)^T$, with each entry having an independent standard normal distribution (that is, mean 0 and variance 1). Derive the probability density function (PDF) and moment-generating function (MGF) of \mathbf{Z} , expressed in vector notation. We say that \mathbf{Z} has a standard multivariate normal distribution.
- (C) A vector-valued random variable $\mathbf{X} = (X_1, \dots, X_p)^T$ has a multivariate normal distribution if and only if every linear combination of its components is univariate normal. That is, for all vectors \mathbf{a} not identically zero, the scalar quantity $Z = \mathbf{a}^T \mathbf{X}$ is normally distributed. From this definition, prove that \mathbf{X} is multivariate normal, written $\mathbf{X} \sim N(\boldsymbol{\mu}, \Sigma)$, if and only if its moment-generating function is of the form $E(\exp\{\mathbf{t}^T \mathbf{X}\}) = \exp(\mathbf{t}^T \boldsymbol{\mu} + \mathbf{t}^T \Sigma \mathbf{t}/2)$. Hint: what are the mean, variance, and moment-generating function of Z , expressed in terms of moments of \mathbf{X} ?
- (D) Another basic theorem is that a random vector is multivariate normal if and only if it is an affine transformation of independent univariate normals. You will first prove the “if” statement. Let \mathbf{Z} have a standard multivariate normal distribution, and define the random vector $\mathbf{X} = \mathbf{L}\mathbf{Z} + \boldsymbol{\mu}$ for some $p \times p$ matrix \mathbf{L} of full column rank. Prove that \mathbf{X} is multivariate normal. In addition, use the moment identities you proved above to compute the expected value and covariance matrix of \mathbf{X} .
- (E) Now for the “only if”, Suppose that \mathbf{X} has a multivariate normal distribution. Prove that \mathbf{X} can be written as an affine transformation of standard normal random variables. (Note: a good way to prove that something can be done is to do it!) Use this insight to propose an algorithm for simulating multivariate normal random variables with a specified mean and covariance matrix.

Appendix A

R code