



REQUIRES SPECIFIC TRAINING - USE UNDER MEDICAL SUPERVISION

EQUIPMENT & SUPPLIES

Item	Description	Use	Quantity
Pulsimeter	PulsoMedical C101B1	Pulse and Oxygen saturation	1
Hemostatic dressing	Hemostatic wadding	Stops bleeding from wounds/nose	1
Gauze, roll	Sterile 7x5 cm	Wound dressing	3
Gauze, pads medium	Sterile 18x40 cm	Wound dressing	6
Gauze, pads large	Sterile 36x40 cm	Wound dressing	4
Medical Tape small	2.5 cm wide	Wound dressing	1
Medical Tape large	5 cm wide	Wound dressing	1
Band Aid large	Sterile 76x100 cm	Wound dressing	2
Band Aid medium	Sterile 50x100 cm	Wound dressing	2
Band Aid small	Sterile 29x70 cm	Wound dressing	4
Self-adhesive bandages		Wound dressing	1
Vented chest seal)		Treat open chest wounds	1
Syringes 5 ml	with needle	Injections	2
Eye dressing	Sterile	Eye wound dressing	2
Surgical blade, sterile	23 and 24	Incision, cutting, debridement	2
Tourniquet	elastic, simple type	Stops bleeding	1
Steri-strip stitches	Sterile bandage stitches	Closing wounds	2
Lister Scissors		Removing induments / dressing	1
Anatomical tweezers		Debridement	1
Surgical forceps		Debridement, bleeding	1

Drug	Class	Purpose	Dosage	Comments	Q.ty
Alovex spray	Disinfectant	Disinfection of wounds and heat burns	Apply thin layer on wound	For severe burns, apply and cover with a plastic bag/wrap. Give electrolytes	1
Amoxicilline ▲ tablets	Antibiotic	Systemic antibiotic to treat ongoing infections	Use 1 table twice a day	CONSULT A MEDIC. Avoid in case of known allergy to antibiotics	6
Bentelan ▲ vials	Cortisone	Anaphylactic shock	Intramuscular injection	CONSULT A MEDIC. Avoid in case of known allergy to corticosteroids	2
Bentelan ▲ tablets	Cortisone	Allergies, asma, inflammations	Use 1 to 4 tablets per day	Avoid in case of known allergy to corticosteroids	20
Betadine 10% solution	Disinfectant	Wound disinfection	Apply directly on the wound		1
Brufen ▲ tablets	Painkiller	Treatment of pain	use 1 to 3 pills per day; 6 h between doses	Avoid in case of known allergy to FANS and painkillers. Might interfere with cardioaspirin	12
Cardioaspirin ▲ tablets	Anticoagulant	Heart attack, stroke	Use in case of ongoing heart attack or stroke	CONSULT A MEDIC. Avoid in case of known allergy to acetylsalicylic acid	15
Cetirizine tablets	Antihistamine	Allergic reactions	Use 1 pill/day	Induces drowsiness	20
Gaviscon solution	Antacid	Stomach acidity	Use orally when needed		4
Gentalin beta ointment	Antibiotic + Cortisone	Preventing / treating local wound infections	Apply directly on the wound		1
Loperamide tablets	Antidiarrheal	To stop diarrhea	Use 2 pills per day		10
Plasil vials	Antiemetic	To stop vomit	Intramuscular injection		1
Plasil tablets	Antiemetic	To stop vomit	1 up to 3 pills per day; 6h between doses		30
Saline solution	Saline solution	Cleaning wounds and eye	directly on the wound		1
Tobradex eye drops	Antibiotic + Cortisone	Eye infections / External ear infection	Apply 2 drops for 5 times per day	CONSULT A MEDIC	1

▲ WARNING: known possible allergies