

Computing cone-constrained singular values of matrices

Giovanni Barbarino ¹ Nicolas Gillis ¹ David Sossa ²



Université de Mons



Universidad
de O'Higgins

06 March 2025

¹Université de Mons, Belgium

²Universidad de O'Higgins, Rancagua, Chile

Class of Computational Complexity

Cones and Notation

P and Q are convex, closed and finitely generated cones

$$P = G(\mathbb{R}_+^p) = \{Gx \mid x \geq 0\}$$

$$Q = H(\mathbb{R}_+^q) = \{Hy \mid y \geq 0\}$$

where the generators $G = [g_1, \dots, g_p]$, $H = [h_1, \dots, h_q]$
have nonnegative independent unit columns


The *faces* of P are the polyhedral cones $F \subseteq P$ such that

$$v_1 + v_2 \in F, \quad v_1, v_2 \in P \implies v_1, v_2 \in F$$

The dimension of a face is equal to the dimension of the
subspace spanned

A *vertex* of P is any nonzero vector of a 1-dimensional
face of P , or any positive multiple of the columns of G

The *facets* of the cone P are the faces of maximal di-
mension different from P



Cones and Notation

P and Q are convex, closed and finitely generated cones

$$P = G(\mathbb{R}_+^p) = \{Gx \mid x \geq 0\}$$

$$Q = H(\mathbb{R}_+^q) = \{Hy \mid y \geq 0\}$$

where the generators $G = [g_1, \dots, g_p]$, $H = [h_1, \dots, h_q]$
have nonnegative independent unit columns


The *faces* of P are the polyhedral cones $F \subseteq P$ such that

$$v_1 + v_2 \in F, \quad v_1, v_2 \in P \implies v_1, v_2 \in F$$

The dimension of a face is equal to the dimension of the
subspace spanned

A *vertex* of P is any nonzero vector of a 1-dimensional
face of P , or any positive multiple of the columns of G

The *facets* of the cone P are the faces of maximal di-
mension different from P



Cones and Notation

P and Q are convex, closed and finitely generated cones

$$P = G(\mathbb{R}_+^p) = \{Gx \mid x \geq 0\}$$

$$Q = H(\mathbb{R}_+^q) = \{Hy \mid y \geq 0\}$$

where the generators $G = [g_1, \dots, g_p]$, $H = [h_1, \dots, h_q]$
have nonnegative independent unit columns


The *faces* of P are the polyhedral cones $F \subseteq P$ such that

$$v_1 + v_2 \in F, \quad v_1, v_2 \in P \implies v_1, v_2 \in F$$

The dimension of a face is equal to the dimension of the
subspace spanned

A *vertex* of P is any nonzero vector of a 1-dimensional
face of P , or any positive multiple of the columns of G

The *facets* of the cone P are the faces of maximal di-
mension different from P



Cones and Notation

P and Q are convex, closed and finitely generated cones

$$P = G(\mathbb{R}_+^p) = \{Gx \mid x \geq 0\}$$

$$Q = H(\mathbb{R}_+^q) = \{Hy \mid y \geq 0\}$$

where the generators $G = [g_1, \dots, g_p]$, $H = [h_1, \dots, h_q]$
have nonnegative independent unit columns


The **faces** of P are the polyhedral cones $F \subseteq P$ such that

$$v_1 + v_2 \in F, \quad v_1, v_2 \in P \implies v_1, v_2 \in F$$

The dimension of a face is equal to the dimension of the
subspace spanned

A **vertex** of P is any nonzero vector of a 1-dimensional
face of P , or any positive multiple of the columns of G

The **facets** of the cone P are the faces of maximal di-
mension different from P



Cones and Notation

P and Q are convex, closed and finitely generated cones

$$P = G(\mathbb{R}_+^p) = \{Gx \mid x \geq 0\}$$

$$Q = H(\mathbb{R}_+^q) = \{Hy \mid y \geq 0\}$$

where the generators $G = [g_1, \dots, g_p]$, $H = [h_1, \dots, h_q]$
have nonnegative independent unit columns


The **faces** of P are the polyhedral cones $F \subseteq P$ such that

$$v_1 + v_2 \in F, \quad v_1, v_2 \in P \implies v_1, v_2 \in F$$

The dimension of a face is equal to the dimension of the
subspace spanned

A **vertex** of P is any nonzero vector of a 1-dimensional
face of P , or any positive multiple of the columns of G

The **facets** of the cone P are the faces of maximal di-
mension different from P



Conical Singular Values

$$\min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1}} u^\top A v \quad P, Q \quad \begin{array}{l} \text{closed convex cones} \\ \text{finitely generated} \end{array}$$

Conical Singular Values

$$\min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1}} u^\top A v \quad P, Q \text{ closed convex cones finitely generated}$$



Pareto Singular Values

$$\min_{\substack{u \geq 0, \|u\| = 1, \\ v \geq 0, \|v\| = 1}} u^\top A v$$

Conical Singular Values

$$\min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1}} u^\top A v \quad P, Q \text{ closed convex cones}$$

finitely generated



Pareto Singular Values

$$\min_{\substack{u \geq 0, \|u\| = 1, \\ v \geq 0, \|v\| = 1}} u^\top A v$$

Conic Angles

$$\min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1}} u^\top v$$

Conical Singular Values

$$\min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1}} u^\top A v \quad P, Q \text{ closed convex cones finitely generated}$$



Pareto Singular Values

$$\min_{\substack{u \geq 0, \|u\| = 1, \\ v \geq 0, \|v\| = 1}} u^\top A v$$

Conic Angles

$$\min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1}} u^\top v$$

Singular Values

$$\min_{\|u\|=\|v\|=1} u^\top A v$$

Conical Singular Values

$$\min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1}} u^\top A v \quad P, Q \text{ closed convex cones finitely generated}$$



Pareto Singular Values

$$\min_{\substack{u \geq 0, \|u\| = 1, \\ v \geq 0, \|v\| = 1}} u^\top A v$$

Conic Angles

$$\min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1}} u^\top v$$

Singular Values

$$\min_{\|u\|=\|v\|=1} u^\top A v$$

Polynomial Time
 $O(mn^2)$ to compute all Singular Values

Conical Singular Values

$$\min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1}} u^\top A v \quad P, Q \text{ closed convex cones}$$

finitely generated



Pareto Singular Values

$$\min_{\substack{u \geq 0, \|u\| = 1, \\ v \geq 0, \|v\| = 1}} u^\top A v$$

?

Conic Angles

$$\min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1}} u^\top v$$

?

Singular Values

$$\min_{\|u\|=\|v\|=1} u^\top A v$$

Polynomial Time
 $O(mn^2)$ to compute
all Singular Values

Pareto Singular Values

$$\min_{\substack{u \geq 0, \|u\| = 1, \\ v \geq 0, \|v\| = 1}} u^\top A v \quad A \in \mathbb{R}^{m \times n} \quad \Xi(A) := \{\text{stationary values of the o.p.}\}$$

"Simple" Case: $A \geq 0$

Theorem (Seeger, S. 2023)

$$A \geq 0 \iff \Xi(A) = \{\|B\| : B \trianglelefteq A\}$$

Theorem

The set of matrices $A \in \mathbb{R}_+^{m \times n}$ for which all the submatrices have different norms, is dense, open and its complementary has measure zero on the space of nonnegative matrices

⇒ A generic nonnegative matrix has exponentially many Pareto Singular Values

Still it does say nothing on the complexity of recovering the minimum, since

$$A \geq 0 \implies \min \Xi(A) = \min_{i,j} A_{i,j}$$

Pareto Singular Values

$$\min_{\substack{u \geq 0, \|u\| = 1, \\ v \geq 0, \|v\| = 1}} u^\top A v \quad A \in \mathbb{R}^{m \times n} \quad \Xi(A) := \{\text{stationary values of the o.p.}\}$$

"Simple" Case: $A \geq 0$

Theorem (Seeger, S. 2023)

$$A \geq 0 \iff \Xi(A) = \{\|B\| : B \trianglelefteq A\}$$

Theorem

The set of matrices $A \in \mathbb{R}_+^{m \times n}$ for which all the submatrices have different norms, is dense, open and its complementary has measure zero on the space of nonnegative matrices

⇒ A generic nonnegative matrix has exponentially many Pareto Singular Values

Still it does say nothing on the complexity of recovering the minimum, since

$$A \geq 0 \implies \min \Xi(A) = \min_{i,j} A_{i,j}$$

Pareto Singular Values

$$\min_{\substack{u \geq 0, \|u\| = 1, \\ v \geq 0, \|v\| = 1}} u^\top A v \quad A \in \mathbb{R}^{m \times n} \quad \Xi(A) := \{\text{stationary values of the o.p.}\}$$

"Simple" Case: $A \geq 0$

Theorem (Seeger, S. 2023)

$$A \geq 0 \iff \Xi(A) = \{\|B\| : B \trianglelefteq A\}$$

Theorem

The set of matrices $A \in \mathbb{R}_+^{m \times n}$ for which all the submatrices have different norms, is dense, open and its complementary has measure zero on the space of nonnegative matrices

⇒ A generic nonnegative matrix has exponentially many Pareto Singular Values

Still it does say nothing on the complexity of recovering the minimum, since

$$A \geq 0 \implies \min \Xi(A) = \min_{i,j} A_{i,j}$$

Pareto Singular Values

$$\min_{\substack{u \geq 0, \|u\| = 1, \\ v \geq 0, \|v\| = 1}} u^\top A v \quad A \in \mathbb{R}^{m \times n} \quad \Xi(A) := \{\text{stationary values of the o.p.}\}$$

"Simple" Case: $A \geq 0$

Theorem (Seeger, S. 2023)

$$A \geq 0 \iff \Xi(A) = \{\|B\| : B \trianglelefteq A\}$$

Theorem

The set of matrices $A \in \mathbb{R}_+^{m \times n}$ for which all the submatrices have different norms, is dense, open and its complementary has measure zero on the space of nonnegative matrices

⇒ A generic nonnegative matrix has exponentially many Pareto Singular Values

Still it does say nothing on the complexity of recovering the minimum, since

$$A \geq 0 \implies \min \Xi(A) = \min_{i,j} A_{i,j}$$

Conic Singular Values

$$\min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1,}} u^\top A v = \min_{\substack{x \geq 0, \|Gx\| = 1, \\ y \geq 0, \|Hy\| = 1,}} x^\top G^\top A H y \quad A \in \mathbb{R}^{m \times n}$$

"Simple" Case: $G^\top A H \geq 0$

Theorem (Seeger, S. 2023)

$$G^\top A H \geq 0 \implies \min_{\substack{x \geq 0, \|Gx\| = 1, \\ y \geq 0, \|Hy\| = 1,}} x^\top G^\top A H y = \min_{i,j} g_i^\top A h_j$$

It is possible to extend the above results to any **Nash Pair**:

- (u^*, v^*) is a Nash Pair of the problem if

$$\langle u, Av^* \rangle \geq \langle u^*, Av^* \rangle, \quad \langle u^*, Av \rangle \geq \langle u^*, Av^* \rangle, \quad \forall (u, v) \in P \times Q$$

where u, v, u^*, v^* have all unit norm

- If (u^*, v^*) is a Nash Pair of the problem, then $u = g_i, v = h_j$ for some i, j

Conic Singular Values

$$\min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1,}} u^\top A v = \min_{\substack{x \geq 0, \|Gx\| = 1, \\ y \geq 0, \|Hy\| = 1,}} x^\top G^\top A H y \quad A \in \mathbb{R}^{m \times n}$$

"Simple" Case: $G^\top A H \geq 0$

Theorem (Seeger, S. 2023)

$$G^\top A H \geq 0 \implies \min_{\substack{x \geq 0, \|Gx\| = 1, \\ y \geq 0, \|Hy\| = 1,}} x^\top G^\top A H y = \min_{i,j} g_i^\top A h_j$$

It is possible to extend the above results to any Nash Pair:

- (u^*, v^*) is a Nash Pair of the problem if

$$\langle u, Av^* \rangle \geq \langle u^*, Av^* \rangle, \quad \langle u^*, Av \rangle \geq \langle u^*, Av^* \rangle, \quad \forall (u, v) \in P \times Q$$

where u, v, u^*, v^* have all unit norm

- If (u^*, v^*) is a Nash Pair of the problem, then $u = g_i, v = h_j$ for some i, j

Conic Singular Values

$$\min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1}} u^\top A v = \min_{\substack{x \geq 0, \|Gx\| = 1, \\ y \geq 0, \|Hy\| = 1}} x^\top G^\top A H y \quad A \in \mathbb{R}^{m \times n}$$

"Simple" Case: $G^\top A H \geq 0$

Theorem (Seeger, S. 2023)

$$G^\top A H \geq 0 \implies \min_{\substack{x \geq 0, \|Gx\| = 1, \\ y \geq 0, \|Hy\| = 1}} x^\top G^\top A H y = \min_{i,j} g_i^\top A h_j$$

It is possible to extend the above results to any **Nash Pair**:

- (u^*, v^*) is a Nash Pair of the problem if

$$\langle u, Av^* \rangle \geq \langle u^*, Av^* \rangle, \quad \langle u^*, Av \rangle \geq \langle u^*, Av^* \rangle, \quad \forall (u, v) \in P \times Q$$

where u, v, u^*, v^* have all unit norm



- If (u^*, v^*) is a Nash Pair of the problem, then $u = g_i, v = h_j$ for some i, j

Conic Angles

$$\min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1,}} u^\top v \quad P, Q \subseteq \mathbb{R}^n \text{ non trivial polyhedral cones}$$

"Simple" Case:

$$\min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1,}} u^\top v \geq 0 \implies u, v \text{ are vertices of } P, Q$$





Conic Angles

$$\min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1,}} u^\top v \quad P, Q \subseteq \mathbb{R}^n \text{ non trivial polyhedral cones}$$

"Simple" Case:

$$\min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1,}} u^\top v \geq 0 \implies u, v \text{ are vertices of } P, Q$$



Conic Angles

If one of u, v in the antipodal pair is a vertex then the problem is
Polynomial in n and the number of generators of P, Q

$$\min_{v \in Q, \|v\|=1} u^T v = - \max_{v \in -Q, \|v\|=1} u^T v \implies v = -\frac{\text{Proj}(u, -Q)}{\|\text{Proj}(u, -Q)\|}$$

$$\text{Proj}(u, -Q) \equiv \min_{y \geq 0} \|u - (-H)y\| \quad \text{NNLS, convex}$$

if $\min_{\substack{u \in P, \|u\|=1 \\ v \in Q, \|v\|=1}} u^T v < 0$, when is it that one among u, v is a vertex?

Theorem (B., G., S. 2024)

Let (u, v) be a stationary point and let $u \in \text{int}(F_u)$, $v \in \text{int}(F_v)$ where F_u, F_v are facets of P, Q . If $\dim(F_u) + \dim(F_v) > n$ and $v \neq \pm u$, then (u, v) is a saddle point

Corollary (B., G., S. 2024)

If (u, v) is a local minimum in dimension $n \leq 3$ with $u \neq -v$, then at least one among u, v is a vertex

Conic Angles

If one of u, v in the antipodal pair is a vertex then the problem is
Polynomial in n and the number of generators of P, Q

$$\min_{v \in Q, \|v\|=1} u^\top v = - \max_{v \in -Q, \|v\|=1} u^\top v \implies v = -\frac{\text{Proj}(u, -Q)}{\|\text{Proj}(u, -Q)\|}$$

$$\text{Proj}(u, -Q) \equiv \min_{y \geq 0} \|u - (-H)y\| \quad \text{NNLS, convex}$$

if $\min_{\substack{u \in P, \|u\|=1 \\ v \in Q, \|v\|=1}} u^\top v < 0$, when is it that one among u, v is a vertex?

Theorem (B., G., S. 2024)

Let (u, v) be a stationary point and let $u \in \text{int}(F_u)$, $v \in \text{int}(F_v)$ where F_u, F_v are facets of P, Q . If $\dim(F_u) + \dim(F_v) > n$ and $v \neq \pm u$, then (u, v) is a saddle point

Corollary (B., G., S. 2024)

If (u, v) is a local minimum in dimension $n \leq 3$ with $u \neq -v$, then at least one among u, v is a vertex

Conic Angles

If one of u, v in the antipodal pair is a vertex then the problem is
Polynomial in n and the number of generators of P, Q

$$\min_{v \in Q, \|v\|=1} u^\top v = - \max_{v \in -Q, \|v\|=1} u^\top v \implies v = -\frac{\text{Proj}(u, -Q)}{\|\text{Proj}(u, -Q)\|}$$

$$\text{Proj}(u, -Q) \equiv \min_{y \geq 0} \|u - (-H)y\| \quad \text{NNLS, convex}$$

if $\min_{\substack{u \in P, \|u\|=1 \\ v \in Q, \|v\|=1}} u^\top v < 0$, when is it that one among u, v is a vertex?

Theorem (B., G., S. 2024)

Let (u, v) be a stationary point and let $u \in \text{int}(F_u)$, $v \in \text{int}(F_v)$ where F_u, F_v are facets of P, Q . If $\dim(F_u) + \dim(F_v) > n$ and $v \neq \pm u$, then (u, v) is a saddle point

Corollary (B., G., S. 2024)

If (u, v) is a local minimum in dimension $n \leq 3$ with $u \neq -v$, then at least one among u, v is a vertex

Conic Angles

If one of u, v in the antipodal pair is a vertex then the problem is
Polynomial in n and the number of generators of P, Q

$$\min_{v \in Q, \|v\|=1} u^\top v = - \max_{v \in -Q, \|v\|=1} u^\top v \implies v = -\frac{\text{Proj}(u, -Q)}{\|\text{Proj}(u, -Q)\|}$$

$$\text{Proj}(u, -Q) \equiv \min_{y \geq 0} \|u - (-H)y\| \quad \text{NNLS, convex}$$

if $\min_{\substack{u \in P, \|u\|=1 \\ v \in Q, \|v\|=1}} u^\top v < 0$, when is it that one among u, v is a vertex?

Theorem (B., G., S. 2024)

Let (u, v) be a stationary point and let $u \in \text{int}(F_u)$, $v \in \text{int}(F_v)$ where F_u, F_v are facets of P, Q . If $\dim(F_u) + \dim(F_v) > n$ and $v \neq \pm u$, then (u, v) is a saddle point

Corollary (B., G., S. 2024)

If (u, v) is a local minimum in dimension $n \leq 3$ with $u \neq -v$, then at least one among u, v is a vertex

Conic Angles

If one of u, v in the antipodal pair is a vertex then the problem is
Polynomial in n and the number of generators of P, Q

$$\min_{v \in Q, \|v\|=1} u^\top v = - \max_{v \in -Q, \|v\|=1} u^\top v \implies v = -\frac{\text{Proj}(u, -Q)}{\|\text{Proj}(u, -Q)\|}$$
$$\text{Proj}(u, -Q) \equiv \min_{y \geq 0} \|u - (-H)y\| \quad \text{NNLS, convex}$$

if $\min_{\substack{u \in P, \|u\|=1 \\ v \in Q, \|v\|=1}} u^\top v < 0$, when is it that one among u, v is a vertex?

Theorem (B., G., S. 2024)

Let (u, v) be a stationary point and let $u \in \text{int}(F_u)$, $v \in \text{int}(F_v)$ where F_u, F_v are facets of P, Q . If $\dim(F_u) + \dim(F_v) > n$ and $v \neq \pm u$, then (u, v) is a saddle point

Corollary (B., G., S. 2024)

If (u, v) is a local minimum in dimension $n \leq 3$ with $u \neq -v$, then at least one among u, v is a vertex

Conic Angles

- $\dim(F_u) + \dim(F_v) > n$ then (u, v) is a saddle point
- (u, v) local minimum, $n \leq 3$, then u or v is a vertex

Idea of proof:


- If $\dim(F_u) + \dim(F_v) > n$ then $0 \neq z \in \text{Span}(F_u) \cap \text{Span}(F_v)$
- u, v are in the interior part of F_u, F_v , so they can move in $\text{Span}(u, z), \text{Span}(v, z)$

Conic Angles

- $\dim(F_u) + \dim(F_v) > n$ then (u, v) is a saddle point
- (u, v) local minimum, $n \leq 3$, then u or v is a vertex

Idea of proof:

- If $\dim(F_u) + \dim(F_v) > n$ then $0 \neq z \in \text{Span}(F_u) \cap \text{Span}(F_v)$
- u, v are in the interior part of F_u, F_v , so they can move in $\text{Span}(u, z), \text{Span}(v, z)$



$$H_{\theta, \varphi}(u_\theta, v_\varphi) \Big|_{\theta=\varphi=0} = \begin{pmatrix} -\langle u, v \rangle & \pm 1 \\ \pm 1 & -\langle u, v \rangle \end{pmatrix}$$

Hessian has negative determinant

since $u \neq \pm v$

↓


(u, v) is a saddle point

Conic Angles

- $\dim(F_u) + \dim(F_v) > n$ then (u, v) is a saddle point
- (u, v) local minimum, $n \leq 3$, then u or v is a vertex

Idea of proof:

- If $\dim(F_u) + \dim(F_v) > n$ then $0 \neq z \in \text{Span}(F_u) \cap \text{Span}(F_v)$
- u, v are in the interior part of F_u, F_v , so they can move in $\text{Span}(u, z), \text{Span}(v, z)$



$$H_{\theta, \varphi}(u_\theta, v_\varphi) \Big|_{\theta=\varphi=0} = \begin{pmatrix} -\langle u, v \rangle & \pm 1 \\ \pm 1 & -\langle u, v \rangle \end{pmatrix}$$

Hessian has negative determinant

since $u \neq \pm v$




(u, v) is a saddle point

Conic Angles

- $\dim(F_u) + \dim(F_v) > n$ then (u, v) is a saddle point
- (u, v) local minimum, $n \leq 3$, then u or v is a vertex

Idea of proof:

- If $\dim(F_u) + \dim(F_v) > n$ then $0 \neq z \in \text{Span}(F_u) \cap \text{Span}(F_v)$
- u, v are in the interior part of F_u, F_v , so they can move in $\text{Span}(u, z), \text{Span}(v, z)$



$$H_{\theta, \varphi}(u_\theta, v_\varphi) \Big|_{\theta=\varphi=0} = \begin{pmatrix} -\langle u, v \rangle & \pm 1 \\ \pm 1 & -\langle u, v \rangle \end{pmatrix}$$

Hessian has negative determinant
since $u \neq \pm v$
 \downarrow
 (u, v) is a saddle point

Conic Angles

- $\dim(F_u) + \dim(F_v) > n$ then (u, v) is a saddle point
- (u, v) local minimum, $n \leq 3$, then u or v is a vertex

Idea of proof:

- If $\dim(F_u) + \dim(F_v) > n$ then $0 \neq z \in \text{Span}(F_u) \cap \text{Span}(F_v)$
- u, v are in the interior part of F_u, F_v , so they can move in $\text{Span}(u, z), \text{Span}(v, z)$

Counterexample for $n \geq 4$:

$$P = \left\langle \begin{pmatrix} 1 & 1 \\ -1 & 1 \\ 0 & 0 \\ 0 & 0 \end{pmatrix} \right\rangle \quad Q = \left\langle \begin{pmatrix} -1 & -1 \\ 0 & 0 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} \right\rangle \quad u = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} \quad v = \frac{1}{\sqrt{2}} \begin{pmatrix} -1 \\ 0 \\ 1 \\ 0 \end{pmatrix}$$

$$\dim(F_u) = \dim(F_v) = 2 \quad u^\top v = -\frac{1}{\sqrt{2}} < 0$$

(u, v) is an antipodal pair in the interior part of P and Q

Conical Singular Values

$$\min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1}} u^\top A v \quad P, Q \quad \begin{array}{l} \text{closed convex cones} \\ \text{finitely generated} \end{array}$$



Pareto Singular Values

$$\min_{\substack{u \geq 0, \|u\| = 1, \\ v \geq 0, \|v\| = 1}} u^\top A v$$

Polynomial Time
for $A \geq 0$ or $A \leq 0$

Conic Angles

$$\min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1}} u^\top v$$

Polynomial Time if
 $u^\top v \geq 0 \forall u \in P, v \in Q$
or for $n \leq 3$

Singular Values

$$\min_{\|u\|=\|v\|=1} u^\top A v$$

Polynomial Time
 $O(mn^2)$ to compute
all Singular Values

Conical Singular Values

$$\min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1,}} u^\top A v$$

P, Q

closed convex cones
finitely generated

Hardest Problem



Pareto Singular Values

$$\min_{\substack{u \geq 0, \|u\| = 1, \\ v \geq 0, \|v\| = 1,}} u^\top A v$$

Polynomial Time
for $A \geq 0$ or $A \leq 0$

Conic Angles

$$\min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1,}} u^\top v$$

Polynomial Time if
 $u^\top v \geq 0 \forall u \in P, v \in Q$
or for $n \leq 3$

Singular Values

$$\min_{\|u\|=\|v\|=1} u^\top A v$$

Polynomial Time
 $O(mn^2)$ to compute
all Singular Values

Lemma (B., G., S. 2024)

Any matrix $A \in \mathbb{R}^{m \times n}$ of spectral norm 1 and $m \geq n$ can be decomposed as $A = U^T V$ where $U \in \mathbb{R}^{(m+n) \times m}$, $V \in \mathbb{R}^{(m+n) \times n}$ have orthonormal columns

Proof: Let

$$A = U^T V \quad U = \begin{pmatrix} I \\ 0 \end{pmatrix} \in \mathbb{R}^{(m+n) \times m} \quad V := \begin{pmatrix} A \\ C \end{pmatrix} \in \mathbb{R}^{(m+n) \times n}$$

Given the SVD $A = W\Sigma Z^T$ let $C = (I - \Sigma^T \Sigma)^{1/2} Z^T$ so that

$$V^T V = A^T A + C^T C = A^T A + Z(I - \Sigma^T \Sigma)Z^T = ZZ^T = I$$

i.e. all columns of V are orthogonal to each other and with unitary norm

Notice:

The columns of U are a subset of the canonical basis, A is the projection of V on \mathbb{R}^m

Lemma (B., G., S. 2024)

Any matrix $A \in \mathbb{R}^{m \times n}$ of spectral norm 1 and $m \geq n$ can be decomposed as $A = U^T V$ where $U \in \mathbb{R}^{(m+n) \times m}$, $V \in \mathbb{R}^{(m+n) \times n}$ have orthonormal columns

Proof: Let

$$A = U^T V \quad U = \begin{pmatrix} I \\ 0 \end{pmatrix} \in \mathbb{R}^{(m+n) \times m} \quad V := \begin{pmatrix} A \\ C \end{pmatrix} \in \mathbb{R}^{(m+n) \times n}$$

Given the SVD $A = W\Sigma Z^\top$ let $C = (I - \Sigma^\top \Sigma)^{1/2} Z^\top$ so that

$$V^\top V = A^\top A + C^\top C = A^\top A + Z(I - \Sigma^\top \Sigma)Z^\top = ZZ^\top = I$$

i.e. all columns of V are orthogonal to each other and with unitary norm

Notice:

The columns of U are a subset of the canonical basis, A is the projection of V on \mathbb{R}^m

Lemma (B., G., S. 2024)

Any matrix $A \in \mathbb{R}^{m \times n}$ of spectral norm 1 and $m \geq n$ can be decomposed as $A = U^T V$ where $U \in \mathbb{R}^{(m+n) \times m}$, $V \in \mathbb{R}^{(m+n) \times n}$ have orthonormal columns

Proof: Let

$$A = U^T V \quad U = \begin{pmatrix} I \\ 0 \end{pmatrix} \in \mathbb{R}^{(m+n) \times m} \quad V := \begin{pmatrix} A \\ C \end{pmatrix} \in \mathbb{R}^{(m+n) \times n}$$

Given the SVD $A = W\Sigma Z^\top$ let $C = (I - \Sigma^\top \Sigma)^{1/2} Z^\top$ so that

$$V^\top V = A^\top A + C^\top C = A^\top A + Z(I - \Sigma^\top \Sigma)Z^\top = ZZ^\top = I$$

i.e. all columns of V are orthogonal to each other and with unitary norm

Notice:

The columns of U are a subset of the canonical basis, A is the projection of V on \mathbb{R}^m

Reduction

$$\min_{u,v} u^T A v : \quad \|u\| = \|v\| = 1 \quad u \in P = \langle G \rangle \subseteq \mathbb{R}^n \quad v \in Q = \langle H \rangle \subseteq \mathbb{R}^n$$

$$\min_{x,y} (Gx)^T A (Hy) : \quad \|Gx\| = \|Hy\| = 1 \quad x, y \geq 0$$

$$\|A\| \min_{x,y} x^T G^T U^T VH y : \quad \|Gx\| = \|Hy\| = 1 \quad x, y \geq 0$$

$$\|A\| \min_{x,y} x^T G^T U^T VH y : \quad \|UGx\| = \|VHy\| = 1 \quad x, y \geq 0$$

$$\|A\| \min_{\tilde{u}, \tilde{v}} \tilde{u}^T \tilde{v} : \quad \|\tilde{u}\| = \|\tilde{v}\| = 1 \quad \tilde{u} \in P' = \langle UG \rangle \subseteq \mathbb{R}^{n+m} \quad \tilde{v} \in Q' = \langle VH \rangle \subseteq \mathbb{R}^{n+m}$$

From Conical SV to Conic Angles

The minimum conical singular value of dimension n with number of generators p, q for G, H reduces polynomially to the maximum angle between cones of dimension $n + m$ with same number of generators for UG, VH

Reduction

$$\min_{u,v} u^\top A v : \|u\| = \|v\| = 1 \quad u \in P = \langle G \rangle \subseteq \mathbb{R}^n \quad v \in Q = \langle H \rangle \subseteq \mathbb{R}^n$$

$$\min_{x,y} (Gx)^\top A (Hy) : \|Gx\| = \|Hy\| = 1 \quad x, y \geq 0$$

$$\|A\| \min_{x,y} x^\top G^\top U^\top VH y : \|Gx\| = \|Hy\| = 1 \quad x, y \geq 0$$

$$\|A\| \min_{x,y} x^\top G^\top U^\top VH y : \|UGx\| = \|VH y\| = 1 \quad x, y \geq 0$$

$$\|A\| \min_{\tilde{u}, \tilde{v}} \tilde{u}^\top \tilde{v} : \|\tilde{u}\| = \|\tilde{v}\| = 1 \quad \tilde{u} \in P' = \langle UG \rangle \subseteq \mathbb{R}^{n+m} \quad \tilde{v} \in Q' = \langle VH \rangle \subseteq \mathbb{R}^{n+m}$$

From Conical SV to Conic Angles

The minimum conical singular value of dimension n with number of generators p, q for G, H reduces polynomially to the maximum angle between cones of dimension $n + m$ with same number of generators for UG, VH

Reduction

$$\min_{u,v} u^\top A v : \|u\| = \|v\| = 1 \quad u \in P = \langle G \rangle \subseteq \mathbb{R}^n \quad v \in Q = \langle H \rangle \subseteq \mathbb{R}^n$$

$$\min_{x,y} (Gx)^\top A (Hy) : \|Gx\| = \|Hy\| = 1 \quad x, y \geq 0$$

$$\|A\| \min_{x,y} x^\top G^\top U^\top VH y : \|Gx\| = \|Hy\| = 1 \quad x, y \geq 0$$

$$\|A\| \min_{x,y} x^\top G^\top U^\top VH y : \|UGx\| = \|VH y\| = 1 \quad x, y \geq 0$$

$$\|A\| \min_{\tilde{u}, \tilde{v}} \tilde{u}^\top \tilde{v} : \|\tilde{u}\| = \|\tilde{v}\| = 1 \quad \tilde{u} \in P' = \langle UG \rangle \subseteq \mathbb{R}^{n+m} \quad \tilde{v} \in Q' = \langle VH \rangle \subseteq \mathbb{R}^{n+m}$$

From Conical SV to Conic Angles

The minimum conical singular value of dimension n with number of generators p, q for G, H reduces polynomially to the maximum angle between cones of dimension $n + m$ with same number of generators for UG, VH

Reduction

$$\min_{u,v} u^\top A v : \|u\| = \|v\| = 1 \quad u \in P = \langle G \rangle \subseteq \mathbb{R}^n \quad v \in Q = \langle H \rangle \subseteq \mathbb{R}^n$$

$$\min_{x,y} (Gx)^\top A (Hy) : \|Gx\| = \|Hy\| = 1 \quad x, y \geq 0$$

$$\|A\| \min_{x,y} x^\top G^\top U^\top VH y : \|Gx\| = \|Hy\| = 1 \quad x, y \geq 0$$

$$\|A\| \min_{x,y} x^\top G^\top U^\top VH y : \|UGx\| = \|VH y\| = 1 \quad x, y \geq 0$$

$$\|A\| \min_{\tilde{u}, \tilde{v}} \tilde{u}^\top \tilde{v} : \|\tilde{u}\| = \|\tilde{v}\| = 1 \quad \tilde{u} \in P' = \langle UG \rangle \subseteq \mathbb{R}^{n+m} \quad \tilde{v} \in Q' = \langle VH \rangle \subseteq \mathbb{R}^{n+m}$$

From Conical SV to Conic Angles

The minimum conical singular value of dimension n with number of generators p, q for G, H reduces polynomially to the maximum angle between cones of dimension $n + m$ with same number of generators for UG, VH

Reduction

$$\min_{u,v} u^\top A v : \|u\| = \|v\| = 1 \quad u \in P = \langle G \rangle \subseteq \mathbb{R}^n \quad v \in Q = \langle H \rangle \subseteq \mathbb{R}^n$$

$$\min_{x,y} (Gx)^\top A (Hy) : \|Gx\| = \|Hy\| = 1 \quad x, y \geq 0$$

$$\|A\| \min_{x,y} x^\top G^\top U^\top VH y : \|Gx\| = \|Hy\| = 1 \quad x, y \geq 0$$

$$\|A\| \min_{x,y} x^\top G^\top U^\top VH y : \|UGx\| = \|VH y\| = 1 \quad x, y \geq 0$$

$$\|A\| \min_{\tilde{u}, \tilde{v}} \tilde{u}^\top \tilde{v} : \|\tilde{u}\| = \|\tilde{v}\| = 1 \quad \tilde{u} \in P' = \langle UG \rangle \subseteq \mathbb{R}^{n+m} \quad \tilde{v} \in Q' = \langle VH \rangle \subseteq \mathbb{R}^{n+m}$$

From Conical SV to Conic Angles

The minimum conical singular value of dimension n with number of generators p, q for G, H reduces polynomially to the maximum angle between cones of dimension $n + m$ with same number of generators for UG, VH

Reduction

$$\min_{u,v} u^\top A v : \|u\| = \|v\| = 1 \quad u \in P = \langle G \rangle \subseteq \mathbb{R}^n \quad v \in Q = \langle H \rangle \subseteq \mathbb{R}^n$$

$$\min_{x,y} (Gx)^\top A (Hy) : \|Gx\| = \|Hy\| = 1 \quad x, y \geq 0$$

$$\|A\| \min_{x,y} x^\top G^\top U^\top VH y : \|Gx\| = \|Hy\| = 1 \quad x, y \geq 0$$

$$\|A\| \min_{x,y} x^\top G^\top U^\top VH y : \|UGx\| = \|VHy\| = 1 \quad x, y \geq 0$$

$$\|A\| \min_{\tilde{u}, \tilde{v}} \tilde{u}^\top \tilde{v} : \|\tilde{u}\| = \|\tilde{v}\| = 1 \quad \tilde{u} \in P' = \langle UG \rangle \subseteq \mathbb{R}^{n+m} \quad \tilde{v} \in Q' = \langle VH \rangle \subseteq \mathbb{R}^{n+m}$$

From Conical SV to Conic Angles

The minimum conical singular value of dimension n with number of generators p, q for G, H reduces polynomially to the maximum angle between cones of dimension $n + m$ with same number of generators for UG, VH

Conical Singular Values

$$\min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1,}} u^\top A v \quad P, Q \text{ closed convex cones}$$

finitely generated



Pareto Singular Values

$$\min_{\substack{u \geq 0, \|u\| = 1, \\ v \geq 0, \|v\| = 1,}} u^\top A v$$

Conic Angles

$$\min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1,}} u^\top v$$

How difficult is the Pareto SV problem?

Conical Singular Values

$$\min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1,}} u^\top A v \quad P, Q \text{ closed convex cones finitely generated}$$



Pareto Singular Values

$$\min_{\substack{u \geq 0, \|u\| = 1, \\ v \geq 0, \|v\| = 1,}} u^\top A v$$

Conic Angles

$$\min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1,}} u^\top v$$



How difficult is the Pareto SV problem?

Conical Singular Values

$$\min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1,}} u^\top A v \quad P, Q \text{ closed convex cones}$$

finitely generated



Pareto Singular Values

$$\min_{\substack{u \geq 0, \|u\| = 1, \\ v \geq 0, \|v\| = 1,}} u^\top A v$$

Conic Angles

$$\min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1,}} u^\top v$$



How difficult is the Pareto SV problem?

Reduction from Maximum Edge Biclique to Minimal Pareto Singular Value

Theorem (Seeger, S. 2023)

Let (σ_0, u, v) be the optimal solution of

$$\sigma_0 = \min_{u, v \geq 0} u^\top A v \quad : \quad \|u\| = \|v\| = 1$$

If A has at least one negative entry then $(x, y) = \sqrt{-\sigma_0}(u, v)$ is optimal for

$$\min_{x, y \geq 0} \| -A - xy^\top \|_F^2$$

This shows that the Minimal Pareto Singular Value is at least as hard as the
Nonnegative Rank 1 Approximation problem

Theorem (G., Glineur 2013)

Let $B \in \{0, 1\}^{m \times n}$ be the bi-adjacency matrix of a bipartite graph (N_1, N_2, E) where $B_{i,j} = 1$ iff node i in N_1 and node j in N_2 are connected and $d \geq \max\{m, n\}$.

$$\min_{x, y \geq 0} \|B - d(1 - B) - xy^\top\|_F^2$$

is solved by binary vectors x, y that identify the fully connected subsets $S_1 \subseteq N_1$ and $S_2 \subseteq N_2$ corresponding to the Maximum Edge Biclique, i.e they maximise $|S_1| \cdot |S_2|$

Reduction from Maximum Edge Biclique to Minimal Pareto Singular Value

Theorem (Seeger, S. 2023)

Let (σ_0, u, v) be the optimal solution of

$$\sigma_0 = \min_{u, v \geq 0} u^\top A v \quad : \quad \|u\| = \|v\| = 1$$

If A has at least one negative entry then $(x, y) = \sqrt{-\sigma_0}(u, v)$ is optimal for

$$\min_{x, y \geq 0} \| -A - xy^\top \|_F^2$$

This shows that the Minimal Pareto Singular Value is **at least as hard** as the
Nonnegative Rank 1 Approximation problem

Theorem (G., Glineur 2013)

Let $B \in \{0, 1\}^{m \times n}$ be the bi-adjacency matrix of a bipartite graph (N_1, N_2, E) where $B_{i,j} = 1$ iff node i in N_1 and node j in N_2 are connected and $d \geq \max\{m, n\}$.

$$\min_{x, y \geq 0} \|B - d(1 - B) - xy^\top\|_F^2$$

is solved by binary vectors x, y that identify the fully connected subsets $S_1 \subseteq N_1$ and $S_2 \subseteq N_2$ corresponding to the **Maximum Edge Biclique**, i.e they maximise $|S_1| \cdot |S_2|$

Reduction from Maximum Edge Biclique to Minimal Pareto Singular Value

Theorem (Seeger, S. 2023)

Let (σ_0, u, v) be the optimal solution of

$$\sigma_0 = \min_{u, v \geq 0} u^\top A v \quad : \quad \|u\| = \|v\| = 1$$

If A has at least one negative entry then $(x, y) = \sqrt{-\sigma_0}(u, v)$ is optimal for

$$\min_{x, y \geq 0} \| -A - xy^\top \|_F^2$$

This shows that the Minimal Pareto Singular Value is **at least as hard** as the
Nonnegative Rank 1 Approximation problem

Theorem (G., Glineur 2013)

Let $B \in \{0, 1\}^{m \times n}$ be the bi-adjacency matrix of a bipartite graph (N_1, N_2, E) where $B_{i,j} = 1$ iff node i in N_1 and node j in N_2 are connected and $d \geq \max\{m, n\}$.

$$\min_{x, y \geq 0} \|B - d(1 - B) - xy^\top\|_F^2$$

is solved by binary vectors x, y that identify the fully connected subsets $S_1 \subseteq N_1$ and $S_2 \subseteq N_2$ corresponding to the **Maximum Edge Biclique**, i.e they maximise $|S_1| \cdot |S_2|$

- $\min_{x,y \geq 0} \|B - d(1 - B) - xy^\top\|_F^2$ identifies the Maximum Edge Biclique

$$M = B - d(1 - B) = \begin{pmatrix} -d & 1 & 1 & -d & 1 & 1 & -d \\ 1 & 1 & -d & -d & 1 & 1 & 1 \\ -d & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & -d & 1 & 1 & 1 & -d \\ 1 & -d & 1 & -d & -d & 1 & 1 \end{pmatrix}$$

Idea of Proof:

All the Maximal Bicliques (S_1, S_2) are local minima of $\|M - xy^\top\|_F^2$ where $x = \chi(S_1)$, $y = \chi(S_2)$ because any extension of $S_1 \times S_2$ gets a $-d$ and

$$(-d - \epsilon)^2 = d^2 + 2d\epsilon + \epsilon^2$$

with the error going up by $d\epsilon$ that is way more than what we gain by removing ϵ from all the ones in a row/column

As a consequence xy^\top has zeros in correspondence of the $-d$ of M and the rest nonzero entries equal to 1, meaning that local minima x, y are indicator for the maximal bicliques

- $\min_{x,y \geq 0} \|B - d(1 - B) - xy^\top\|_F^2$ identifies the Maximum Edge Biclique

$$M = B - d(1 - B) = \begin{pmatrix} -d & 1 & 1 & -d & 1 & 1 & -d \\ 1 & 1 & -d & -d & 1 & 1 & 1 \\ -d & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & -d & 1 & 1 & 1 & -d \\ 1 & -d & 1 & -d & -d & 1 & 1 \end{pmatrix}$$

Idea of Proof:

All the Maximal Bicliques (S_1, S_2) are local minima of $\|M - xy^\top\|_F^2$ where $x = \chi(S_1)$, $y = \chi(S_2)$ because any extension of $S_1 \times S_2$ gets a $-d$ and

$$(-d - \epsilon)^2 = d^2 + 2d\epsilon + \epsilon^2$$

with the error going up by $d\epsilon$ that is way more than what we gain by removing ϵ from all the ones in a row/column

As a consequence xy^\top has zeros in correspondence of the $-d$ of M and the rest nonzero entries equal to 1, meaning that local minima x, y are indicator for the maximal bicliques

- $\min_{x,y \geq 0} \|B - d(1 - B) - xy^\top\|_F^2$ identifies the Maximum Edge Biclique

$$M = B - d(1 - B) = \begin{pmatrix} -d & 1 & 1 & -d & 1 & 1 & -d \\ 1 & 1 & -d & -d & 1 & 1 & 1 \\ -d & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & -d & 1 & 1 & 1 & -d \\ 1 & -d & 1 & -d & -d & 1 & 1 \end{pmatrix}$$

Idea of Proof:

All the Maximal Bicliques (S_1, S_2) are local minima of $\|M - xy^\top\|_F^2$ where $x = \chi(S_1)$, $y = \chi(S_2)$ because any extension of $S_1 \times S_2$ gets a $-d$ and

$$(-d - \epsilon)^2 = d^2 + 2d\epsilon + \epsilon^2$$

with the error going up by $d\epsilon$ that is way more than what we gain by removing ϵ from all the ones in a row/column

As a consequence xy^\top has zeros in correspondence of the $-d$ of M and the rest nonzero entries equal to 1, meaning that local minima x, y are indicator for the maximal bicliques

- $\min_{x,y \geq 0} \|B - d(1 - B) - xy^\top\|_F^2$ identifies the Maximum Edge Biclique

$$M = B - d(1 - B) = \begin{pmatrix} -d & 1 & 1 & -d & 1 & 1 & -d \\ 1 & 1 & -d & -d & 1 & 1 & 1 \\ -d & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & -d & 1 & 1 & 1 & -d \\ 1 & -d & 1 & -d & -d & 1 & 1 \end{pmatrix}$$

Idea of Proof:

All the Maximal Bicliques (S_1, S_2) are local minima of $\|M - xy^\top\|_F^2$ where $x = \chi(S_1)$, $y = \chi(S_2)$ because any extension of $S_1 \times S_2$ gets a $-d$ and

$$(-d - \epsilon)^2 = d^2 + 2d\epsilon + \epsilon^2$$

with the error going up by $d\epsilon$ that is way more than what we gain by removing ϵ from all the ones in a row/column

As a consequence xy^\top has zeros in correspondence of the $-d$ of M and the rest nonzero entries equal to 1, meaning that local minima x, y are indicator for the maximal bicliques

Everything is Hard

Theorem (Peeters 2003)

The Maximal Edge Biclique problem is NP-hard

Maximal Edge Biclique

Maximum Number of
Edges in a Bipartite
Connected Subgraph

NP-hard

Everything is Hard

Theorem (Peeters 2003)

The Maximal Edge Biclique problem is NP-hard

Maximal Edge Biclique

Maximum Number of
Edges in a Bipartite
Connected Subgraph

NP-hard



Nonnegative Rank 1

$$\min_{x,y \geq 0} \|M - xy^\top\|$$

NP-hard

Everything is Hard

Theorem (Peeters 2003)

The Maximal Edge Biclique problem is NP-hard

Maximal Edge Biclique

Maximum Number of
Edges in a Bipartite
Connected Subgraph

NP-hard



**Pareto Singular
Values**

$$\begin{array}{ll} \min & u^\top A v \\ \text{s.t.} & u \geq 0, \|u\| = 1, \\ & v \geq 0, \|v\| = 1, \end{array}$$

**Nonnegative
Rank 1**

$$\min_{x,y \geq 0} \|M - xy^\top\|$$

NP-hard



NP-hard

Everything is Hard

Theorem (Peeters 2003)

The Maximal Edge Biclique problem is NP-hard

Maximal Edge Biclique

Maximum Number of Edges in a Bipartite Connected Subgraph

NP-hard



Nonnegative Rank 1

$$\min_{x,y \geq 0} \|M - xy^\top\|$$

NP-hard

Pareto Singular Values

$$\begin{array}{ll} \min & u^\top Av \\ \begin{array}{l} u \geq 0, \|u\| = 1, \\ v \geq 0, \|v\| = 1 \end{array} & \end{array}$$



NP-hard

Conic Singular Values

$$\begin{array}{ll} \min & u^\top Av \\ \begin{array}{l} u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1, \end{array} & \end{array}$$

NP-hard



Conic Angles

$$\begin{array}{ll} \min & u^\top v \\ \begin{array}{l} u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1, \end{array} & \end{array}$$

NP-hard



Recall: from $A = \|A\|U^\top V$

$$\min_{\substack{x \geq 0, \|x\| = 1, \\ y \geq 0, \|y\| = 1}} \langle x, Ay \rangle = \|A\| \quad \min_{\substack{x \geq 0, \|Ux\| = 1, \\ y \geq 0, \|Vy\| = 1}} \langle Ux, Vy \rangle = \|A\| \quad \min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1}} \langle u, v \rangle,$$

with $U^\top = \begin{pmatrix} I & 0 \end{pmatrix}$

Theorem (B., G., S. 2024)

The maximum angle between convex closed cones problem

$$\min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1}} \langle u, v \rangle$$

with P being generated by a subset of the canonical basis is NP-hard

Conjecture (B., G., S. 2024)

The maximum angle between the positive orthant and another convex closed cone

$$\min_{\substack{x \geq 0, \|x\| = 1, \\ v \in Q, \|v\| = 1}} \langle x, v \rangle = - \max_{v \in Q, \|v\|=1} \|v^\perp\|$$

is NP-hard

Recall: from $A = \|A\|U^\top V$

$$\min_{\substack{x \geq 0, \|x\| = 1, \\ y \geq 0, \|y\| = 1}} \langle x, Ay \rangle = \|A\| \quad \min_{\substack{x \geq 0, \|Ux\| = 1, \\ y \geq 0, \|Vy\| = 1}} \langle Ux, Vy \rangle = \|A\| \quad \min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1}} \langle u, v \rangle,$$

with $U^\top = \begin{pmatrix} I & 0 \end{pmatrix}$

Theorem (B., G., S. 2024)

The maximum angle between convex closed cones problem

$$\min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1}} \langle u, v \rangle$$

with P being generated by a subset of the canonical basis is NP-hard

Conjecture (B., G., S. 2024)

The maximum angle between the positive orthant and another convex closed cone

$$\min_{\substack{x \geq 0, \|x\| = 1, \\ v \in Q, \|v\| = 1}} \langle x, v \rangle = - \max_{v \in Q, \|v\|=1} \|v^\perp\|$$

is NP-hard

Recall: from $A = \|A\|U^\top V$

$$\min_{\substack{x \geq 0, \|x\| = 1, \\ y \geq 0, \|y\| = 1}} \langle x, Ay \rangle = \|A\| \quad \min_{\substack{x \geq 0, \|Ux\| = 1, \\ y \geq 0, \|Vy\| = 1}} \langle Ux, Vy \rangle = \|A\| \quad \min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1}} \langle u, v \rangle,$$

with $U^\top = \begin{pmatrix} I & 0 \end{pmatrix}$

Theorem (B., G., S. 2024)

The maximum angle between convex closed cones problem

$$\min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1}} \langle u, v \rangle$$

with P being generated by a subset of the canonical basis is NP-hard

Conjecture (B., G., S. 2024)

The maximum angle between the positive orthant and another convex closed cone

$$\min_{\substack{x \geq 0, \|x\| = 1, \\ v \in Q, \|v\| = 1}} \langle x, v \rangle = - \max_{v \in Q, \|v\|=1} \|v^- \|$$

is NP-hard

Algorithms

Active Set

$$\lambda^* = \min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1}} u^\top A v = \min_{\substack{x \geq 0, \|Gx\| = 1, \\ y \geq 0, \|Hy\| = 1}} x^\top G^\top A H y$$

Idea: If we know the sets \mathcal{I} , \mathcal{J} of indices for which $x_i^*, y_j^* > 0$, called **Active Sets**, then a direct gradient computation solves the problem

KKT Conditions:

$$\begin{cases} P \ni u^* \perp (A v^* - \lambda^* u^*) \in P^* \\ Q \ni v^* \perp (A^\top u^* - \lambda^* v^*) \in Q^* \\ \|u^*\| = \|v^*\| = 1 \end{cases} \rightarrow \begin{cases} 0 \leq x^* \perp G^\top A H y^* - \lambda^* G^\top G x^* \geq 0 \\ 0 \leq y^* \perp H^\top A^\top G x^* - \lambda^* H^\top H y^* \geq 0 \\ \|Gx^*\| = \|Hy^*\| = 1 \end{cases}$$

$$\begin{cases} 0 < \bar{x} \quad \bar{G}^\top A \bar{H} \bar{y} - \lambda^* \bar{G}^\top \bar{G} \bar{x} = 0 \\ 0 < \bar{y} \quad \bar{H}^\top A^\top \bar{G} \bar{x} - \lambda^* \bar{H}^\top \bar{H} \bar{y} = 0 \\ \bar{G}^\top A \bar{H} \bar{y} - \lambda^* \bar{G}^\top \bar{G} \bar{x} \geq 0 \\ \bar{H}^\top A^\top \bar{G} \bar{x} - \lambda^* \bar{H}^\top \bar{H} \bar{y} \geq 0 \\ \|\bar{G} \bar{x}\| = \|\bar{H} \bar{y}\| = 1 \end{cases}$$

where $\begin{cases} \bar{x} := x_{\mathcal{I}}^*, \bar{y} := y_{\mathcal{J}}^* \\ \bar{G} := G_{:, \mathcal{I}}, \bar{H} := H_{:, \mathcal{J}} \\ \tilde{G} := G_{:, \mathcal{I}^c}, \tilde{H} := H_{:, \mathcal{J}^c} \\ \lambda^* = \bar{x}^\top \bar{G}^\top A \bar{H} \bar{y} = (u^*)^\top A v^* \end{cases}$

Active Set

$$\lambda^* = \min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1}} u^\top A v = \min_{\substack{x \geq 0, \|Gx\| = 1, \\ y \geq 0, \|Hy\| = 1}} x^\top G^\top A H y$$

Idea: If we know the sets \mathcal{I} , \mathcal{J} of indices for which $x_i^*, y_j^* > 0$, called **Active Sets**, then a direct gradient computation solves the problem

KKT Conditions:

$$\begin{cases} P \ni u^* \perp (A v^* - \lambda^* u^*) \in P^* \\ Q \ni v^* \perp (A^\top u^* - \lambda^* v^*) \in Q^* \\ \|u^*\| = \|v^*\| = 1 \end{cases} \rightarrow \begin{cases} 0 \leq x^* \perp G^\top A H y^* - \lambda^* G^\top G x^* \geq 0 \\ 0 \leq y^* \perp H^\top A^\top G x^* - \lambda^* H^\top H y^* \geq 0 \\ \|Gx^*\| = \|Hy^*\| = 1 \end{cases}$$

$$\begin{cases} 0 < \bar{x} \quad \bar{G}^\top A \bar{H} \bar{y} - \lambda^* \bar{G}^\top \bar{G} \bar{x} = 0 \\ 0 < \bar{y} \quad \bar{H}^\top A^\top \bar{G} \bar{x} - \lambda^* \bar{H}^\top \bar{H} \bar{y} = 0 \\ \bar{G}^\top A \bar{H} \bar{y} - \lambda^* \bar{G}^\top \bar{G} \bar{x} \geq 0 \\ \bar{H}^\top A^\top \bar{G} \bar{x} - \lambda^* \bar{H}^\top \bar{H} \bar{y} \geq 0 \\ \|\bar{G} \bar{x}\| = \|\bar{H} \bar{y}\| = 1 \end{cases}$$

where $\begin{cases} \bar{x} := x_{\mathcal{I}}^*, \bar{y} := y_{\mathcal{J}}^* \\ \bar{G} := G_{:, \mathcal{I}}, \bar{H} := H_{:, \mathcal{J}} \\ \tilde{G} := G_{:, \mathcal{I}^c}, \tilde{H} := H_{:, \mathcal{J}^c} \\ \lambda^* = \bar{x}^\top \bar{G}^\top A \bar{H} \bar{y} = (u^*)^\top A v^* \end{cases}$

Active Set

$$\lambda^* = \min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1}} u^\top A v = \min_{\substack{x \geq 0, \|Gx\| = 1, \\ y \geq 0, \|Hy\| = 1}} x^\top G^\top A H y$$

Idea: If we know the sets \mathcal{I} , \mathcal{J} of indices for which $x_i^*, y_j^* > 0$, called **Active Sets**, then a direct gradient computation solves the problem

KKT Conditions:

$$\begin{cases} P \ni u^* \perp (A v^* - \lambda^* u^*) \in P^* \\ Q \ni v^* \perp (A^\top u^* - \lambda^* v^*) \in Q^* \\ \|u^*\| = \|v^*\| = 1 \end{cases} \rightarrow \begin{cases} 0 \leq x^* \perp G^\top A H y^* - \lambda^* G^\top G x^* \geq 0 \\ 0 \leq y^* \perp H^\top A^\top G x^* - \lambda^* H^\top H y^* \geq 0 \\ \|Gx^*\| = \|Hy^*\| = 1 \end{cases}$$

$$\begin{cases} 0 < \bar{x} \quad \bar{G}^\top A \bar{H} \bar{y} - \lambda^* \bar{G}^\top \bar{G} \bar{x} = 0 \\ 0 < \bar{y} \quad \bar{H}^\top A^\top \bar{G} \bar{x} - \lambda^* \bar{H}^\top \bar{H} \bar{y} = 0 \\ \bar{G}^\top A \bar{H} \bar{y} - \lambda^* \bar{G}^\top \bar{G} \bar{x} \geq 0 \\ \bar{H}^\top A^\top \bar{G} \bar{x} - \lambda^* \bar{H}^\top \bar{H} \bar{y} \geq 0 \\ \|\bar{G} \bar{x}\| = \|\bar{H} \bar{y}\| = 1 \end{cases}$$

where $\begin{cases} \bar{x} := x_{\mathcal{I}}^*, \bar{y} := y_{\mathcal{J}}^* \\ \bar{G} := G_{:, \mathcal{I}}, \bar{H} := H_{:, \mathcal{J}} \\ \tilde{G} := G_{:, \mathcal{I}^c}, \tilde{H} := H_{:, \mathcal{J}^c} \\ \lambda^* = \bar{x}^\top \bar{G}^\top A \bar{H} \bar{y} = (u^*)^\top A v^* \end{cases}$

Active Set

$$\lambda^* = \min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1}} u^\top A v = \min_{\substack{x \geq 0, \|Gx\| = 1, \\ y \geq 0, \|Hy\| = 1}} x^\top G^\top A H y$$

Idea: If we know the sets \mathcal{I} , \mathcal{J} of indices for which $x_i^*, y_j^* > 0$, called **Active Sets**, then a direct gradient computation solves the problem

KKT Conditions:

$$\begin{cases} P \ni u^* \perp (A v^* - \lambda^* u^*) \in P^* \\ Q \ni v^* \perp (A^\top u^* - \lambda^* v^*) \in Q^* \\ \|u^*\| = \|v^*\| = 1 \end{cases} \rightarrow \begin{cases} 0 \leq x^* \perp G^\top A H y^* - \lambda^* G^\top G x^* \geq 0 \\ 0 \leq y^* \perp H^\top A^\top G x^* - \lambda^* H^\top H y^* \geq 0 \\ \|Gx^*\| = \|Hy^*\| = 1 \end{cases}$$

$$\begin{cases} 0 < \bar{x} \quad \bar{G}^\top A \bar{H} \bar{y} - \lambda^* \bar{G}^\top \bar{G} \bar{x} = 0 \\ 0 < \bar{y} \quad \bar{H}^\top A^\top \bar{G} \bar{x} - \lambda^* \bar{H}^\top \bar{H} \bar{y} = 0 \\ \bar{G}^\top A \bar{H} \bar{y} - \lambda^* \bar{G}^\top \bar{G} \bar{x} \geq 0 \\ \bar{H}^\top A^\top \bar{G} \bar{x} - \lambda^* \bar{H}^\top \bar{H} \bar{y} \geq 0 \\ \|\bar{G} \bar{x}\| = \|\bar{H} \bar{y}\| = 1 \end{cases}$$

where $\begin{cases} \bar{x} := x_{\mathcal{I}}^*, \bar{y} := y_{\mathcal{J}}^* \\ \bar{G} := G_{:, \mathcal{I}}, \bar{H} := H_{:, \mathcal{J}} \\ \tilde{G} := G_{:, \mathcal{I}^c}, \tilde{H} := H_{:, \mathcal{J}^c} \\ \lambda^* = \bar{x}^\top \bar{G}^\top A \bar{H} \bar{y} = (u^*)^\top A v^* \end{cases}$

Active Set

$$\lambda^* = \min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1}} u^\top A v = \min_{\substack{x \geq 0, \|Gx\| = 1, \\ y \geq 0, \|Hy\| = 1}} x^\top G^\top A H y$$

Idea: If we know the sets \mathcal{I} , \mathcal{J} of indices for which $x_i^*, y_j^* > 0$, called **Active Sets**, then a direct gradient computation solves the problem

KKT Conditions:

$$\begin{cases} P \ni u^* \perp (A v^* - \lambda^* u^*) \in P^* \\ Q \ni v^* \perp (A^\top u^* - \lambda^* v^*) \in Q^* \\ \|u^*\| = \|v^*\| = 1 \end{cases} \rightarrow \begin{cases} 0 \leq x^* \perp G^\top A H y^* - \lambda^* G^\top G x^* \geq 0 \\ 0 \leq y^* \perp H^\top A^\top G x^* - \lambda^* H^\top H y^* \geq 0 \\ \|Gx^*\| = \|Hy^*\| = 1 \end{cases}$$

$$\begin{cases} 0 < \bar{x} \quad \bar{G}^\top A \bar{H} \bar{y} - \lambda^* \bar{G}^\top \bar{G} \bar{x} = 0 \\ 0 < \bar{y} \quad \bar{H}^\top A^\top \bar{G} \bar{x} - \lambda^* \bar{H}^\top \bar{H} \bar{y} = 0 \\ \bar{G}^\top A \bar{H} \bar{y} - \lambda^* \bar{G}^\top \bar{G} \bar{x} \geq 0 \\ \bar{H}^\top A^\top \bar{G} \bar{x} - \lambda^* \bar{H}^\top \bar{H} \bar{y} \geq 0 \\ \|\bar{G} \bar{x}\| = \|\bar{H} \bar{y}\| = 1 \end{cases}$$

where $\begin{cases} \bar{x} := x_{\mathcal{I}}^*, \bar{y} := y_{\mathcal{J}}^* \\ \bar{G} := G_{:, \mathcal{I}}, \bar{H} := H_{:, \mathcal{J}} \\ \tilde{G} := G_{:, \mathcal{I}^c}, \tilde{H} := H_{:, \mathcal{J}^c} \\ \lambda^* = \bar{x}^\top \bar{G}^\top A \bar{H} \bar{y} = (u^*)^\top A v^* \end{cases}$

Active Set

$$\begin{cases} 0 < \bar{x} & \bar{G}^\top A \bar{H} \bar{y} - \lambda^* \bar{G}^\top \bar{G} \bar{x} = 0 \\ 0 < \bar{y} & \bar{H}^\top A^\top \bar{G} \bar{x} - \lambda^* \bar{H}^\top \bar{H} \bar{y} = 0 \\ \bar{G}^\top A \bar{H} \bar{y} - \lambda^* \bar{G}^\top \bar{G} \bar{x} \geq 0 \\ \bar{H}^\top A^\top \bar{G} \bar{x} - \lambda^* \bar{H}^\top \bar{H} \bar{y} \geq 0 \\ \|\bar{G} \bar{x}\| = \|\bar{H} \bar{y}\| = 1 \end{cases}$$

where $\begin{cases} \bar{x} := x_{\mathcal{I}}^*, \bar{y} := y_{\mathcal{J}}^* \\ \bar{G} := G_{:, \mathcal{I}}, \bar{H} := H_{:, \mathcal{J}} \\ \tilde{G} := G_{:, \mathcal{I}^c}, \tilde{H} := H_{:, \mathcal{J}^c} \\ \lambda^* = \bar{x}^\top \bar{G}^\top A \bar{H} \bar{y} = (u^*)^\top A v^* \end{cases}$

Theorem (B., G., S. 2024)

For the optimal solution $(u^*, v^*) = (Gx^*, Hy^*) = (\bar{G}\bar{x}, \bar{H}\bar{y})$, and $\lambda^* = (u^*)^\top A v^*$ we have that

$$M^* := \begin{pmatrix} 0 & \bar{H}^\top A^\top \bar{G} \\ \bar{G}^\top A \bar{H} & 0 \end{pmatrix} \implies M^* \begin{pmatrix} \bar{y} \\ \bar{x} \end{pmatrix} = \lambda^* \begin{pmatrix} \bar{y} \\ \bar{x} \end{pmatrix}$$

and λ^* is the least eigenvalue of M^* (from 2° order KKT)

Viceversa, if M is generated by random sets \mathcal{I}, \mathcal{J} , λ is its least eigenvalue and it admits a nonnegative eigenvector, then $\lambda \geq \lambda^*$.

The Active Set algorithm cycles over all subsets of indices \mathcal{I}, \mathcal{J} and tests if the least eigenvalue of M has a nonnegative eigenvector, giving us upper bounds on λ^* , and one of them will be the exact solution

Active Set

$$\begin{cases} 0 < \bar{x} & \bar{G}^\dagger A \bar{H} \bar{y} - \lambda^* \bar{x} = 0 \\ 0 < \bar{y} & \bar{H}^\dagger A^\top \bar{G} \bar{x} - \lambda^* \bar{y} = 0 \\ \tilde{G}^\top A \bar{H} \bar{y} - \lambda^* \tilde{G}^\top \bar{G} \bar{x} \geq 0 \\ \tilde{H}^\top A^\top \bar{G} \bar{x} - \lambda^* \tilde{H}^\top \bar{H} \bar{y} \geq 0 \\ \|\bar{G} \bar{x}\| = \|\bar{H} \bar{y}\| = 1 \end{cases}$$

where $\begin{cases} \bar{x} := x_{\mathcal{I}}^*, \bar{y} := y_{\mathcal{J}}^* \\ \bar{G} := G_{:, \mathcal{I}}, \bar{H} := H_{:, \mathcal{J}} \\ \tilde{G} := G_{:, \mathcal{I}^c}, \tilde{H} := H_{:, \mathcal{J}^c} \\ \lambda^* = \bar{x}^\top \bar{G}^\top A \bar{H} \bar{y} = (u^*)^\top A v^* \end{cases}$

Theorem (B., G., S. 2024)

For the optimal solution $(u^*, v^*) = (Gx^*, Hy^*) = (\bar{G}\bar{x}, \bar{H}\bar{y})$, and $\lambda^* = (u^*)^\top A v^*$ we have that

$$M^* := \begin{pmatrix} 0 & \bar{H}^\dagger A^\top \bar{G} \\ \bar{G}^\dagger A \bar{H} & 0 \end{pmatrix} \implies M^* \begin{pmatrix} \bar{y} \\ \bar{x} \end{pmatrix} = \lambda^* \begin{pmatrix} \bar{y} \\ \bar{x} \end{pmatrix}$$

and λ^* is the least eigenvalue of M^* (from 2° order KKT)

Viceversa, if M is generated by random sets \mathcal{I}, \mathcal{J} , λ is its least eigenvalue and it admits a nonnegative eigenvector, then $\lambda \geq \lambda^*$.

The Active Set algorithm cycles over all subsets of indices \mathcal{I}, \mathcal{J} and tests if the least eigenvalue of M has a nonnegative eigenvector, giving us upper bounds on λ^* , and one of them will be the exact solution

Active Set

$$\begin{cases} 0 < \bar{x} & \bar{G}^\dagger A \bar{H} \bar{y} - \lambda^* \bar{x} = 0 \\ 0 < \bar{y} & \bar{H}^\dagger A^\top \bar{G} \bar{x} - \lambda^* \bar{y} = 0 \\ \tilde{G}^\top A \bar{H} \bar{y} - \lambda^* \tilde{G}^\top \bar{G} \bar{x} \geq 0 \\ \tilde{H}^\top A^\top \bar{G} \bar{x} - \lambda^* \tilde{H}^\top \bar{H} \bar{y} \geq 0 \\ \|\bar{G} \bar{x}\| = \|\bar{H} \bar{y}\| = 1 \end{cases}$$

where $\begin{cases} \bar{x} := x_{\mathcal{I}}^*, \bar{y} := y_{\mathcal{J}}^* \\ \bar{G} := G_{:, \mathcal{I}}, \bar{H} := H_{:, \mathcal{J}} \\ \tilde{G} := G_{:, \mathcal{I}^c}, \tilde{H} := H_{:, \mathcal{J}^c} \\ \lambda^* = \bar{x}^\top \bar{G}^\top A \bar{H} \bar{y} = (u^*)^\top A v^* \end{cases}$

Theorem (B., G., S. 2024)

For the optimal solution $(u^*, v^*) = (Gx^*, Hy^*) = (\bar{G}\bar{x}, \bar{H}\bar{y})$, and $\lambda^* = (u^*)^\top A v^*$ we have that

$$M^* := \begin{pmatrix} 0 & \bar{H}^\dagger A^\top \bar{G} \\ \bar{G}^\dagger A \bar{H} & 0 \end{pmatrix} \implies M^* \begin{pmatrix} \bar{y} \\ \bar{x} \end{pmatrix} = \lambda^* \begin{pmatrix} \bar{y} \\ \bar{x} \end{pmatrix}$$

and λ^* is the least eigenvalue of M^* (from 2° order KKT)

Viceversa, if M is generated by random sets \mathcal{I}, \mathcal{J} , λ is its least eigenvalue and it admits a nonnegative eigenvector, then $\lambda \geq \lambda^*$.

The Active Set algorithm cycles over all subsets of indices \mathcal{I}, \mathcal{J} and tests if the least eigenvalue of M has a nonnegative eigenvector, giving us upper bounds on λ^* , and one of them will be the exact solution

Active Set

$$\begin{cases} 0 < \bar{x} & \bar{G}^\dagger A \bar{H} \bar{y} - \lambda^* \bar{x} = 0 \\ 0 < \bar{y} & \bar{H}^\dagger A^\top \bar{G} \bar{x} - \lambda^* \bar{y} = 0 \\ \tilde{G}^\top A \bar{H} \bar{y} - \lambda^* \tilde{G}^\top \bar{G} \bar{x} \geq 0 \\ \tilde{H}^\top A^\top \bar{G} \bar{x} - \lambda^* \tilde{H}^\top \bar{H} \bar{y} \geq 0 \\ \|\bar{G} \bar{x}\| = \|\bar{H} \bar{y}\| = 1 \end{cases}$$

where $\begin{cases} \bar{x} := x_{\mathcal{I}}^*, \bar{y} := y_{\mathcal{J}}^* \\ \bar{G} := G_{:, \mathcal{I}}, \bar{H} := H_{:, \mathcal{J}} \\ \tilde{G} := G_{:, \mathcal{I}^c}, \tilde{H} := H_{:, \mathcal{J}^c} \\ \lambda^* = \bar{x}^\top \bar{G}^\top A \bar{H} \bar{y} = (u^*)^\top A v^* \end{cases}$

Theorem (B., G., S. 2024)

For the optimal solution $(u^*, v^*) = (Gx^*, Hy^*) = (\bar{G}\bar{x}, \bar{H}\bar{y})$, and $\lambda^* = (u^*)^\top A v^*$ we have that

$$M^* := \begin{pmatrix} 0 & \bar{H}^\dagger A^\top \bar{G} \\ \bar{G}^\dagger A \bar{H} & 0 \end{pmatrix} \implies M^* \begin{pmatrix} \bar{y} \\ \bar{x} \end{pmatrix} = \lambda^* \begin{pmatrix} \bar{y} \\ \bar{x} \end{pmatrix}$$

and λ^* is the least eigenvalue of M^* (from 2° order KKT)

Viceversa, if M is generated by random sets \mathcal{I}, \mathcal{J} , λ is its least eigenvalue and it admits a nonnegative eigenvector, then $\lambda \geq \lambda^*$.

The Active Set algorithm cycles over all subsets of indices \mathcal{I}, \mathcal{J} and tests if the least eigenvalue of M has a nonnegative eigenvector, giving us upper bounds on λ^* , and one of them will be the exact solution

Active Set

Input: $A \in \mathbb{R}^{m \times n}$, $G \in \mathbb{R}^{m \times p}$, $H \in \mathbb{R}^{n \times q}$, $P = \langle G \rangle$, $Q = \langle H \rangle$

Output: $\lambda = \min u^\top A v$ such that $\|u\| = \|v\| = 1$, $u \in P$, $v \in Q$

- 1: $\lambda = g_i^\top A h_j = \min_{k,\ell} (G^\top A H)_{k,\ell}$, $u = g_i$, $v = h_j$, $r = \text{Null}(A^\top A - \|A\|^2 I_n)$
- 2: $\mathcal{I} := \{(\mathcal{I}, \mathcal{J}) : 2 < |\mathcal{I}| + |\mathcal{J}| \leq m+n-r, \bar{G} := G_{:, \mathcal{I}}$ and $\bar{H} := H_{:, \mathcal{J}}$ full column rank}
- 3: **for** $(\mathcal{I}, \mathcal{J}) \in \mathcal{I}$, **do**
- 4: $A_x = \bar{G}^\dagger A^\top \bar{H}$, $A_y = \bar{H}^\dagger A \bar{G}$
- 5: $A_\lambda = A_y A_x$, $\tilde{A}_\lambda = A_x$ (or $A_\lambda = A_x A_y$, $\tilde{A}_\lambda = A_y$ if $|\mathcal{I}| > |\mathcal{J}|$)
- 6: **if** $\rho(A_\lambda) \leq \lambda^2$ **then** Skip to the next $(\mathcal{I}, \mathcal{J}) \in \mathcal{I}$
- 7: U right eigenspace of $\rho(A_\lambda)$ in A_λ , $\mu = -\sqrt{\rho(A_\lambda)}$, $W = \begin{pmatrix} \tilde{A}_\lambda U / \mu \\ U \end{pmatrix}$
- 8: Compute the reduced QR of $W = VR$
- 9: **if** $(VV^\top - I)z = 0$, $z \geq 0$, $e^\top z = 1$ admits a solution **then**
- 10: $\lambda = \mu$, $z = [y^\top \ x^\top]^\top$ (or $z = [x^\top \ y^\top]^\top$ if $|\mathcal{I}| > |\mathcal{J}|$)
- 11: $u = \bar{G}x / \|\bar{G}x\|$, $v = \bar{H}y / \|\bar{H}y\|$
- 12: **end if**
- 13: **end for**

(Not) a Gurobi AD



(Not) a Gurobi AD



minimize $x^T Qx + c^T x + \text{alpha}$

subject to $Ax = b$ (linear constraints)

$\ell \leq x \leq u$ (bound constraints)

some x_j integral (integrality constraints)

$x^T Qc x + q^T x \leq \text{beta}$ (quadratic constraints)

some x_i in SOS (special ordered set constraints)

min, max, abs, or, ... (general constraints)

- ✓ Interfaces: C, C++, Python, Java, Matlab, .NET, R
- ✗ Proprietary, not Open Source
- ✓ Free for Academic use
- Slower than approximating iterative solvers, but has decent Heuristics
- ✓ Solve the problem exactly even in the indefinite case

Uses McCormick Envelope Relaxation:

$$\min_{(u,v) \in K} \langle u, Av \rangle = \min_{(u,v) \in K} \sum_{i,j} A_{i,j} u_i v_j = \min_{(u,v) \in K, u_i v_j = w_{i,j}} \sum_{i,j} A_{i,j} w_{i,j} \geq \min_{(u_i, v_j, w_{i,j}) \in K_{i,j}} \sum_{i,j} A_{i,j} w_{i,j}$$

where $K_{i,j} = \text{Conv}((u_i, v_j, w_{i,j}) : w_{i,j} = u_i v_j, \underline{u}_i \leq u_i \leq \bar{u}_i, \underline{v}_j \leq v_j \leq \bar{v}_j, u, v \in K)$

Behaves great when A is Sparse since it has less slack variables $w_{i,j}$

(Not) a Gurobi AD



minimize $x^T Qx + c^T x + \text{alpha}$

subject to $Ax = b$ (linear constraints)

$\ell \leq x \leq u$ (bound constraints)

some x_j integral (integrality constraints)

$x^T Q_c x + q^T x \leq \text{beta}$ (quadratic constraints)

some x_i in SOS (special ordered set constraints)

min, max, abs, or, ... (general constraints)

- ✓ Interfaces: C, C++, Python, Java, Matlab, .NET, R
- ✗ Proprietary, not Open Source
- ✓ Free for Academic use
- Slower than approximating iterative solvers, but has decent Heuristics
- ✓ Solve the problem exactly even in the indefinite case

Uses McCormick Envelope Relaxation:

$$\min_{(u,v) \in K} \langle u, Av \rangle = \min_{(u,v) \in K} \sum_{i,j} A_{i,j} u_i v_j = \min_{(u,v) \in K, u_i v_j = w_{i,j}} \sum_{i,j} A_{i,j} w_{i,j} \geq \min_{(u_i, v_j, w_{i,j}) \in K_{i,j}} \sum_{i,j} A_{i,j} w_{i,j}$$

where $K_{i,j} = \text{Conv}((u_i, v_j, w_{i,j}) : w_{i,j} = u_i v_j, \underline{u}_i \leq u_i \leq \bar{u}_i, \underline{v}_j \leq v_j \leq \bar{v}_j, u, v \in K)$

Behaves great when A is **Sparse** since it has less slack variables $w_{i,j}$

(Not) a Gurobi AD



minimize $x^T Qx + c^T x + \text{alpha}$

subject to $Ax = b$ (linear constraints)

$\ell \leq x \leq u$ (bound constraints)

some x_j integral (integrality constraints)

$x^T Q_c x + q^T x \leq \text{beta}$ (quadratic constraints)

some x_i in SOS (special ordered set constraints)

min, max, abs, or, ... (general constraints)

- ✓ Interfaces: C, C++, Python, Java, Matlab, .NET, R
- ✗ Proprietary, not Open Source
- ✓ Free for Academic use
- Slower than approximating iterative solvers, but has decent Heuristics
- ✓ Solve the problem exactly even in the indefinite case

Uses McCormick Envelope Relaxation:

$$\min_{(u,v) \in K} \langle u, Av \rangle = \min_{(u,v) \in K} \sum_{i,j} A_{i,j} u_i v_j = \min_{(u,v) \in K, u_i v_j = w_{i,j}} \sum_{i,j} A_{i,j} w_{i,j} \geq \min_{(u_i, v_j, w_{i,j}) \in K_{i,j}} \sum_{i,j} A_{i,j} w_{i,j}$$

where $K_{i,j} = \text{Conv}((u_i, v_j, w_{i,j}) : w_{i,j} = u_i v_j, \underline{u}_i \leq u_i \leq \bar{u}_i, \underline{v}_j \leq v_j \leq \bar{v}_j, u, v \in K)$

Behaves great when A is **Sparse** since it has less slack variables $w_{i,j}$

(Not) a Gurobi AD



minimize $x^T Qx + c^T x + \text{alpha}$

subject to $Ax = b$ (linear constraints)

$\ell \leq x \leq u$ (bound constraints)

some x_j integral (integrality constraints)

$x^T Qc x + q^T x \leq \text{beta}$ (quadratic constraints)

some x_i in SOS (special ordered set constraints)

min, max, abs, or, ... (general constraints)

- ✓ Interfaces: C, C++, Python, Java, Matlab, .NET, R
- ✗ Proprietary, not Open Source
- ✓ Free for Academic use
- Slower than approximating iterative solvers, but has decent Heuristics
- ✓ Solve the problem exactly even in the indefinite case

Uses McCormick Envelope Relaxation:

$$\min_{(u,v) \in K} \langle u, Av \rangle = \min_{(u,v) \in K} \sum_{i,j} A_{i,j} u_i v_j = \min_{(u,v) \in K, u_i v_j = w_{i,j}} \sum_{i,j} A_{i,j} w_{i,j} \geq \min_{(u_i, v_j, w_{i,j}) \in K_{i,j}} \sum_{i,j} A_{i,j} w_{i,j}$$

where $K_{i,j} = \text{Conv}((u_i, v_j, w_{i,j}) : w_{i,j} = u_i v_j, \underline{u}_i \leq u_i \leq \bar{u}_i, \underline{v}_j \leq v_j \leq \bar{v}_j, u, v \in K)$

Behaves great when A is **Sparse** since it has less slack variables $w_{i,j}$

Alternating projection with extrapolation

$$\lambda^* = \min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1}} u^\top A v$$

Idea: We have seen that if we know u^* or v^* , then finding the other is equivalent to solve an easy convex problem

Alternate Projection: starting from an initial feasible point (u_0, v_0) and $k = 0$

- $u_{k+1} = \arg \min_{x \in P} x^\top A v_k$ such that $\|x\|_2 = 1$
- $v_{k+1} = \arg \min_{y \in Q} u_{k+1}^\top A y$ such that $\|y\|_2 = 1$
- $k = k + 1$

To accelerate the convergence, we add an **Extrapolation step** after each update

- $u_{k+1} = u_{k+1} + \beta(u_{k+1} - u_k)$
- $v_{k+1} = v_{k+1} + \beta(v_{k+1} - v_k)$
- If the objective increases then we decrease β and go back to (u_k, v_k) , otherwise we increase β

The method converges to a stationary point, that may not be optimal

Alternating projection with extrapolation

$$\lambda^* = \min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1}} u^\top A v$$

Idea: We have seen that if we know u^* or v^* , then finding the other is equivalent to solve an easy convex problem

Alternate Projection: starting from an initial feasible point (u_0, v_0) and $k = 0$

- $u_{k+1} = \arg \min_{x \in P} x^\top A v_k$ such that $\|x\|_2 = 1$
- $v_{k+1} = \arg \min_{y \in Q} u_{k+1}^\top A y$ such that $\|y\|_2 = 1$
- $k = k + 1$

To accelerate the convergence, we add an **Extrapolation step** after each update

- $u_{k+1} = u_{k+1} + \beta(u_{k+1} - u_k)$
- $v_{k+1} = v_{k+1} + \beta(v_{k+1} - v_k)$
- If the objective increases then we decrease β and go back to (u_k, v_k) , otherwise we increase β

The method converges to a stationary point, that may not be optimal

Alternating projection with extrapolation

$$\lambda^* = \min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1}} u^\top A v$$

Idea: We have seen that if we know u^* or v^* , then finding the other is equivalent to solve an easy convex problem

Alternate Projection: starting from an initial feasible point (u_0, v_0) and $k = 0$

- $u_{k+1} = \arg \min_{x \in P} x^\top A v_k$ such that $\|x\|_2 = 1$
- $v_{k+1} = \arg \min_{y \in Q} u_{k+1}^\top A y$ such that $\|y\|_2 = 1$
- $k = k + 1$

To accelerate the convergence, we add an **Extrapolation step** after each update

- $u_{k+1} = u_{k+1} + \beta(u_{k+1} - u_k)$
- $v_{k+1} = v_{k+1} + \beta(v_{k+1} - v_k)$
- If the objective increases then we decrease β and go back to (u_k, v_k) , otherwise we increase β

The method converges to a stationary point, that may not be optimal

Alternating projection with extrapolation

$$\lambda^* = \min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1,}} u^\top A v$$

Idea: We have seen that if we know u^* or v^* , then finding the other is equivalent to solve an easy convex problem

Alternate Projection: starting from an initial feasible point (u_0, v_0) and $k = 0$

- $u_{k+1} = \arg \min_{x \in P} x^\top A v_k$ such that $\|x\|_2 = 1$
- $v_{k+1} = \arg \min_{y \in Q} u_{k+1}^\top A y$ such that $\|y\|_2 = 1$
- $k = k + 1$

To accelerate the convergence, we add an **Extrapolation step** after each update

- $u_{k+1} = u_{k+1} + \beta(u_{k+1} - u_k)$
- $v_{k+1} = v_{k+1} + \beta(v_{k+1} - v_k)$
- If the objective increases then we decrease β and go back to (u_k, v_k) , otherwise we increase β

The method converges to a stationary point, that may not be optimal

Alternating projection with extrapolation

$$\lambda^* = \min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1,}} u^\top A v$$

Idea: We have seen that if we know u^* or v^* , then finding the other is equivalent to solve an easy convex problem

Alternate Projection: starting from an initial feasible point (u_0, v_0) and $k = 0$

- $u_{k+1} = \arg \min_{x \in P} x^\top A v_k$ such that $\|x\|_2 = 1$
- $v_{k+1} = \arg \min_{y \in Q} u_{k+1}^\top A y$ such that $\|y\|_2 = 1$
- $k = k + 1$

To accelerate the convergence, we add an **Extrapolation step** after each update

- $u_{k+1} = u_{k+1} + \beta(u_{k+1} - u_k)$
- $v_{k+1} = v_{k+1} + \beta(v_{k+1} - v_k)$
- If the objective increases then we decrease β and go back to (u_k, v_k) , otherwise we increase β

The method **converges** to a stationary point, that **may not be optimal**

Alternating projection with extrapolation

Input: $A \in \mathbb{R}^{m \times n}$, cones $P \subseteq \mathbb{R}^m$ and $Q \subseteq \mathbb{R}^n$

Output: An approximate solution to $\min_{u \in P, v \in Q} u^\top A v$ such that $\|u\|_2 = \|v\|_2 = 1$.

```
1:  $u = 0, v = 0, v_e = v_0, k = 1.$ 
2: while  $k \leq K$  and ( $\|u - u_p\|_2 \geq \delta$  or  $\|v - v_p\|_2 \geq \delta$ ) do
3:    $u_p = u$ . % Keep previous iterate in memory
4:    $u = \arg \min_{x \in P} x^\top A v_e$  such that  $\|x\|_2 = 1.$ 
5:    $u_e = u + \beta(u - u_p)$ . % Extrapolated point
6:    $v_p = v$ . % Keep previous iterate in memory
7:    $v = \arg \min_{y \in Q} u_e^\top A y$  such that  $\|y\|_2 = 1.$ 
8:    $v_e = v + \beta(v - v_p)$ . % Extrapolated point
9:    $e_k \leftarrow u^\top A v.$ 
10:  if  $k \geq 2$  and  $e_k > e_{k-1}$  then
11:     $u = u_p, v = v_p, \beta = \frac{\beta}{\eta}.$ 
12:  else
13:     $\beta \leftarrow \min(1, \gamma\beta).$ 
14:  end if
15:   $k \leftarrow k + 1.$ 
16: end while
```

Sequential Regularized Partial Linearization

$$\lambda^* = \min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1}} u^\top A v = \min_{\substack{u \in P, u \neq 0, \\ v \in Q, v \neq 0}} \frac{u^\top A v}{\|u\| \|v\|} = \min_{\substack{e^\top x = 1, x \geq 0, \\ e^\top y = 1, y \geq 0}} \frac{x^\top G^\top A H y}{\|Gx\| \|Hy\|}$$

Idea: If the minimum of $f_\delta(x, y) := x^\top G^\top A H y - \delta \|Gx\| \|Hy\|$ over $(x, y) \in \Delta_p \times \Delta_q$ is $\mu < 0$ then we get a decrease in the objective function

$$\frac{x^\top G^\top A H y}{\|Gx\| \|Hy\|} = \delta + \frac{\mu}{\|Gx\| \|Hy\|} < \delta$$

Partial Linearization: starting from an initial feasible point (x_0, y_0) and $k = 0$,

- $\delta = \frac{x_k^\top G^\top A H y_k}{\|Gx_k\| \|Hy_k\|}$
- Linearize wrt x the function $f_\delta(x, y_k)$, penalize it with $\|x - x_k\|^2$ and minimize it to obtain x_{k+1}
- Linearize wrt y the function $f_\delta(x_{k+1}, y)$, penalize it with $\|y - y_k\|^2$ and minimize it to obtain y_{k+1}
- $k = k + 1$

To accelerate the convergence, we add an **Extrapolation step**

Sequential Regularized Partial Linearization

$$\lambda^* = \min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1}} u^\top A v = \min_{\substack{u \in P, u \neq 0, \\ v \in Q, v \neq 0}} \frac{u^\top A v}{\|u\| \|v\|} = \min_{\substack{e^\top x = 1, x \geq 0, \\ e^\top y = 1, y \geq 0}} \frac{x^\top G^\top A H y}{\|Gx\| \|Hy\|}$$

Idea: If the minimum of $f_\delta(x, y) := x^\top G^\top A H y - \delta \|Gx\| \|Hy\|$ over $(x, y) \in \Delta_p \times \Delta_q$ is $\mu < 0$ then we get a decrease in the objective function

$$\frac{x^\top G^\top A H y}{\|Gx\| \|Hy\|} = \delta + \frac{\mu}{\|Gx\| \|Hy\|} < \delta$$

Partial Linearization: starting from an initial feasible point (x_0, y_0) and $k = 0$,

- $\delta = \frac{x_k^\top G^\top A H y_k}{\|Gx_k\| \|Hy_k\|}$
- Linearize wrt x the function $f_\delta(x, y_k)$, penalize it with $\|x - x_k\|^2$ and minimize it to obtain x_{k+1}
- Linearize wrt y the function $f_\delta(x_{k+1}, y)$, penalize it with $\|y - y_k\|^2$ and minimize it to obtain y_{k+1}
- $k = k + 1$

To accelerate the convergence, we add an **Extrapolation step**

Sequential Regularized Partial Linearization

$$\lambda^* = \min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1}} u^\top A v = \min_{\substack{u \in P, u \neq 0, \\ v \in Q, v \neq 0}} \frac{u^\top A v}{\|u\| \|v\|} = \min_{\substack{e^\top x = 1, x \geq 0, \\ e^\top y = 1, y \geq 0}} \frac{x^\top G^\top A H y}{\|Gx\| \|Hy\|}$$

Idea: If the minimum of $f_\delta(x, y) := x^\top G^\top A H y - \delta \|Gx\| \|Hy\|$ over $(x, y) \in \Delta_p \times \Delta_q$ is $\mu < 0$ then we get a decrease in the objective function

$$\frac{x^\top G^\top A H y}{\|Gx\| \|Hy\|} = \delta + \frac{\mu}{\|Gx\| \|Hy\|} < \delta$$

Partial Linearization: starting from an initial feasible point (x_0, y_0) and $k = 0$,

- $\delta = \frac{x_k^\top G^\top A H y_k}{\|Gx_k\| \|Hy_k\|}$
- Linearize wrt x the function $f_\delta(x, y_k)$, penalize it with $\|x - x_k\|^2$ and minimize it to obtain x_{k+1}
- Linearize wrt y the function $f_\delta(x_{k+1}, y)$, penalize it with $\|y - y_k\|^2$ and minimize it to obtain y_{k+1}
- $k = k + 1$

To accelerate the convergence, we add an **Extrapolation step**

Sequential Regularized Partial Linearization

$$\lambda^* = \min_{\substack{u \in P, \|u\| = 1, \\ v \in Q, \|v\| = 1}} u^\top A v = \min_{\substack{u \in P, u \neq 0, \\ v \in Q, v \neq 0}} \frac{u^\top A v}{\|u\| \|v\|} = \min_{\substack{e^\top x = 1, x \geq 0, \\ e^\top y = 1, y \geq 0}} \frac{x^\top G^\top A H y}{\|Gx\| \|Hy\|}$$

Idea: If the minimum of $f_\delta(x, y) := x^\top G^\top A H y - \delta \|Gx\| \|Hy\|$ over $(x, y) \in \Delta_p \times \Delta_q$ is $\mu < 0$ then we get a decrease in the objective function

$$\frac{x^\top G^\top A H y}{\|Gx\| \|Hy\|} = \delta + \frac{\mu}{\|Gx\| \|Hy\|} < \delta$$

Partial Linearization: starting from an initial feasible point (x_0, y_0) and $k = 0$,

- $\delta = \frac{x_k^\top G^\top A H y_k}{\|Gx_k\| \|Hy_k\|}$
- Linearize wrt x the function $f_\delta(x, y_k)$, penalize it with $\|x - x_k\|^2$ and minimize it to obtain x_{k+1}
- Linearize wrt y the function $f_\delta(x_{k+1}, y)$, penalize it with $\|y - y_k\|^2$ and minimize it to obtain y_{k+1}
- $k = k + 1$

To accelerate the convergence, we add an **Extrapolation step**

Input: $A \in \mathbb{R}^{m \times n}$, cones $P \subseteq \mathbb{R}^m$ and $Q \subseteq \mathbb{R}^n$

Output: An approximate solution to $\min_{u \in P, v \in Q} \langle u, Av \rangle$ such that $\|u\| = \|v\| = 1$

1: Set

$$\delta_k := \frac{\langle Gx^k, AHy^k \rangle}{\|Gx^k\| \|Hy^k\|}$$

2: Let $L_1^k(x) := \langle Gx, AHy^k - \delta_k \|Gx^k\|^{-1} \|Hy^k\| Gx^k \rangle$

Compute a solution \tilde{x}^k to the convex program

$$\min L_1^k(x) + \frac{\mu_1}{2} \|x - x^k\|^2 \quad \text{such that } x \in \Delta_p$$

3: Let $L_2^k(y) := \langle Hy, A^\top Gx^k - \delta_k \|Gx^k\| \|Hy^k\|^{-1} Hy^k \rangle$

Compute a solution \tilde{y}^k to the convex program

$$\min L_2^k(y) + \frac{\mu_2}{2} \|y - y^k\|^2 \quad \text{such that } y \in \Delta_q$$

4: Let $d_1^k := \tilde{x}^k - x^k$ and $d_2^k := \tilde{y}^k - y^k$

5: If $(|L_1^k(d_1^k)| < \delta \text{ and } |L_2^k(d_2^k)| < \delta)$ or $k \geq K$ terminate

Otherwise, let $t_k := \beta \rho^{\ell_k}$, where ℓ_k is the smallest nonnegative integer ℓ such that

$$\Phi(x^k + t^k d_1^k, y^k + t^k d_2^k) \leq \Phi(x^k, y^k) + \alpha t_k \frac{L_1^k(d_1^k) + L_2^k(d_2^k)}{\|Gx^k\| \|Hy^k\|}$$

Set $(x^{k+1}, y^{k+1}) := (x^k, y^k) + t_k(d_1^k, d_2^k)$ and $k = k + 1$. Go to step 1

Experiments

An Example: Schur Cone

The **Schur Cone** is generated by the matrix

$$H = \begin{pmatrix} 1 & 0 & \dots & 0 \\ -1 & 1 & \dots & 0 \\ 0 & -1 & \dots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \dots & 1 \\ 0 & 0 & \dots & -1 \end{pmatrix} \in \mathbb{R}^{n \times n-1} \quad \langle H \rangle \subseteq e^\perp$$

One can prove that the maximum angle between the Schur cone Q and the positive orthant P is achieved by

$$y = e_n \in P \quad x = (aa \dots ab) \in Q \quad a = \sqrt{\frac{1}{n(n-1)}} \quad b = -\sqrt{1 - \frac{1}{n}} = x^\top y$$

that can be proved being the maximum angle as

$$\min_{\substack{x \in Q, y \in P \\ \|x\| = \|y\| = 1}} x^\top y \geq \min_{\substack{x \in e^\perp, y \geq 0 \\ \|x\| = \|y\| = 1}} x^\top y = \min_{\substack{y \geq 0, \|y\|=1}} -\|P_{e^\perp}(y)\| = -\sqrt{1 - \frac{1}{n}}$$

An Example: Schur Cone

The **Schur Cone** is generated by the matrix

$$H = \begin{pmatrix} 1 & 0 & \dots & 0 \\ -1 & 1 & \dots & 0 \\ 0 & -1 & \dots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \dots & 1 \\ 0 & 0 & \dots & -1 \end{pmatrix} \in \mathbb{R}^{n \times n-1} \quad \langle H \rangle \subseteq e^\perp$$

One can prove that the maximum angle between the Schur cone Q and the positive orthant P is achieved by

$$y = e_n \in P \quad x = (a \ a \ \dots \ a \ b) \in Q \quad a = \sqrt{\frac{1}{n(n-1)}} \quad b = -\sqrt{1 - \frac{1}{n}} = x^\top y$$

that can be proved being the maximum angle as

$$\min_{\substack{x \in Q, y \in P \\ \|x\| = \|y\| = 1}} x^\top y \geq \min_{\substack{x \in e^\perp, y \geq 0 \\ \|x\| = \|y\| = 1}} x^\top y = \min_{\substack{y \geq 0, \|y\|=1}} -\|P_{e^\perp}(y)\| = -\sqrt{1 - \frac{1}{n}}$$

Schur Cone and Positive Orthant

Table 1: Numerical comparison for Gur and BFAS for different dimensions for the problem of finding the maximum angle between the Schur cone and the positive orthant cone. The table reports the optimal objective functions values found in the timelimit (60 seconds) and the actual elapsed time. We also report the exact value for each problem.

n	5	10	20	50
exact	0.852416π	0.897584π	0.928217π	0.954833π
Gur	0.852416π	0.897584π	0.928218π	0.954833π
	0.1134 s	0.2016 s	20.1493 s	60^* s
BFAS	0.852416π	0.897584π	0.750000π	0.750000π
	0.3310 s	48.3153 s	60^* s	60^* s

n	100	200	500
exact	0.968116π	0.977473π	0.985760π
Gur	0.968116π	0.977473π	0.985756π
	60^* s	60^* s	60^* s
BFAS	0.750000π	0.750000π	0.750000π
	60^* s	60^* s	60^* s

Schur Cone and Positive Orthant


Table 1: Numerical comparison for Gurobi and BFAS for different dimensions for the problem of finding the maximum angle between the Schur cone and itself. The table reports the optimal objective functions values found in the timelimit (60 seconds) and the actual elapsed time. We also report the exact value for each problem.

n	5	10	20	50
exact	0.800000π	0.900000π	0.950000π	0.980000π
Gur	0.800001π 0.2508 s	0.900000π 60* s	0.950000π 60* s	0.980000π 60* s
BFAS	0.800000π 0.3856 s	0.900000π 60* s	0.859157π 60* s	0.804087π 60* s


n	100	200	500
exact	0.990000π	0.995000π	0.998000π
Gur	0.936315π 60* s	0.994996π 60* s	0.998011π 60* s
BFAS	0.750000π 60* s	0.750000π 60* s	0.750000π 60* s

Schur Cone and Positive Orthant

Schur - Nonnegative Orthant, $n = 200$



Schur - Schur, $n = 200$



Maximum Edge Biclique Problem

Recall that solving the Pareto singular value problem is equivalent to solve the maximum edge biclique problem.

Here we thus test all four algorithms on four bipartite graphs taken from a benchmark dataset¹. All graphs have been randomly generated with a fixed edge density, and then a biclique has been added to them. In particular,

- the first graph is a 100×100 graph with density 0.2 and planted biclique of size $50 \times 50 = 2500$,
- the second graph is a 300×300 graph with density 0.3 and planted biclique of size $2 \times 55 = 110$,
- the third graph is a 100×100 graph with density 0.71 and planted biclique of size $80 \times 80 = 6400$,
- the fourth graph is a 10000×300 graph with density 0.03 and planted biclique of size $22 \times 2 = 44$.

¹Shaham, E.: maximum biclique benchmark. <https://github.com/shahamer/maximum-biclique-benchmark> (2019)

Maximum Edge Biclique Problem

Table 1: Numerical comparison for Gurobi, BFAS, E-AO and SRPL for the problem of finding the maximum edge biclique in four different bipartite graphs. The table reports the maximum edge biclique found in the timelimit (10 seconds) for Gurobi and BFAS. The reported number for E-AO and SRPL are instead the average value found at 10 seconds for 100 runs, and in parentheses the best value found throughout all 100 runs when it differs from the average one. Gurobi cannot be executed on the last graph due to its excessive size.

n	100×100	300×300	100×100	10000×300
Gur	2500	0	310	NA
BFAS	3	2	2	2
E-AO	66	114	87	12
SRPL	2500	114	6400	46(358)

Maximum Angle between PSD and Nonnegative Symmetric Matrices

Given the inner product $\langle A, B \rangle = \text{Tr}(A^\top B)$ on the space of $n \times n$ real symmetric matrices \mathcal{S}^n an open question is the maximum angle between the cone of PSD matrices \mathcal{P}^n and the cone of nonnegative symmetric matrices \mathcal{N}^n for $n \geq 5$

$$\gamma_n := \min_{\substack{A \in \mathcal{P}^n, B \in \mathcal{N}^n \\ \|A\|_F = \|B\|_F = 1}} \langle A, B \rangle = - \max_{A \in \mathcal{P}^n, \|A\|_F = 1} \|A^\top\|_F = -\frac{1}{2} \max_{B \in \mathcal{N}^n, \|B\|_F = 1} \|B - \sqrt{B^2}\|_F$$

It is known that

$$n = 2, 3, 4 \implies \gamma_n = -\frac{1}{\sqrt{2}} = \cos\left(\frac{3}{4}\pi\right) \quad \lim_{n \rightarrow \infty} \gamma_n \downarrow -1 = \cos(\pi)$$

This is a lower bound on the maximum angle in the cone of copositive matrices

$$\mathcal{C}^n := \{A \in \mathcal{S}^n : x^\top A x \geq 0 \ \forall x \geq 0\}$$

All the algorithms to compute γ_n are iteratively converging to a critical angle, i.e. a stationary point of the optimization problem

For $n \geq 5$ we only have lower bounds on the exact angle

Maximum Angle between PSD and Nonnegative Symmetric Matrices

Given the inner product $\langle A, B \rangle = \text{Tr}(A^\top B)$ on the space of $n \times n$ real symmetric matrices \mathcal{S}^n an open question is the maximum angle between the cone of PSD matrices \mathcal{P}^n and the cone of nonnegative symmetric matrices \mathcal{N}^n for $n \geq 5$

$$\gamma_n := \min_{\substack{A \in \mathcal{P}^n, B \in \mathcal{N}^n \\ \|A\|_F = \|B\|_F = 1}} \langle A, B \rangle = - \max_{A \in \mathcal{P}^n, \|A\|_F=1} \|A^\top\|_F = -\frac{1}{2} \max_{B \in \mathcal{N}^n, \|B\|_F=1} \|B - \sqrt{B^2}\|_F$$

It is known that

$$n = 2, 3, 4 \implies \gamma_n = -\frac{1}{\sqrt{2}} = \cos\left(\frac{3}{4}\pi\right) \quad \lim_{n \rightarrow \infty} \gamma_n \downarrow -1 = \cos(\pi)$$

This is a lower bound on the maximum angle in the cone of copositive matrices

$$\mathcal{C}^n := \{A \in \mathcal{S}^n : x^\top A x \geq 0 \ \forall x \geq 0\}$$

All the algorithms to compute γ_n are iteratively converging to a critical angle, i.e. a stationary point of the optimization problem

For $n \geq 5$ we only have lower bounds on the exact angle

Maximum Angle between PSD and Nonnegative Symmetric Matrices

Given the inner product $\langle A, B \rangle = \text{Tr}(A^\top B)$ on the space of $n \times n$ real symmetric matrices \mathcal{S}^n an open question is the maximum angle between the cone of PSD matrices \mathcal{P}^n and the cone of nonnegative symmetric matrices \mathcal{N}^n for $n \geq 5$

$$\gamma_n := \min_{\substack{A \in \mathcal{P}^n, B \in \mathcal{N}^n \\ \|A\|_F = \|B\|_F = 1}} \langle A, B \rangle = - \max_{A \in \mathcal{P}^n, \|A\|_F=1} \|A^\top\|_F = -\frac{1}{2} \max_{B \in \mathcal{N}^n, \|B\|_F=1} \|B - \sqrt{B^2}\|_F$$

It is known that

$$n = 2, 3, 4 \implies \gamma_n = -\frac{1}{\sqrt{2}} = \cos\left(\frac{3}{4}\pi\right) \quad \lim_{n \rightarrow \infty} \gamma_n \downarrow -1 = \cos(\pi)$$

This is a lower bound on the maximum angle in the cone of copositive matrices

$$\mathcal{C}^n := \{A \in \mathcal{S}^n : x^\top A x \geq 0 \ \forall x \geq 0\}$$

All the algorithms to compute γ_n are iteratively converging to a critical angle, i.e. a stationary point of the optimization problem

For $n \geq 5$ we only have lower bounds on the exact angle

A Conjecture

Known Antipodal Couples:

$$n=1: A_1 = B_1 = 1$$

$$n=2: A_2 = \frac{1}{2} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \quad B_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

$$n=3, 4: A_n = \begin{pmatrix} A_2 & 0 \\ 0 & 0 \end{pmatrix} \quad B_n = \begin{pmatrix} B_2 & 0 \\ 0 & 0 \end{pmatrix}$$

$n=5$: (Best Known Stationary Point) F_5 is the Fourier matrix

$$A_n = F_5 \begin{pmatrix} 0 & & & & \\ & 0 & & & \\ & & \frac{1}{\sqrt{2}} & & \\ & & & \frac{1}{\sqrt{2}} & \\ & & & & 0 \end{pmatrix} F_5^H \quad B_n = \frac{1}{\sqrt{10}} \begin{pmatrix} 0 & 1 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 & 0 \end{pmatrix}$$

Conjecture

Every antipodal pair is block circulant

A Conjecture

Known Antipodal Couples:

$$n = 1: A_1 = B_1 = 1$$

$$n = 2: A_2 = \frac{1}{2} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \quad B_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

$$n = 3, 4: A_n = \begin{pmatrix} A_2 & 0 \\ 0 & 0 \end{pmatrix} \quad B_n = \begin{pmatrix} B_2 & 0 \\ 0 & 0 \end{pmatrix}$$

$n = 5$: (Best Known Stationary Point) F_5 is the Fourier matrix

$$A_n = F_5 \begin{pmatrix} 0 & & & & \\ & 0 & & & \\ & & \frac{1}{\sqrt{2}} & & \\ & & & \frac{1}{\sqrt{2}} & \\ & & & & 0 \end{pmatrix} F_5^H \quad B_n = \frac{1}{\sqrt{10}} \begin{pmatrix} 0 & 1 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 & 0 \end{pmatrix}$$

Conjecture

Every antipodal pair is block circulant

A Conjecture

Known Antipodal Couples:

$$n = 1: A_1 = B_1 = 1$$

$$n = 2: A_2 = \frac{1}{2} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \quad B_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

$$n = 3, 4: A_n = \begin{pmatrix} A_2 & 0 \\ 0 & 0 \end{pmatrix} \quad B_n = \begin{pmatrix} B_2 & 0 \\ 0 & 0 \end{pmatrix}$$

n = 5: (Best Known Stationary Point) F_5 is the Fourier matrix

$$A_n = F_5 \begin{pmatrix} 0 & & & & \\ & 0 & & & \\ & & \frac{1}{\sqrt{2}} & & \\ & & & \frac{1}{\sqrt{2}} & \\ & & & & 0 \end{pmatrix} F_5^H \quad B_n = \frac{1}{\sqrt{10}} \begin{pmatrix} 0 & 1 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 & 0 \end{pmatrix}$$

Conjecture

Every antipodal pair is block circulant

A Conjecture

Known Antipodal Couples:

$$n = 1: A_1 = B_1 = 1$$

$$n = 2: A_2 = \frac{1}{2} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \quad B_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

$$n = 3, 4: A_n = \begin{pmatrix} A_2 & 0 \\ 0 & 0 \end{pmatrix} \quad B_n = \begin{pmatrix} B_2 & 0 \\ 0 & 0 \end{pmatrix}$$

n = 5: (Best Known Stationary Point) F_5 is the Fourier matrix

$$A_n = F_5 \begin{pmatrix} 0 & & & & \\ & 0 & & & \\ & & \frac{1}{\sqrt{2}} & & \\ & & & \frac{1}{\sqrt{2}} & \\ & & & & 0 \end{pmatrix} F_5^H \quad B_n = \frac{1}{\sqrt{10}} \begin{pmatrix} 0 & 1 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 & 0 \end{pmatrix}$$

Conjecture

Every antipodal pair is block circulant

A Conjecture

Known Antipodal Couples:

$$n = 1: A_1 = B_1 = 1$$

$$n = 2: A_2 = \frac{1}{2} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \quad B_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

$$n = 3, 4: A_n = \begin{pmatrix} A_2 & 0 \\ 0 & 0 \end{pmatrix} \quad B_n = \begin{pmatrix} B_2 & 0 \\ 0 & 0 \end{pmatrix}$$

$n = 5$: (Best Known Stationary Point) F_5 is the Fourier matrix

$$A_n = F_5 \begin{pmatrix} 0 & & & & \\ & 0 & & & \\ & & \frac{1}{\sqrt{2}} & & \\ & & & \frac{1}{\sqrt{2}} & \\ & & & & 0 \end{pmatrix} F_5^H \quad B_n = \frac{1}{\sqrt{10}} \begin{pmatrix} 0 & 1 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 & 0 \end{pmatrix}$$

Conjecture

Every antipodal pair is block circulant

A Conjecture

Known Antipodal Couples:

$$n = 1: A_1 = B_1 = 1$$

$$n = 2: A_2 = \frac{1}{2} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \quad B_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

$$n = 3, 4: A_n = \begin{pmatrix} A_2 & 0 \\ 0 & 0 \end{pmatrix} \quad B_n = \begin{pmatrix} B_2 & 0 \\ 0 & 0 \end{pmatrix}$$

$n = 5$: (Best Known Stationary Point) F_5 is the Fourier matrix

$$A_n = F_5 \begin{pmatrix} 0 & & & & \\ & 0 & & & \\ & & \frac{1}{\sqrt{2}} & & \\ & & & \frac{1}{\sqrt{2}} & \\ & & & & 0 \end{pmatrix} F_5^H \quad B_n = \frac{1}{\sqrt{10}} \begin{pmatrix} 0 & 1 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 & 0 \end{pmatrix}$$

Conjecture

Every antipodal pair is block circulant

Circulant Symmetric Matrices

The algebra of circulant real symmetric matrices \mathcal{SC}^n is the set of

$$\begin{pmatrix} a_0 & a_1 & a_2 & \dots & a_1 \\ a_1 & a_0 & a_1 & \ddots & \vdots \\ a_2 & a_1 & \ddots & \ddots & a_2 \\ \vdots & \ddots & \ddots & a_0 & a_1 \\ a_1 & \dots & a_2 & a_1 & a_0 \end{pmatrix} = a_0 I_n + F_n \operatorname{diag} \left(\sum_{j>0} 2a_j \cos(2\pi ij/n) \right)_{i=0:n-1} F_n^H$$

Properties:

- Both $\mathcal{SC}^n \cap \mathcal{P}^n$ and $\mathcal{SC}^n \cap \mathcal{N}^n$ are finitely generated cones with $\lceil \frac{n+1}{2} \rceil$ generators
- Given $C \in \mathcal{SC}^n \cap \mathcal{P}^n$ its projection C' onto $-\mathcal{N}^n$ is still in \mathcal{SC}^n , and the angle between $C, -C'$ is the maximum angle between C and \mathcal{N}^n
- Given $C \in \mathcal{SC}^n \cap \mathcal{N}^n$ its projection C' onto $-\mathcal{P}^n$ is still in \mathcal{SC}^n and the angle between $C, -C'$ is the maximum angle between C and \mathcal{P}^n
- An alternating algorithm using projections to minimize γ_n starting from a $C \in \mathcal{SC}^n$ will converge to a stationary point of the problem that is still in \mathcal{SC}^n

Circulant Symmetric Matrices

The algebra of circulant real symmetric matrices \mathcal{SC}^n is the set of

$$\begin{pmatrix} a_0 & a_1 & a_2 & \dots & a_1 \\ a_1 & a_0 & a_1 & \ddots & \vdots \\ a_2 & a_1 & \ddots & \ddots & a_2 \\ \vdots & \ddots & \ddots & a_0 & a_1 \\ a_1 & \dots & a_2 & a_1 & a_0 \end{pmatrix} = a_0 I_n + F_n \operatorname{diag} \left(\sum_{j>0} 2a_j \cos(2\pi ij/n) \right)_{i=0:n-1} F_n^H$$

Properties:

- Both $\mathcal{SC}^n \cap \mathcal{P}^n$ and $\mathcal{SC}^n \cap \mathcal{N}^n$ are finitely generated cones with $\lceil \frac{n+1}{2} \rceil$ generators
- Given $C \in \mathcal{SC}^n \cap \mathcal{P}^n$ its projection C' onto $-\mathcal{N}^n$ is still in \mathcal{SC}^n , and the angle between $C, -C'$ is the maximum angle between C and \mathcal{N}^n
- Given $C \in \mathcal{SC}^n \cap \mathcal{N}^n$ its projection C' onto $-\mathcal{P}^n$ is still in \mathcal{SC}^n and the angle between $C, -C'$ is the maximum angle between C and \mathcal{P}^n
- An alternating algorithm using projections to minimize γ_n starting from a $C \in \mathcal{SC}^n$ will converge to a stationary point of the problem that is still in \mathcal{SC}^n

Circulant Symmetric Matrices

The algebra of circulant real symmetric matrices \mathcal{SC}^n is the set of

$$\begin{pmatrix} a_0 & a_1 & a_2 & \dots & a_1 \\ a_1 & a_0 & a_1 & \ddots & \vdots \\ a_2 & a_1 & \ddots & \ddots & a_2 \\ \vdots & \ddots & \ddots & a_0 & a_1 \\ a_1 & \dots & a_2 & a_1 & a_0 \end{pmatrix} = a_0 I_n + F_n \operatorname{diag} \left(\sum_{j>0} 2a_j \cos(2\pi ij/n) \right)_{i=0:n-1} F_n^H$$

Properties:

- Both $\mathcal{SC}^n \cap \mathcal{P}^n$ and $\mathcal{SC}^n \cap \mathcal{N}^n$ are finitely generated cones with $\lceil \frac{n+1}{2} \rceil$ generators
- Given $C \in \mathcal{SC}^n \cap \mathcal{P}^n$ its projection C' onto $-\mathcal{N}^n$ is still in \mathcal{SC}^n , and the angle between $C, -C'$ is the maximum angle between C and \mathcal{N}^n
- Given $C \in \mathcal{SC}^n \cap \mathcal{N}^n$ its projection C' onto $-\mathcal{P}^n$ is still in \mathcal{SC}^n and the angle between $C, -C'$ is the maximum angle between C and \mathcal{P}^n
- An alternating algorithm using projections to minimize γ_n starting from a $C \in \mathcal{SC}^n$ will converge to a stationary point of the problem that is still in \mathcal{SC}^n

Circulant Symmetric Matrices

The algebra of circulant real symmetric matrices \mathcal{SC}^n is the set of

$$\begin{pmatrix} a_0 & a_1 & a_2 & \dots & a_1 \\ a_1 & a_0 & a_1 & \ddots & \vdots \\ a_2 & a_1 & \ddots & \ddots & a_2 \\ \vdots & \ddots & \ddots & a_0 & a_1 \\ a_1 & \dots & a_2 & a_1 & a_0 \end{pmatrix} = a_0 I_n + F_n \operatorname{diag} \left(\sum_{j>0} 2a_j \cos(2\pi ij/n) \right)_{i=0:n-1} F_n^H$$

Properties:

- Both $\mathcal{SC}^n \cap \mathcal{P}^n$ and $\mathcal{SC}^n \cap \mathcal{N}^n$ are finitely generated cones with $\lceil \frac{n+1}{2} \rceil$ generators
- Given $C \in \mathcal{SC}^n \cap \mathcal{P}^n$ its projection C' onto $-\mathcal{N}^n$ is still in \mathcal{SC}^n , and the angle between $C, -C'$ is the maximum angle between C and \mathcal{N}^n
- Given $C \in \mathcal{SC}^n \cap \mathcal{N}^n$ its projection C' onto $-\mathcal{P}^n$ is still in \mathcal{SC}^n and the angle between $C, -C'$ is the maximum angle between C and \mathcal{P}^n
- An alternating algorithm using projections to minimize γ_n starting from a $C \in \mathcal{SC}^n$ will converge to a stationary point of the problem that is still in \mathcal{SC}^n

Circulant Symmetric Matrices

The algebra of circulant real symmetric matrices \mathcal{SC}^n is the set of

$$\begin{pmatrix} a_0 & a_1 & a_2 & \dots & a_1 \\ a_1 & a_0 & a_1 & \ddots & \vdots \\ a_2 & a_1 & \ddots & \ddots & a_2 \\ \vdots & \ddots & \ddots & a_0 & a_1 \\ a_1 & \dots & a_2 & a_1 & a_0 \end{pmatrix} = a_0 I_n + F_n \operatorname{diag} \left(\sum_{j>0} 2a_j \cos(2\pi ij/n) \right)_{i=0:n-1} F_n^H$$

Properties:

- Both $\mathcal{SC}^n \cap \mathcal{P}^n$ and $\mathcal{SC}^n \cap \mathcal{N}^n$ are finitely generated cones with $\lceil \frac{n+1}{2} \rceil$ generators
- Given $C \in \mathcal{SC}^n \cap \mathcal{P}^n$ its projection C' onto $-\mathcal{N}^n$ is still in \mathcal{SC}^n , and the angle between $C, -C'$ is the maximum angle between C and \mathcal{N}^n
- Given $C \in \mathcal{SC}^n \cap \mathcal{N}^n$ its projection C' onto $-\mathcal{P}^n$ is still in \mathcal{SC}^n and the angle between $C, -C'$ is the maximum angle between C and \mathcal{P}^n
- An alternating algorithm using projections to minimize γ_n starting from a $C \in \mathcal{SC}^n$ will converge to a stationary point of the problem that is still in \mathcal{SC}^n

Circulant Symmetric Matrices

The algebra of circulant real symmetric matrices \mathcal{SC}^n is the set of

$$\begin{pmatrix} a_0 & a_1 & a_2 & \dots & a_1 \\ a_1 & a_0 & a_1 & \ddots & \vdots \\ a_2 & a_1 & \ddots & \ddots & a_2 \\ \vdots & \ddots & \ddots & a_0 & a_1 \\ a_1 & \dots & a_2 & a_1 & a_0 \end{pmatrix} = a_0 I_n + F_n \operatorname{diag} \left(\sum_{j>0} 2a_j \cos(2\pi ij/n) \right)_{i=0:n-1} F_n^H$$

Properties:

- Both $\mathcal{SC}^n \cap \mathcal{P}^n$ and $\mathcal{SC}^n \cap \mathcal{N}^n$ are finitely generated cones with $\lceil \frac{n+1}{2} \rceil$ generators
- Given $C \in \mathcal{SC}^n \cap \mathcal{P}^n$ its projection C' onto $-\mathcal{N}^n$ is still in \mathcal{SC}^n , and the angle between $C, -C'$ is the maximum angle between C and \mathcal{N}^n
- Given $C \in \mathcal{SC}^n \cap \mathcal{N}^n$ its projection C' onto $-\mathcal{P}^n$ is still in \mathcal{SC}^n and the angle between $C, -C'$ is the maximum angle between C and \mathcal{P}^n
- An alternating algorithm using projections to minimize γ_n starting from a $C \in \mathcal{SC}^n$ will converge to a stationary point of the problem that is still in \mathcal{SC}^n

Check with Gurobi

If n is odd and $n = 1 + 2m$

$$\min_{\substack{A \in \mathcal{SC}^n \cap \mathcal{P}^n, \|A\|_F = 1 \\ B \in \mathcal{SC}^n \cap \mathcal{N}^n, \|B\|_F = 1}} \langle A, B \rangle = \min_{\substack{x \geq 0, \|x\|_F = 1 \\ y \geq 0, \|y\|_F = 1}} \langle x, My \rangle \quad M = \frac{2}{\sqrt{n}} \left[\cos \left(\frac{2\pi}{n} ij \right) \right]_{i,j=1:m}$$

A similar reduction holds for n even

Check with Gurobi

If n is odd and $n = 1 + 2m$

$$\min_{\substack{A \in \mathcal{SC}^n \cap \mathcal{P}^n, \|A\|_F = 1 \\ B \in \mathcal{SC}^n \cap \mathcal{N}^n, \|B\|_F = 1}} \langle A, B \rangle = \min_{\substack{x \geq 0, \|x\|_F = 1 \\ y \geq 0, \|y\|_F = 1}} \langle x, My \rangle \quad M = \frac{2}{\sqrt{n}} \left[\cos \left(\frac{2\pi}{n} ij \right) \right]_{i,j=1:m}$$

A similar reduction holds for n even

5	0.7575π	0.7575π	18	0.7699π	$\textcolor{blue}{0.7670 \pi}$
6	0.7575π	$\textcolor{blue}{0.7575 \pi}$	19	0.7703π	0.7681π
7	0.7575π	$\textcolor{blue}{0.7575 \pi}$	20	0.7719π	0.7719π
8	0.7608π	0.7608π	21	0.7719π	$\textcolor{blue}{0.7719 \pi}$
9	0.7608π	$\textcolor{blue}{0.7608 \pi}$	22	0.7719π	$\textcolor{blue}{0.7719 \pi}$
10	0.7609π	$\textcolor{blue}{0.7608 \pi}$	23	0.7722π	$\textcolor{blue}{0.7719 \pi}$
11	0.7627π	0.7627π	24	0.7735π	$\textcolor{red}{0.7730 \pi}$
12	0.7649π	0.7649π	25	0.7735π	$\textcolor{blue}{0.7730 \pi}$
13	0.7649π	$\textcolor{blue}{0.7649 \pi}$	26	0.7735π	$\textcolor{blue}{0.7730 \pi}$
14	0.7659π	$\textcolor{blue}{0.7649 \pi}$	27	0.7739π	$\textcolor{blue}{0.7730 \pi}$
15	0.7678π	$\textcolor{blue}{0.7649 \pi}$	28	0.7750π	$\textcolor{blue}{0.7730 \pi}$
16	0.7699π	0.7670π	29	0.7750π	$\textcolor{blue}{0.7741 \pi}$
17	0.7699π	$\textcolor{blue}{0.7670 \pi}$	30	0.7757π	$\textcolor{red}{0.7741 \pi}$

Left: Lower bounds on γ_n

Right:

- In **black** the exact angle $\mathcal{SC}^n \cap \mathcal{P}^n \angle \mathcal{SC}^n \cap \mathcal{N}^n$
- In **blue** if a previous angle was bigger then the exact solution
- In **red** if it is a lower bound

Table 2: Numerical comparison of Gur and BFAS for different dimensions for the problem of finding the maximum angle between the PSD cone and the nonnegative symmetric cone, both restricted to the subalgebra of circulant matrices. Timelimit: 60 seconds

n	13	15	17	19	21	23
exact	0.762950π	0.757765π	0.764971π	0.768062π	0.768769π	0.766370π
Gur	0.762950π 0.854 s	0.757765π 25.061 s	0.764971π 60* s	0.767876π 60* s	0.765409π 60* s	0.766370π 60* s
BFAS	0.762950π 0.333 s	0.757765π 0.356 s	0.764971π 1.114 s	0.768062π 4.418 s	0.768768π 19.953 s	0.766370π 60* s

Table 3: Numerical comparison of Gur, BFAS, E-AO and SRPL for the same problem. Timelimit: 10 seconds. We also report the exact value for each problem, and the best known lower bound when the exact value is not available, indicated with an asterisk

n	17	19	21	23	25	27
exact	0.764971π	0.768062π	0.768769π	0.766370π	$0.767385\pi^*$	$0.768258\pi^*$
Gur	0.764971π	0.759309π	0.765409π	0.766370π	0.767385π	0.760879π
BFAS	0.764971π	0.768062π	0.768768π	0.766370π	0.762620π	0.756841π
E-AO	0.764971π	0.768062π	0.768768π	0.766370π	0.767385π	0.768258π
SRPL	0.764970π	0.768062π	0.768768π	0.766369π	0.767384π	0.768257π

PSD and SNN matrices

Since E-AO and SRPL main steps are projections, they can be adapted to the case of NON-polyhedral cones, as long as we know how to compute the projection on such cones

We can thus test them on the task to find the maximum angle between the cone of Positive Semi-Definite matrices and the con of Symmetric Nonnegative matrices

Table 4: Numerical comparison for E-AO and SRPL for different dimensions for the problem of finding the maximum angle between the PSD cone and the nonnegative symmetric cone. The table reports the best and average value found over 10000 random initializations, together with the average elapsed time. We also report the best known value for each dimension.

n	30	40	50	60
best known	0.7757π	0.7789π	0.7812π	0.7837π
EAO _b	0.7755π	0.7782π	0.7802π	0.7811π
EAO _a	0.7717π	0.7734π	0.7745π	0.7760π
	0.006 ± 0.002 s	0.056 ± 0.019 s	0.030 ± 0.011 s	0.057 ± 0.019 s
SRPL _b	0.7757π	0.7789π	0.7812π	0.7837π
SRPL _a	0.7739π	0.7766π	0.7787π	0.7802π
	0.062 ± 0.025 s	0.155 ± 0.060 s	0.319 ± 0.130 s	0.565 ± 0.229 s

PSD and SNN matrices

Since E-AO and SRPL main steps are projections, they can be adapted to the case of NON-polyhedral cones, as long as we know how to compute the projection on such cones

We can thus test them on the task to find the maximum angle between the cone of Positive Semi-Definite matrices and the con of Symmetric Nonnegative matrices

Table 4: Numerical comparison for E-AO and SRPL for different dimensions for the problem of finding the maximum angle between the PSD cone and the nonnegative symmetric cone. The table reports the best and average value found over 10000 random initializations, together with the average elapsed time. We also report the best known value for each dimension.

n	30	40	50	60
best known	0.7757π	0.7789π	0.7812π	0.7837π
EAO _b	0.7755π	0.7782π	0.7802π	0.7811π
EAO _a	0.7717π	0.7734π	0.7745π	0.7760π
	0.006 ± 0.002 s	0.056 ± 0.019 s	0.030 ± 0.011 s	0.057 ± 0.019 s
SRPL _b	0.7757π	0.7789π	0.7812π	0.7837π
SRPL _a	0.7739π	0.7766π	0.7787π	0.7802π
	0.062 ± 0.025 s	0.155 ± 0.060 s	0.319 ± 0.130 s	0.565 ± 0.229 s

-  Wellington de Oliveira, Valentina Sessa, and David Sossa. **Computing critical angles between two convex cones.** *Journal of Optimization Theory and Applications*, 201(2):866–898, 2024.
-  Alberto Seeger and David Sossa. **Cone-constrained singular value problems.** *J. Convex Anal.*, 30:1285–1306, 2023.
-  Alberto Seeger and David Sossa. **Singular value problems under nonnegativity constraints.** *Positivity*, 27, 2023.
-  Nicolas Gillis and Francois Glineur. **A continuous characterization of the maximum-edge biclique problem.** *J. Glob. Optim.*, 58:439–464, 2014.
-  Felix Goldberg and Naomi Shaked-Monderer. **On the maximum angle between copositive matrices.** *The Electronic Journal of Linear Algebra*, 27(1), 2014.
-  Alberto Seeger and Alfredo Iusem. **Axiomatization of the index of pointedness for closed convex cones.** *Computational and Applied Mathematics*, 24(2):245–283, 2005.
-  R. Peeters. **The maximum edge biclique problem is np-complete.** *Discret. Appl. Math.*, 131(3):651–654, 2003.
-  G. P. McCormick. **Computability of global solutions to factorable nonconvex solutions: Part i: convex underestimating problems.** *Mathematical Programming*, 10:147–175, 1976.

Workshop on Low-Rank Models and Applications (LRMA)

11-12 September 2025, Mons, Belgium



Plenary speakers: Stanislav Budzinskiy, Luca Calatroni, Alice Cortinovis, Mariya Ishteva, Paul Magron, Margherita Porcelli, Bertrand Rivet, and Lawrence Saul.

<https://sites.google.com/view/lrma25>

Thank You!