

Climate-driven unprecedented events: impacts on Monterrey urban air pollution

Adriana Ipiña^{1*}, Constanza Zúñiga-Villarreal²,
Gamaliel López-Padilla³, Jair Rafael Carrillo Ávila⁴,
Michel Grutter⁵

^{1*} Instituto de Física Rosario (CONICET-UNR), 27 de Febrero 210 BIS,
Rosario, 2000, Santa Fe, Argentina, .

² Centro de Investigación Científica y de Educación Superior de
Ensenada (CONAHCYT), Carretera Ensenada - Tijuana 3918,
Ensenada, 22860, Baja California, México .

³ Centro de Investigación en Matemáticas A.C (CONAHCYT), De
Jalisco s/n, Valenciana, Guanajuato, 36023, Guanajuato, México .

⁴ Agencia de la Calidad del Aire, Secretaría de Medio Ambiente de Nuevo
León, Washington 2000 Ote, Monterrey, 64010, Nuevo León, México .

⁵ Instituto de Ciencias de la Atmósfera y Cambio Climático,
Universidad Nacional Autónoma de México, Investigación Científica s/n,
CU, Ciudad de México, 04510, Ciudad de México, México .

*Corresponding author(s). E-mail(s): ipina@ifir-conicet.gov.ar;
Contributing authors: constanza@email; giovanni.lopez@cimat.mx;
jair@email; michel@email;

Abstract

Extreme weather episodes have increased in frequency and intensity due to climate change. Record-breaking temperatures, prolonged tropical or subtropical storms, and drought have greatly impacted many large urban areas, changing air pollutants. The present work analyzes the pollutant concentrations affected by the passage of unprecedented events such as Hurricane Hanna, cold fronts, wildfires, and African dust clouds over the Monterrey urban area in the period 2015-2022. The average percentage changes with respect to the baseline of the primary and secondary pollutants during the most strict pandemic lockdown were mainly attributed to exceptional climate phenomena. The PM₁₀ percentage differences calculated with the random forest deweather method as well as

the anomalies formula had opposite changes to raw data without the presence of extraordinary occurrences. These findings are of particular interest in establishing methods for further quantifying the influence factors and formulating a better air quality policy in large cities facing new climate challenges.

Keywords: Extreme weather events, Air pollutants, Deseasonalized, Deweather, Climate change

1 Introduction

The World Health Organization (WHO) recognizes that the disease burden associated with air pollution is similar to that of the biggest health threats in large metropolitan areas [1]. Moreover, assessing the changes in atmospheric photochemistry, meteorology, and pollution is essential for proposing effective policies on air quality. The COVID-19 pandemic provided an opportunity to study nitrogen dioxide (NO_2), ozone (O_3), particulate matter (PM), and sulfur dioxide (SO_2) worldwide [2], as most countries implemented a mandatory lockdown to stop infection, unintentionally benefiting the environment [3, 4]. The reduction in vehicle traffic and the closure of industrial operations led to a drop in atmospheric pollution [5–8], especially in the major metropolises [9, 10]. In particular, NO_2 decreased along the full lockdown period in at least 63 cities around the world [11]. In a similar way, the low $\text{PM}_{2.5}$ values reflected some degree of the partial and total activity shutdown in most cities [9, 12], such as a decrease ranging from 5.93% [13] up to 46.5% [14] in China and nearly 200% in India [15]. These changes were generally much lower in some urban areas of Europe than in Asia [6, 16] and a slight increase in some US cities [17, 18]. Since the COVID-19 disease was identified, China, India, and the USA lead the list of countries with publications about lockdown and air pollution, while Mexico is one of the most underexplored [19]. Comparative studies with respect to prior periods have predominated, but few have examined the meteorological and environmental causes [20–22], and even less in the post-pandemic period [23]. Although the pandemic restrictions provided great data to fill the knowledge gap about air pollutants, there are still unresolved questions about cities where the reduction was not significant. Monterrey Metropolitan Area (MMA) is an industrial development hub in northeastern Mexico, with more than 4000 foreign-capitalized companies accelerating urban growth [24]. Furthermore, its proximity to the United States border maintains one of the principal commercial routes and an intense flow of heavy transportation [25]. The Pan American Health Organization has recognized PM_{10} in MMA as one of the highest in Latin America [26]. Its particle composition includes black carbon (BC) [27] sulfates, organic carbon (OC) [28] and Volatile Organic Compounds (VOCs), with a seasonal dependence on maximum photolysis and temperature [29]. Carbonyl compounds increase in the morning and decrease in the afternoon [30], which are sensitive to the NO_x – O_3 relationship [31]. Likewise O_3 has a sensitive response to the changes in NO_x and VOCs emissions, meteorology, and photochemistry [32]. Mobile sources are the main regional source of $\text{PM}_{2.5}$ and its precursors (NO_x , VOCs, and NH_3) [33]. The state vehicle register

enlists about 2 million, 700 thousand auto motors [34], that substantially contribute to metropolitan emissions. Recently, Tesla announced the construction of its largest plant in the world on the Monterrey outskirts [35]. This urban area is a key point for the investigation of economic growth, increasing emissions, and the energy transition. Consequently, the aim of this work is to analyze eight years (2015-2022) of air pollutant observations affected by unprecedented events in Monterrey, one of the largest cities in the Western Hemisphere [36].

2 Materials and Methods

2.1 Mobility and COVID-19 cases

The *Healthy Distance Program* was a countrywide strategy in Mexico based on social distancing measures and the closure of all non-essential activities [37]. From the beginning of the pandemic, some local governments disagreed with the national recommendations, as was in the MMA [38]. It implemented a mixture of measures from March 2020 to October 2021 (Table S1). Social media information was widely used as a measure of mobility during the lockdown. Figure S1 shows the COVID-19 cases [39], the mobility from a *Python* library [40], and the Unified Mobility Index (MIU) [41].

2.2 Ground data

The Integral Environmental Monitoring System (SIMA) records in the MMA the criteria pollutants $PM_{2.5}$, PM_{10} , NO_2 , O_3 , SO_2 , and CO . Additionally, this network also collected meteorological data on humidity, temperature, global solar irradiance, precipitation, wind speed and direction. The available data for the period 2015–2022 were analyzed, except for the SE3 station that started operations in 2017. The CO measurements were not included because of a lack of calibration from 2016 to 2021. Figure 1 shows and lists the selected stations that comply with at least 75% of the available monthly measurements per pollutant [42].




Fig. 1 SIMA stations (red dots) in the MMA.

2.2.1 Deseasonalized air pollutants data and anomaly calculation

A function derived from the linear term and truncated Fourier series was fitted with a frequency of $K = 2$, and its coefficients were obtained [20] for the baseline (January 2015 to December 2019). The air pollution anomalies $A(t)$ were calculated as follows:

$$A(t) = D(t) - \left[c_0 + c_1 t + \sum_{k=1}^K (a_k \cos(2\pi t k) + b_k \sin(2\pi t k)) \right] \quad (1)$$

By subtracting the linear trend ($c_0 + c_1 t$) and the Fourier fitting from the original air pollutant measurements $D(t)$, the influence of seasonality and long-term trends were reduced. Considering meteorological conditions, the pollutant measurements in the presence of daily accumulated precipitation of 10 mm and wind speed values of 60 km/h were filtered.

2.3 Deweather model for PM₁₀ data

Machine learning methods to determine changes in air quality have gained popularity over the past decades. Random Forest (RF) is a regression model that has been implemented to extract meteorological influences in air pollution [43–45]. It involves applying a normalization technique to meteorology based on pollutant emissions [46–48]. The RF model was trained using measurements in the period 2015–2022, with resampling for each monitoring station. The algorithm inputs were the meteorological (relative humidity, temperature, pressure, direction, and velocity wind) and temporal (hour and Julian day) variables to predict the PM₁₀. The RF process repeats hundreds to thousands of times until it obtains an average regression called Deweather. In this work, the Deweather model was a method to evaluate the monthly changes in PM₁₀ during lockdown, comparing the results with anomalies analysis.

2.4 Satellite data

2.4.1 OMI

The Total Ozone Column (TOC) [49], NO₂ Vertical Column Density (VCD) [50], and maps of Aerosol Optical Depth at 500nm (AOD₅₀₀) are provided by the Ozone Monitoring Instrument (OMI) on board the AURA-NASA satellite. The satellite overpass time is between 13:00 and 14:00 h CST for the coordinates of Monterrey downtown (25.40°N, 100.28°W, and 560m asl). The products used were OMTO3 (L2) v8.5 for TOC, OMNO2 (L2) v4.0 for VCD, and OMAERUVd (L3) v3 for AOD₅₀₀. Gases information was utilized to determine concentration patterns. For maps of the incoming aerosols from Africa, the Giovanni platform was employed [51].

2.4.2 VIIRS

The satellite instrument of the Visible Infrared Imaging Radiometer Suite (VIIRS) [52] was used for tracking wildfires. Near Real-Time NOAA-20 (formally JPSS-1) VIIRS Active Fire detection product is derived collaboratively by NASA and National

Oceanic and Atmospheric Administration (NOAA). VIIRS has local overpass times of approximately 01:30 -13:30 h with a spatial resolution of 375 m. The data product Fire Radiative Power [53] was processed to localize the active fire and estimate the number of hotspots related to wildfires on the mountain range (the Sierra Madre Oriental, henceforth SMO) nearby MMA.

2.5 HYSPLIT model

The Hybrid-Single Particle Lagrangian Integrated Trajectory (HYSPLIT) model from NOAA - Air Resources Laboratory (ARL) [54] was executed on the Realtime Environmental Applications and Display System website to demonstrate dust transport. Data from reanalysis 2015–2022, stored on the HYSPLIT website, were applied for wind backward-trajectories simulations during African dust events.

3 Results and Discussion

3.1 Unprecedented events

Affected pollutant concentrations from the passage of an unprecedented cold front, large African dust clouds, a hurricane, record wildfires, an atypical drought, the COVID-19 lockdown, and even controls prior were analyzed.

3.1.1 Cold fronts

The frontal systems crossing the northern hemisphere considerably affect the air quality along their transport pathways. As the cold front enters an urban area, the contaminated air mass from other regions can cause a sharp increase in PM_{10} [55]. Figure S2 shows an extreme PM_{10} value of $430\mu g m^{-3}$ after cold front arrival. At the end of 2019 and the beginning of 2020, the long persistence of the jet stream with westerly winds and widespread displacement toward Mexican territory increased the number of cold fronts [56]. This phenomenon caused low raw averages of PM_{10} , $PM_{2.5}$, and NO_2 (Figures S3 and S4). Mexico City also recorded this decline in January and February 2020, prior to the pandemic restrictions [57]. In mid-February 2021, an unprecedented winter storm crossed the United States, causing failures in the electricity system supply in Texas [58, 59] at the same time as a new closure (L2) in the MMA. The cold snap whipped as far as northeastern Mexico, decreasing temperatures by below $10^\circ C$ along with particle matter (Figure S2). The air pollutants measured during the passage of the cold fronts were filtered for the analysis of anomalies.

3.1.2 African dust

In June 2020, the strongest dust storm over the last two decades (dubbed *Godzilla*) transported massive amounts of dust from Africa to the Americas [60]. The origin, transport, and deposition of dust associated with the PM_{10} increase can be identified by comparing the AOD_{500} and HYSPLIT model [61]. Figure S5 illustrates the back trajectories from HYSPLIT about the main dust transportation routes from northwest Africa to the southeast part of the MMA. This was aligned with the AOD_{500} records

from the OMI satellite before and during the COVID-19 lockdown. Part of the satellite grid cells cover a mountain bend that acts as a barrier by accumulating particles [62, 63]. On the other hand, SIMA sensors registered the arrival of the dust storm modeled by HYSPLIT, exhibiting ridges on 16 July 2018 and 17 July 2020 (Figure S6). The dusty episodes during the June-July period have also been recorded in Puerto Rico (18.22°N, 66.52°W) [64] and Houston (667 km northeast of Monterrey) coincident with the large-scale Saharan dust intrusion [65] when the dry deposition peak over the eastern Gulf of Mexico [66]. The dust is present year-round due to erosion of soil, unpaved roads, particle resuspension, and quarry activities. Notwithstanding, PM₁₀ values rise midyear when African dust arrives, exceeding the WHO air quality guidelines [67]. As at other sites affected by Saharan dust outbreaks, these PM₁₀ measurements were removed [68] before applying the anomaly formula.

3.1.3 Hurricane Hanna

The number of systems that at least qualified as tropical or subtropical storms during the 2020 Atlantic hurricane season exceeded that of the previous record-holder (in 2005) [69, 70]. Tropical Storm Hanna was the earliest 8th storm on record, and becoming the season's first hurricane. The eye made landfall 500 km from Monterrey (Figure S7) with maximum sustained winds of 150 km/h and a minimum central pressure of 973 hPa [71]. It was the last significant rainfall in the MMA, crossing the region at the end of July 2020 and dropping the daily PM₁₀ for four days (Figure S6). The maximum precipitation accumulated was 489 mm at SE3 station and 604 mm in the vicinity of NE station (25.73°N, 100.30°W). Table S2 shows the PM₁₀ averages in July for the years 2020 and 2019, with the effects of African dust and Hurricane Hanna and without both events, respectively. The mean percentage relative differences (RD) without measurements affected by these episodes was -0.4%.

3.1.4 Wildfires

From March to May in the central region of Mexico, wildfires occur [72]. Early in the spring of 2021, one of the most devastating forest fires in the last decade raged over the SMO. Wildfires destroyed almost 1000 hectares and forced the evacuation of roughly 400 people. The VIIRS and the Moderate Resolution Imaging Spectroradiometer (MODIS) on board the Terra-NASA satellite detected at least eight clusters of hotspots (active fires) visible in the forested region, with smoke flowing northeast [73]. In 2022, a second historical record of wildfires was observed in the same area. Figure S8 depicts the daily number of hotspots between 2015-2022 detected by VIIRS and a map of active fires during the most critical period. For assessing impact, a weighted kernel function was applied to the PM₁₀ and the wind speed and direction measurements in a specific radius [74–76]. Figure S9 shows the non-parametric regression for five stations from 13 to 18 March 2021. Wildfire increased the PM₁₀ values higher than 100 μgm^{-3} with the dominant southeast wind. Smoke is partially responsible for the increase of monthly average PM₁₀ in March 2021 and 2022 (Figure S2). In Mexico City outskirts, it produces over 70% of the principal fine particle mass [77] and several trace gases in northeast Mexico [78]. This PM₁₀ increase does not seem

to have a relationship between O_3 monthly mean and the number of fire detections, as in other regions of Mexico [79].

3.1.5 COVID-19 lockdown

In general, the particle matter decline in the first two months of 2020 was due to the atypical number of cold fronts. Furthermore, the meteorological conditions led to high-ozone episodes and driven the dispersion and accumulation of pollutants during the lockdown period [80–82]. Despite 54.6% of the population using their own vehicles as the main mode of transportation to work [83] and more than 90% of them having mobile internet connection [84], PM and NO_2 did not show a proportional reduction to the MIU in 2020. This result was more in line with the transport of air masses and meteorological parameters than the vehicular flux reduction, similar to those of Rio de Janeiro [85] and Ireland [86]. The predominance of the decrease in pollutants without taking into account environmental and meteorological factors could be falsely attributed to the COVID-19 lockdown. The environmental episodes should be taken into account in lockstep [87] and they can help other cities identify natural and anthropogenic influences.

3.2 Satellite NO_2 and O_3 changes

The NO_2 VCD monthly averages derived from OMI data [50] were used to calculate the RD for Monterrey city ($25.67^\circ N$ and $100.30^\circ W$). The pixel size is $1^\circ \times 1^\circ$ with grid boxes drawn around the city coordinates. Figure S10 shows the main NO_2 VCD reductions in April and October 2020, around -21% and -19%, respectively. The values are lower than half of those reported in other large cities [88]. The lowest VCD was 31% in August 2022, during a total opening of activities. A deseasonalized analysis from platform Air Quality/NASA based on satellite data for Monterrey indicated a slight change from March to June 2020 during L1. Indeed, two months before the COVID-19 lockdown, the values were lower than in that period [89]. In the case of the O_3 column from OMI, peaks appeared on December 2021 and November 2022, within variations lower than $\pm 10\%$ along the period, implicating the main changes were in the tropospheric ozone.

3.3 Air Pollutants long term and anomalies

The air pollutants were not spatially homogeneous in the MMA due to the orography, urban sprawl, and a wide variety of anthropogenic sources. At least three years before the pandemic outbreak, PM_{10} , $PM_{2.5}$ and NO_2 (at SE3) concentrations decreased. This fact could have caused the O_3 formation from increased UV radiation [90]. Figure 2 shows the truncated Fourier series and the linear trends for the monthly means of criteria pollutants in the period 2015–2022. The highest values for PM_{10} , $PM_{2.5}$ and NO_2 occur in wintertime when O_3 reaches the minimum values. By contrast, SO_2 had no clear annual pattern. The solar UV radiation is around two times higher in the summer than in the winter and it promotes the formation of ozone and the aging of organic aerosols. The PM_{10} and $PM_{2.5}$ trends exhibited clear decreases on practically all stations, following prior findings [91], attributed to the government’s corresponding

policy related to the 2017 controls on quarries and industries. Higher levels of particle concentration were recorded in the western zone, consistent with the predominant wind direction from east to west [62]. A drastic drop in SO_2 at SW station was caused by ceasing in a factory as the main source of emission. In the post-pandemic period (2021–2022), a regional extreme drought promoted severe pollution accumulation along with low wind speeds and the second opening stage (N2). Excluding ozone, an increase in the daily pollutant concentrations was noticed in the first half of 2022 at almost all stations. The downtown station (near the water supply service of Monterrey) recorded a significant increase in NO_2 linked to the passage of heavy trucks used as water pipes during the drought-related crisis.




Fig. 2 Monthly averages of criteria pollutant concentrations recorded from 2015 to 2022 in the MMA, the linear trend, the Fourier fitting, and the anomaly $A(t)$. The main lockdown were L1 (pink dotted line), L2 (violet dotted line), and the complete systematic phase (light yellow area).

The trends decreased on average up to 2019 for PM_{10} (-4 %/yr), $\text{PM}_{2.5}$ (-7 %/yr) and NO_2 (-1 %/yr), while O_3 and SO_2 had an increase of approximately 8%/yr and 14%/yr, respectively (Table S3). The daily data set was filtered, considering meteorological and environmental factors. Subsequently, Equation 1 was applied to observe the net variation. The anomalies over the complete period exhibited values of $A(t) < 0$ not only in the lockdown context (Figure 2). The baseline anomaly and overlapping

years of 2020, 2021, and 2022 revealed the most noticeable changes were in PM_{10} during L1, as shown in Figure 3. The lowest anomaly was at NW, near the subway, encompassing high mobility and unpaved roads.




Fig. 3 Monthly average anomalies of PM_{10} , $PM_{2.5}$, NO_2 , O_3 , and SO_2 for the years 2020, 2021, and 2022, and the period 2015-2019 with its standard deviation.

The real average percentage changes during strict lockdown (L1) with respect to baseline in PM_{10} , $PM_{2.5}$ and SO_2 declined up to -14%, with an increase of 20% for O_3 and 10% for NO_2 (Table S4). The low reductions for PM_{10} , $PM_{2.5}$ and NO_2 are consistent with the uninterrupted passage of heavy transport, which was considered an essential activity along with most industries. The ascending behavior at the end of 2021 and 2022 was present in the majority of pollutants, due to an exceptional drought period. The O_3 reached their maximum in the spring, followed by a prolonged slump until the early autumn coincident with the maximum UV solar intensity. In the cold dry season, the low photolysis rate might lead to a low conversion of SO_2 to sulfate ions and thus accumulate higher concentrations [92]. At least since September 2020, the decreasing of SO_2 could be inferred from the fact that several companies and industries in Sta. Catarina (SW) did not work during L1. The Cadereyta refinery near SE3 station had SO_2 and NO_2 levels below the baseline between 2021 and 2022 due to an improvement in the refinery emissions system in spite of uninterrupted activity and an intense drought.

3.4 Deweather model performance

A special normalization of the meteorological effects on the PM_{10} concentrations was carried out by means of the deweather-RF method. The results of ten metrics (Table S5) used to evaluate the deweather model performance aligned with the PM_{10} values [48]. The non-linearity between predicted and observed values generated a mean Pearson correlation coefficient of 0.6, which is evidence of large differences due to extreme events. Figure S11 displays the behavior during the months involving relevant events.

3.5 PM_{10} changes in extraordinary events

The extraordinary events most significant were the COVID-19 restrictions (March 2020 to October 2021), storms (April 2020), hurricane Hanna, African dust (July 2020), cold fronts (February 2021), and wildfires (March 2021). Figure 4 shows the monthly mean PM_{10} changes with respect to the baseline using raw data, the deweather model and the anomaly formula in the absence of environmental and meteorological episodes. An apparent reduction occurred by subtracting only the absolute values, except in the month of wildfires. Instead, the net changes calculated by Eq. 1 were minor compared to the raw data, with differences more than twice and even opposite. Despite the fact that the deweathering method did not account environmental phenomena such as wildfires and African dust, the results were more similar to anomalies with filtered data. It demonstrates the relevance of considering these factors because they can inverse the changes in air pollution.

4 Conclusion

This study seeks to provide a broader picture of changes in air pollution due to the impact of unprecedented events beyond the COVID-19 lockdown effects. In this region with an industrial heritage and continuous growth, the opening-closure of activities was not really more relevant than the extraordinary phenomena. There was no "background atmosphere" in the MMA due to the pandemic restrictions, as has been reported elsewhere. Since the heavy transport, the larger factories and companies worked as usual. PM_{10} assessing by three methods revealed that the main episodes in the context of the health crisis influenced the differences with respect to baseline and even produced opposite changes. Climate change could intensify extreme events and consequently have a growing impact on atmospheric pollution, to which a large number of people are exposed. It is important to consider in pollution models the long drought periods that increase the likelihood of wildfires and dust episodes. This analysis advances the understanding of the natural contributions needed to formulate better air quality policies in large cities.

5 Acknowledgments

The authors would like to thank the Nuevo León State government's Sistema Integral de Monitoreo Ambiental for providing the measurements. AI thanks Armandina Valdez of the Air Quality Agency for the management support, Mario Graff for the advisory




Fig. 4 Monthly mean changes of PM_{10} with respect to baseline using raw data, the deweather model, and the anomaly formula. The last two take into account the passage of storms (April 2020), hurricane Hanna and African dust (July 2020), cold fronts (February 2021), and wildfires (March 2021).

about the IM method, Lok Lamsal (NO_2) and Michael Yan (O_3) for satellite data. CZV and GLP thank CONAHCYT for their fellowships.

References

- [1] WHO: Global air quality guidelines: Particulate matter ($PM_{2.5}$ and PM_{10}), ozone, nitrogen dioxide, sulfur dioxide and carbon monoxide (2021)
- [2] Kumari, P., Toshniwal, D.: Impact of lockdown on air quality over major cities across the globe during COVID-19 pandemic. *Urban Climate* **34**, 100719 (2020) <https://doi.org/10.1016/j.uclim.2020.100719>
- [3] Han, J., Yin, J., Wu, X., Wang, D., Li, C.: Environment and COVID-19 incidence: A critical review. *Journal of Environmental Sciences* (2022) <https://doi.org/>

- [4] Ravindra, K., Singh, T., Vardhan, S., Shrivastava, A., Singh, S., Kumar, P., Mor, S.: Covid-19 pandemic: What can we learn for better air quality and human health? *Journal of Infection and Public Health* **15**(2), 187–198 (2022) <https://doi.org/10.1016/j.jiph.2021.12.001>
- [5] Barré, J., Petetin, H., Colette, A., Guevara, M., Peuch, V.-H., Rouil, L., Engelen, R., Inness, A., Flemming, J., García-Pando, C.P., Bowdalo, D., Meleux, F., Geels, C., Christensen, J.H., Gauss, M., Benedictow, A., Tsyro, S., Friese, E., Struzewska, J., Kaminski, J.W., Douros, J., Timmermans, R., Robertson, L., Adani, M., Jorba, O., Joly, M., Kouznetsov, R.: Estimating lockdown-induced european no₂ changes using satellite and surface observations and air quality models. *Atmospheric Chemistry and Physics* **21**(9), 7373–7394 (2021) <https://doi.org/10.5194/acp-21-7373-2021>
- [6] Sicard, P., De Marco, A., Agathokleous, E., Feng, Z., Xu, X., Paoletti, E., Rodriguez, J.J.D., Calatayud, V.: Amplified ozone pollution in cities during the covid-19 lockdown. *Science of The Total Environment* **735**, 139542 (2020) <https://doi.org/10.1016/j.scitotenv.2020.139542>
- [7] Sharma, S., Zhang, M., Anshika, Gao, J., Zhang, H., Kota, S.H.: Effect of restricted emissions during COVID-19 on air quality in India. *Science of The Total Environment* **728**, 138878 (2020) <https://doi.org/10.1016/j.scitotenv.2020.138878>
- [8] Venter, Z.S., Aunan, K., Chowdhury, S., Lelieveld, J.: COVID-19 lockdowns cause global air pollution declines with implications for public health risk. *PNAS* **117**(32), 18984–18990 (2020) <https://doi.org/10.1073/pnas.2006853117>
- [9] Saha, L., Kumar, A., Kumar, S., Korstad, J., Srivastava, S., Baudhh, K.: The impact of the COVID-19 lockdown on global air quality: A review. *Environmental Sustainability* (2022) <https://doi.org/10.1007/s42398-021-00213-6>
- [10] Slezakova, K., Pereira, M.C.: 2020 COVID-19 lockdown and the impacts on air quality with emphasis on urban suburban and rural zones. *Scientific Reports* **11**(1) (2021) <https://doi.org/10.1038/s41598-021-99491-7>
- [11] Sokhi, R.S., Singh, V., Querol, X., Finardi, S., Targino, A.C., Fatima Andrade, M., Pavlovic, R., Garland, R.M., Massagué, J., Kong, S., Baklanov, A., Ren, L., Tarasova, O., Carmichael, G., Peuch, V.-H., Anand, V., Arbilla, G., Badali, K., Beig, G., Belalcazar, L.C., Bolignano, A., Brimblecombe, P., Camacho, P., Casaslas, A., Charland, J.-P., Choi, J., Chourdakis, E., Coll, I., Collins, M., Cyrus, J., da Silva, C.M., Di Giosa, A.D., Di Leo, A., Ferro, C., Gavidia-Calderon, M., Gayen, A., Ginzburg, A., Godefroy, F., Gonzalez, Y.A., Guevara-Luna, M., Haque, S.M., Havenga, H., Herod, D., Hörrak, U., Hussein, T., Ibarra, S.,

- Jaimes, M., Kaasik, M., Khaiwal, R., Kim, J., Kousa, A., Kukkonen, J., Kulmala, M., Kuula, J., La Violette, N., Lanzani, G., Liu, X., MacDougall, S., Manseau, P.M., Marchegiani, G., McDonald, B., Mishra, S.V., Molina, L.T., Mooibroek, D., Mor, S., Moussiopoulos, N., Murena, F., Niemi, J.V., Noe, S., Nogueira, T., Norman, M., Pérez-Camaño, J.L., Petäjä, T., Piketh, S., Rathod, A., Reid, K., Retama, A., Rivera, O., Rojas, N.Y., Rojas-Quincho, J.P., San José, R., Sánchez, O., Seguel, R.J., Sillanpää, S., Su, Y., Tapper, N., Terrazas, A., Timonen, H., Toscano, D., Tsegas, G., Velders, G.J.M., Vlachokostas, C., von Schneidmesser, E., VPM, R., Yadav, R., Zalakeviciute, R., Zavala, M.: A global observational analysis to understand changes in air quality during exceptionally low anthropogenic emission conditions. *Environment International* **157**, 106818 (2021) <https://doi.org/10.1016/j.envint.2021.106818>
- [12] Chauhan, A., Singh, R.P.: Decline in PM_{2.5} concentrations over major cities around the world associated with COVID-19. *Environmental Research* **187**, 109634 (2020) <https://doi.org/10.1016/j.envres.2020.109634>
- [13] Bao, R., Zhang, A.: Does lockdown reduce air pollution? Evidence from 44 cities in northern China. *Science of the Total Environment* **731**, 139052 (2020)
- [14] Xu, K., Cui, K., Young, L.-H., Wang, Y.-F., Hsieh, Y.-K., Wan, S., Zhang, J.: Air quality index indicator air pollutants and impact of covid-19 event on the air quality near central china. *Aerosol and Air Quality Research* **20**(6), 1204–1221 (2020) <https://doi.org/10.4209/aaqr.2020.04.0139>
- [15] Kotnala, G., Mandal, T.K., Sharma, S.K., Kotnala, R.K.: Emergence of Blue Sky Over Delhi Due to Coronavirus Disease (COVID-19) Lockdown Implications. *Aerosol Science and Engineering* **4**(3), 228–238 (2020) <https://doi.org/10.1007/s41810-020-00062-6>
- [16] Fu, F., Purvis-Roberts, K.L., Williams, B.: Impact of the COVID-19 Pandemic Lockdown on Air Pollution in 20 Major Cities around the World. *Atmosphere* **11**(11), 1189 (2020) <https://doi.org/10.3390/atmos11111189>
- [17] Zangari, S., Hill, D.T., Charette, A.T., Mirowsky, J.E.: Air quality changes in New York City during the COVID-19 pandemic. *Science of The Total Environment* **742**, 140496 (2020) <https://doi.org/10.1016/j.scitotenv.2020.140496>
- [18] Archer, C.L., Cervone, G., Golbazi, M., Fahel, N.A., Hultquist, C.: Changes in air quality and human mobility in the USA during the COVID-19 pandemic. *Bulletin of Atmospheric Science and Technology* **1**(3-4), 491–514 (2020) <https://doi.org/10.1007/s42865-020-00019-0>
- [19] Aboagye, E.M., Effah, N.A.A., Effah, K.O.: A bibliometric analysis of the impact of covid-19 social lockdowns on air quality: research trends and future directions. *Environmental Science and Pollution Research* **30**, 74500–74520 (2023) <https://doi.org/10.1007/s11356-023-27699-3>

- [20] Hernández-Paniagua, I.Y., Valdez, S.I., Almanza, V., Rivera-Cárdenas, C., Grutter, M., Stremme, W., García-Reynoso, A., Ruiz-Suárez, L.G.: Impact of the covid-19 lockdown on air quality and resulting public health benefits in the mexico city metropolitan area. *Frontiers in Public Health* **9**, 642630 (2021) <https://doi.org/10.3389/fpubh.2021.642630>
- [21] Bontempi, E., Carnevale, C., Cornelio, A., Volta, M., Zanoletti, A.: Analysis of the lockdown effects due to the COVID-19 on air pollution in Brescia (Lombardy). *Environmental Research* **212**, 113193 (2022) <https://doi.org/10.1016/j.envres.2022.113193>
- [22] Zoran, M.A., Savastru, R.S., Savastru, D.M., Tautan, M.N.: Peculiar weather patterns effects on air pollution and covid-19 spread in tokyo metropolis. *Environmental Research* **228**, 115907 (2023) <https://doi.org/10.1016/j.envres.2023.115907>
- [23] Priya, S., Iqbal, J.: Assessment of no(2) concentrations over industrial state jharkhand, at the time frame of pre, concurrent, and post-covid-19 lockdown along with the meteorological behaviour: an overview from satellite and ground approaches. *Environmental Science and Pollution Research* **30**, 68591–68608 (2023) <https://doi.org/10.1007/s11356-023-27236-2>
- [24] Nuevo Leon, S.: Monterrey, Nuevo León Is One of the Most Attractive Regions to Invest and do Business in North America. (2022). https://www.nl.gob.mx/sites/default/files/nuevo_leon_2s2022_para_pdf_sept_22_1.pdf
- [25] CMM: PROYECTO: “Propuestas para el desarrollo sustentable de una ciudad mexicana” Estudio del Área Metropolitana de Monterrey. (2019). http://aire.nl.gob.mx/docs/reportes/Propuestas_de_sustentabilidad_Monterrey.pdf
- [26] Riojas-Rodríguez, H., Silva, A.S., Texcalac-Sangrador, J.L., Moreno-Banda, G.L.: Air pollution management and control in Latin America and the Caribbean: implications for climate change. *Pan American Journal of Public Health* **40**, 150–159 (2016)
- [27] Peralta, O., Ortíz-Alvarez, A., Basaldud, R., Santiago, N., Alvarez-Ospina, H., Cruz, K., Barrera, V., Luz Espinosa, M., Saavedra, I., Castro, T., Martínez-Arroyo, A., Páramo, V.H., Ruíz-Suárez, L.G., Vazquez-Gálvez, F.A., Gavilán, A.: Atmospheric black carbon concentrations in Mexico. *Atmospheric Research* **230**, 104626 (2019) <https://doi.org/10.1016/j.atmosres.2019.104626>
- [28] Mancilla, Y., Paniagua, I.Y.H., Mendoza, A.: Spatial differences in ambient coarse and fine particles in the Monterrey metropolitan area Mexico: Implications for source contribution. *Journal of the Air & Waste Management Association* **69**(5), 548–564 (2019) <https://doi.org/10.1080/10962247.2018.1549121>

- [29] Menchaca-Torre, H.L., Mercado-Hernández, R., Mendoza-Domínguez, A.: Diurnal and seasonal variation of volatile organic compounds in the atmosphere of Monterrey Mexico. *Atmospheric Pollution Research* **6**(6), 1073–1081 (2015) <https://doi.org/10.1016/j.apr.2015.06.004>
- [30] D.M., F.-T., E., R.-L., J.G., C.-B., R.M., C.-B., Y., G.-V., R., M.-G., J., R.D.L.R.: Measurement of Carbonyls and its relation with Criteria Pollutants (O₃, NO, NO₂, NO_x, CO and SO₂) in an Urban Site within the Metropolitan Area of Monterrey, in Nuevo León, México. *International Journal of Energy and Environment* **6** (2012)
- [31] Menchaca-Torre, H.L., Mercado-Hernández, R., Rodríguez-Rodríguez, J., Mendoza-Domínguez, A.: Diurnal and seasonal variations of carbonyls and their effect on ozone concentrations in the atmosphere of Monterrey Mexico. *Journal of the Air & Waste Management Association* **65**(4), 500–510 (2015) <https://doi.org/10.1080/10962247.2015.1005849>
- [32] Paniagua, I.Y.H., K.C.C., Mendoza, A.: Observed trends in ground-level o₃ in monterrey, mexico, during 1993–2014: comparison with mexico city and guadalajara. *Atmospheric Chemistry and Physics* **17**(14), 9163–9185 (2017) <https://doi.org/10.5194/acp-17-9163-2017>
- [33] Martínez-Cinco, M., Santos-Guzmán, J., Mejía-Velázquez, G.: Source apportionment of PM_{2.5} for supporting control strategies in the Monterrey Metropolitan Area, Mexico. *Journal of the Air & Waste Management Association* **66**, 631–42 (2016)
- [34] Instituto de Control Vehicular-Gobierno del Estado de Nuevo León. Accessed on Thu, January 19, 2023 (2022). <https://www.nl.gob.mx/boletines-comunicados-y-avisos/anuncia-instituto-de-control-vehicular-baja-de-automoviles-con-placas>
- [35] Elon Musk announces plan to build the world’s largest Tesla plant in Mexico. Accessed on Sat, Oct 7, 2023 (2023). <https://www.gob.mx/sre/prensa/elon-musk-announces-plan-to-build-the-largest-tesla-plant-in-the-world-in-mexico>
- [36] Ni, J., Zhao, Y., Li, B., Liu, J., Zhou, Y., Zhang, P., Shao, J., Chen, Y., Jin, J., He, C.: Investigation of the impact mechanisms and patterns of meteorological factors on air quality and atmospheric pollutant concentrations during extreme weather events in zhengzhou city, henan province. *Atmospheric Pollution Research* **14**(12), 101932 (2023) <https://doi.org/10.1016/j.apr.2023.101932>
- [37] Secretaría de Gobernación: Presidential decree to establish measures to mitigate and control health risks involved in the disease caused by the SARS-CoV2 virus (COVID-19) (2020). https://dof.gob.mx/nota_detalle.php?codigo=5590340&fecha=24/03/2020#gsc.tab=0
- [38] Official Journal of the Nuevo León State: Official Publications about COVID 19

- (2020). <https://saludnl.gob.mx/regulacion-sanitaria/index.php/acuerdos/>
- [39] COVID-19 México. Accessed on Mon, January 30, 2023 (2023). <https://datos.covid-19.conacyt.mx/#COMNac>
- [40] Graff, M., Moctezuma, D., Miranda-Jiménez, S., Tellez, E.S.: A Python library for exploratory data analysis on twitter data based on tokens and aggregated origin-destination information. *Computers & Geosciences* **159**, 105012 (2022) <https://doi.org/10.1016/j.cageo.2021.105012>
- [41] índice de Movilidad — CONACYT. Accessed on Wed, December 14, 2022 (2022). <https://salud.conacyt.mx/coronavirus/investigacion/>
- [42] aire.nl.gob.mx. Accessed on Fri, March 17, 2023 (2015). http://aire.nl.gob.mx/nor_metodos.html
- [43] Breiman, L. *Machine Learning* **45**(1), 5–32 (2001) <https://doi.org/10.1023/a:1010933404324>
- [44] Vu, T.V., Shi, Z., Cheng, J., Zhang, Q., He, K., Wang, S., Harrison, R.M.: Assessing the impact of clean air action on air quality trends in beijing using a machine learning technique. *Atmos. Chem. Phys.* **19**(17), 11303–11314 (2019) <https://doi.org/10.5194/acp-19-11303-2019>
- [45] Cole, M.A., Elliott, R.J.R., Liu, B.: The impact of the wuhan covid-19 lockdown on air pollution and health: A machine learning and augmented synthetic control approach. *Environ Resource Econ* **76**(4), 553–580 (2020) <https://doi.org/10.1007/s10640-020-00483-4>
- [46] Grange, S.K., Carslaw, D.C.: Using meteorological normalisation to detect interventions in air quality time series. *Science of The Total Environment* **653**, 578–588 (2019) <https://doi.org/10.1016/j.scitotenv.2018.10.344>
- [47] Shi, Z., Song, C., Liu, B., Lu, G., Xu, J., Vu, T.V., Elliott, R.J.R., Li, W., Bloss, W.J., Harrison, R.M.: Abrupt but smaller than expected changes in surface air quality attributable to covid-19 lockdowns. *Science Advances* **7**(3), 6696 (2021) <https://doi.org/10.1126/sciadv.abd6696>
- [48] Lv, Y., Tian, H., Luo, L., Liu, S., Bai, X., Zhao, H., Lin, S., Zhao, S., Guo, Z., Xiao, Y., Yang, J.: Meteorology-normalized variations of air quality during the covid-19 lockdown in three chinese megacities. *Atmospheric Pollution Research* **13**(6), 101452 (2022) <https://doi.org/10.1016/j.apr.2022.101452>
- [49] NASA EOS/Aura Validation Data Center (AVDC) - Correlative data, Field of View Predictions, Data Subsets, GEOMS, DCIO. Accessed on Fri, March 17, 2023 (2010). https://avdc.gsfc.nasa.gov/pub/most_popular/overpass/OMI/

- [50] Lamsal, L.N., Krotkov, N.A., Vasilkov, A., Marchenko, S., Qin, W., Yang, E.-S., Fasnacht, Z., Joiner, J., Choi, S., Haffner, D., Swartz, W.H., Fisher, B., Bucsela, E.: OMI/Aura Nitrogen Dioxide Standard Product with Improved Surface and Cloud Treatments. *Atmospheric Measurement Techniques* **14**(1), 455–479 (2021) <https://doi.org/10.5194/amt-2020-200>
- [51] Torres, O.O.: OMI/Aura Near UV Aerosol Optical Depth and Single Scattering Albedo L3 1 day 1.0 degree x 1.0 degree V3. NASA Goddard Space Flight Center, Goddard Earth Sciences Data and Information Services Center (GES DISC) **10.5067/Aura/OMI/DATA3003** (2008)
- [52] Schroeder, W., Oliva, P., Giglio, L., Csiszar, I.A.: The New VIIRS 375 m active fire detection data product: Algorithm description and initial assessment. *Remote Sensing of Environment* **143**, 85–96 (2014) <https://doi.org/10.1016/j.rse.2013.12.008>
- [53] VJ114IMGTDL_NRT Earthdata. Accessed on Wed, January 25, 2023 (2021). https://doi.org/10.5067/FIRMS/VIIRS/VNP14IMGT_NRT.002
- [54] Stein, A.F., Draxler, R.R., Rolph, G.D., Stunder, B.J.B., Cohen, M.D., Ngan, F.: NOAA’s HYSPLIT Atmospheric Transport and Dispersion Modeling System. *Bulletin of the American Meteorological Society* **96**(12), 2059–2077 (2015) <https://doi.org/10.1175/bams-d-14-00110.1>
- [55] Kang, H., Zhu, B., Gao, J., He, Y., Wang, H., Su, J., Pan, C., Zhu, T., Yu, B.: Potential impacts of cold frontal passage on air quality over the Yangtze River Delta China. *Atmospheric Chemistry and Physics* **19**(6), 3673–3685 (2019) <https://doi.org/10.5194/acp-19-3673-2019>
- [56] Preven evento de Norte con rachas mayores a 70 km/h y oleaje de 2 a 4 m en costas de Campeche, Tabasco, Tamaulipas, Veracruz y Yucatán. Servicio Meteorológico Nacional. Accessed on Sat, February 04, 2023 (2020). <http://www.gob.mx/smn/prensa/preven-evento-de-norte-con-rachas-mayores-a-70-km-h-y-oleaje-de-2-a-4-m-en-costas-de-campeche-tabasco-tamaulipas-veracruz-y-yucatan-234066?idiom=es>
- [57] Vega, E., Namdeo, A., Bramwell, L., Miquelajauregui, Y., Resendiz-Martinez, C.G., Jaimes-Palomera, M., Luna-Falfan, F., Terrazas-Ahumada, A., Maji, K.J., Entwistle, J., Enríquez, J.C.N., Mejia, J.M., Portas, A., Hayes, L., McNally, R.: Changes in air quality in Mexico City London and Delhi in response to various stages and levels of lockdowns and easing of restrictions during COVID-19 pandemic. *Environmental Pollution* **285**, 117664 (2021) <https://doi.org/10.1016/j.envpol.2021.117664>
- [58] Doss-Gollin, J., Farnham, D.J., Lall, U., Modi, V.: How unprecedented was the February 2021 Texas cold snap? *Environmental Research Letters* **16**(6), 064056 (2021) <https://doi.org/10.1088/1748-9326/ac0278>

- [59] U.S. had its coldest February in more than 30 years. Accessed on Thu, February 02, 2023 (2021). <https://www.noaa.gov/news/us-had-its-coldest-february-in-more-than-30-years>
- [60] Francis, D., Fonseca, R., Nelli, N., Cuesta, J., Weston, M., Evan, A., Temimi, M.: The Atmospheric Drivers of the Major Saharan Dust Storm in June 2020. *Geophysical Research Letters* **47**(24) (2020) <https://doi.org/10.1029/2020gl090102>
- [61] Yassin, M.F., Almutairi, S.K., Al-Hemoud, A.: Dust storms backward Trajectories' and source identification over Kuwait. *Atmospheric Research* **212**, 158–171 (2018) <https://doi.org/10.1016/j.atmosres.2018.05.020>
- [62] González-Santiago, O., Badillo-Castañeda, C.T., Kahl, J.D.W., Ramírez-Lara, E., Balderas-Renteria, I.: Temporal analysis of pm₁₀ in metropolitan monterrey méxico. *Journal of the Air & Waste Management Association* **61**(5), 573–579 (2011) <https://doi.org/10.3155/1047-3289.61.5.573>
- [63] Martinez, M.A., Caballero, P., Carrillo, O., Mendoza, A., Mejia, G.M.: Chemical characterization and factor analysis of pm_{2.5} in two sites of monterrey mexico. *Journal of the Air & Waste Management Association* **62**(7), 817–827 (2012) <https://doi.org/10.1080/10962247.2012.681421>
- [64] Euphrasie-Clotilde, L., Plocoste, T., Feuillard, T., Velasco-Merino, C., Mateos, D., Toledano, C., Brute, F.-N., Bassette, C., Gobinddass, M.: Assessment of a new detection threshold for pm₁₀ concentrations linked to african dust events in the caribbean basin. *Atmospheric Environment* **224**, 117354 (2020) <https://doi.org/10.1016/j.atmosenv.2020.117354>
- [65] Bozlaker, A., Prospero, J.M., Fraser, M.P., Chellam, S.: Quantifying the Contribution of Long-Range Saharan Dust Transport on Particulate Matter Concentrations in Houston Texas, Using Detailed Elemental Analysis. *Environmental Science & Technology*, 130909083424001 (2013) <https://doi.org/10.1021/es4015663>
- [66] Lenes, J.M., Prospero, J.M., Landing, W.M., Virmani, J.I., Walsh, J.J.: A model of Saharan dust deposition to the eastern Gulf of Mexico. *Marine Chemistry* **134–135**, 1–9 (2012) <https://doi.org/10.1016/j.marchem.2012.02.007>
- [67] Prospero, J.M., Collard, F.-X., Molinié, J., Jeannot, A.: Characterizing the annual cycle of African dust transport to the Caribbean Basin and South America and its impact on the environment and air quality. *Global Biogeochemical Cycles* **28**(7), 757–773 (2014) <https://doi.org/10.1002/2013gb004802>
- [68] Álvaro Clemente and Eduardo Yubero and Jose F. Nicolás and Sandra Caballero and Javier Crespo and Nuria Galindo: Changes in the concentration and composition of urban aerosols during the COVID-19 lockdown. *Environmental Research* **203**, 111788 (2022) <https://doi.org/10.1016/j.envres.2021.111788>

- [69] Probst, P., Annunziato, A., Chiara Proietti, S.P.: 2020 Atlantic hurricane season: a record-breaking season. Joint Research Centre (European Commission) (2021) <https://doi.org/10.2760/00114,JRC123923>
- [70] Beven, J.L.: The 2020 Atlantic Hurricane Season: The Most Active Season on Record. *Weatherwise* **74**(5), 33–43 (2021) <https://doi.org/10.1080/00431672.2021.1953905>
- [71] CONAGUA: Reseña del huracán Hanna del Océano Atlántico (22 al 27 de julio de 2020). Accessed on Tue, March 07, 2023 (2020). <https://smn.conagua.gob.mx/tools/DATA/Ciclones%20Tropicales/Ciclones/2020-Hanna.pdf>
- [72] Bravo, A.H., Sosa, E.R., Sánchez, A.P., Jaimes, P.M., Saavedra, R.M.I.: Impact of wildfires on the air quality of Mexico City 1992-1999. *Environmental Pollution* **117**(2), 243–253 (2002) [https://doi.org/10.1016/S0269-7491\(01\)00277-9](https://doi.org/10.1016/S0269-7491(01)00277-9)
- [73] MODIS Web. Accessed on Wed, March 08, 2023 (2021). https://modis.gsfc.nasa.gov/gallery/individual.php?db_date=2021-03-30
- [74] Henry, R., Norris, G.A., Vedantham, R., Turner, J.R.: Source region identification using kernel smoothing. *Environmental Science Technology* **43**, 4090–7 (2009)
- [75] Petit, J.-E., Favez, O., Albinet, A., Canonaco, F.: A user-friendly tool for comprehensive evaluation of the geographical origins of atmospheric pollution: Wind and trajectory analyses. *Environmental Modelling & Software* **88**, 183–187 (2017) <https://doi.org/10.1016/j.envsoft.2016.11.022>
- [76] Ji, D., Gao, W., Maenhaut, W., He, J., Wang, Z., Li, J., Du, W., Wang, L., Sun, Y., Xin, J., Hu, B., Wang, Y.: Impact of air pollution control measures and regional transport on carbonaceous aerosols in fine particulate matter in urban Beijing China: insights gained from long-term measurement. *Atmospheric Chemistry and Physics* **19**(13), 8569–8590 (2019) <https://doi.org/10.5194/acp-19-8569-2019>
- [77] Yokelson, R.J., Urbanski, S.P., Atlas, E.L., Toohey, D.W., Alvarado, E.C., Crounse, J.D., Wennberg, P.O., Fisher, M.E., Wold, C.E., Campos, T.L., Adachi, K., Buseck, P.R., Hao, W.M.: Emissions from forest fires near Mexico City. *Atmospheric Chemistry and Physics* **7**(21), 5569–5584 (2007) <https://doi.org/10.5194/acp-7-5569-2007>
- [78] Mendoza, A., Garcia, M.R., Vela, P., Lozano, D.F., Allen, D.: Trace Gases and Particulate Matter Emissions from Wildfires and Agricultural Burning in Northeastern Mexico during the 2000 Fire Season. *Journal of the Air & Waste Management Association* **55**(12), 1797–1808 (2005) <https://doi.org/10.1080/10473289.2005.10464778>
- [79] Carbajal, N., Pineda-Martinez, L.F., Vicente, F.B.: Air Quality Deterioration

- of Urban Areas Caused by Wildfires in a Natural Reservoir Forest of Mexico. *Advances in Meteorology* **2015**, 1–13 (2015) <https://doi.org/10.1155/2015/912946>
- [80] Tello-Leal, E., Macías-Hernández, B.A.: Association of environmental and meteorological factors on the spread of COVID-19 in Victoria Mexico, and air quality during the lockdown. *Environmental Research* **196**, 110442 (2021) <https://doi.org/10.1016/j.envres.2020.110442>
 - [81] Bera, B., Bhattacharjee, S., Sengupta, N., Saha, S.: Variation and dispersal of PM10 and PM2.5 during COVID-19 lockdown over Kolkata metropolitan city India investigated through HYSPLIT model. *Geoscience Frontiers* **13**(1), 101291 (2022) <https://doi.org/10.1016/j.gsf.2021.101291>
 - [82] Sulaymon, I.D., Zhang, Y., Hopke, P.K., Guo, S., Ye, F., Sun, J., Zhu, Y., Hu, J.: Using the covid-19 lockdown to identify atmospheric processes and meteorology influences on regional pm2.5 pollution episodes in the beijing-tianjin-hebei, china. *Atmospheric Research* **294**, 106940 (2023) <https://doi.org/10.1016/j.atmosres.2023.106940>
 - [83] Monterrey: Economía, empleo, equidad, calidad de vida, educación, salud y seguridad pública — Data México. Accessed on Mon, January 30, 2023 (2023). <https://datamexico.org/es/profile/geo/monterrey>
 - [84] INEGI: Censo de Población y Vivienda 2020 (Cuestionario Básico y Cuestionario Ampliado). Technical report (2021)
 - [85] Dantas, G., Siciliano, B., França, B.B., Silva, C.M., Arbilla, G.: The impact of COVID-19 partial lockdown on the air quality of the city of Rio de Janeiro Brazil. *Science of The Total Environment* **729**, 139085 (2020) <https://doi.org/10.1016/j.scitotenv.2020.139085>
 - [86] Spohn, T.K., Martin, D., Geever, M., O’Dowd, C.: Effect of COVID-19 lockdown on regional pollution in ireland. *Air Quality Atmosphere & Health* **15**(2), 221–234 (2021) <https://doi.org/10.1007/s11869-021-01098-4>
 - [87] Liu, Y., Zhou, Y., Lu, J.: Exploring the relationship between air pollution and meteorological conditions in China under environmental governance. *Scientific Reports* **10**, 14518 (2020) <https://doi.org/10.1038/s41598-020-71338-7>
 - [88] Bauwens, M., Compernelle, S., Stavrou, T., Müller, J.-F., Gent, J., Eskes, H., Levelt, P.F., A, R., Veefkind, J.P., Vlietinck, J., Yu, H., Zehner, C.: Impact of coronavirus outbreak on NO2 pollution assessed using TROPOMI and OMI observations. *Geophysical Research Letters*, 2020–087978 (2020) <https://doi.org/10.1029/2020GL087978>
 - [89] Monterrey — Air Quality. Accessed on Wed, March 01, 2023 (2021). <https://>

airquality.gsfc.nasa.gov/no2/world/south-and-central-america/monterrey

- [90] Ipiña, A., López-Padilla, G., Retama, A., Piacentini, R.D., Madronich, S.: Ultra-violet Radiation Environment of a Tropical Megacity in Transition: Mexico City 2000-2019. *Environmental Science & Technology* **55**(16), 10946–10956 (2021) <https://doi.org/10.1021/acs.est.0c08515>
- [91] Aguirre-López, M.A., Rodríguez-González, M.A., Soto-Villalobos, R., Gómez-Sánchez, L.E., Benavides-Ríos, Á.G., Benavides-Bravo, F.G., Walle-García, O., Pamanés-Aguilar, M.G.: Statistical Analysis of PM10 Concentration in the Monterrey Metropolitan Area Mexico (2010-2018). *Atmosphere* **13**(2), 297 (2022) <https://doi.org/10.3390/atmos13020297>
- [92] González, L.T., Longoria-Rodríguez, F.E., Sánchez-Domínguez, M., Leyva-Porras, C., Acuña-Askar, K., Kharissov, B.I., Arizpe-Zapata, A., Alfaro-Barbosa, J.M.: Seasonal variation and chemical composition of particulate matter: A study by XPS ICP-AES and sequential microanalysis using Raman with SEM/EDS. *Journal of Environmental Sciences* **74**, 32–49 (2018) <https://doi.org/10.1016/j.jes.2018.02.002>