



The dispersal of compact protoplanetary discs

Giovanni Picogna^{1*} and Barbara Ercolano^{1,2,3}

¹*Universitäts-Sternwarte, Ludwig-Maximilians-Universität München, Scheinerstr. 1, D-81679 München, Germany*

²*Excellence Cluster Origins, Boltzmannstrasse 2, D-85748 Garching bei München, Germany*

³*Max-Planck-Institut für Extraterrestrische Physik, Giessenbachstraße 1, 85748 Garching, Germany*

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

Compact protoplanetary discs are becoming increasingly prominent in observations. Their dispersal pathways may differ substantially from those of extended discs. We aim to quantify the role of the disc outer radius in internal photoevaporation, provide a simple scaling relation for compact discs, and test whether the resulting evolutionary tracks reproduce the observed inside-out clearing of young stellar populations. We performed radiation-hydrodynamic simulations of X-ray-driven photoevaporation for discs with different outer radii, and derived the dependence of the total mass-loss rate on the critical radius. We incorporated this scaling into disc population synthesis models. We find that the surface mass-loss profiles are nearly independent of disc size, but their integrated wind rates are reduced according to the cumulative mass-loss rate distribution. Compact discs disperse more slowly, evolving via inside-out clearing consistent with observational diagnostics. Population synthesis shows that the scaled prescription reproduces both disc fractions and the persistence of highly accreting systems at late times ($t > 4$ Myr) better than unscaled models. Disc dispersal prescriptions must include the dependence on disc outer radius to capture the evolution of compact discs. The proposed scaling provides a simple, physically motivated correction that better predicts the growing observational evidence for compact discs and inside-out dispersal.

Key words: keyword1 – keyword2 – keyword3

1 INTRODUCTION

Protoplanetary discs are the birthplaces of planets, and their lifetimes and dispersal pathways set essential constraints on planet formation models. Most discs dissipate within a few Myr, with only a minority surviving to ~ 10 Myr (Haisch et al. 2001; Fedele et al. 2010). Observations also show that dispersal proceeds preferentially from the inside-out: colour evolution studies of young stellar populations (Ercolano et al. 2011; Koepferl et al. 2013; Ercolano et al. 2015) reveal clear signatures of inner clearing, inconsistent with outside-in erosion. Any successful theoretical model must therefore reproduce this inside-out behaviour.

Mounting evidence points to magnetically driven winds and photoevaporation (EUV, X-ray, FUV) as the dominant agents of angular momentum removal and mass loss, respectively (Ercolano & Pascucci 2017; Lesur et al. 2023; Pascucci et al. 2023). X-ray photoevaporation models in particular have been refined in recent years to yield dispersal prescriptions consistent with several independent observational diagnostics (Ercolano & Owen 2016; Weber et al. 2020; Flaischlen et al. 2021; Rab et al. 2023; Ercolano et al. 2023).

Most existing models, however, assume discs with outer radii of order hundreds of au – well beyond the gravitational radius R_g , defined as the location where the gas sound speed becomes comparable with the Keplerian speed – around 5 au for a Solar mass star, creating a natural “sweet spot” for photoevaporation (Hollenbach et al. 1994; Liffman 2003). Yet compact discs, with outer radii comparable to

or smaller than this scale, are increasingly recognised as common. ALMA surveys show that a substantial fraction of discs are indeed very compact (e.g. Trapman et al. 2025), with critical radii much smaller than typically assumed. Such compact configurations may result from internal evolution (mass and angular momentum loss via winds) or from external influences such as stellar fly-bys or external photoevaporation.

In this Letter, we investigate the dispersal of compact discs by explicitly accounting for their finite outer radii in photoevaporation models. Using radiation-hydrodynamic simulations, we derive the dependence of the total wind mass-loss rate on the critical radius and implement this scaling into one-dimensional viscous evolution and population synthesis models. Our aim is to provide a simple prescription valid in the compact-disc regime and to test whether the resulting evolutionary pathways are consistent with observed disc fractions, accretion rates, and the observationally inferred inside-out dispersal of young stellar populations.

2 METHODS

We performed a series of radiation hydrodynamic simulations following the approach described in Picogna et al. (2019), which we briefly summarise here for completeness.

* E-mail: picogna@usm.lmu.de

2.1 Radiative Transfer

We focused on a $1 M_{\odot}$ star, where the stellar properties were taken from [Siess et al. \(2000\)](#), assuming an age of 1 Myr and metallicity $Z = 0.02$ without convective overshooting. The initial disc structure was based on the gas densities and dust temperatures from the hydrostatic disc models of the DAlessio Irradiated Accretion disc (DIAD) radiative transfer framework ([D'Alessio et al. 1998, 1999, 2001, 2005, 2006](#)). These models provide the best fits to the median spectral energy distribution (SED) observed in Taurus. To compute the thermal structure of the upper layers of the disc, we used the gas photoionisation and dust radiative transfer code MOCASSIN ([Ercolano et al. 2003, 2005, 2008](#)). This code self-consistently solves for heating and cooling processes under thermal equilibrium, allowing us to derive temperature prescriptions up to the maximum penetration depth of X-rays ($\sim 10^{22}$ pp/cm²) for discs irradiated by X-ray and EUV stellar spectra. The resulting temperature prescription depends on the spectral hardness as detailed in ([Ercolano et al. 2021](#)). It relates the local gas temperature to both the column density towards the central star (from 5×10^{20} pp/cm² to 2×10^{22} pp/cm²), and the local ionisation parameter ([Tarter et al. 1969](#)) $\xi = L_X / (nr^2)$, where L_X is the stellar X-ray luminosity, n the local number density, and r the spherical radius.

2.2 Hydrodynamical model

Having the temperature prescription at thermal equilibrium, we then performed hydrodynamical simulations using a modified version of the PLUTO code ([Mignone et al. 2007](#)), as described in [Picogna et al. \(2019\)](#). In this setup, the MOCASSIN-derived prescription was applied at column densities below the X-ray penetration depth ($\sim 2 \times 10^{22}$ pp/cm²), while at larger column densities we assumed perfect thermal coupling between gas and dust, adopting the DIAD models. Each simulation was evolved until the disc structure and wind streamlines reached steady state. We explored 4 different initial critical radii for the circumstellar disc: 10, 50, 100, 200 au, where we exponentially cut off the gas surface densities from the unperturbed disc. We adopted a spherical coordinate system centred on the star. The computational grid is logarithmically spaced in the radial direction with 500 cells, ensuring enhanced resolution in the innermost disc regions where photoevaporation is most effective, while still extending to large radii ($R_{\text{out}} = 300$ au) without excessive computational cost. In the polar direction, the grid is linearly spaced with 250 cells between 0.01 rad and $\pi/2$. The system was evolved for 500 – 1000 orbits at 10 au. A quasi-steady state is typically reached after a few hundred orbital periods, at which point the cumulative mass-loss rate and the gas streamlines in the wind stabilise.

2.3 Disc population synthesis

In order to model the long-term evolution of compact protoplanetary discs, we adopted a 1D viscous evolution code ([Ercolano & Rosotti 2015](#)) where the 1D surface density is evolved according to

$$\frac{\partial \Sigma}{\partial t} = \frac{1}{R} \frac{\partial}{\partial R} \left[3R^{1/2} \frac{\partial}{\partial R} \left(\nu \Sigma R^{1/2} \right) \right] - \dot{\Sigma}_w(R, t), \quad (1)$$

where the first term describes the disc viscous evolution ([Lynden-Bell & Pringle 1974](#)) and the last term the mass loss due to photoevaporation derived from our hydrodynamical models ([Picogna et al.](#)

2021)

$$\dot{\Sigma}_w(R) = \ln(10) \left(\frac{6a \ln(R)^5}{R \ln(10)^6} + \frac{5b \ln(R)^4}{R \ln(10)^5} + \frac{4c \ln(R)^3}{R \ln(10)^4} + \frac{3d \ln(R)^2}{R \ln(10)^3} + \frac{2e \ln(R)}{R \ln(10)^2} + \frac{f}{R \ln(10)} \right) \frac{\dot{M}_w(R)}{2\pi R} [\text{M}_{\odot} \text{au}^{-2} \text{yr}^{-1}], \quad (2)$$

where

$$\dot{M}_w(R) = 10^{a \log R^6 + b \log R^5 + c \log R^4 + d \log R^3 + e \log R^2 + f \log R + g} \quad (3)$$

is the cumulative mass-loss rate, and the parameters a, b, c, d, e, f, g are stellar-mass dependant and given in [Picogna et al. \(2021, Table 2\)](#). The discs are initialized with a self-similar solution (for $\gamma = 1$):

$$\Sigma = \frac{M_0}{2\pi r_1 R} T^{-3/2} \exp \left[-\frac{(R/r_1)}{T} \right], \quad (4)$$

where M_0 is the initial disc mass, $T = t/t_s$ is the dimensionless time with t_s the viscous timescale, and r_1 is the disc characteristic or critical radius.

We performed a parameter space analysis by probing a large set of disc and stellar properties. We assumed a stellar initial mass function following [Kroupa \(2001\)](#), $\xi(m) \propto m^{-\alpha}$, where $\alpha = 1.3 \pm 0.5$ for $0.08 \leq m/M_{\odot} < 0.5$ and $\alpha = 2.3 \pm 0.3$ for $0.5 < m/M_{\odot} < 1$. For the X-ray luminosities we followed the same approach outlined in [Ercolano et al. \(2023\)](#), deriving an X-ray luminosity function in three stellar mass bins from a subsample of the Chandra Orion Ultra-deep Project (COUP, cf. [Getman et al. 2005](#)) as shown in [Kuhn & Hillenbrand \(2019\)](#). For the disc properties, we derive the disc dust mass from observational constraint in young star forming regions ([Pascucci et al. 2016; Ansdell et al. 2017](#)):

$$\log(M_{\text{dust}}[M_{\oplus}]) = (1.2 \pm 0.2) + (1.8 \pm 0.4) \log(M_{\star}[M_{\odot}]) \quad (5)$$

with an associated uncertainty of $\delta = 0.9 \pm 0.1$. We then consider a standard 100 conversion factor between the dust and the gas mass, and derive the critical radius by inverting the observed dependence of the critical radius as a function of the gas mass ([Trapman et al. 2025](#)), $M_{\text{gas}} \propto r_1^{1.7}$. Finally, the disc viscosity is the less constrained parameter, thus we linearly sampled its values between $\alpha = 10^{-4}$ and 10^{-2} . The resulting parameter space sampled is shown in Fig. 1.

3 RESULTS

3.1 Hydrodynamical models

We computed the cumulative mass-loss rate from our hydrodynamical models, as shown in Fig. 2. The four cases exhibit very similar radial behaviours, with $\dot{M}_w(R)$ rising steeply in the inner disc and flattening towards large radii. This outcome is expected, since the incident radiation field at smaller radii is not directly affected by the details of the density distribution further out. The main effect of reducing the critical radius is to truncate the radial extent of the mass-loss distribution (equation 3), thereby lowering the total integrated mass-loss rate. In Fig. 2 we show also that the complex fit of equation 3 can be actually broken in two linear regimes, in order to find a simpler (and physically motivated) way to describe the wind mass-loss, where for radii comparable with the gravitational radius it grows as $\dot{M}_w \propto R^3$, while for larger radii like $\dot{M}_w \propto R^{0.5}$. This shows how important it is to consider the critical radius truncated mass-loss rate for compact discs, where the bulk of the disc fits inside the fast growth regime of the wind mass-loss rate.





Figure 1. Parameter space sampled in the population synthesis. In panel a, stellar mass as a function of X-ray luminosity sampled. In panel b, critical radius as a function of disc mass sampled. The observationally derived fits from Güdel et al. (2007) and Trapman et al. (2025) are overplotted for comparison.

We can now provide a prescription for the long-term evolution of compact discs. The normalization factor to be used is given by substituting in equation 3 the critical radius r_1

$$\dot{M}_{w,\text{norm}} = \dot{M}_w(r_1). \quad (6)$$

The total mass-loss rate for a compact disc will then be:

$$\dot{M}_w = \dot{M}_{w,\text{norm}} \dot{M}_w(M_*) \dot{M}_w(L_X), \quad (7)$$

where the mass-loss rate as a function of stellar mass (for a mean X-ray luminosity, Picogna et al. (2021)) is

$$\dot{M}_w(M_*) = 3.93 \times 10^{-8} \left(\frac{M_*}{M_\odot} \right) \quad (8)$$

and the dependence on the X-ray luminosity (Ercolano et al. 2021) is

$$\begin{aligned} \dot{M}_w(L_X) &= \frac{\dot{M}_w(L_{X,\text{soft}})}{\dot{M}_w(L_{X,\text{soft,mean}})} \\ &= 10^{a_L \left[\exp\left(\frac{(\ln(\log(L_{X,\text{soft}}) - b_L)^2}{c_L})\right) - \exp\left(\frac{(\ln(\log(L_{X,\text{soft,mean}}) - b_L)^2}{c_L})\right) \right]}, \end{aligned} \quad (9)$$

with $a_L = -1.947 \cdot 10^{17}$, $b_L = -1.572 \cdot 10^{-4}$, $c_L = -0.2866$, the soft component of the X-ray luminosity is

$$L_{X,\text{soft}} = 10^{0.95 \log(L_X) + 1.19}, \quad (10)$$

and $L_{X,\text{soft,mean}}$ is the soft component for a star with total X-ray luminosity given by the observational relation between stellar mass and mean X-ray luminosity (Güdel et al. 2007)

$$\log L_{X,\text{mean}} = 1.54 \log(M_*) + 30.31 \quad (11)$$

The consistency of the surface mass loss rate distribution however is reassuring, as it demonstrates that existing mass-loss prescriptions remain valid for compact discs, provided they are appropriately scaled to the total wind rate according to the relation above. In the following section, we normalize the mass-loss rates to the value of the cumulative mass-loss rate at the critical radius.




Figure 2. Cumulative mass loss rate as a function of radius for different critical radii.

3.2 Single disc evolution

We investigated the disc long-term evolution using our 1-D viscous evolution code SPOCK for discs with $r_1 = 5, 20$, and 80 au, $M_d = 0.01 M_\odot$, $\alpha = 10^{-2}$ around a $1 M_\odot$ star with $L_X = 2.29 \cdot 10^{30}$ erg/s, evolving the surface density according to equations 1-3, 6-11. In Figure 3, the left-hand panels show the surface density evolution obtained using our new, radius-scaled mass-loss prescription, while the right-hand panels display the corresponding results using the standard, unscaled prescription. The two approaches lead to strikingly different evolutionary outcomes. With the unscaled prescription, compact discs tend to disperse from the outside-in, gradually eroding their outer edges until the disc vanishes. In contrast, when the radius dependence is included, the discs evolve via inside-out dispersal, with gap opening driving the final dispersal. The difference in global lifetimes is also substantial, and the scaled prescription predicts significantly longer dispersal for the most compact discs. This distinction is not only physically motivated, but also observationally significant. The inside-out dispersal predicted by the scaled prescription is consistent with the colour evolution of young stellar objects (YSOs) observed in large surveys




Figure 3. Surface density evolution for discs with a critical radius of 5 au (top row), 20 au (middle row) and 80 au (bottom row) for the old prescription (right column) and the critical radius dependence (left column). The line colours indicate the different evolutionary stage as explained in the legend on top.

(Koepferl et al. 2013), which strongly disfavour outside-in clearing. The improved treatment of compact discs therefore provides a natural explanation for the observational evidence, reconciling theory and data.

3.3 Population synthesis models

We performed a population synthesis of $N_{\text{disc}} = 10,000$ discs, with initial critical radii and masses drawn from the observationally derived distribution described in Section 2. The resulting population naturally reproduces the observed disc fraction as a function of cluster age (e.g. Mamajek 2009) without the need of tuning the parameters as shown in Figure 4. We scaled the initial disc fraction to be 86%, considering that a fraction of discs would not form due to close binary interaction (see e.g. Haisch et al. 2001). The resulting e-folding time for the critical radius-dependent population synthesis is 2.99 ± 0.12 Myr, which is compatible with the observationally derived one (2.96 ± 0.43 Myr) and considerably larger than the previous prescription (1.64 ± 0.07 Myr). We then compared the accretion rate distributions as a function of age predicted by our population synthesis models with observational data from several stellar clusters, including Taurus, Lupus, Upper Scorpius, Chameleon I, Chameleon II, ρ Ophiuchi, Corona Australis, Orionis Cloud A, Orionis Cloud B, γ Vel, σ Orionis, OriOB1a, OriOB1b (Manara et al. 2023; Thanathibodee et al. 2022). We evaluated the predictions obtained using both the standard (unscaled) prescription and the new, radius-scaled formulation of the photoevaporation-driven mass-loss




Figure 4. Disc fraction as a function of time for the old (gray solid line) and new critical radius prescription (black solid line). The observational data from Mamajek (2009) and their exponential fit are shown with red dots and a red dashed line respectively. The e-folding time is overplotted with vertical dotted lines for each model and the observations.

rate. The two approaches yield systematically different evolutionary trends. As shown in Fig. 5, the unscaled prescription (panel a) results in a more rapid decline of accretion activity, underpredicting the number of systems with high accretion rates at ages beyond ~ 4 Myr. In contrast, the new radius-dependent prescription (panel b) produces a broader distribution of accretion rates and a more extended tail of strongly accreting discs persisting to older ages. This tail is a natural outcome of the weaker dependence of the dispersal timescale on stellar and disc parameters when the critical radius is accounted for explicitly. The resulting accretion-rateage distribution from the scaled model aligns more closely with the observations, which consistently reveal a minority population of long-lived, highly accreting discs in older clusters. Furthermore, the peak in the accretion rate is now shifted from 10^{-9} to 10^{-12} , which is below our current detection threshold (Thanathibodee et al. 2023) but it could be an interesting observable for future campaigns. This improved agreement suggests that incorporating the radius dependence into the photoevaporation prescription captures an essential physical ingredient missing from the standard approach, namely, the coupling between disc structure and the location of the flow launching region, which regulates both the efficiency and timescale of disc dispersal.

4 CONCLUSIONS

We have shown that compact discs, with a critical radius comparable with the gravitational radius for photoevaporation, follow the same local mass-loss profile as extended discs, but their integrated wind rates are reduced according to the previously found cumulative mass-loss rate distribution, given in equations 6, 7, 8, 9, 10. Incorporating this scaling into viscous evolution models leads to longer disc lifetimes and inside-out dispersal, in agreement with observational constraints from colour-colour diagrams of young stellar populations. Population synthesis demonstrates that the corrected prescription reproduces both observed disc fractions and the presence of highly accreting discs at late ages ($t > 4$ Myr). The dispersal of compact discs cannot be captured by standard unscaled models, and radius-dependent photoevaporation prescriptions are essential to connect disc evolution theory with observations.




Figure 5. Mass accretion rate as a function of time for the old prescription (panel a) and the critical radius criteria (panel b). The corner plots show the density distribution in the accretion rates and disc age, and the difference as a gray shadow region. The disc observed properties (Manara et al. 2023; Thanathibodee et al. 2022) are overplotted as box plots.

ACKNOWLEDGEMENTS

We acknowledge the support of the DFG (German Research Foundation), Research Unit "Transition discs" - 325594231 and of the Excellence Cluster ORIGINS - EXC-2094 - 390783311. We acknowledge support from the Max Planck Society.

DATA AVAILABILITY

The data underlying this article will be shared on reasonable request to the corresponding author.

REFERENCES

- Ansdell M., Williams J. P., Manara C. F., Miotello A., Facchini S., van der Marel N., Testi L., van Dishoeck E. F., 2017, *AJ*, **153**, 240
- D'Alessio P., Cantó J., Calvet N., Lizano S., 1998, *ApJ*, **500**, 411
- D'Alessio P., Calvet N., Hartmann L., Lizano S., Cantó J., 1999, *ApJ*, **527**, 893
- D'Alessio P., Calvet N., Hartmann L., 2001, *ApJ*, **553**, 321
- D'Alessio P., et al., 2005, *ApJ*, **621**, 461
- D'Alessio P., Calvet N., Hartmann L., Franco-Hernández R., Servín H., 2006, *ApJ*, **638**, 314
- Ercolano B., Owen J. E., 2016, *MNRAS*, **460**, 3472
- Ercolano B., Pascucci I., 2017, *Royal Society Open Science*, **4**, 170114
- Ercolano B., Rosotti G., 2015, *MNRAS*, **450**, 3008
- Ercolano B., Barlow M. J., Storey P. J., Liu X. W., 2003, *MNRAS*, **340**, 1136
- Ercolano B., Barlow M. J., Storey P. J., 2005, *MNRAS*, **362**, 1038
- Ercolano B., Young P. R., Drake J. J., Raymond J. C., 2008, *ApJS*, **175**, 534
- Ercolano B., Clarke C. J., Hall A. C., 2011, *MNRAS*, **410**, 671
- Ercolano B., Koepferl C., Owen J., Robitaille T., 2015, *MNRAS*, **452**, 3689
- Ercolano B., Picogna G., Monsch K., Drake J. J., Preibisch T., 2021, *MNRAS*, **508**, 1675
- Ercolano B., Picogna G., Monsch K., 2023, *MNRAS*, **526**, L105
- Fedele D., van den Ancker M. E., Henning T., Jayawardhana R., Oliveira J. M., 2010, *A&A*, **510**, A72
- Flaischlen S., Preibisch T., Manara C. F., Ercolano B., 2021, *A&A*, **648**, A121
- Getman K. V., et al., 2005, *ApJS*, **160**, 319
- Güdel M., et al., 2007, *A&A*, **468**, 353
- Haisch Jr. K. E., Lada E. A., Lada C. J., 2001, *ApJ*, **553**, L153
- Hollenbach D., Johnstone D., Lizano S., Shu F., 1994, *ApJ*, **428**, 654
- Koepferl C. M., Ercolano B., Dale J., Teixeira P. S., Ratzka T., Spezzi L., 2013, *MNRAS*, **428**, 3327
- Kroupa P., 2001, *MNRAS*, **322**, 231
- Kuhn M. A., Hillenbrand L. A., 2019, *ApJ*, **883**, 117
- Lesur G., et al., 2023, in Inutsuka S., Aikawa Y., Muto T., Tomida K., Tamura M., eds, *Astronomical Society of the Pacific Conference Series Vol. 534, Protostars and Planets VII*. p. 465 ([arXiv:2203.09821](https://arxiv.org/abs/2203.09821)), [doi:10.48550/arXiv.2203.09821](https://doi.org/10.48550/arXiv.2203.09821)
- Liffman K., 2003, *Publ. Astron. Soc. Australia*, **20**, 337
- Lynden-Bell D., Pringle J. E., 1974, *MNRAS*, **168**, 603
- Mamajek E. E., 2009, in Usuda T., Tamura M., Ishii M., eds, *American In-*

- stitute of Physics Conference Series Vol. 1158, Exoplanets and Disks: Their Formation and Diversity. AIP, pp 3–10 ([arXiv:0906.5011](#)), [doi:10.1063/1.3215910](#)
- Manara C. F., Ansdell M., Rosotti G. P., Hughes A. M., Armitage P. J., Lodato G., Williams J. P., 2023, in Inutsuka S., Aikawa Y., Muto T., Tomida K., Tamura M., eds, Astronomical Society of the Pacific Conference Series Vol. 534, Protostars and Planets VII. p. 539 ([arXiv:2203.09930](#)), [doi:10.48550/arXiv.2203.09930](#)
- Mignone A., Bodo G., Massaglia S., Matsakos T., Tesileanu O., Zanni C., Ferrari A., 2007, *ApJS*, **170**, 228
- Pascucci I., et al., 2016, *ApJ*, **831**, 125
- Pascucci I., Cabrit S., Edwards S., Gorti U., Gressel O., Suzuki T. K., 2023, in Inutsuka S., Aikawa Y., Muto T., Tomida K., Tamura M., eds, Astronomical Society of the Pacific Conference Series Vol. 534, Protostars and Planets VII. p. 567 ([arXiv:2203.10068](#)), [doi:10.48550/arXiv.2203.10068](#)
- Picogna G., Ercolano B., Owen J. E., Weber M. L., 2019, *MNRAS*, **487**, 691
- Picogna G., Ercolano B., Espaillat C. C., 2021, *MNRAS*, **508**, 3611
- Rab C., Weber M. L., Picogna G., Ercolano B., Owen J. E., 2023, *ApJ*, **955**, L11
- Siess L., Dufour E., Forestini M., 2000, *A&A*, **358**, 593
- Tarter C. B., Tucker W. H., Salpeter E. E., 1969, *ApJ*, **156**, 943
- Thanathibodee T., Calvet N., Hernandez J., Mauco K., Briceno C., 2022, VizieR Online Data Catalog: A census of the low accretors. I. The catalog (Thanathibodee+, 2022), VizieR On-line Data Catalog: J/AJ/163/74. Originally published in: 2022AJ....163...74T, [doi:10.26093/cds/vizier.51630074](#)
- Thanathibodee T., Molina B., Serna J., Calvet N., Hernández J., Muzerolle J., Franco-Hernández R., 2023, *ApJ*, **944**, 90
- Trapman L., et al., 2025, *The Astrophysical Journal*, 989, 5
- Weber M. L., Ercolano B., Picogna G., Hartmann L., Rodenkirch P. J., 2020, *MNRAS*, **496**, 223

This paper has been typeset from a $\text{\LaTeX}$ file prepared by the author.