

Non-zero Constraints in PDEs and Applications to Hybrid Inverse Problems

Giovanni S. Alberti

Machine Learning Genoa Center, Department of Mathematics, University of Genoa

International Zoom Inverse Problems seminar, 9 October 2025

Internal data in quantitative hybrid imaging problems

- Hybrid conductivity imaging [Widlak, Scherzer, 2012]

$$\begin{cases} -\operatorname{div}(a \nabla u_i) = 0 & \text{in } \Omega, \\ u_i = \varphi_i & \text{on } \partial\Omega. \end{cases}$$

$$u_i(x) \quad \text{or} \quad a(x) \nabla u_i(x) \quad \text{or} \quad a(x) |\nabla u_i|^2(x) \quad \xrightarrow{?} \quad a$$

- Quantitative thermoacoustic tomography [Bal et al., 2011, Ammari et al., 2013]

$$\begin{cases} \Delta u_i + (\omega^2 + i\omega\sigma) u_i = 0 & \text{in } \Omega, \\ u_i = \varphi_i & \text{on } \partial\Omega. \end{cases}$$

$$\sigma(x) |u_i|^2(x) \quad \xrightarrow{?} \quad \sigma$$

- MREIT [Seo et al., 2012, Bal and Guo, 2013]

$$\begin{cases} \operatorname{curl} E^i = i\omega H^i & \text{in } \Omega, \\ \operatorname{curl} H^i = -i(\omega\varepsilon + i\sigma) E^i & \text{in } \Omega, \\ E^i \times \nu = \varphi_i \times \nu & \text{on } \partial\Omega. \end{cases}$$

$$H^i(x) \quad \xrightarrow{?} \quad \varepsilon, \sigma$$

Internal data in quantitative hybrid imaging problems

- Hybrid conductivity imaging [Widlak, Scherzer, 2012], quantitative photoacoustic tomography

$$\begin{cases} -\operatorname{div}(\textcolor{red}{a} \nabla u_i) + \mu u_i = 0 & \text{in } \Omega, \\ u_i = \varphi_i & \text{on } \partial\Omega. \end{cases}$$

$$u_i(x) \quad \text{or} \quad a(x) \nabla u_i(x) \quad \text{or} \quad a(x) |\nabla u_i|^2(x) \quad \text{or} \quad \mu(x) u_i(x) \quad \xrightarrow{?} \quad \textcolor{red}{a}, \mu$$

- Quantitative thermoacoustic tomography [Bal et al., 2011, Ammari et al., 2013]

$$\begin{cases} \Delta u_i + (\omega^2 + i\omega \textcolor{red}{\sigma}) u_i = 0 & \text{in } \Omega, \\ u_i = \varphi_i & \text{on } \partial\Omega. \end{cases}$$

$$\sigma(x) |u_i|^2(x) \quad \xrightarrow{?} \quad \textcolor{red}{\sigma}$$

- MREIT [Seo et al., 2012, Bal and Guo, 2013]

$$\begin{cases} \operatorname{curl} E^i = i\omega H^i & \text{in } \Omega, \\ \operatorname{curl} H^i = -i(\omega \textcolor{red}{\varepsilon} + i\textcolor{red}{\sigma}) E^i & \text{in } \Omega, \\ E^i \times \nu = \varphi_i \times \nu & \text{on } \partial\Omega. \end{cases}$$

$$H^i(x) \quad \xrightarrow{?} \quad \textcolor{red}{\varepsilon}, \textcolor{red}{\sigma}$$

Internal data in quantitative hybrid imaging problems

- Hybrid conductivity imaging [Widlak, Scherzer, 2012], quantitative photoacoustic tomography

$$\begin{cases} -\operatorname{div}(\mathbf{a} \nabla u_i) + \mu u_i = 0 & \text{in } \Omega, \\ u_i = \varphi_i & \text{on } \partial\Omega. \end{cases}$$

$$u_i(x) \quad \text{or} \quad a(x) \nabla u_i(x) \quad \text{or} \quad a(x) |\nabla u_i|^2(x) \quad \text{or} \quad \mu(x) u_i(x) \quad \xrightarrow{?} \quad \mathbf{a}, \mu$$

- Quantitative thermoacoustic tomography [Bal et al., 2011, Ammari et al., 2013]

$$\begin{cases} \Delta u_i + (\omega^2 + i\omega\sigma) u_i = 0 & \text{in } \Omega, \\ u_i = \varphi_i & \text{on } \partial\Omega. \end{cases}$$

$$\sigma(x) |u_i|^2(x) \quad \xrightarrow{?} \quad \sigma$$

- MREIT [Seo et al., 2012, Bal and Guo, 2013]

$$\begin{cases} \operatorname{curl} E^i = i\omega H^i & \text{in } \Omega, \\ \operatorname{curl} H^i = -i(\omega\epsilon + i\sigma) E^i & \text{in } \Omega, \\ E^i \times \nu = \varphi_i \times \nu & \text{on } \partial\Omega. \end{cases}$$

$$H^i(x) \quad \xrightarrow{?} \quad \epsilon, \sigma$$

Internal data in quantitative hybrid imaging problems

- Hybrid conductivity imaging [Widlak, Scherzer, 2012], quantitative photoacoustic tomography

$$\begin{cases} -\operatorname{div}(\textcolor{red}{a} \nabla u_i) + \mu u_i = 0 & \text{in } \Omega, \\ u_i = \varphi_i & \text{on } \partial\Omega. \end{cases}$$

$$u_i(x) \quad \text{or} \quad a(x) \nabla u_i(x) \quad \text{or} \quad a(x) |\nabla u_i|^2(x) \quad \text{or} \quad \mu(x) u_i(x) \quad \xrightarrow{?} \quad \textcolor{red}{a}, \mu$$

- Quantitative thermoacoustic tomography [Bal et al., 2011, Ammari et al., 2013]

$$\begin{cases} \Delta u_i + (\omega^2 + i\omega \textcolor{red}{\sigma}) u_i = 0 & \text{in } \Omega, \\ u_i = \varphi_i & \text{on } \partial\Omega. \end{cases}$$

$$\sigma(x) |u_i|^2(x) \quad \xrightarrow{?} \quad \textcolor{red}{\sigma}$$

- MREIT [Seo et al., 2012, Bal and Guo, 2013]

$$\begin{cases} \operatorname{curl} E^i = i\omega H^i & \text{in } \Omega, \\ \operatorname{curl} H^i = -i(\omega \varepsilon + i\textcolor{red}{\sigma}) E^i & \text{in } \Omega, \\ E^i \times \nu = \varphi_i \times \nu & \text{on } \partial\Omega. \end{cases}$$

$$H^i(x) \quad \xrightarrow{?} \quad \varepsilon, \textcolor{red}{\sigma}$$

Non-vanishing gradients and Jacobians

- Consider for simplicity the hybrid conductivity problem with internal data ∇u and unknown a :

$$\begin{cases} -\operatorname{div}(a \nabla u) = 0 & \text{in } \Omega, \\ u = \varphi & \text{on } \partial\Omega. \end{cases}$$

- With 1 measurement:

$$\nabla a \cdot \nabla u = -a \Delta u \quad \implies \quad \nabla(\log a) \cdot \nabla u = -\Delta u$$

This equation may be solved in a if a is known on $\partial\Omega$ and if

$$\nabla u(x) \neq 0, \quad x \in \Omega.$$

- With d measurements:

$$\begin{aligned} \nabla(\log a) \cdot (\nabla u_1, \dots, \nabla u_d) &= -(\Delta u_1, \dots, \Delta u_d) \\ \implies \nabla(\log a) &= -(\Delta u_1, \dots, \Delta u_d)(\nabla u_1, \dots, \nabla u_d)^{-1} \end{aligned}$$

This equation may be solved in a if a is known at $x_0 \in \partial\Omega$ and

$$\det [\nabla u_1(x) \quad \dots \quad \nabla u_d(x)] \neq 0, \quad x \in \Omega.$$

Non-vanishing gradients and Jacobians

- Consider for simplicity the hybrid conductivity problem with internal data ∇u and unknown a :

$$\begin{cases} -\operatorname{div}(a \nabla u) = 0 & \text{in } \Omega, \\ u = \varphi & \text{on } \partial\Omega. \end{cases}$$

- With 1 measurement:

$$\nabla a \cdot \nabla u = -a \Delta u \quad \implies \quad \nabla(\log a) \cdot \nabla u = -\Delta u$$

This equation may be solved in a if a is known on $\partial\Omega$ and if

$$\nabla u(x) \neq 0, \quad x \in \Omega.$$

- With d measurements:

$$\begin{aligned} \nabla(\log a) \cdot (\nabla u_1, \dots, \nabla u_d) &= -(\Delta u_1, \dots, \Delta u_d) \\ \implies \nabla(\log a) &= -(\Delta u_1, \dots, \Delta u_d)(\nabla u_1, \dots, \nabla u_d)^{-1} \end{aligned}$$

This equation may be solved in a if a is known at $x_0 \in \partial\Omega$ and

$$\det [\nabla u_1(x) \quad \dots \quad \nabla u_d(x)] \neq 0, \quad x \in \Omega.$$

Non-vanishing gradients and Jacobians

- Consider for simplicity the hybrid conductivity problem with internal data ∇u and unknown a :

$$\begin{cases} -\operatorname{div}(a \nabla u) = 0 & \text{in } \Omega, \\ u = \varphi & \text{on } \partial\Omega. \end{cases}$$

- With 1 measurement:

$$\nabla a \cdot \nabla u = -a \Delta u \quad \implies \quad \nabla(\log a) \cdot \nabla u = -\Delta u$$

This equation may be solved in a if a is known on $\partial\Omega$ and if

$$\nabla u(x) \neq 0, \quad x \in \Omega.$$

- With d measurements:

$$\begin{aligned} \nabla(\log a) \cdot (\nabla u_1, \dots, \nabla u_d) &= -(\Delta u_1, \dots, \Delta u_d) \\ \implies \nabla(\log a) &= -(\Delta u_1, \dots, \Delta u_d)(\nabla u_1, \dots, \nabla u_d)^{-1} \end{aligned}$$

This equation may be solved in a if a is known at $x_0 \in \partial\Omega$ and

$$\det [\nabla u_1(x) \quad \dots \quad \nabla u_d(x)] \neq 0, \quad x \in \Omega.$$

Non-vanishing gradients and Jacobians

- Consider for simplicity the hybrid conductivity problem with internal data ∇u and unknown a :

$$\begin{cases} -\operatorname{div}(a \nabla u) = 0 & \text{in } \Omega, \\ u = \varphi & \text{on } \partial\Omega. \end{cases}$$

- With 1 measurement:

$$\nabla a \cdot \nabla u = -a \Delta u \quad \implies \quad \nabla(\log a) \cdot \nabla u = -\Delta u$$

This equation may be solved in a if a is known on $\partial\Omega$ and if

$$\nabla u(x) \neq 0, \quad x \in \Omega.$$

- With d measurements:

$$\begin{aligned} \nabla(\log a) \cdot (\nabla u_1, \dots, \nabla u_d) &= -(\Delta u_1, \dots, \Delta u_d) \\ \implies \nabla(\log a) &= -(\Delta u_1, \dots, \Delta u_d)(\nabla u_1, \dots, \nabla u_d)^{-1} \end{aligned}$$

This equation may be solved in a if a is known at $x_0 \in \partial\Omega$ and

$$\det [\nabla u_1(x) \quad \dots \quad \nabla u_d(x)] \neq 0, \quad x \in \Omega.$$

Non-vanishing gradients and Jacobians

- Consider for simplicity the hybrid conductivity problem with internal data ∇u and unknown a :

$$\begin{cases} -\operatorname{div}(a \nabla u) = 0 & \text{in } \Omega, \\ u = \varphi & \text{on } \partial\Omega. \end{cases}$$

- With 1 measurement:

$$\nabla a \cdot \nabla u = -a \Delta u \quad \implies \quad \nabla(\log a) \cdot \nabla u = -\Delta u$$

This equation may be solved in a if a is known on $\partial\Omega$ and if

$$\nabla u(x) \neq 0, \quad x \in \Omega.$$

- With d measurements:

$$\begin{aligned} \nabla(\log a) \cdot (\nabla u_1, \dots, \nabla u_d) &= -(\Delta u_1, \dots, \Delta u_d) \\ \implies \nabla(\log a) &= -(\Delta u_1, \dots, \Delta u_d)(\nabla u_1, \dots, \nabla u_d)^{-1} \end{aligned}$$

This equation may be solved in a if a is known at $x_0 \in \partial\Omega$ and

$$\det [\nabla u_1(x) \quad \dots \quad \nabla u_d(x)] \neq 0, \quad x \in \Omega.$$

Main question

Is it possible to find suitable boundary values φ_i so that the corresponding solutions u_i satisfy certain non-zero constraints, such as a **non-vanishing Jacobian**

$$\det [\nabla u_1(x) \quad \cdots \quad \nabla u_d(x)] \neq 0?$$

- ▶ In other words, this ensures that the internal data are **rich enough**
- ▶ Ideally, we would like to construct the φ_i s **independently of the unknown coefficients**

Main question

Is it possible to find suitable boundary values φ_i so that the corresponding solutions u_i satisfy certain non-zero constraints, such as a **non-vanishing Jacobian**

$$\det [\nabla u_1(x) \quad \cdots \quad \nabla u_d(x)] \neq 0?$$

- ▶ In other words, this ensures that the internal data are **rich enough**
- ▶ Ideally, we would like to construct the φ_i s **independently of the unknown coefficients**

Main question

Is it possible to find suitable boundary values φ_i so that the corresponding solutions u_i satisfy certain non-zero constraints, such as a **non-vanishing Jacobian**

$$\det [\nabla u_1(x) \quad \cdots \quad \nabla u_d(x)] \neq 0?$$

- ▶ In other words, this ensures that the internal data are **rich enough**
- ▶ Ideally, we would like to construct the φ_i s **independently of the unknown coefficients**

Outline of the talk

- 1 The Radó-Kneser-Choquet theorem
- 2 Runge approximation & Whitney embedding
- 3 Using random boundary conditions

Outline of the talk

- 1 The Radó-Kneser-Choquet theorem
- 2 Runge approximation & Whitney embedding
- 3 Using random boundary conditions

The Radó-Kneser-Choquet theorem


Theorem (Alessandrini Magnanini 1994, Bauman et al. 2000, Alessandrini Nesi 2015)

Let $\Omega \subseteq \mathbb{R}^2$ be bounded and convex and $a \in C^{0,\alpha}(\overline{\Omega}; \mathbb{R}^{2 \times 2})$ be uniformly elliptic. Let $u_i \in H^1(\Omega)$ solve

$$-\operatorname{div}(a \nabla u_i) = 0 \quad \text{in } \Omega, \quad u_i = x_i \quad \text{on } \partial\Omega.$$

Then

$$\det [\nabla u_1(x) \quad \nabla u_2(x)] \neq 0, \quad x \in \Omega.$$



- ▶ $\alpha \nabla u_1(x_0) + \beta \nabla u_2(x_0) = 0$
- ▶ Set $v(x) = \alpha u_1(x) + \beta u_2(x)$:
- ▶ $\nabla v(x_0) = 0$
- ▶ Thus, v has a saddle point in x_0

The Radó-Kneser-Choquet theorem


Theorem (Alessandrini Magnanini 1994, Bauman et al. 2000, Alessandrini Nesi 2015)

Let $\Omega \subseteq \mathbb{R}^2$ be bounded and convex and $a \in C^{0,\alpha}(\overline{\Omega}; \mathbb{R}^{2 \times 2})$ be uniformly elliptic. Let $u_i \in H^1(\Omega)$ solve

$$-\operatorname{div}(a \nabla u_i) = 0 \quad \text{in } \Omega, \quad u_i = x_i \quad \text{on } \partial\Omega.$$

Then

$$\det [\nabla u_1(x) \quad \nabla u_2(x)] \neq 0, \quad x \in \Omega.$$



- ▶ $\alpha \nabla u_1(x_0) + \beta \nabla u_2(x_0) = 0$
- ▶ Set $v(x) = \alpha u_1(x) + \beta u_2(x)$:
- ▶ $\nabla v(x_0) = 0$
- ▶ Thus, v has a saddle point in x_0

The failure in 3D and for other elliptic PDEs

- ▶ In 3D, the above result fails. Counterexamples by Laugesen 1996, Briane et al 2004 and Capdeboscq 2015: it is not possible to find $(\varphi^1, \varphi^2, \varphi^3)$ independently of a so that

$$\det [\nabla u_1(x) \quad \nabla u_2(x) \quad \nabla u_3(x)] \neq 0, \quad x \in \Omega.$$

- ▶ This result fails for Helmholtz type problems or for eigenvalue problems

$$\operatorname{div}(a \nabla u) + k^2 q u = 0$$

since solutions oscillate.

The failure in 3D and for other elliptic PDEs

- ▶ In 3D, the above result fails. Counterexamples by Laugesen 1996, Briane et al 2004 and Capdeboscq 2015: it is not possible to find $(\varphi^1, \varphi^2, \varphi^3)$ independently of a so that

$$\det [\nabla u_1(x) \quad \nabla u_2(x) \quad \nabla u_3(x)] \neq 0, \quad x \in \Omega.$$

- ▶ This result fails for Helmholtz type problems or for eigenvalue problems

$$\operatorname{div}(a \nabla u) + k^2 q u = 0$$

since solutions oscillate.

Critical points in 3D

What about critical points: can we find φ independently of a so that

$$\nabla u(x) \neq 0, \quad x \in \Omega?$$

Theorem (GSA, Bal, Di Cristo, ARMA 2017)

Let $\Omega \subseteq \mathbb{R}^3$ be a bounded Lipschitz domain. Take $\varphi \in C(\partial X) \cap H^{\frac{1}{2}}(\partial X)$. There exists a (nonempty open set of) $a \in C^\infty(\overline{X})$ such that the solution $u \in H^1(X)$ to

$$\begin{cases} -\operatorname{div}(a \nabla u) = 0 & \text{in } \Omega, \\ u = \varphi & \text{on } \partial\Omega, \end{cases}$$

has a critical point in Ω , namely $\nabla u(x) = 0$ for some $x \in \Omega$.

Critical points in 3D

What about critical points: can we find φ independently of a so that


$$\nabla u(x) \neq 0, \quad x \in \Omega?$$

Theorem (GSA, Bal, Di Cristo, ARMA 2017)

Let $\Omega \subseteq \mathbb{R}^3$ be a bounded Lipschitz domain. Take $\varphi \in C(\partial X) \cap H^{\frac{1}{2}}(\partial X)$. There exists a (nonempty open set of) $a \in C^\infty(\overline{X})$ such that the solution $u \in H^1(X)$ to

$$\begin{cases} -\operatorname{div}(a \nabla u) = 0 & \text{in } \Omega, \\ u = \varphi & \text{on } \partial\Omega, \end{cases}$$

has a critical point in Ω , namely $\nabla u(x) = 0$ for some $x \in \Omega$.



Alternative approaches

- Complex geometrical optics solutions [Bal and Uhlmann, IP 2010]

$$\cos^2 x + \sin^2 x = 1 \neq 0$$

- Use multiple frequencies [GSA, Commun. PDE 2015]

$$\cos^2(k_1 x) + \cos^2(k_2 x) \neq 0$$

Also with Neumann eigenfunctions (bypassing the “hot spots” conjecture):

$$\sum_{k=1}^K |\nabla e_k(x)|^2 \geq c, \quad x \in \overline{\Omega'}$$

[GSA, Barnes, Jambhale and Nickl, Math. Stat. Learn. 2025]

- Runge approximation & Whitney embedding
- Random boundary conditions

Alternative approaches

- Complex geometrical optics solutions [Bal and Uhlmann, IP 2010]

$$\cos^2 x + \sin^2 x = 1 \neq 0$$

- Use multiple frequencies [GSA, Commun. PDE 2015]

$$\cos^2(k_1 x) + \cos^2(k_2 x) \neq 0$$

Also with Neumann eigenfunctions (bypassing the “hot spots” conjecture):

$$\sum_{k=1}^K |\nabla e_k(x)|^2 \geq c, \quad x \in \overline{\Omega'}$$

[GSA, Barnes, Jambhale and Nickl, Math. Stat. Learn. 2025]

- Runge approximation & Whitney embedding
- Random boundary conditions

Alternative approaches

- Complex geometrical optics solutions [Bal and Uhlmann, IP 2010]

$$\cos^2 x + \sin^2 x = 1 \neq 0$$

- Use multiple frequencies [GSA, Commun. PDE 2015]

$$\cos^2(k_1 x) + \cos^2(k_2 x) \neq 0$$

Also with Neumann eigenfunctions (bypassing the “hot spots” conjecture):

$$\sum_{k=1}^K |\nabla e_k(x)|^2 \geq c, \quad x \in \overline{\Omega'}$$

[GSA, Barnes, Jambhale and Nickl, Math. Stat. Learn. 2025]

- Runge approximation & Whitney embedding
- Random boundary conditions

Alternative approaches

- Complex geometrical optics solutions [Bal and Uhlmann, IP 2010]

$$\cos^2 x + \sin^2 x = 1 \neq 0$$

- Use multiple frequencies [GSA, Commun. PDE 2015]

$$\cos^2(k_1 x) + \cos^2(k_2 x) \neq 0$$

Also with Neumann eigenfunctions (bypassing the “hot spots” conjecture):

$$\sum_{k=1}^K |\nabla e_k(x)|^2 \geq c, \quad x \in \overline{\Omega'}$$

[GSA, Barnes, Jambhale and Nickl, Math. Stat. Learn. 2025]

- Runge approximation & Whitney embedding
- Random boundary conditions

Outline of the talk

- 1 The Radó-Kneser-Choquet theorem
- 2 Runge approximation & Whitney embedding
- 3 Using random boundary conditions

The model problem

- ▶ Let $\Omega \subseteq \mathbb{R}^d$, $d \geq 2$ be a smooth bounded domain. Consider the elliptic PDE

$$Lu := -\operatorname{div}(a\nabla u) + b \cdot \nabla u + cu = 0 \quad \text{in } \Omega,$$

with a , b and c smooth enough so that $u \in C^{1,\alpha}$ and the **unique continuation property** (UCP) holds

- ▶ Consider, for simplicity, the **non-vanishing Jacobian** constraint: look for φ_i such that

$$\det [\nabla u_1 \quad \cdots \quad \nabla u_d](x) \neq 0$$

possibly locally, where

$$\begin{cases} Lu_i = 0 & \text{in } \Omega, \\ u_i = \varphi_i & \text{on } \partial\Omega. \end{cases}$$

The model problem

- ▶ Let $\Omega \subseteq \mathbb{R}^d$, $d \geq 2$ be a smooth bounded domain. Consider the elliptic PDE

$$Lu := -\operatorname{div}(a \nabla u) + b \cdot \nabla u + cu = 0 \quad \text{in } \Omega,$$

with a , b and c smooth enough so that $u \in C^{1,\alpha}$ and the **unique continuation property** (UCP) holds


- ▶ Consider, for simplicity, the **non-vanishing Jacobian** constraint: look for φ_i such that

$$\det [\nabla u_1 \quad \cdots \quad \nabla u_d](x) \neq 0$$

possibly locally, where

$$\begin{cases} Lu_i = 0 & \text{in } \Omega, \\ u_i = \varphi_i & \text{on } \partial\Omega. \end{cases}$$

Main tool: the Runge Approximation [Lax 1956]



- ▶ Let $\Omega' \subseteq \Omega$ be simply connected and $v \in H^1(\Omega')$ be a local solution:

$$Lv = 0 \quad \text{in } \Omega'.$$

In general, v cannot be extended to a global solution u , BUT:

- ▶ Runge approximation: there exist global solutions u_n to

$$Lu_n = 0 \quad \text{in } \Omega$$


such that

$$\|u_n|_{\Omega'} - v\|_{L^2(\Omega')} \rightarrow 0.$$

- ▶ By elliptic regularity, we get for $\Omega'' \Subset \Omega'$:

$$\|u_n|_{\Omega'} - v\|_{C^1(\overline{\Omega''})} \rightarrow 0.$$

Main tool: the Runge Approximation [Lax 1956]



- ▶ Let $\Omega' \subseteq \Omega$ be simply connected and $v \in H^1(\Omega')$ be a local solution:

$$Lv = 0 \quad \text{in } \Omega'.$$

In general, v cannot be extended to a global solution u , BUT:

- ▶ Runge approximation: there exist global solutions u_n to

$$Lu_n = 0 \quad \text{in } \Omega$$


such that

$$\|u_n|_{\Omega'} - v\|_{L^2(\Omega')} \rightarrow 0.$$

- ▶ By elliptic regularity, we get for $\Omega'' \Subset \Omega'$:

$$\|u_n|_{\Omega'} - v\|_{C^1(\overline{\Omega''})} \rightarrow 0.$$

Main tool: the Runge Approximation [Lax 1956]



- ▶ Let $\Omega' \subseteq \Omega$ be simply connected and $v \in H^1(\Omega')$ be a local solution:

$$Lv = 0 \quad \text{in } \Omega'.$$

In general, v cannot be extended to a global solution u , BUT:

- ▶ Runge approximation: there exist global solutions u_n to

$$Lu_n = 0 \quad \text{in } \Omega$$


such that

$$\|u_n|_{\Omega'} - v\|_{L^2(\Omega')} \rightarrow 0.$$

- ▶ By elliptic regularity, we get for $\Omega'' \Subset \Omega'$:

$$\|u_n|_{\Omega'} - v\|_{C^1(\overline{\Omega''})} \rightarrow 0.$$

Main tool: the Runge Approximation [Lax 1956]



- ▶ Let $\Omega' \subseteq \Omega$ be simply connected and $v \in H^1(\Omega')$ be a local solution:

$$Lv = 0 \quad \text{in } \Omega'.$$

In general, v cannot be extended to a global solution u , BUT:

- ▶ Runge approximation: there exist global solutions u_n to

$$Lu_n = 0 \quad \text{in } \Omega$$


such that

$$\|u_n|_{\Omega'} - v\|_{L^2(\Omega')} \rightarrow 0.$$

- ▶ By elliptic regularity, we get for $\Omega'' \Subset \Omega'$:

$$\|u_n|_{\Omega'} - v\|_{C^1(\overline{\Omega''})} \rightarrow 0.$$

Main tool: the Runge Approximation [Lax 1956]



- ▶ Let $\Omega' \subseteq \Omega$ be simply connected and $v \in H^1(\Omega')$ be a local solution:

$$Lv = 0 \quad \text{in } \Omega'.$$

In general, v cannot be extended to a global solution u , BUT:

- ▶ Runge approximation: there exist global solutions u_n to

$$Lu_n = 0 \quad \text{in } \Omega$$

such that

$$\|u_n|_{\Omega'} - v\|_{L^2(\Omega')} \rightarrow 0.$$

- ▶ By elliptic regularity, we get for $\Omega'' \Subset \Omega'$:

$$\|u_n|_{\Omega'} - v\|_{C^1(\overline{\Omega''})} \rightarrow 0.$$

The Runge approximation and non-zero constraints [Bal and Uhlmann, CPAM 2013]

1. Fix $x_0 \in \overline{\Omega}$ and $r > 0$. Consider local solutions $v_i^0 = x_i$:

$$-\operatorname{div}(a(x_0)\nabla v_i^0) = 0 \quad \text{in } B(x_0, r)$$

such that $\det [\nabla v_1^0 \quad \cdots \quad \nabla v_d^0] \neq 0$ in $B(x_0, r)$.

2. Find $\tilde{r} \in (0, r]$ and v_i such that $Lv_i = 0$ in $B(x_0, \tilde{r})$ and


$$\|v_i^0 - v_i\|_{C^1(\overline{B(x_0, \tilde{r})})}$$

is arbitrarily small.


3. Runge approximation: find u_i such that $Lu_i = 0$ in Ω and $\|v_i - u_i\|_{C^1(\overline{B(x_0, \tilde{r}/2)})}$ is arbitrarily small. Thus

$$\det [\nabla u_1 \quad \cdots \quad \nabla u_d](x) \neq 0, \quad x \in B(x_0, \tilde{r}/2).$$

4. Covering of $\overline{\Omega}$ with N balls: $N \cdot d$ boundary conditions.



The Runge approximation and non-zero constraints [Bal and Uhlmann, CPAM 2013]



1. Fix $x_0 \in \overline{\Omega}$ and $r > 0$. Consider local solutions $v_i^0 = x_i$:

$$-\operatorname{div}(a(x_0)\nabla v_i^0) = 0 \quad \text{in } B(x_0, r)$$

such that $\det [\nabla v_1^0 \ \cdots \ \nabla v_d^0] \neq 0$ in $B(x_0, r)$.

2. Find $\tilde{r} \in (0, r]$ and v_i such that $L v_i = 0$ in $B(x_0, \tilde{r})$ and

$$\|v_i^0 - v_i\|_{C^1(\overline{B(x_0, \tilde{r})})}$$


is arbitrarily small.

3. Runge approximation: find u_i such that $L u_i = 0$ in Ω and $\|v_i - u_i\|_{C^1(\overline{B(x_0, \tilde{r}/2)})}$ is arbitrarily small. Thus

$$\det [\nabla u_1 \ \cdots \ \nabla u_d](x) \neq 0, \quad x \in B(x_0, \tilde{r}/2).$$

4. Covering of $\overline{\Omega}$ with N balls: $N \cdot d$ boundary conditions.

The Runge approximation and non-zero constraints [Bal and Uhlmann, CPAM 2013]



1. Fix $x_0 \in \overline{\Omega}$ and $r > 0$. Consider local solutions $v_i^0 = x_i$:

$$-\operatorname{div}(a(x_0)\nabla v_i^0) = 0 \quad \text{in } B(x_0, r)$$

such that $\det [\nabla v_1^0 \ \cdots \ \nabla v_d^0] \neq 0$ in $B(x_0, r)$.

2. Find $\tilde{r} \in (0, r]$ and v_i such that $Lv_i = 0$ in $B(x_0, \tilde{r})$ and

$$\|v_i^0 - v_i\|_{C^1(\overline{B(x_0, \tilde{r})})}$$


is arbitrarily small.

3. Runge approximation: find u_i such that $Lu_i = 0$ in Ω and $\|v_i - u_i\|_{C^1(\overline{B(x_0, \tilde{r}/2)})}$ is arbitrarily small. Thus

$$\det [\nabla u_1 \ \cdots \ \nabla u_d] (x) \neq 0, \quad x \in B(x_0, \tilde{r}/2).$$

4. Covering of $\overline{\Omega}$ with N balls: $N \cdot d$ boundary conditions.

The Runge approximation and non-zero constraints [Bal and Uhlmann, CPAM 2013]



1. Fix $x_0 \in \overline{\Omega}$ and $r > 0$. Consider local solutions $v_i^0 = x_i$:

$$-\operatorname{div}(a(x_0)\nabla v_i^0) = 0 \quad \text{in } B(x_0, r)$$

such that $\det [\nabla v_1^0 \ \cdots \ \nabla v_d^0] \neq 0$ in $B(x_0, r)$.

2. Find $\tilde{r} \in (0, r]$ and v_i such that $Lv_i = 0$ in $B(x_0, \tilde{r})$ and

$$\|v_i^0 - v_i\|_{C^1(\overline{B(x_0, \tilde{r})})}$$


is arbitrarily small.

3. Runge approximation: find u_i such that $Lu_i = 0$ in Ω and $\|v_i - u_i\|_{C^1(\overline{B(x_0, \tilde{r}/2)})}$ is arbitrarily small. Thus

$$\det [\nabla u_1 \ \cdots \ \nabla u_d] (x) \neq 0, \quad x \in B(x_0, \tilde{r}/2).$$

4. Covering of $\overline{\Omega}$ with N balls: $N \cdot d$ boundary conditions.

The Runge approximation and non-zero constraints [Bal and Uhlmann, CPAM 2013]



1. Fix $x_0 \in \overline{\Omega}$ and $r > 0$. Consider local solutions $v_i^0 = x_i$:

$$-\operatorname{div}(a(x_0)\nabla v_i^0) = 0 \quad \text{in } B(x_0, r)$$

such that $\det [\nabla v_1^0 \quad \cdots \quad \nabla v_d^0] \neq 0$ in $B(x_0, r)$.

2. Find $\tilde{r} \in (0, r]$ and v_i such that $Lv_i = 0$ in $B(x_0, \tilde{r})$ and

$$\|v_i^0 - v_i\|_{C^1(\overline{B(x_0, \tilde{r})})}$$

is arbitrarily small.

3. Runge approximation: find u_i such that $Lu_i = 0$ in Ω and $\|v_i - u_i\|_{C^1(\overline{B(x_0, \tilde{r}/2)})}$ is arbitrarily small. Thus

$$\det [\nabla u_1 \quad \cdots \quad \nabla u_d](x) \neq 0, \quad x \in B(x_0, \tilde{r}/2).$$

4. Covering of $\overline{\Omega}$ with N balls: $N \cdot d$ boundary conditions.

Two main issues

- ▶ You need a **large number of measurements** to satisfy the constraint

$$\text{rank} \begin{bmatrix} \nabla u_1 & \nabla u_2 & \cdots & \nabla u_{Nd} \end{bmatrix} = d$$

everywhere.

- ▶ The suitable solutions, and so their boundary values, are **not explicitly constructed** (axiom of choice).

Two main issues

- ▶ You need a **large number of measurements** to satisfy the constraint

$$\text{rank} \begin{bmatrix} \nabla u_1 & \nabla u_2 & \cdots & \nabla u_{Nd} \end{bmatrix} = d$$

everywhere.

- ▶ The suitable solutions, and so their boundary values, are **not explicitly constructed** (axiom of choice).

Whitney projection argument

Lemma (Greene and Wu 1975)

Take $k > 2d$ (possibly large). Let u_1, \dots, u_k be solutions to $Lu_i = 0$ in Ω such that

$$\text{rank} \begin{bmatrix} \nabla u_1 & \cdots & \nabla u_{k-1} & \nabla u_k \end{bmatrix} (x) = d, \quad x \in \overline{\Omega}.$$

Then, for almost every $a \in \mathbb{R}^{k-1}$, we have

$$\text{rank} \begin{bmatrix} \nabla (u_1 - a_1 u_k) & \nabla (u_2 - a_2 u_k) & \cdots & \nabla (u_{k-1} - a_{k-1} u_k) \end{bmatrix} (x) = d, \quad x \in \overline{\Omega}.$$

In other words: we can almost always reduce the number of solutions (until $2d$) and keep the constraint. In particular, arbitrarily small weights a can be used.

Whitney projection argument

Lemma (Greene and Wu 1975)

Take $k > 2d$ (possibly large). Let u_1, \dots, u_k be solutions to $Lu_i = 0$ in Ω such that

$$\text{rank} \begin{bmatrix} \nabla u_1 & \cdots & \nabla u_{k-1} & \nabla u_k \end{bmatrix} (x) = d, \quad x \in \overline{\Omega}.$$

Then, for almost every $a \in \mathbb{R}^{k-1}$, we have

$$\text{rank} \begin{bmatrix} \nabla (u_1 - a_1 u_k) & \nabla (u_2 - a_2 u_k) & \cdots & \nabla (u_{k-1} - a_{k-1} u_k) \end{bmatrix} (x) = d, \quad x \in \overline{\Omega}.$$

In other words: we can almost always reduce the number of solutions (until $2d$) and keep the constraint. In particular, arbitrarily small weights a can be used.

Whitney projection argument

Lemma (Greene and Wu 1975)

Take $k > 2d$ (possibly large). Let u_1, \dots, u_k be solutions to $Lu_i = 0$ in Ω such that

$$\text{rank} \begin{bmatrix} \nabla u_1 & \cdots & \nabla u_{k-1} & \nabla u_k \end{bmatrix} (x) = d, \quad x \in \overline{\Omega}.$$

Then, for almost every $a \in \mathbb{R}^{k-1}$, we have

$$\text{rank} \begin{bmatrix} \nabla (u_1 - a_1 u_k) & \nabla (u_2 - a_2 u_k) & \cdots & \nabla (u_{k-1} - a_{k-1} u_k) \end{bmatrix} (x) = d, \quad x \in \overline{\Omega}.$$

In other words: we can almost always reduce the number of solutions (until $2d$) and keep the constraint.
In particular, arbitrarily small weights a can be used.

Whitney projection argument

Lemma (Greene and Wu 1975)

Take $k > 2d$ (possibly large). Let u_1, \dots, u_k be solutions to $Lu_i = 0$ in Ω such that

$$\text{rank} \begin{bmatrix} \nabla u_1 & \cdots & \nabla u_{k-1} & \nabla u_k \end{bmatrix} (x) = d, \quad x \in \overline{\Omega}.$$

Then, for almost every $a \in \mathbb{R}^{k-1}$, we have

$$\text{rank} \begin{bmatrix} \nabla (u_1 - a_1 u_k) & \nabla (u_2 - a_2 u_k) & \cdots & \nabla (u_{k-1} - a_{k-1} u_k) \end{bmatrix} (x) = d, \quad x \in \overline{\Omega}.$$

In other words: we can almost always reduce the number of solutions (until $2d$) and keep the constraint. In particular, arbitrarily small weights a can be used.

Runge & Whitney: main result

Theorem (GSA and Capdeboscq, IMRN 2019)

The set of $2d$ solutions u_1, \dots, u_{2d} to $Lu_i = 0$ in Ω such that

$$\text{rank} \begin{bmatrix} \nabla u_1 & \cdots & \nabla u_{2d} \end{bmatrix} (x) = d, \quad x \in \overline{\Omega},$$

is open and dense in the set of $2d$ solutions to $Lu_i = 0$ in Ω .

Proof.

Open. The rank is stable under small perturbations of u_i .

Dense. Take $\tilde{u}_1, \dots, \tilde{u}_{2d}$ solutions to $L\tilde{u}_i = 0$. By Runge, we have a large number of solutions so that

$$\text{rank} \begin{bmatrix} \nabla u_1 & \cdots & \nabla u_{Nd} \end{bmatrix} (x) = d, \quad x \in \overline{\Omega}.$$

In particular

$$\text{rank} \begin{bmatrix} \nabla \tilde{u}_1 & \cdots & \nabla \tilde{u}_{2d} & \nabla u_1 & \cdots & \nabla u_{Nd} \end{bmatrix} (x) = d, \quad x \in \overline{\Omega}.$$

Apply Whitney reduction lemma Nd times with small weights a , until you reach $2d$ solutions.

Runge & Whitney: main result

Theorem (GSA and Capdeboscq, IMRN 2019)

The set of $2d$ solutions u_1, \dots, u_{2d} to $Lu_i = 0$ in Ω such that

$$\text{rank} \begin{bmatrix} \nabla u_1 & \cdots & \nabla u_{2d} \end{bmatrix} (x) = d, \quad x \in \overline{\Omega},$$

is open and dense in the set of $2d$ solutions to $Lu_i = 0$ in Ω .

Proof.

Open. The rank is stable under small perturbations of u_i .

Dense. Take $\tilde{u}_1, \dots, \tilde{u}_{2d}$ solutions to $L\tilde{u}_i = 0$. By Runge, we have a large number of solutions so that

$$\text{rank} \begin{bmatrix} \nabla u_1 & \cdots & \nabla u_{Nd} \end{bmatrix} (x) = d, \quad x \in \overline{\Omega}.$$

In particular

$$\text{rank} \begin{bmatrix} \nabla \tilde{u}_1 & \cdots & \nabla \tilde{u}_{2d} & \nabla u_1 & \cdots & \nabla u_{Nd} \end{bmatrix} (x) = d, \quad x \in \overline{\Omega}.$$

Apply Whitney reduction lemma Nd times with small weights a , until you reach $2d$ solutions.

Runge & Whitney: main result

Theorem (GSA and Capdeboscq, IMRN 2019)

The set of $2d$ solutions u_1, \dots, u_{2d} to $Lu_i = 0$ in Ω such that

$$\text{rank} \begin{bmatrix} \nabla u_1 & \cdots & \nabla u_{2d} \end{bmatrix} (x) = d, \quad x \in \overline{\Omega},$$

is *open and dense* in the set of $2d$ solutions to $Lu_i = 0$ in Ω .

Proof.

Open. The rank is stable under small perturbations of u_i .

Dense. Take $\tilde{u}_1, \dots, \tilde{u}_{2d}$ solutions to $L\tilde{u}_i = 0$. By Runge, we have a large number of solutions so that

$$\text{rank} \begin{bmatrix} \nabla u_1 & \cdots & \nabla u_{Nd} \end{bmatrix} (x) = d, \quad x \in \overline{\Omega}.$$

In particular

$$\text{rank} \begin{bmatrix} \nabla \tilde{u}_1 & \cdots & \nabla \tilde{u}_{2d} & \nabla u_1 & \cdots & \nabla u_{Nd} \end{bmatrix} (x) = d, \quad x \in \overline{\Omega}.$$

Apply Whitney reduction lemma Nd times with small weights a , until you reach $2d$ solutions.

Runge & Whitney: main result

Theorem (GSA and Capdeboscq, IMRN 2019)

The set of $2d$ solutions u_1, \dots, u_{2d} to $Lu_i = 0$ in Ω such that

$$\text{rank} \begin{bmatrix} \nabla u_1 & \cdots & \nabla u_{2d} \end{bmatrix} (x) = d, \quad x \in \overline{\Omega},$$

is *open and dense* in the set of $2d$ solutions to $Lu_i = 0$ in Ω .

Proof.

Open. The rank is stable under small perturbations of u_i .

Dense. Take $\tilde{u}_1, \dots, \tilde{u}_{2d}$ solutions to $L\tilde{u}_i = 0$. By [Runge](#), we have a large number of solutions so that

$$\text{rank} \begin{bmatrix} \nabla u_1 & \cdots & \nabla u_{Nd} \end{bmatrix} (x) = d, \quad x \in \overline{\Omega}.$$

In particular

$$\text{rank} \begin{bmatrix} \nabla \tilde{u}_1 & \cdots & \nabla \tilde{u}_{2d} & \nabla u_1 & \cdots & \nabla u_{Nd} \end{bmatrix} (x) = d, \quad x \in \overline{\Omega}.$$

Apply [Whitney reduction](#) lemma Nd times with small weights a , until you reach $2d$ solutions.

Runge & Whitney: main result

Theorem (GSA and Capdeboscq, IMRN 2019)

The set of $2d$ solutions u_1, \dots, u_{2d} to $Lu_i = 0$ in Ω such that

$$\text{rank} \begin{bmatrix} \nabla u_1 & \cdots & \nabla u_{2d} \end{bmatrix} (x) = d, \quad x \in \overline{\Omega},$$

is *open and dense* in the set of $2d$ solutions to $Lu_i = 0$ in Ω .

Proof.

Open. The rank is stable under small perturbations of u_i .

Dense. Take $\tilde{u}_1, \dots, \tilde{u}_{2d}$ solutions to $L\tilde{u}_i = 0$. By [Runge](#), we have a large number of solutions so that

$$\text{rank} \begin{bmatrix} \nabla u_1 & \cdots & \nabla u_{Nd} \end{bmatrix} (x) = d, \quad x \in \overline{\Omega}.$$

In particular

$$\text{rank} \begin{bmatrix} \nabla \tilde{u}_1 & \cdots & \nabla \tilde{u}_{2d} & \nabla u_1 & \cdots & \nabla u_{Nd} \end{bmatrix} (x) = d, \quad x \in \overline{\Omega}.$$

Apply [Whitney reduction](#) lemma Nd times with small weights a , until you reach $2d$ solutions.

Runge & Whitney: main result

Theorem (GSA and Capdeboscq, IMRN 2019)

The set of $2d$ solutions u_1, \dots, u_{2d} to $Lu_i = 0$ in Ω such that

$$\text{rank} \begin{bmatrix} \nabla u_1 & \cdots & \nabla u_{2d} \end{bmatrix} (x) = d, \quad x \in \overline{\Omega},$$

is *open and dense* in the set of $2d$ solutions to $Lu_i = 0$ in Ω .

Proof.

Open. The rank is stable under small perturbations of u_i .

Dense. Take $\tilde{u}_1, \dots, \tilde{u}_{2d}$ solutions to $L\tilde{u}_i = 0$. By [Runge](#), we have a large number of solutions so that

$$\text{rank} \begin{bmatrix} \nabla u_1 & \cdots & \nabla u_{Nd} \end{bmatrix} (x) = d, \quad x \in \overline{\Omega}.$$

In particular

$$\text{rank} \begin{bmatrix} \nabla \tilde{u}_1 & \cdots & \nabla \tilde{u}_{2d} & \nabla u_1 & \cdots & \nabla u_{Nd} \end{bmatrix} (x) = d, \quad x \in \overline{\Omega}.$$

Apply [Whitney reduction](#) lemma Nd times with small weights a , until you reach $2d$ solutions.

Remarks on the result

- ▶ As a corollary, the set of $2d$ boundary conditions whose solutions satisfy the constraint everywhere is open and dense.
- ▶ The approach is very general, and works with many other constraints, like

$ u_1 (x) > 0$ (nodal set)	$d + 1$ solutions
$ \det [\nabla u_1 \ \cdots \ \nabla u_d] (x) > 0$ (Jacobian)	$2d$ solutions
$ \det \begin{bmatrix} u_1 & \cdots & u_{d+1} \\ \nabla u_1 & \cdots & \nabla u_{d+1} \end{bmatrix} (x) > 0$ (“augmented” Jacobian)	$2d + 1$ solutions

which appear in several hybrid problems.

- ▶ However, open and dense sets may have arbitrarily small measure...

Remarks on the result

- ▶ As a corollary, the set of $2d$ boundary conditions whose solutions satisfy the constraint everywhere is open and dense.
- ▶ The approach is very general, and works with many other constraints, like

$ u_1 (x) > 0$ (nodal set)	$d + 1$ solutions
$ \det [\nabla u_1 \ \cdots \ \nabla u_d] (x) > 0$ (Jacobian)	$2d$ solutions
$ \det \begin{bmatrix} u_1 & \cdots & u_{d+1} \\ \nabla u_1 & \cdots & \nabla u_{d+1} \end{bmatrix} (x) > 0$ (“augmented” Jacobian)	$2d + 1$ solutions

which appear in several hybrid problems.

- ▶ However, open and dense sets may have arbitrarily small measure...

Remarks on the result

- ▶ As a corollary, the set of $2d$ boundary conditions whose solutions satisfy the constraint everywhere is open and dense.
- ▶ The approach is very general, and works with many other constraints, like

$ u_1 (x) > 0$ (nodal set)	$d + 1$ solutions
$ \det [\nabla u_1 \ \cdots \ \nabla u_d] (x) > 0$ (Jacobian)	$2d$ solutions
$ \det \begin{bmatrix} u_1 & \cdots & u_{d+1} \\ \nabla u_1 & \cdots & \nabla u_{d+1} \end{bmatrix} (x) > 0$ (“augmented” Jacobian)	$2d + 1$ solutions

which appear in several hybrid problems.

- ▶ However, open and dense sets may have arbitrarily **small measure**...

Outline of the talk

- 1 The Radó-Kneser-Choquet theorem
- 2 Runge approximation & Whitney embedding
- 3 Using random boundary conditions

Using random boundary conditions

- ▶ ν : subgaussian probability distribution on $H^{1/2}(\partial\Omega)$
- ▶ $\varphi \sim \nu$ is of the form

$$\varphi = \sum_{k \in \mathbb{N}} a_k e_k$$

where $\{e_k\}$ is an ONB of $H^{1/2}(\partial\Omega)$ and $\{a_k\}_k$ are uncorrelated real random variables

Theorem (GSA, IP 2022)

Take $N \in \mathbb{N}$. Let $\varphi_1^l, \dots, \varphi_d^l \sim \nu$ be sampled i.i.d. in $H^{1/2}(\partial\Omega)$ for $l = 1, \dots, N$. Then

$$\max_{l=1, \dots, N} |\det [\nabla u_1^l \quad \dots \quad \nabla u_d^l](x)| \geq C_1, \quad x \in \overline{\Omega'},$$

where

$$\begin{cases} -\operatorname{div}(a \nabla u_i^l) + q u_i^l = 0 & \text{in } \Omega, \\ u_i^l = \varphi_i^l & \text{on } \partial\Omega, \end{cases}$$

with probability greater than

$$1 - C_2 N^{d/2} \exp(-C_3 N^{1/d}).$$

Using random boundary conditions

- ▶ ν : subgaussian probability distribution on $H^{1/2}(\partial\Omega)$
- ▶ $\varphi \sim \nu$ is of the form

$$\varphi = \sum_{k \in \mathbb{N}} a_k e_k$$

where $\{e_k\}$ is an ONB of $H^{1/2}(\partial\Omega)$ and $\{a_k\}_k$ are uncorrelated real random variables

Theorem (GSA, IP 2022)

Take $N \in \mathbb{N}$. Let $\varphi_1^l, \dots, \varphi_d^l \sim \nu$ be sampled i.i.d. in $H^{1/2}(\partial\Omega)$ for $l = 1, \dots, N$. Then

$$\max_{l=1, \dots, N} |\det [\nabla u_1^l \quad \dots \quad \nabla u_d^l](x)| \geq C_1, \quad x \in \overline{\Omega'},$$

where

$$\begin{cases} -\operatorname{div}(a \nabla u_i^l) + q u_i^l = 0 & \text{in } \Omega, \\ u_i^l = \varphi_i^l & \text{on } \partial\Omega, \end{cases}$$

with probability greater than

$$1 - C_2 N^{d/2} \exp(-C_3 N^{1/d}).$$

Using random boundary conditions

- ▶ ν : subgaussian probability distribution on $H^{1/2}(\partial\Omega)$
- ▶ $\varphi \sim \nu$ is of the form

$$\varphi = \sum_{k \in \mathbb{N}} a_k e_k$$

where $\{e_k\}$ is an ONB of $H^{1/2}(\partial\Omega)$ and $\{a_k\}_k$ are uncorrelated real random variables

Theorem (GSA, IP 2022)

Take $N \in \mathbb{N}$. Let $\varphi_1^l, \dots, \varphi_d^l \sim \nu$ be sampled i.i.d. in $H^{1/2}(\partial\Omega)$ for $l = 1, \dots, N$. Then

$$\max_{l=1, \dots, N} |\det [\nabla u_1^l \quad \dots \quad \nabla u_d^l](x)| \geq C_1, \quad x \in \overline{\Omega'},$$

where

$$\begin{cases} -\operatorname{div}(a \nabla u_i^l) + q u_i^l = 0 & \text{in } \Omega, \\ u_i^l = \varphi_i^l & \text{on } \partial\Omega, \end{cases}$$

with probability greater than

$$1 - C_2 N^{d/2} \exp(-C_3 N^{1/d}).$$

Using random boundary conditions

- ▶ ν : subgaussian probability distribution on $H^{1/2}(\partial\Omega)$
- ▶ $\varphi \sim \nu$ is of the form

$$\varphi = \sum_{k \in \mathbb{N}} a_k e_k$$

where $\{e_k\}$ is an ONB of $H^{1/2}(\partial\Omega)$ and $\{a_k\}_k$ are uncorrelated real random variables

Theorem (GSA, IP 2022)

Take $N \in \mathbb{N}$. Let $\varphi_1^l, \dots, \varphi_d^l \sim \nu$ be sampled i.i.d. in $H^{1/2}(\partial\Omega)$ for $l = 1, \dots, N$. Then

$$\max_{l=1, \dots, N} |\det [\nabla u_1^l \quad \dots \quad \nabla u_d^l](x)| \geq C_1, \quad x \in \overline{\Omega'},$$

where

$$\begin{cases} -\operatorname{div}(a \nabla u_i^l) + q u_i^l = 0 & \text{in } \Omega, \\ u_i^l = \varphi_i^l & \text{on } \partial\Omega, \end{cases}$$

with probability greater than

$$1 - C_2 N^{d/2} \exp(-C_3 N^{1/d}).$$

Sketch of the proof

Theorem (GSA, IP 2022)

Take $N \in \mathbb{N}$. Let $\varphi_1^l, \dots, \varphi_d^l \sim \nu$ be sampled i.i.d. in $H^{1/2}(\partial\Omega)$ for $l = 1, \dots, N$. Then

$$\max_{l=1, \dots, N} |\det [\nabla u_1^l \ \cdots \ \nabla u_d^l](x)| \geq C_1, \quad x \in \overline{\Omega'}$$

with probability greater than

$$1 - C_2 N^{d/2} \exp(-C_3 N^{1/d}).$$

Two steps:

1. By quantitative Runge approximation (as in Salo-Rüland-2018, but with arbitrary norms on $\partial\Omega$):

$$\mathbb{E}(|\det [\nabla u_1 \ \cdots \ \nabla u_d](x)|) \geq C$$

2. Concentration inequalities

Sketch of the proof

Theorem (GSA, IP 2022)

Take $N \in \mathbb{N}$. Let $\varphi_1^l, \dots, \varphi_d^l \sim \nu$ be sampled i.i.d. in $H^{1/2}(\partial\Omega)$ for $l = 1, \dots, N$. Then

$$\max_{l=1, \dots, N} |\det [\nabla u_1^l \ \cdots \ \nabla u_d^l](x)| \geq C_1, \quad x \in \overline{\Omega'}$$

with probability greater than

$$1 - C_2 N^{d/2} \exp(-C_3 N^{1/d}).$$

Two steps:

1. By quantitative Runge approximation (as in Salo-Rüland-2018, but with arbitrary norms on $\partial\Omega$):

$$\mathbb{E}(|\det [\nabla u_1 \ \cdots \ \nabla u_d](x)|) \geq C$$

2. Concentration inequalities

Random boundary values: simulations [GSA, Cen and Zhou, preprint 2025]




Figure: Boundary values and the non-zero region $\max_{\ell=1, \dots, N} |\partial_{x_1} u^{(\ell)}(x)| \geq 0.1$, $N = 1, 2, 3, 4, 5$.

Conclusions

- ▶ The inversion in **quantitative hybrid imaging** often requires the solutions to the direct problem to satisfy certain **non-zero constraints**.
- ▶ Available methods:
 - ▶ The Radó-Kneser-Choquet theorem and its generalizations (only in 2D, not for Helmholtz)
 - ▶ CGO solutions
 - ▶ multiple frequencies
 - ▶ Runge & Whitney
 - ▶ Random boundary values
- ▶ Future perspectives:
 - ▶ combine Runge & Whitney with random boundary values
 - ▶ other PDEs (Maxwell, elasticity, etc.)
 - ▶ numerical experiments

Slides:

