


Phase transitions for Gödel incompleteness

Andreas Weiermann
Ghent University

Ghent Logic Seminar, Ghent, September 28th, 2023


Welcome to our logic group



Aim of the talk

- ▶ Talk is of expository nature
- ▶ Addressed to non proof-theorists.
- ▶ Will keep examples simple.
- ▶ Several fairytales which are independent from each other.
- ▶ Will highlight some applications of mathematics within logic.
- ▶ Project is open ended.
- ▶ Interested students, PhD students, and Postdocs may join in.
- ▶ Search for international collaboration is intended.

Phase transition in real life



- ▶ Phase transitions in logic?
- ▶ First or second order phase transitions?
- ▶ Which methods can be used to classify them?
- ▶ Can physical methods like renormalization be useful here?

Outline

- ▶ Peano arithmetic
- ▶ Historical overview: Hilbert, Gödel,
- ▶ The big picture behind phase transitions for incompleteness
- ▶ Ordinals for beginners
- ▶ Ackermann function
- ▶ Goodstein sequences
- ▶ Ramseyan theorems
- ▶ Slowly well orderedness
- ▶ Kruskal's tree theorem
- ▶ Bolzano Weierstrass and Abel's theorem about integrals

Peano arithmetic

How to model the natural numbers?

Via Peano arithmetic PA .

Have symbols for $0, S, +$ and \cdot together with the axioms:

- ▶ $\forall x[\neg Sx = 0]$.
- ▶ $\forall x, y[Sx = Sy \rightarrow x = y]$.
- ▶ $\forall x[x + 0 = x]$.
- ▶ $\forall x, y[x + Sy = S(x + y)]$.
- ▶ $\forall x[x \cdot 0 = 0]$.
- ▶ $\forall x, y[x \cdot Sy = x \cdot y + x]$.
- ▶ For all formulas φ : $(\varphi(0) \wedge \forall x[\varphi(x) \rightarrow \varphi(Sx)]) \rightarrow \forall y\varphi(y)$.

PA is good for developing discrete mathematics.

Compactness

Is every countable model of PA isomorphic to the standard model?

Compactness

Is every countable model of PA isomorphic to the standard model?
No, by compactness.

Inconsistency in naïve set theory

Comprehension [Cantor]: For all properties φ there is a set s such that $s = \{x : \varphi(x)\}$.

Inconsistency in naïve set theory

Comprehension [Cantor]: For all properties φ there is a set s such that $s = \{x : \varphi(x)\}$.

Russell: Let $s := \{x : x \notin x\}$.

Inconsistency in naïve set theory

Comprehension [Cantor]: For all properties φ there is a set s such that $s = \{x : \varphi(x)\}$.

Russell: Let $s := \{x : x \notin x\}$. Then $s \in s \iff s \notin s$.

Inconsistency in naïve set theory

Comprehension [Cantor]: For all properties φ there is a set s such that $s = \{x : \varphi(x)\}$.

Russell: Let $s := \{x : x \notin x\}$. Then $s \in s \iff s \notin s$.

Possible way out: Intuitionism. Predicativism.

ZFC: Replace comprehension by separation/replacement.

Hilbert: Let's save mathematics via proof theory



- ▶ Formalize all of mathematics:
Is every assertion true in the standard model provable in *PA*?
- ▶ Prove the consistency of mathematics by finitary means.
Is the consistency of *PA* provable in *PA*?

Enter Gödel



Is every assertion true in the standard model provable in *PA*?

Gödel one simplified: No

Is the consistency of *PA* provable in *PA*?

Gödel two simplified: No

Phase transitions in logic

Assumptions:

- ▶ $A(h)$ is true for all values of the order parameter h .
- ▶ $A(h)$ is provable for small values of h .
- ▶ $A(h)$ is unprovable for large enough values of h .
- ▶ Unprovability of $A(h)$ is monotone in h .

Then we are looking for the threshold for the transition from provability to unprovability.

Ordinals for beginners

Following Cantor ordinals model counting into the infinite:

$$0 < 1 < 2 < \dots < \omega < \omega + 1 < \dots < \omega + \omega \dots$$

Ordinals for beginners

Following Cantor ordinals model counting into the infinite:

$$0 < 1 < 2 < \dots < \omega < \omega + 1 < \dots < \omega + \omega \dots$$

In our talk we model an interesting segment of ordinals by unary number-theoretic functions under eventual domination.

Ordinals for beginners

Following Cantor ordinals model counting into the infinite:

$$0 < 1 < 2 < \dots < \omega < \omega + 1 < \dots < \omega + \omega \dots$$

In our talk we model an interesting segment of ordinals by unary number-theoretic functions under eventual domination.

Let E be the least class of unary functions such that

- ▶ $m \mapsto 0 \in E$
- ▶ If $g, h \in E$ then $m \mapsto m^{g(m)} + h(m) \in E$.

Ordinals for beginners

Following Cantor ordinals model counting into the infinite:

$$0 < 1 < 2 < \dots < \omega < \omega + 1 < \dots < \omega + \omega \dots$$

In our talk we model an interesting segment of ordinals by unary number-theoretic functions under eventual domination.

Let E be the least class of unary functions such that

- ▶ $m \mapsto 0 \in E$
- ▶ If $g, h \in E$ then $m \mapsto m^{g(m)} + h(m) \in E$.

As for Hardy's orders of infinity define $g < h$ if there exists a K such that for all $m > K$ we have $g(m) < h(m)$.

Ordinals for beginners

Following Cantor ordinals model counting into the infinite:

$$0 < 1 < 2 < \dots < \omega < \omega + 1 < \dots < \omega + \omega \dots$$

In our talk we model an interesting segment of ordinals by unary number-theoretic functions under eventual domination.

Let E be the least class of unary functions such that

- ▶ $m \mapsto 0 \in E$
- ▶ If $g, h \in E$ then $m \mapsto m^{g(m)} + h(m) \in E$.

As for Hardy's orders of infinity define $g < h$ if there exists a K such that for all $m > K$ we have $g(m) < h(m)$.

Then $\langle E, < \rangle$ is a well order. So $<$ is a linear ordering and there is no sequence $f_0, f_1, \dots \in E$ such that $f_{i+1} < f_i$ for all i .

Ordinals for beginners

How can we model the elements of E with natural numbers?

Ordinals for beginners

How can we model the elements of E with natural numbers?

Let c_k be the function $m \mapsto k$. Then

$$c_0 < c_1 < \dots < id < id + 1 < \dots < id + id < \dots$$

Ordinals for beginners

How can we model the elements of E with natural numbers?

Let c_k be the function $m \mapsto k$. Then

$$c_0 < c_1 < \dots < id < id + 1 < \dots < id + id < \dots$$

Write ω^f for the function $m \mapsto m^{f(m)}$.

Ordinals for beginners

How can we model the elements of E with natural numbers?

Let c_k be the function $m \mapsto k$. Then

$c_0 < c_1 < \dots < id < id + 1 < \dots < id + id < \dots$

Write ω^f for the function $m \mapsto m^{f(m)}$.

Theorem

- ▶ Every $\alpha \in E$ can be written uniquely as $\alpha = \omega^{\alpha_1} + \dots + \omega^{\alpha_n}$ where $\alpha > \alpha_1 \geq \dots \geq \alpha_n$.

Ordinals for beginners

How can we model the elements of E with natural numbers?

Let c_k be the function $m \mapsto k$. Then

$$c_0 < c_1 < \dots < id < id + 1 < \dots < id + id < \dots$$

Write ω^f for the function $m \mapsto m^{f(m)}$.

Theorem

- ▶ Every $\alpha \in E$ can be written uniquely as $\alpha = \omega^{\alpha_1} + \dots + \omega^{\alpha_n}$ where $\alpha > \alpha_1 \geq \dots \geq \alpha_n$.
- ▶ If $\alpha = \omega^{\alpha_1} + \dots + \omega^{\alpha_n}$ where $\alpha > \alpha_1 \geq \dots \geq \alpha_n$ and $\beta = \omega^{\beta_1} + \dots + \omega^{\beta_m}$ where $\beta > \beta_1 \geq \dots \geq \beta_m$ then $\alpha < \beta$

Ordinals for beginners

How can we model the elements of E with natural numbers?

Let c_k be the function $m \mapsto k$. Then

$$c_0 < c_1 < \dots < id < id + 1 < \dots < id + id < \dots$$

Write ω^f for the function $m \mapsto m^{f(m)}$.

Theorem

- ▶ Every $\alpha \in E$ can be written uniquely as $\alpha = \omega^{\alpha_1} + \dots + \omega^{\alpha_n}$ where $\alpha > \alpha_1 \geq \dots \geq \alpha_n$.
- ▶ If $\alpha = \omega^{\alpha_1} + \dots + \omega^{\alpha_n}$ where $\alpha > \alpha_1 \geq \dots \geq \alpha_n$ and $\beta = \omega^{\beta_1} + \dots + \omega^{\beta_m}$ where $\beta > \beta_1 \geq \dots \geq \beta_m$ then $\alpha < \beta$ iff either $n < m$ and $(\forall i \leq n)[\alpha_i = \beta_i]$, or there exists an $i \leq \min(m, n)$ such that $\alpha_i < \beta_i$ and $(\forall j < i)[\alpha_j = \beta_j]$.

Gentzen

Assume that $\langle E, < \rangle$ is order isomorphic to $\langle \mathbb{N}, \prec \rangle$.

$$I(\alpha) := \forall x[(\forall y(y \prec x \varphi(y)) \rightarrow \varphi(x)] \rightarrow \forall x \prec \alpha \varphi(x)$$

$$I := \forall x[(\forall y(y \prec x \varphi(y)) \rightarrow \varphi(x)] \rightarrow \forall x \varphi(x)$$

Gentzen:

- ▶ PA proves $I(\alpha)$ for any $\alpha \in E$.
- ▶ PA does not prove I .

This is first example of a (meta mathematical) phase transition for Gödel incompleteness.

$PA + I$ proves the consistency of PA .

Phase transition for the Ackermann function

$$A_0(n) := n + 1.$$

$$A_{d+1}(n) := A_d^n(n) = \overbrace{A_d(\dots A_d(n))}^{n\text{-times}}.$$

$$A(n) := A_n(n).$$

Then A_d is primitive recursive for any fixed d but the function A is not prim rec.

Phase transition for the Ackermann function

$$\begin{aligned}
 A_0(n) &:= n + 1. \\
 A_{d+1}(n) &:= \overbrace{A_d(\dots A_d(n))}^{n\text{-times}}. \\
 A(n) &:= A_n(n).
 \end{aligned}$$

Then A_d is primitive recursive for any fixed d but the function A is not prim rec. Introduce a parameter $h : \mathbb{N} \rightarrow \mathbb{N}$.

$$\begin{aligned}
 A(h)_0(n) &:= n + 1. \\
 A(h)_{d+1}(n) &:= A(h)_d^{h(n)}(n). \\
 A(h)(n) &:= A(h)_n(n).
 \end{aligned}$$

Then for $h = id$ the function $A(h)$ is not prim rec but $A(h)$ is prim rec for $h = const.$ Is there a non trivial phase transition?

Phase transition for Ackermann function

We round h up to natural number values.

Theorem

- ▶ If $h(n) = \sqrt[d]{n}$ then $A(h)$ is not prim rec.
- ▶ If $h(n) = \log(n)$ then $A(h)$ is prim rec.

Phase transition for Ackermann function

We round h up to natural number values.

Theorem

- ▶ If $h(n) = \sqrt[d]{n}$ then $A(h)$ is not prim rec.
- ▶ If $h(n) = \log(n)$ then $A(h)$ is prim rec.

For a non decreasing unbounded function h let

$h^{-1}(n) := \min\{m : h(m) \geq n\}$. Then for quickly growing functions h the induced function h^{-1} is slowly growing.

Theorem (Omri and W.)

- ▶ If $h(n) = \sqrt[{}^{A^{-1}(n)}]{n}$ then $A(h)$ is not prim rec.
- ▶ If $h(n) = \sqrt[{}^{A_d^{-1}(n)}]{n}$ for some fixed d then $A(h)$ is prim rec.

Proof idea: $A^{-1}(n)$ behaves like a constant for the branches of the Ackermann function.

Phase transition for the generalized Ackermann function

Let g be strictly increasing such that $g(n) \geq n + 1$.

$$A(g, h)_0(n) := g(n).$$

$$A(g, h)_{d+1}(n) := A(g, h)_d^{h(n)}(n).$$

$$A(g, h)(n) := A(g, h)_n(n).$$

A classification result for a general function g seems out of reach

Phase transition for the generalized Ackermann function

Let g be strictly increasing such that $g(n) \geq n + 1$.

$$A(g, h)_0(n) := g(n).$$

$$A(g, h)_{d+1}(n) := A(g, h)_d^{h(n)}(n).$$

$$A(g, h)(n) := A(g, h)_n(n).$$

A classification result for a general function g seems out of reach but our search was successful. Let $g'(n) := g^n(n)$.

Theorem

- ▶ If $h(n) = \sqrt[A^{-1}(n)]{(g')^{-1}(n)}$ then $A(g, h)$ is not prim rec.
- ▶ If $h(n) = \sqrt[A_d^{-1}(n)]{(g')^{-1}(n)}$ for some fixed d then $A(g, h)$ is prim rec.

This theorem has applications to the pigeonhole principle.

Moreover this theorem extends to transfinite extensions $A_\alpha(g, h)$ but we will skip further details here.

Phase transition for the Goodstein principle

Fix a natural number $k \geq 2$.

Phase transition for the Goodstein principle

Fix a natural number $k \geq 2$. Every positive integer m can be written uniquely as $m = k^r \cdot p + q$ where $k^r \leq m < k^{r+1}$ and $p < k$ and $q < k^r$.

Phase transition for the Goodstein principle

Fix a natural number $k \geq 2$. Every positive integer m can be written uniquely as $m = k^r \cdot p + q$ where $k^r \leq m < k^{r+1}$ and $p < k$ and $q < k^r$. Define $0\{k\} := 0$ and $m\{k\} := (k+1)^{r\{k\}} \cdot p + q\{k\}$.

Phase transition for the Goodstein principle

Fix a natural number $k \geq 2$. Every positive integer m can be written uniquely as $m = k^r \cdot p + q$ where $k^r \leq m < k^{r+1}$ and $p < k$ and $q < k^r$. Define $0\{k\} := 0$ and $m\{k\} := (k+1)^{r\{k\}} \cdot p + q\{k\}$. Define $m_0 := m$ and if $m_l > 0$ let $m_{l+1} := m_l\{2+l\} - 1$. If $m_l = 0$ then $m_{l+1} := 0$.

Phase transition for the Goodstein principle

Fix a natural number $k \geq 2$. Every positive integer m can be written uniquely as $m = k^r \cdot p + q$ where $k^r \leq m < k^{r+1}$ and $p < k$ and $q < k^r$. Define $0\{k\} := 0$ and $m\{k\} := (k+1)^{r\{k\}} \cdot p + q\{k\}$. Define $m_0 := m$ and if $m_l > 0$ let $m_{l+1} := m_l\{2+l\} - 1$. If $m_l = 0$ then $m_{l+1} := 0$.

These numbers grow quickly in the beginning and every laptop will make you believe that m_l is diverging for not too small starting values of m .

Example: $m = 2^2 + 2$. $m_0 = m$. $m_1 = 3^3 + 3 - 1 = 3^3 + 2$. etc.

Theorem

Let G stand for $\forall m \exists l (m_l = 0)$.

Phase transition for the Goodstein principle

Fix a natural number $k \geq 2$. Every positive integer m can be written uniquely as $m = k^r \cdot p + q$ where $k^r \leq m < k^{r+1}$ and $p < k$ and $q < k^r$. Define $0\{k\} := 0$ and $m\{k\} := (k+1)^{r\{k\}} \cdot p + q\{k\}$. Define $m_0 := m$ and if $m_l > 0$ let $m_{l+1} := m_l\{2+l\} - 1$. If $m_l = 0$ then $m_{l+1} := 0$.

These numbers grow quickly in the beginning and every laptop will make you believe that m_l is diverging for not too small starting values of m .

Example: $m = 2^2 + 2$. $m_0 = m$. $m_1 = 3^3 + 3 - 1 = 3^3 + 2$. etc.

Theorem

Let G stand for $\forall m \exists l (m_l = 0)$.

- ▶ The assertion G is true.
- ▶ But $PA \not\vdash G$.

Mentioned by Penrose.

Phase transition for the Goodstein principle

Let us introduce a function parameter h for G where $h : \mathbb{N} \rightarrow \mathbb{N}$ is non decreasing.

Every positive integer m can be written uniquely as

$m = (2 + h(k))^r \cdot p + q$ where $(2 + h(k))^r \leq m < (2 + h(k))^{r+1}$

and $p < 2 + h(k)$ and $q < (2 + h(k))^r$. Define $0\{k\}^h := 0$ and

$m\{k\}^h := (2 + h(k+1))^{r\{k\}^h} \cdot p + q\{k\}^h$. Define $m_0^h := m$ and if
 $m_I^h > 0$ let $m_{I+1}^h := m_I^h\{I\}^h - 1$. If $m_I = 0$ then $m_{I+1}^h := 0$.

Phase transition for the Goodstein principle

Let us introduce a function parameter h for G where $h : \mathbb{N} \rightarrow \mathbb{N}$ is non decreasing.

Every positive integer m can be written uniquely as

$m = (2 + h(k))^r \cdot p + q$ where $(2 + h(k))^r \leq m < (2 + h(k))^{r+1}$

and $p < 2 + h(k)$ and $q < (2 + h(k))^r$. Define $0\{k\}^h := 0$ and

$m\{k\}^h := (2 + h(k+1))^{r\{k\}^h} \cdot p + q\{k\}^h$. Define $m_0^h := m$ and if $m_I^h > 0$ let $m_{I+1}^h := m_I^h\{I\}^h - 1$. If $m_I = 0$ then $m_{I+1}^h := 0$.

Theorem

Let $G(h)$ stand for $\forall m \exists I (m_I^h = 0)$. Let \log stand for the binary length function. $\log^1 := \log$ and $\log^{d+1} := \log(\log^d)$. Let $\log^*(m)$ be the least number d such that $\log^d(m) \leq 1$.

- ▶ The assertion $G(h)$ is true for all h .
- ▶ $PA \vdash G(\log^*)$.
- ▶ But for any fixed d $PA \not\vdash G(\log^d)$.

Phase transition for the Paris Harrington principle

We identify R with the set $\{0, 1, \dots, R - 1\}$. With $[X]^d$ we denote the set of d -element subsets of X .

The finite Ramsey theorem RT states:

$\forall d, c, m \exists R \forall P : [R]^d \rightarrow c \exists H \subseteq R [P \upharpoonright [H]^d = \text{const} \wedge |H| \geq m]$.
The least R depending on d, c, m is denoted by $R_c^d(m)$.

Theorem

- ▶ *RT is true.*
- ▶ *PA ⊢ RT.*
- ▶ *There exists a constant const such that $R_c^d(m) \leq 2_{d-1}(\text{const} \cdot c \cdot m)$.*

No phase transition yet.

Phase transition for the Paris Harrington principle

The Paris Harrington thm PH states: $\forall d, c, m \exists R \forall P : [R]^d \rightarrow c \exists H \subseteq R [P \upharpoonright [H]^d = \text{const} \wedge |H| \geq \max\{\min(H), m\}]$.

Theorem (Paris Harrington)

- ▶ PH is true.
- ▶ $PA \not\vdash PH$.

This indicates that a phase transition might be possible.

Phase transition for the Paris Harrington principle

The Paris Harrington theorem $PH(h)$ states: $\forall d, c, m \exists R \forall P : [R]^d \rightarrow c \exists H \subseteq R[P \upharpoonright [H]^d = \text{const} \wedge |H| \geq \max\{h(\min(H)), m\}]$.

Theorem (W.)

- ▶ $PH(h)$ is true.
- ▶ $PA \vdash PH(\log^*)$.
- ▶ If d is fixed then $PA \not\vdash PH(\log^d)$.

Similar results hold for the regressive Ramsey theorem and the canonical Ramsey theorem. Threshold is similar to the one for Goodstein. Further refinements are possible.

Phase transition for the Paris Harrington principle

Proof of $PA \vdash PH(\log^*)$.

Let m, d, c be given and put $R := R_c^d(m \cdot 2)$. W.l.o.g $m = d = c$.

Let $P : [R]^d \rightarrow c$ be given. Then by RT there exists H such that $P \upharpoonright [H]^d = \text{const}$ and $2m \leq |H|$. But now we have the following insightful conclusion using the Erdős Rado bound $\log^*(\min H) \leq \log^*(R) \leq \log^*(2_{d-1}(\text{const} \cdot m \cdot c)) \leq \log^*(2_{2m}) = 2m \leq |H|$.

Phase transition for the Paris Harrington principle

Proof of $PA \vdash PH(\log^*)$.

Let m, d, c be given and put $R := R_c^d(m \cdot 2)$. W.l.o.g $m = d = c$. Let $P : [R]^d \rightarrow c$ be given. Then by RT there exists H such that $P \upharpoonright [H]^d = \text{const}$ and $2m \leq |H|$. But now we have the following insightful conclusion using the Erdős Rado bound $\log^*(\min H) \leq \log^*(R) \leq \log^*(2_{d-1}(\text{const} \cdot m \cdot c)) \leq \log^*(2_{2m}) = 2m \leq |H|$.

Insight: Finite combinatorics may indicate the threshold function. The threshold emerges as the **functional inverse** of the bounding function.

Ordinal counting

Let us consider ordinals in E . Let $N\alpha := n + N\alpha_1 + \dots + N\alpha_n$ if $\alpha = \omega^{\alpha_1} + \dots + \omega^{\alpha_n}$ where $\alpha > \alpha_1 \geq \dots \geq \alpha_n$.

Then for all d the set of all $\alpha \in E$ with $N\alpha \leq d$ is finite (bounded by an exponential upper bound in n).

Let $c_\alpha(n) = \#\{\beta < \alpha : N\beta = n\}$.

Then $c_{\omega^d}(n) \sim \text{const} \cdot n^{d-1}$.

What about other values of $c_\alpha(n)$?

Ordinal counting

Let us consider ordinals in E . Let $N\alpha := n + N\alpha_1 + \dots + N\alpha_n$ if $\alpha = \omega^{\alpha_1} + \dots + \omega^{\alpha_n}$ where $\alpha > \alpha_1 \geq \dots \geq \alpha_n$.

Then for all d the set of all $\alpha \in E$ with $N\alpha \leq d$ is finite (bounded by an exponential upper bound in n).

Let $c_\alpha(n) = \#\{\beta < \alpha : N\beta = n\}$.

Then $c_{\omega^d}(n) \sim \text{const} \cdot n^{d-1}$.

What about other values of $c_\alpha(n)$?

Theorem

Let $\omega_1 = \omega$ and $\omega_{d+1} = \omega^{\omega_d}$.

► $c_{\omega^\omega}(n) \sim$

Ordinal counting

Let us consider ordinals in E . Let $N\alpha := n + N\alpha_1 + \dots + N\alpha_n$ if $\alpha = \omega^{\alpha_1} + \dots + \omega^{\alpha_n}$ where $\alpha > \alpha_1 \geq \dots \geq \alpha_n$.

Then for all d the set of all $\alpha \in E$ with $N\alpha \leq d$ is finite (bounded by an exponential upper bound in n).

Let $c_\alpha(n) = \#\{\beta < \alpha : N\beta = n\}$.

Then $c_{\omega^d}(n) \sim \text{const} \cdot n^{d-1}$.

What about other values of $c_\alpha(n)$?

Theorem

Let $\omega_1 = \omega$ and $\omega_{d+1} = \omega^{\omega_d}$.

- ▶ $c_{\omega^\omega}(n) \sim \frac{e^{\pi\sqrt{\frac{2}{3}\cdot n}}}{4\sqrt{3n}}$ following Hardy and Ramanujan.

Ordinal counting

Let us consider ordinals in E . Let $N\alpha := n + N\alpha_1 + \dots + N\alpha_n$ if $\alpha = \omega^{\alpha_1} + \dots + \omega^{\alpha_n}$ where $\alpha > \alpha_1 \geq \dots \geq \alpha_n$.

Then for all d the set of all $\alpha \in E$ with $N\alpha \leq d$ is finite (bounded by an exponential upper bound in n).

Let $c_\alpha(n) = \#\{\beta < \alpha : N\beta = n\}$.

Then $c_{\omega^d}(n) \sim \text{const} \cdot n^{d-1}$.

What about other values of $c_\alpha(n)$?

Theorem

Let $\omega_1 = \omega$ and $\omega_{d+1} = \omega^{\omega_d}$.

- ▶ $c_{\omega^\omega}(n) \sim \frac{e^{\pi\sqrt{\frac{2}{3}\cdot n}}}{4\sqrt{3n}}$ following Hardy and Ramanujan.
- ▶ $\ln(c_{\omega_{d+2}}(n)) \sim$

Ordinal counting

Let us consider ordinals in E . Let $N\alpha := n + N\alpha_1 + \dots + N\alpha_n$ if $\alpha = \omega^{\alpha_1} + \dots + \omega^{\alpha_n}$ where $\alpha > \alpha_1 \geq \dots \geq \alpha_n$.

Then for all d the set of all $\alpha \in E$ with $N\alpha \leq d$ is finite (bounded by an exponential upper bound in n).

Let $c_\alpha(n) = \#\{\beta < \alpha : N\beta = n\}$.

Then $c_{\omega^d}(n) \sim \text{const} \cdot n^{d-1}$.

What about other values of $c_\alpha(n)$?

Theorem

Let $\omega_1 = \omega$ and $\omega_{d+1} = \omega^{\omega_d}$.

- ▶ $c_{\omega^\omega}(n) \sim \frac{e^{\pi\sqrt{\frac{2}{3}\cdot n}}}{4\sqrt{3n}}$ following Hardy and Ramanujan.
- ▶ $\ln(c_{\omega_{d+2}}(n)) \sim \frac{\pi^2}{6} \cdot \frac{n}{\ln^d(n)}$.

Proof by additive number theory, generating functions (Odlyzko).
 Spin off to zero one laws for ordinals.

Slowly well orderedness a la Friedman



Let SWO be the following assertion.

$$\forall K \exists M \forall \alpha_1, \dots, \alpha_M [(\forall i \leq M (N\alpha_i \leq K + i)) \rightarrow \exists i < M (\alpha_i \leq \alpha_{i+1})].$$

Theorem (Friedman)

- ▶ SWO is true.
- ▶ $PA \not\vdash SWO$.

Slowly well orderedness a la Friedman

Let $SWO(h)$ be the following assertion.

$$\forall K \exists M \forall \alpha_1, \dots, \alpha_M [(\forall i \leq M (N\alpha_i \leq K + h(i))) \rightarrow \exists i < M (\alpha_i \leq \alpha_{i+1})].$$

Theorem (W.)

- ▶ $PA \vdash SWO(\log \cdot \log^*)$.
- ▶ $PA \not\vdash SWO(\log \cdot \log^d)$.

Slowly well orderedness a la Friedman

Let $SWO(h)$ be the following assertion.

$$\forall K \exists M \forall \alpha_1, \dots, \alpha_M [(\forall i \leq M (N\alpha_i \leq K + h(i))) \rightarrow \exists i < M (\alpha_i \leq \alpha_{i+1})].$$

Theorem (W.)

- ▶ $PA \vdash SWO(\log \cdot \log^*)$.
- ▶ $PA \not\vdash SWO(\log \cdot \log^d)$.

Proof: $f = \log \cdot \log^d$ is the inverse of the counting function $c_{\omega_{d+2}}$ which yields the provability part by counting. For the unprovability part we apply "renormalization" to Friedman's result for $f = id$.

Slowly well orderedness a la Friedman

Let $SWO(h)$ be the following assertion.

$$\forall K \exists M \forall \alpha_1, \dots, \alpha_M [(\forall i \leq M (N\alpha_i \leq K + h(i))) \rightarrow \exists i < M (\alpha_i \leq \alpha_{i+1})].$$

Theorem (W.)

- ▶ $PA \vdash SWO(\log \cdot \log^*)$.
- ▶ $PA \not\vdash SWO(\log \cdot \log^d)$.

Proof: $f = \log \cdot \log^d$ is the inverse of the counting function $c_{\omega_{d+2}}$ which yields the provability part by counting. For the unprovability part we apply "renormalization" to Friedman's result for $f = id$.

We can also vary the count function. If $N\alpha$ is given by a canonical Gödel numbering we can apply multiplicative number theory (Dirichlet generating functions) and Tauberian theory.

Kruskal's theorem

A finite tree is a finite partial order $\langle T, \leq_T \rangle$ such that

- ▶ T has a minimum r such that $\forall t \in T (r \leq_T t)$ and
- ▶ such that for all $t \in T$ the set $\{s \in T : s \leq t\}$ is totally (linearly) ordered.

Kruskal's theorem

A finite tree is a finite partial order $\langle T, \leq_T \rangle$ such that

- ▶ T has a minimum r such that $\forall t \in T (r \leq_T t)$ and
- ▶ such that for all $t \in T$ the set $\{s \in T : s \leq t\}$ is totally (linearly) ordered.

Then for $s, t \in T$ there exists $\inf(s, t) \in T$.

A one to one function $e : T \rightarrow T'$ is called a homeomorphic embedding if for all $s, t \in T$ we have

$e(\inf_T(s, t)) = \inf_{T'}(e(s), e(t))$. We say $T \trianglelefteq T'$ if there exists a homeomorphic embedding of T into T' .

Kruskal's theorem

A finite tree is a finite partial order $\langle T, \leq_T \rangle$ such that

- ▶ T has a minimum r such that $\forall t \in T (r \leq_T t)$ and
- ▶ such that for all $t \in T$ the set $\{s \in T : s \leq t\}$ is totally (linearly) ordered.

Then for $s, t \in T$ there exists $\inf(s, t) \in T$.

A one to one function $e : T \rightarrow T'$ is called a homeomorphic embedding if for all $s, t \in T$ we have

$e(\inf_T(s, t)) = \inf_{T'}(e(s), e(t))$. We say $T \trianglelefteq T'$ if there exists a homeomorphic embedding of T into T' .

Theorem (Kruskal's theorem KT)

For every ω -sequence of finite trees T_i there exists $i, j < \omega$ such that $i < j$ and $T_i \trianglelefteq T_j$.

Friedman showed that KT is not provable in ATR_0 .

Friedman's miniaturization of Kruskal's theorem FKT



Let $N(T)$ denote the number of nodes in T . Let FKT stand for:
 $\forall K \exists M \forall T_1, \dots, T_M [(\forall i \leq M (N(T_i) \leq K + i)) \rightarrow \exists i < j < M T_i \trianglelefteq T_j]$.

Theorem (Friedman)

- ▶ *FKT is true*
- ▶ *$PA \nvdash FKT$. Even stronger $ATR_0 \nvdash FKT$*

Friedman's miniaturization of Kruskal's theorem FKT

Let $h : \mathbb{N} \rightarrow \mathbb{N}$. The assertion $\text{FKT}(h)$ stands for:

$$\forall K \exists M \forall T_1, \dots, T_M [(\forall i \leq M)[(N(T_i) \leq K + h(i))] \rightarrow (\exists i, j)[i < j < M \wedge T_i \trianglelefteq T_j]].$$

Theorem

- ▶ $\text{FKT}(h)$ is true.
- ▶ If $h = \text{const}$ then $\text{PA} \vdash \text{FKT}(h)$.
- ▶ $\text{PA} \nvdash \text{FKT}(\text{id})$.

Theorem (Matousek and Loebl)

Let $h_r(i) := r \cdot \log_2(i)$.

- ▶ If $r \leq 0.5$ then $\text{PA} \vdash \text{FKT}(h_r)$.
- ▶ If $r \geq 4$ then $\text{PA} \nvdash \text{FKT}(h_r)$.

Any guesses for the threshold?

Let $c = 0.63957768999472013311\dots$

Let $c = 0.63957768999472013311\dots$ and $h_r(i) := r \cdot \log(i)$.

Theorem (W.)

- ▶ If $r \leq c$ then $PA \vdash FKT(h_r)$.
- ▶ If $r > c$ then $PA \not\vdash FKT(h_r)$.

What is c and how did I find it?

Let $c = 0.63957768999472013311\dots$ and $h_r(i) := r \cdot \log(i)$.

Theorem (W.)

- ▶ If $r \leq c$ then $PA \vdash FKT(h_r)$.
- ▶ If $r > c$ then $PA \not\vdash FKT(h_r)$.

What is c and how did I find it?

With a search machine I looked around 2000 for "finite tree". I found Otter's constant $\alpha = 2.955765\dots$ on Stephen Finch's pages about mathematical constants. Voila: $c = \frac{1}{\log_2(\alpha)}$.

Let $c = 0.63957768999472013311\dots$ and $h_r(i) := r \cdot \log(i)$.

Theorem (W.)

- ▶ If $r \leq c$ then $PA \vdash FKT(h_r)$.
- ▶ If $r > c$ then $PA \not\vdash FKT(h_r)$.

What is c and how did I find it?

With a search machine I looked around 2000 for "finite tree". I found Otter's constant $\alpha = 2.955765\dots$ on Stephen Finch's pages about mathematical constants. Voila: $c = \frac{1}{\log_2(\alpha)}$. What is α ?

Let $c = 0.63957768999472013311\dots$ and $h_r(i) := r \cdot \log(i)$.

Theorem (W.)

- ▶ If $r \leq c$ then $PA \vdash FKT(h_r)$.
- ▶ If $r > c$ then $PA \not\vdash FKT(h_r)$.

What is c and how did I find it?

With a search machine I looked around 2000 for "finite tree". I found Otter's constant $\alpha = 2.955765\dots$ on Stephen Finch's pages about mathematical constants. Voila: $c = \frac{1}{\log_2(\alpha)}$. What is α ? Let $T(z) = \sum_{n=0}^{\infty} t_n z^n$ be a solution of $T(z) = z \cdot \exp(\sum_{i=1}^{\infty} T(z^i)/i)$. Let ρ be the radius of convergence of T . Then $\alpha = 1/\rho$.

Let $c = 0.63957768999472013311\dots$ and $h_r(i) := r \cdot \log(i)$.

Theorem (W.)

- ▶ If $r \leq c$ then $PA \vdash FKT(h_r)$.
- ▶ If $r > c$ then $PA \not\vdash FKT(h_r)$.

What is c and how did I find it?

With a search machine I looked around 2000 for "finite tree". I found Otter's constant $\alpha = 2.955765\dots$ on Stephen Finch's pages about mathematical constants. Voila: $c = \frac{1}{\log_2(\alpha)}$. What is α ? Let $T(z) = \sum_{n=0}^{\infty} t_n z^n$ be a solution of $T(z) = z \cdot \exp(\sum_{i=1}^{\infty} T(z^i)/i)$. Let ρ be the radius of convergence of T . Then $\alpha = 1/\rho$. What is t_n ? We have $t_n \sim \text{const} \cdot \frac{\alpha^n}{n^{3/2}}$.

Let $c = 0.63957768999472013311\dots$ and $h_r(i) := r \cdot \log(i)$.

Theorem (W.)

- ▶ If $r \leq c$ then $PA \vdash FKT(h_r)$.
- ▶ If $r > c$ then $PA \not\vdash FKT(h_r)$.

What is c and how did I find it?

With a search machine I looked around 2000 for "finite tree". I found Otter's constant $\alpha = 2.955765\dots$ on Stephen Finch's pages about mathematical constants. Voila: $c = \frac{1}{\log_2(\alpha)}$. What is α ? Let $T(z) = \sum_{n=0}^{\infty} t_n z^n$ be a solution of $T(z) = z \cdot \exp(\sum_{i=1}^{\infty} T(z^i)/i)$. Let ρ be the radius of convergence of T . Then $\alpha = 1/\rho$. What is t_n ? We have $t_n \sim \text{const} \cdot \frac{\alpha^n}{n^{3/2}}$. What is t_n ?
 $t_n = \#\{T : N(T) = n\}$.

Let $c = 0.63957768999472013311\dots$ and $h_r(i) := r \cdot \log(i)$.

Theorem (W.)

- ▶ If $r \leq c$ then $PA \vdash FKT(h_r)$.
- ▶ If $r > c$ then $PA \not\vdash FKT(h_r)$.

What is c and how did I find it?

With a search machine I looked around 2000 for "finite tree". I found Otter's constant $\alpha = 2.955765\dots$ on Stephen Finch's pages about mathematical constants. Voila: $c = \frac{1}{\log_2(\alpha)}$. What is α ? Let $T(z) = \sum_{n=0}^{\infty} t_n z^n$ be a solution of $T(z) = z \cdot \exp(\sum_{i=1}^{\infty} T(z^i)/i)$. Let ρ be the radius of convergence of T . Then $\alpha = 1/\rho$.

What is t_n ? We have $t_n \sim \text{const} \cdot \frac{\alpha^n}{n^{3/2}}$. What is t_n ?

$$t_n = \#\{T : N(T) = n\}.$$

The threshold function h_c is morally the same as the inverse function of $n \mapsto t_n$. The concrete digits of c after 5 entries behind the . have been calculated by Moritz Firsching.

The scaling window for FKT

$$A_0(x) := x + 1.$$

$$A_{\alpha+1}(x) := A_\alpha^x(x).$$

$A_\lambda(x) := A_{\lambda[x]}(x)$ if λ is of limit type .

$$A_E(x) := A_{\omega_x}(x).$$

The scaling window for FKT

$$A_0(x) := x + 1.$$

$$A_{\alpha+1}(x) := A_\alpha^x(x).$$

$A_\lambda(x) := A_{\lambda[x]}(x)$ if λ is of limit type .

$$A_E(x) := A_{\omega_x}(x).$$

Here $\lambda[x]$ converges to λ canonically. So $\omega^{\beta+1}[x] = \omega^\beta \cdot x$ and $\lambda[x]$ is defined by continuity otherwise.

The scaling window for FKT

$$\begin{aligned}
 A_0(x) &:= x + 1. \\
 A_{\alpha+1}(x) &:= A_\alpha^x(x). \\
 A_\lambda(x) &:= A_{\lambda[x]}(x) \text{ if } \lambda \text{ is of limit type .} \\
 A_E(x) &:= A_{\omega_x}(x).
 \end{aligned}$$

Here $\lambda[x]$ converges to λ canonically. So $\omega^{\beta+1}[x] = \omega^\beta \cdot x$ and $\lambda[x]$ is defined by continuity otherwise.

The functions A_α are quickly growing.

Let $h^\alpha(i) = (c + 1/A_\alpha^{-1}(i)) \cdot \log_2(i)$ and

$h^E(i) = (c + 1/A_E^{-1}(i)) \cdot \log_2(i)$. Using Goh and Schmutz (Random struct. and alg.) Gordeev and I could show the following result.

The scaling window for FKT

$$\begin{aligned}
 A_0(x) &:= x + 1. \\
 A_{\alpha+1}(x) &:= A_\alpha^x(x). \\
 A_\lambda(x) &:= A_{\lambda[x]}(x) \text{ if } \lambda \text{ is of limit type .} \\
 A_E(x) &:= A_{\omega_x}(x).
 \end{aligned}$$

Here $\lambda[x]$ converges to λ canonically. So $\omega^{\beta+1}[x] = \omega^\beta \cdot x$ and $\lambda[x]$ is defined by continuity otherwise.

The functions A_α are quickly growing.

Let $h^\alpha(i) = (c + 1/A_\alpha^{-1}(i)) \cdot \log_2(i)$ and
 $h^E(i) = (c + 1/A_E^{-1}(i)) \cdot \log_2(i)$. Using Goh and Schmutz (Random struct. and alg.) Gordeev and I could show the following result.

Theorem

- ▶ If $\alpha \in E$ then $PA \vdash FKT(h^\alpha)$.
- ▶ $PA \not\vdash FKT(h^E)$.



Theorem (Bolzano Weierstraß)

Let $\langle x_i : i < \omega \rangle$ be an infinite sequence of real numbers from the closed unit interval $[0, 1]$. Then there exist $k_1 < k_2 < \dots$ such that the subsequence $\langle x_{k_i} : i < \omega \rangle$ converges.

Theorem (Bolzano Weierstraß)

Let $\langle x_i : i < \omega \rangle$ be an infinite sequence of real numbers from the closed unit interval $[0, 1]$. Then there exist $k_1 < k_2 < \dots$ such that the subsequence $\langle x_{k_i} : i < \omega \rangle$ converges.

Theorem (Monotone convergence theorem)

*Let $\langle x_i : i < \omega \rangle$ be a **non decreasing** infinite sequence of real numbers from the closed unit interval $[0, 1]$.*

Then the sequence $\langle x_i : i < \omega \rangle$ converges.

Friedman's BW

Theorem (modified Bolzano Weierstraß (Friedman))

Let $f : \omega \rightarrow \mathbb{R}_{>0}$. Let $\langle x_i : i < \omega \rangle$ be an infinite sequence from the closed unit interval $[0, 1]$.

Friedman's BW

Theorem (modified Bolzano Weierstraß (Friedman))

Let $f : \omega \rightarrow \mathbb{R}_{>0}$. Let $\langle x_i : i < \omega \rangle$ be an infinite sequence from the closed unit interval $[0, 1]$.

Then there exist $k_1 < k_2 < \dots$ such that $|x_{k_{i+1}} - x_{k_{i+2}}| < f(k_i)$ holds for all $i \geq 1$.

Friedman's BW

Theorem (modified Bolzano Weierstraß (Friedman))

Let $f : \omega \rightarrow \mathbb{R}_{>0}$. Let $\langle x_i : i < \omega \rangle$ be an infinite sequence from the closed unit interval $[0, 1]$.

Then there exist $k_1 < k_2 < \dots$ such that $|x_{k_{i+1}} - x_{k_{i+2}}| < f(k_i)$ holds for all $i \geq 1$.

Proof: By BW pick a subsequence $\langle x_{m_i} : i < \omega \rangle$ which converges to a limit c .

Friedman's BW

Theorem (modified Bolzano Weierstraß (Friedman))

Let $f : \omega \rightarrow \mathbb{R}_{>0}$. Let $\langle x_i : i < \omega \rangle$ be an infinite sequence from the closed unit interval $[0, 1]$.

Then there exist $k_1 < k_2 < \dots$ such that $|x_{k_{i+1}} - x_{k_{i+2}}| < f(k_i)$ holds for all $i \geq 1$.

Proof: By BW pick a subsequence $\langle x_{m_i} : i < \omega \rangle$ which converges to a limit c . Then construct the indices k_i by recursion.

Friedman's BW

Theorem (modified Bolzano Weierstraß (Friedman))

Let $f : \omega \rightarrow \mathbb{R}_{>0}$. Let $\langle x_i : i < \omega \rangle$ be an infinite sequence from the closed unit interval $[0, 1]$.

Then there exist $k_1 < k_2 < \dots$ such that $|x_{k_{i+1}} - x_{k_{i+2}}| < f(k_i)$ holds for all $i \geq 1$.

Proof: By BW pick a subsequence $\langle x_{m_i} : i < \omega \rangle$ which converges to a limit c . Then construct the indices k_i by recursion.

What is the connection with the following?

Friedman's BW

Theorem (modified Bolzano Weierstraß (Friedman))

Let $f : \omega \rightarrow \mathbb{R}_{>0}$. Let $\langle x_i : i < \omega \rangle$ be an infinite sequence from the closed unit interval $[0, 1]$.

Then there exist $k_1 < k_2 < \dots$ such that $|x_{k_{i+1}} - x_{k_{i+2}}| < f(k_i)$ holds for all $i \geq 1$.

Proof: By BW pick a subsequence $\langle x_{m_i} : i < \omega \rangle$ which converges to a limit c . Then construct the indices k_i by recursion.

What is the connection with the following?

Theorem (Abel)

Let $d \geq 0$. Let $\log^d(\exp^d)$ be the d -fold iteration of $\log(\exp)$ resp.).



Friedman's BW

Theorem (modified Bolzano Weierstraß (Friedman))

Let $f : \omega \rightarrow \mathbb{R}_{>0}$. Let $\langle x_i : i < \omega \rangle$ be an infinite sequence from the closed unit interval $[0, 1]$.

Then there exist $k_1 < k_2 < \dots$ such that $|x_{k_{i+1}} - x_{k_{i+2}}| < f(k_i)$ holds for all $i \geq 1$.

Proof: By BW pick a subsequence $\langle x_{m_i} : i < \omega \rangle$ which converges to a limit c . Then construct the indices k_i by recursion.

What is the connection with the following?

Theorem (Abel)

Let $d \geq 0$. Let $\log^d(\exp^d)$ be the d -fold iteration of $\log(\exp)$ resp.).

► $\int_{\exp^d(1)}^{\infty} \frac{1}{x \cdot \log(x) \cdot \dots \cdot \log^d(x)} dx = +\infty$.



Friedman's BW

Theorem (modified Bolzano Weierstraß (Friedman))

Let $f : \omega \rightarrow \mathbb{R}_{>0}$. Let $\langle x_i : i < \omega \rangle$ be an infinite sequence from the closed unit interval $[0, 1]$.

Then there exist $k_1 < k_2 < \dots$ such that $|x_{k_{i+1}} - x_{k_{i+2}}| < f(k_i)$ holds for all $i \geq 1$.

Proof: By BW pick a subsequence $\langle x_{m_i} : i < \omega \rangle$ which converges to a limit c . Then construct the indices k_i by recursion.

What is the connection with the following?

Theorem (Abel)

Let $d \geq 0$. Let $\log^d(\exp^d)$ be the d -fold iteration of $\log(\exp)$ resp.).

- ▶ $\int_{\exp^d(1)}^{\infty} \frac{1}{x \cdot \log(x) \cdots \log^d(x)} dx = +\infty$.
- ▶ $\int_{\exp^d(1)}^{\infty} \frac{1}{x \cdot \log(x) \cdots (\log^d(x))^{1+\varepsilon}} dx < +\infty$ for $\varepsilon > 0$.



Friedman's miniaturization of BW



Friedman's miniaturization of BW



Theorem (Friedman)

For all $K \geq 3$ there exists M such for all sequences x_1, x_2, \dots, x_M from the closed unit interval $[0, 1]$ there exist $k_1 < \dots < k_K \leq M$ such that $|x_{k_{i+1}} - x_{k_i}| < 1/(k_i)^2$ holds for all $i \leq K - 2$.

Friedman's miniaturization of BW

Theorem (Friedman)

Let $f : \omega \rightarrow \mathbb{R}_{>0}$. The following assertion $FBW(f)$ is true. For all $K \geq 3$ there exists M such for all sequences x_1, x_2, \dots, x_M from the closed unit interval $[0, 1]$ there exist $k_1 < \dots < k_K \leq M$ such that $|x_{k_{i+1}} - x_{k_{i+2}}| < f(k_i)$ holds for all $i \leq K - 2$.

Friedman's miniaturization of BW

Theorem (Friedman)

Let $f : \omega \rightarrow \mathbb{R}_{>0}$. The following assertion $FBW(f)$ is true. For all $K \geq 3$ there exists M such for all sequences x_1, x_2, \dots, x_M from the closed unit interval $[0, 1]$ there exist $k_1 < \dots < k_K \leq M$ such that $|x_{k_{i+1}} - x_{k_{i+2}}| < f(k_i)$ holds for all $i \leq K - 2$.

Friedman's Proof:

Apply the compactness of the Hilbert cube $[0, 1]^\mathbb{N}$ (Tychonov).

Connections with the Ackermann function

Define by recursion on m and subsidiary recursion on n :

Connections with the Ackermann function

Define by recursion on m and subsidiary recursion on n :

$$Ack_0(m) = m + 1. \quad Ack_{n+1}(0) = Ack_n(1) \text{ and}$$

$$Ack_{n+1}(m + 1) = Ack_n(Ack_{n+1}(m)).$$

Connections with the Ackermann function

Define by recursion on m and subsidiary recursion on n :

$$Ack_0(m) = m + 1. \quad Ack_{n+1}(0) = Ack_n(1) \text{ and}$$

$$Ack_{n+1}(m + 1) = Ack_n(Ack_{n+1}(m)).$$

Theorem (Friedman)

Let $f(i) = \frac{1}{i^2}$. For $K \geq 10$, the least $M(K)$ for $FBW(f)$ exceeds $Ack_{K-8}(64)$.

Connections with the Ackermann function

Define by recursion on m and subsidiary recursion on n :

$$Ack_0(m) = m + 1. \quad Ack_{n+1}(0) = Ack_n(1) \text{ and}$$

$$Ack_{n+1}(m + 1) = Ack_n(Ack_{n+1}(m)).$$

Theorem (Friedman)

Let $f(i) = \frac{1}{i^2}$. For $K \geq 10$, the least $M(K)$ for $FBW(f)$ exceeds $Ack_{K-8}(64)$. In particular, the least $M(13)$ exceeds $Ack_5(64)$.

Connections with the Ackermann function

Define by recursion on m and subsidiary recursion on n :

$$Ack_0(m) = m + 1. \quad Ack_{n+1}(0) = Ack_n(1) \text{ and}$$

$$Ack_{n+1}(m + 1) = Ack_n(Ack_{n+1}(m)).$$

Theorem (Friedman)

Let $f(i) = \frac{1}{i^2}$. For $K \geq 10$, the least $M(K)$ for $FBW(f)$ exceeds $Ack_{K-8}(64)$. In particular, the least $M(13)$ exceeds $Ack_5(64)$.

There are also neat Ackermannian upper bounds on $M(K)$.

Connections with the Ackermann function

Define by recursion on m and subsidiary recursion on n :

$$Ack_0(m) = m + 1. \quad Ack_{n+1}(0) = Ack_n(1) \text{ and}$$

$$Ack_{n+1}(m + 1) = Ack_n(Ack_{n+1}(m)).$$

Theorem (Friedman)

Let $f(i) = \frac{1}{i^2}$. For $K \geq 10$, the least $M(K)$ for $FBW(f)$ exceeds $Ack_{K-8}(64)$. In particular, the least $M(13)$ exceeds $Ack_5(64)$.

There are also neat Ackermannian upper bounds on $M(K)$.

What happens when we modify the estimate $1/(k_i)^2$? Obviously any strictly positive function of k_i will make these statements hold, but the issue is the size of the associated constants $M(K)$.

Connections with the Ackermann function

Define by recursion on m and subsidiary recursion on n :

$$Ack_0(m) = m + 1. \quad Ack_{n+1}(0) = Ack_n(1) \text{ and}$$

$$Ack_{n+1}(m+1) = Ack_n(Ack_{n+1}(m)).$$

Theorem (Friedman)

Let $f(i) = \frac{1}{i^2}$. For $K \geq 10$, the least $M(K)$ for $FBW(f)$ exceeds $Ack_{K-8}(64)$. In particular, the least $M(13)$ exceeds $Ack_5(64)$.

There are also neat Ackermannian upper bounds on $M(K)$.

What happens when we modify the estimate $1/(k_i)^2$? Obviously any strictly positive function of k_i will make these statements hold, but the issue is the size of the associated constants $M(K)$.

If we use the estimate $\frac{1}{(k_i)^{1+\varepsilon}}$, where $\varepsilon > 0$, then (following

Friedman unpublished) we obtain roughly the same Ackermannian upper and lower bounds.

Friedman's miniaturization of BW

Theorem (Friedman)

Let $f(i) := \log(i)/i$. Then the least $M(K)$ for $FBW(f)$ is bounded by an exponential function.

Friedman's miniaturization of BW

Theorem (Friedman)

Let $f(i) := \log(i)/i$. Then the least $M(K)$ for $FBW(f)$ is bounded by an exponential function.

Proof. By a simple bisection argument.

Friedman's miniaturization of BW

Theorem (Friedman)

Let $f(i) := \log(i)/i$. Then the least $M(K)$ for $FBW(f)$ is bounded by an exponential function.

Proof. By a simple bisection argument.

Friedman's challenge from 1999: How does $M(K)$ behave for $f(i) = \frac{1}{i}$ and $f(i) = \frac{1}{i \log(i)}$?

Friedman's miniaturization of BW

Theorem (Friedman)

Let $f(i) := \log(i)/i$. Then the least $M(K)$ for $FBW(f)$ is bounded by an exponential function.

Proof. By a simple bisection argument.

Friedman's challenge from 1999: How does $M(K)$ behave for $f(i) = \frac{1}{i}$ and $f(i) = \frac{1}{i \log(i)}$?

Question: Can we improve on the Ackermannian upper bounds for $f(i) = \frac{1}{i}$ and $f(i) = \frac{1}{i \log(i)}$?

Friedman's monotone convergence principle

Assume that f is strictly positive. Let $FMC(f)$ be the following assertion: For all $K \geq 3$ there exists M such for all weakly increasing sequences $x_1 \leq x_2 \leq \dots \leq x_M$ from the closed unit interval $[0, 1]$ there exist $k_1 < \dots < k_K \leq M$ such that $|x_{k_{i+1}} - x_{k_{i+2}}| < f(k_i)$ holds for all $i \leq K - 2$.

Friedman's monotone convergence principle

Assume that f is strictly positive. Let $FMC(f)$ be the following assertion: For all $K \geq 3$ there exists M such for all weakly increasing sequences $x_1 \leq x_2 \leq \dots \leq x_M$ from the closed unit interval $[0, 1]$ there exist $k_1 < \dots < k_K \leq M$ such that $|x_{k_{i+1}} - x_{k_{i+2}}| < f(k_i)$ holds for all $i \leq K - 2$.

Theorem (W.)

- ▶ Let $f(i) = \frac{1}{i \log(i)}$. Then the least $M(K)$ in $FMC(f)$ is bounded from above by a double exponential function.

Friedman's monotone convergence principle

Assume that f is strictly positive. Let $FMC(f)$ be the following assertion: For all $K \geq 3$ there exists M such for all weakly increasing sequences $x_1 \leq x_2 \leq \dots \leq x_M$ from the closed unit interval $[0, 1]$ there exist $k_1 < \dots < k_K \leq M$ such that $|x_{k_{i+1}} - x_{k_{i+2}}| < f(k_i)$ holds for all $i \leq K - 2$.

Theorem (W.)

- ▶ Let $f(i) = \frac{1}{i \log(i)}$. Then the least $M(K)$ in $FMC(f)$ is bounded from above by a double exponential function.
- ▶ Let $f(i) = \frac{1}{i(\log(i))^{1+\varepsilon}}$. Then the least $M(K)$ in $FMC(f)$ is Ackermannian.

Friedman's miniaturization of BW

Theorem (W.)

Fix $d < \omega$. Let $\log^d(i)$ denote the d -th iterate of \log .

- ▶ Let $f(i) = \frac{1}{i \cdot \log(i) \cdot \dots \cdot \log^d(i)}$. Then the least $M(K)$ in $FMC(f)$ is bounded by a $d + 1$ -times iterated exponential function.

Friedman's miniaturization of BW

Theorem (W.)

Fix $d < \omega$. Let $\log^d(i)$ denote the d -th iterate of \log .

- ▶ Let $f(i) = \frac{1}{i \cdot \log(i) \cdot \dots \cdot \log^d(i)}$. Then the least $M(K)$ in $FMC(f)$ is bounded by a $d + 1$ -times iterated exponential function.
- ▶ Let $f_d(i) = A_d^{-1}(i) - A_d^{-1}(i - 1)$ and where A_d refers to the d -th branch of the Ackermann function, and $^{-1}$ refers to taking functional inverses.

Friedman's miniaturization of BW

Theorem (W.)

Fix $d < \omega$. Let $\log^d(i)$ denote the d -th iterate of \log .

- ▶ Let $f(i) = \frac{1}{i \cdot \log(i) \cdot \dots \cdot \log^d(i)}$. Then the least $M(K)$ in $FMC(f)$ is bounded by a $d + 1$ -times iterated exponential function.
- ▶ Let $f_d(i) = A_d^{-1}(i) - A_d^{-1}(i - 1)$ and where A_d refers to the d -th branch of the Ackermann function, and $^{-1}$ refers to taking functional inverses. Then the least $M(K)$ in $FMC(f)$ is bounded by a primitive recursive function.

Friedman's miniaturization of BW

Theorem (W.)

Fix $d < \omega$. Let $\log^d(i)$ denote the d -th iterate of \log .

- ▶ Let $f(i) = \frac{1}{i \cdot \log(i) \cdot \dots \cdot \log^d(i)}$. Then the least $M(K)$ in $FMC(f)$ is bounded by a $d + 1$ -times iterated exponential function.
- ▶ Let $f_d(i) = A_d^{-1}(i) - A_d^{-1}(i - 1)$ and where A_d refers to the d -th branch of the Ackermann function, and $^{-1}$ refers to taking functional inverses. Then the least $M(K)$ in $FMC(f)$ is bounded by a primitive recursive function.
- ▶ Let $f(i) = \frac{1}{i \cdot \log(i) \cdot \dots \cdot (\log^d(i))^{1+\varepsilon}}$. Then the least $M(K)$ in $FMC(f)$ is Ackermannian.

Friedman's miniaturization of BW

Theorem (W.)

Fix $d < \omega$. Let $\log^d(i)$ denote the d -th iterate of \log .

- ▶ Let $f(i) = \frac{1}{i \cdot \log(i) \cdot \dots \cdot \log^d(i)}$. Then the least $M(K)$ in $FMC(f)$ is bounded by a $d + 1$ -times iterated exponential function.
- ▶ Let $f_d(i) = A_d^{-1}(i) - A_d^{-1}(i - 1)$ and where A_d refers to the d -th branch of the Ackermann function, and $^{-1}$ refers to taking functional inverses. Then the least $M(K)$ in $FMC(f)$ is bounded by a primitive recursive function.
- ▶ Let $f(i) = \frac{1}{i \cdot \log(i) \cdot \dots \cdot (\log^d(i))^{1+\varepsilon}}$. Then the least $M(K)$ in $FMC(f)$ is Ackermannian.



Solution to Friedman's problem

Theorem (W.)

Fix $d < \omega$. Let $\log^d(i)$ denote the d -th iterate of \log .

- ▶ Let $f(i) = \frac{1}{i \cdot \log(i) \cdot \dots \cdot \log^d(i)}$. Then the least $M(K)$ in $FBW(f)$ is bounded by a $d + 2$ -times iterated exponential function. This already answers Friedman's question.

Solution to Friedman's problem

Theorem (W.)

Fix $d < \omega$. Let $\log^d(i)$ denote the d -th iterate of \log .

- ▶ Let $f(i) = \frac{1}{i \cdot \log(i) \cdot \dots \cdot \log^d(i)}$. Then the least $M(K)$ in $FBW(f)$ is bounded by a $d + 2$ -times iterated exponential function. This already answers Friedman's question.
- ▶ Let $f_d(i) = A_d^{-1}(i) - A_d^{-1}(i - 1)$ and where A_d refers to the d -th branch of the Ackermann function, and $^{-1}$ refers to taking functional inverses.

Solution to Friedman's problem

Theorem (W.)

Fix $d < \omega$. Let $\log^d(i)$ denote the d -th iterate of \log .

- ▶ Let $f(i) = \frac{1}{i \cdot \log(i) \cdot \dots \cdot \log^d(i)}$. Then the least $M(K)$ in $FBW(f)$ is bounded by a $d + 2$ -times iterated exponential function. This already answers Friedman's question.
- ▶ Let $f_d(i) = A_d^{-1}(i) - A_d^{-1}(i - 1)$ and where A_d refers to the d -th branch of the Ackermann function, and $^{-1}$ refers to taking functional inverses. Then the least $M(K)$ in $FBW(f)$ is bounded by a primitive recursive function. This answers Friedman's questions completely for the lower bounds.

Solution to Friedman's problem

Theorem (W.)

Fix $d < \omega$. Let $\log^d(i)$ denote the d -th iterate of \log .

- ▶ Let $f(i) = \frac{1}{i \cdot \log(i) \cdot \dots \cdot \log^d(i)}$. Then the least $M(K)$ in $FBW(f)$ is bounded by a $d + 2$ -times iterated exponential function. This already answers Friedman's question.
- ▶ Let $f_d(i) = A_d^{-1}(i) - A_d^{-1}(i-1)$ and where A_d refers to the d -th branch of the Ackermann function, and $^{-1}$ refers to taking functional inverses. Then the least $M(K)$ in $FBW(f)$ is bounded by a primitive recursive function. This answers Friedman's questions completely for the lower bounds.
- ▶ Let $f(i) = \frac{1}{i \cdot \log(i) \cdot \dots \cdot \log_d(i)^{1+\varepsilon}}$. Then the least $M(K)$ in $FBW(f)$ is Ackermannian.

Solution to Friedman's problem

Theorem (W.)

Fix $d < \omega$. Let $\log^d(i)$ denote the d -th iterate of \log .

- ▶ Let $f(i) = \frac{1}{i \cdot \log(i) \cdot \dots \cdot \log^d(i)}$. Then the least $M(K)$ in $FBW(f)$ is bounded by a $d + 2$ -times iterated exponential function. This already answers Friedman's question.
- ▶ Let $f_d(i) = A_d^{-1}(i) - A_d^{-1}(i-1)$ and where A_d refers to the d -th branch of the Ackermann function, and $^{-1}$ refers to taking functional inverses. Then the least $M(K)$ in $FBW(f)$ is bounded by a primitive recursive function. This answers Friedman's questions completely for the lower bounds.
- ▶ Let $f(i) = \frac{1}{i \cdot \log(i) \cdot \dots \cdot \log_d(i)^{1+\varepsilon}}$. Then the least $M(K)$ in $FBW(f)$ is Ackermannian.

Hint for the proof of the first two assertions: Do the counting argument for $FMC(f)$ but now with nested layers.

Mathematical spin off

General question: What is $A_d^{-1}(l) - A_d^{-1}(l-1)$?

Mathematical spin off

General question: What is $A_d^{-1}(I) - A_d^{-1}(I - 1)$? Thinking of the intermediate value theorem one might ask: What is the derivative of (a smoothed version of) $I \mapsto A_d^{-1}(I)$?

Mathematical spin off

General question: What is $A_d^{-1}(I) - A_d^{-1}(I - 1)$? Thinking of the intermediate value theorem one might ask: What is the derivative of (a smoothed version of) $I \mapsto A_d^{-1}(I)$?

Let $F_0(x) = 2^x$ and $F_{d+1}(x) = F_d^x(x)$.

Mathematical spin off

General question: What is $A_d^{-1}(I) - A_d^{-1}(I-1)$? Thinking of the intermediate value theorem one might ask: What is the derivative of (a smoothed version of) $I \mapsto A_d^{-1}(I)$?

Let $F_0(x) = 2^x$ and $F_{d+1}(x) = F_d^x(x)$.

Let $D_0(x) = 1/x$ and $D_{d+1}(x) = \prod_{i=1}^{F_{d+1}^{-1}(x)} D_d((F_d^{-1})^i(x))$.

Mathematical spin off

General question: What is $A_d^{-1}(I) - A_d^{-1}(I-1)$? Thinking of the intermediate value theorem one might ask: What is the derivative of (a smoothed version of) $I \mapsto A_d^{-1}(I)$?

Let $F_0(x) = 2^x$ and $F_{d+1}(x) = F_d^x(x)$.

Let $D_0(x) = 1/x$ and $D_{d+1}(x) = \prod_{i=1}^{F_{d+1}^{-1}(x)} D_d((F_d^{-1})^i(x))$.

Theorem (Strong extension of Abel's result)

Fix $d < \omega$ and $\varepsilon > 0$.

Mathematical spin off

General question: What is $A_d^{-1}(I) - A_d^{-1}(I-1)$? Thinking of the intermediate value theorem one might ask: What is the derivative of (a smoothed version of) $I \mapsto A_d^{-1}(I)$?

Let $F_0(x) = 2^x$ and $F_{d+1}(x) = F_d^x(x)$.

Let $D_0(x) = 1/x$ and $D_{d+1}(x) = \prod_{i=1}^{F_{d+1}^{-1}(x)} D_d((F_d^{-1})^i(x))$.

Theorem (Strong extension of Abel's result)

Fix $d < \omega$ and $\varepsilon > 0$.

- ▶ $\sum_i^\infty D_d(i) = \infty$.

Mathematical spin off

General question: What is $A_d^{-1}(I) - A_d^{-1}(I-1)$? Thinking of the intermediate value theorem one might ask: What is the derivative of (a smoothed version of) $I \mapsto A_d^{-1}(I)$?

Let $F_0(x) = 2^x$ and $F_{d+1}(x) = F_d^x(x)$.

Let $D_0(x) = 1/x$ and $D_{d+1}(x) = \prod_{i=1}^{F_{d+1}^{-1}(x)} D_d((F_d^{-1})^i(x))$.

Theorem (Strong extension of Abel's result)

Fix $d < \omega$ and $\varepsilon > 0$.

- ▶ $\sum_i^\infty D_d(i) = \infty$.
- ▶ $\sum_i^\infty (D_d(i) \cdot \frac{1}{(F_{d+1}^{-1}(i))^{1+\varepsilon}}) < \infty$.

Mathematical spin off

General question: What is $A_d^{-1}(I) - A_d^{-1}(I-1)$? Thinking of the intermediate value theorem one might ask: What is the derivative of (a smoothed version of) $I \mapsto A_d^{-1}(I)$?

Let $F_0(x) = 2^x$ and $F_{d+1}(x) = F_d^x(x)$.

Let $D_0(x) = 1/x$ and $D_{d+1}(x) = \prod_{i=1}^{F_{d+1}^{-1}(x)} D_d((F_d^{-1})^i(x))$.

Theorem (Strong extension of Abel's result)

Fix $d < \omega$ and $\varepsilon > 0$.

- ▶ $\sum_i^\infty D_d(i) = \infty$.
- ▶ $\sum_i^\infty (D_d(i) \cdot \frac{1}{(F_{d+1}^{-1}(i))^{1+\varepsilon}}) < \infty$.

Conjecture

$A_d^{-1}(I) - A_d^{-1}(I-1)$ will be of order $D_d(I)$ as I tends to infinity.
This expression will approximate the derivative of A_d^{-1} .

We are almost there

Let $F_0(x) = 2^x$ and $F_{d+1}(x) = F_d^x(x)$.

Let $D_0(x) = 1/x$ and $D_{d+1}(x) = \prod_{i=1}^{F_{d+1}^{-1}(x)} D_d((F_d^{-1})^i(x))$.

We are almost there

Let $F_0(x) = 2^x$ and $F_{d+1}(x) = F_d^x(x)$.

Let $D_0(x) = 1/x$ and $D_{d+1}(x) = \prod_{i=1}^{F_{d+1}^{-1}(x)} D_d((F_d^{-1})^i(x))$.

For $d^* : \omega \rightarrow \omega$ non decreasing let $D^*(x) := D_{d^*(x)}(x)$.

We are almost there

Let $F_0(x) = 2^x$ and $F_{d+1}(x) = F_d^x(x)$.

Let $D_0(x) = 1/x$ and $D_{d+1}(x) = \prod_{i=1}^{F_{d+1}^{-1}(x)} D_d((F_d^{-1})^i(x))$.

For $d^* : \omega \rightarrow \omega$ non decreasing let $D^*(x) := D_{d^*(x)}(x)$.

Conjecture

Let $d^ : \omega \rightarrow \omega$ be non decreasing and definable in one quantifier induction. Then $FBW(D^*)$ is unprovable with one quantifier induction ($I\Sigma_1$) iff d^* is unbounded.*

We are almost there

Let $F_0(x) = 2^x$ and $F_{d+1}(x) = F_d^x(x)$.

Let $D_0(x) = 1/x$ and $D_{d+1}(x) = \prod_{i=1}^{F_{d+1}^{-1}(x)} D_d((F_d^{-1})^i(x))$.

For $d^* : \omega \rightarrow \omega$ non decreasing let $D^*(x) := D_{d^*(x)}(x)$.

Conjecture

Let $d^ : \omega \rightarrow \omega$ be non decreasing and definable in one quantifier induction. Then $\text{FBW}(D^*)$ is unprovable with one quantifier induction ($I\Sigma_1$) iff d^* is unbounded.*

Left to be done: Classify the phase transition for FBW version of strength PA. FOM 913.

We are almost there

Let $F_0(x) = 2^x$ and $F_{d+1}(x) = F_d^x(x)$.

Let $D_0(x) = 1/x$ and $D_{d+1}(x) = \prod_{i=1}^{F_{d+1}^{-1}(x)} D_d((F_d^{-1})^i(x))$.

For $d^* : \omega \rightarrow \omega$ non decreasing let $D^*(x) := D_{d^*(x)}(x)$.

Conjecture

Let $d^ : \omega \rightarrow \omega$ be non decreasing and definable in one quantifier induction. Then $\text{FBW}(D^*)$ is unprovable with one quantifier induction ($I\Sigma_1$) iff d^* is unbounded.*

Left to be done: Classify the phase transition for FBW version of strength PA. FOM 913.

Also the phase transition for Abel's theorem could be refined by looking at a suitably defined D_α .

New exiting developments

By recent developments (emerging from a discussion with Aschenbrenner) the question on the derivative A_d^{-1} seems to be related to the world of transseries and surreal numbers and also to the work by Écalle. For transseries we have $\log_\alpha = \int \frac{1}{\prod_{\beta < \alpha} \log_\beta} dx$ where α is an ordinal. Our D_d seems to correspond roughly to \log_{ω^d} .

Logic/Mathematics at UGent

If you want to learn more details please send me an email under
Andreas.Weiermann at UGent.be.

Logic/Mathematics at UGent

If you want to learn more details please send me an email under Andreas.Weiermann at UGent.be.

Students, PhD students and Postdocs are welcome in our team in Ghent! For funding options please consult:

<https://www.research.ugent.be>

Logic/Mathematics at UGent

If you want to learn more details please send me an email under Andreas.Weiermann at UGent.be.

Students, PhD students and Postdocs are welcome in our team in Ghent! For funding options please consult:

<https://www.research.ugent.be>

Also institutional collaboration is welcome!

Logic/Mathematics at UGent

If you want to learn more details please send me an email under Andreas.Weiermann at UGent.be.

Students, PhD students and Postdocs are welcome in our team in Ghent! For funding options please consult:

<https://www.research.ugent.be>

Also institutional collaboration is welcome!

Here is by the way our logic group:

