


# Modeled estimates of the health outcomes and economic value of improving the social determinants of mental health

Received: 17 December 2024

Accepted: 4 June 2025

Published online: 24 July 2025

 Check for updates

Paul Crosland<sup>1</sup>✉, Nicholas Ho<sup>1</sup>, Kim-Huong Nguyen<sup>1,2,3</sup>, Kristen Tran<sup>1</sup>, Seyed Hossein Hosseini<sup>1</sup>, Catherine Vacher<sup>1</sup>, Adam Skinner<sup>1</sup>, Jordan van Rosmalen<sup>1</sup>, Sebastian Rosenberg<sup>1,4</sup>, Frank Iorfino<sup>1</sup>, Victoria Loblay<sup>1</sup>, Olivia Iannelli<sup>1</sup>, Sarah Piper<sup>1</sup>, Yun Ju Christine Song<sup>1</sup>, Sophie Morson<sup>5</sup>, Judith M. G. Piccone<sup>6</sup>, Adam Connell<sup>7</sup>, Deborah A. Marshall<sup>8</sup>, Ian B. Hickie<sup>1,10</sup> & Jo-An Occhipinti<sup>1,9,10</sup>

The prevalence and burden of mental disorders have worsened despite increased community awareness. Enhanced access to treatments alone is unlikely to deliver improvements in population mental health, so more attention needs to be paid to social and environmental influences. Here we estimate the health benefits and economic value of improving the social determinants of mental health within Brisbane South, a diverse population of 1.2 million people, in Australia. The incremental net monetary benefit (combining costs and monetized health outcomes) derived from 5% improvements in the average yearly change of social cohesion, childhood difficulties, substance misuse and unemployment over 11 years from 2024 to 2034 was projected to be AUD\$146.64 million, AUD\$234.50 million, AUD\$281.67 million and AUD\$100.43 million, respectively. Quality-adjusted life years, suicide deaths, emergency department presentations and self-harm hospitalizations were also improved. This study demonstrates the health and economic value of investing in the social determinants of mental health.

Despite greater community awareness about mental health, the prevalence and burden associated with mental disorders are deteriorating<sup>1–4</sup>. Depressive disorders were the second-highest cause of nonfatal health burden globally in 2021, an increase in years lived with disability of 36.5% from 2010<sup>5</sup>. Anxiety disorders were the sixth-highest cause of nonfatal burden<sup>5</sup>. Young people, in particular, have experienced a serious deterioration in mental health in the past 14 years. For example, a longitudinal survey found that the prevalence of depression and anxiety more than doubled between 2009 and 2021 in Australian young people aged 15–34 years (from 6.1% to 14.4% for males and from 12.7% to 29.3% for females)<sup>6</sup>. The prevalence of psychological distress also more than doubled between 2011 and 2021 in young people in Australia aged 15–24 years (from 18.4% to 42.3%)<sup>6</sup>. Suicide was the leading cause

of death among people aged 15–44 in 2022 in Australia<sup>7</sup>. Although the decline in the mental health of younger age groups has been underway for more than a decade, the coronavirus disease 2019 pandemic and associated lockdowns have influenced these estimates in recent years<sup>8</sup>. Increases in the provision of mental health care have been outweighed by a concurrent increase in the incidence of high to very high levels of distress, as demonstrated by simulation modeling<sup>9</sup>. The Australian experience is replicated across most high-income countries<sup>10,11</sup> and is also an emerging threat in developing nations<sup>12</sup>.

There are long-term consequences of young people experiencing mental disorders, for individuals and their families, for communities and for the whole economy. These disorders are a risk factor for a wide range of chronic physical illnesses<sup>13</sup> and affect their ability to participate

A full list of affiliations appears at the end of the paper. ✉e-mail: [paul.crosland@sydney.edu.au](mailto:paul.crosland@sydney.edu.au)

in the labor force later in life<sup>14</sup>. Some of this burden could be averted if young people had full access to optimal treatment<sup>15</sup>. However, expanding access to mental health treatments alone is insufficient to reduce the burden of poor mental health due to the rising incidence of disorders<sup>9,16</sup>, which occur in—and interact with—the context of wider social, economic, commercial and environmental factors. More meaningful, long-term, intergenerational reductions in mental illness are thus likely to be achieved by ‘moving upstream’, targeting the ‘causes of the causes’ and preventing mental ill health before it develops<sup>17,18</sup>. This could be achieved by a more intentional approach to choosing mental health and well-being as a key policy objective (which can be formally conceptualized as Mental Wealth<sup>19</sup>) and designing and implementing evidence-based economic and social systems and environments that foster this objective as the norm<sup>17,20</sup>.

Economic implications are interwoven through the causes and consequences of mental disorders. The global economic burden of mental disorders was estimated to be between US\$2.5 trillion and US\$8.5 trillion in 2010 depending on the methodological approach used to monetize health impacts<sup>21</sup>. The more conservative approach is expected to reach annual costs of US\$6 trillion by 2030<sup>21</sup>. In the USA, more health expenditure is dedicated to treating mental health disorders than any other disease area<sup>22</sup>. The persistent nature of mental health and substance use disorders in the USA, and other countries, suggests this expenditure is not currently allocated efficiently. Furthermore, most of the economic burden of poor mental health is incurred outside the health care system<sup>23</sup>. For example, the Australian Productivity Commission estimated there were AUD\$39 billion in productivity costs associated with poor mental health and suicide in 2018–2019 compared with AUD\$16 billion of mental health-related healthcare expenditure<sup>24</sup>. Estimates of the economic burden of mental illness are a part of the prioritization problem facing decision-makers (that is ‘How big is the problem?’); another is economic evidence supporting interventions (that is, ‘What can be done about it?’). Although much evidence exists on the cost-effectiveness of treatments<sup>25,26</sup> and the prevention of mental ill health and promotion of mental health<sup>27,28</sup>, many gaps still exist<sup>29–31</sup>, including strategies that target the social determinants of mental health. Evidence on the economic credentials of current expenditure is also lacking<sup>32</sup>, and accountability in mental health decision-making and planning is poor<sup>33</sup>. Generating more and better economic evidence is critical for guiding better investment decisions and helping decision-makers prioritize the areas and strategies that can have the biggest impact for improving mental health<sup>34</sup>.

The Lancet Commission on Global Mental Health and Sustainable Development recognized that treatments alone are unlikely to achieve sufficient improvements in population mental health and that more attention needs to be paid to social and environmental influences, particularly during important developmental periods in the early life course, childhood and adolescence<sup>12</sup>. The social determinants of mental health refer to the social, economic and physical environments that directly influence the incidence, prevalence and severity of mental illness<sup>35,36</sup>. Examples of the structural risk factors that influence mental health include low educational attainment, unemployment and underemployment, poverty, food insecurity, unstable housing, social isolation and loneliness, discrimination, early life and childhood adversity and trauma, neighborhood social and physical conditions, and access to best practice and affordable health care<sup>37,38</sup>. These determinants can play out in regionally specific ways to impact population mental health and well-being. They often have a bidirectional relationship with mental health and are interlinked and dynamic, interacting with each other in a complex causal web, highlighting the importance of systems thinking when developing strategies to overcome them<sup>17</sup>. Most of these risk factors are associated with a social gradient based on socioeconomic disadvantage and inequality, whereby “the greater the inequality the higher the inequality in risk”<sup>35</sup>. There is also increasing awareness of the role of ‘global megatrends’ in harming the mental health of young

people<sup>11</sup> as well as the ongoing shadow cast by the coronavirus disease pandemic. These megatrends are broader concepts than social determinants, referring to long-lasting societal upheaval in environmental, social, economic, political or technological domains. Nevertheless, they overlap with the social determinants discussed here.

Evidence exists supporting the causal link between social determinants and later mental health outcomes<sup>37</sup>. The importance of different social determinants varies across the life course<sup>37</sup>. An umbrella review of 46 meta-analyses found that numerous social determinants, such as midlife unemployment, homelessness and interaction with the criminal justice system, had consistent associations with suicide-related outcomes<sup>39</sup>. Another umbrella review included 289 systematic reviews on the evidence for potential mechanisms by which social determinants affect mental health conditions<sup>16</sup>. This evidence was mapped to the United Nations Sustainable Development Goals using five domains: demographic, economic, neighborhood, environmental events, and social and cultural domains<sup>16</sup>. The World Health Organization recently issued new guidance on mental health policy and strategic action plans that has a strong emphasis on social and structural determinants<sup>40</sup>. Despite this evidence, social determinants of mental health are rarely addressed within child and youth health policies and services planning<sup>41</sup>, and the social determinants of mental health are either not improving or getting worse over time<sup>11,32</sup>. Investing in child and youth mental health has the potential to optimize the return on investment of scarce healthcare resources, given the potential for major intergenerational impacts that stretch into adulthood and beyond.

The Lancet Psychiatry Commission on Youth Mental Health specify enhancing awareness and advocacy of the social and economic determinants of mental ill health as the first key element of optimal youth mental health care<sup>11</sup>. However, in the context of scarcity and budgetary constraint, decision-makers require guidance on which social determinants should be prioritized, and what specific interventions can be implemented to improve them. This is one of the reasons why prominent reviews of the social determinants of mental health call for the generation of more economic evidence in this area<sup>36,42</sup>.

The objective of this Analysis was to estimate the health benefits and economic value of improving the social determinants of mental health for the Brisbane South region in Australia using system dynamics simulation modeling. To enhance the validity, transparency and usability of the Analysis, participatory model-building processes were followed, including meaningful participation of young people with lived experience of mental health conditions. This makes the Analysis relevant to the multifaceted nature of the problem, which involves diverse stakeholders engaging with the issue in real-world contexts. Results were reported for a variety of health outcomes and economic summary measures, and the cost impacts were disaggregated by payer. Only a selection of social determinants have been included in this Analysis. These were priority determinants identified by stakeholders of the region as key drivers of youth mental health outcomes and where sufficient data could be obtained.


## Results

All costs are reported in 2020–2021 Australian dollars. All estimates relate to all ages of the Brisbane South Primary Health Network region, a population of 1.2 million people (4% of the Australian population), and are cumulative over the 11-year time horizon of the model (2024–2034) unless otherwise specified. Estimates are incremental compared with business as usual. Costs and quality-adjusted life years (QALYs) were discounted at 5%. Table 1 contains a definition of each social determinant and the specific changes that were applied for each scenario of improvement. ‘Improvement’ in this context could mean a reduction in the rate at which people develop substance misuse disorder or an increase in social cohesion, for example (Table 1).


The greatest increase in health outcomes in terms of QALYs was attained by a 5% improvement in childhood difficulties (2,621 QALYs), followed by social cohesion (1,309 QALYs) and substance misuse (82

**Table 1 | Defining the social determinants and scenarios of improvement**

Social determinant	Description as defined for the present model	Source of historical data	What a '5% improvement' means for each scenario (additional detail provided per age group provided in Supplementary Information part B)
Social cohesion	Based on the Scanlon–Monash Index of Social Cohesion, this represents the willingness of members of society to cooperate with each other in order to survive and prosper. There are five domains this index seeks to capture: belonging; worth; social justice and equity; political participation; and acceptance and rejection <sup>75</sup> . The five domains are transformed to a single composite index using factor analysis and assigning weights to each indicator, with 2007 as the reference year assigned the starting index of 100. Further detail on the methods used by the Scanlon–Monash research group can be found in a report by Markus et al. <sup>76</sup> , and <a href="https://www.monash.edu/mapping-population">https://www.monash.edu/mapping-population</a> . The average annual decline between 2011 and 2023 in the Scanlon–Monash Index was by 0.788 units, and this was assumed to continue each year of the model timeframe in a linear fashion as part of the business-as-usual scenario.	Scanlon–Monash Index of Social Cohesion <sup>75</sup>	The social cohesion index continues to decline in future years but at a reduced rate by applying a multiplier of 0.95 to the baseline average annual rate of change from January 2024 onward. Each year, the index will decrease by 0.7486 ( $0.788 \times 0.95$ ).
Early life and childhood behavioral and emotional difficulties (shortened to 'childhood difficulties' in text, tables and figures)	Childhood difficulties (early life and childhood behavioral and emotional difficulties) refers to the outcome of a brief screening questionnaire (SDQ) that measures behavioral and emotional difficulties in children and young people <sup>67</sup> . In our model, the population aged less than 12 years is partitioned to one of three levels of distress: 'close to average', 'slightly raised' or 'high' SDQ levels. Higher levels of SDQ can be an indicator of early life difficulties, adverse experiences or trauma. Rates by which people flow between the three levels of SDQ are dependent on age, rates of engagement with the mental health services system, rates of treatment and the population levels of social cohesion.	The Longitudinal Study of Australian Children, Growing up in Australia <sup>77,78</sup>	The rate at which children transition to higher levels of SDQ was reduced by applying a multiplier of 0.95 from January 2024 onward.
Homelessness	Following the Australian Bureau of Statistics (ABS), homelessness refers to a state in which an individual does not have a permanent living arrangement, no fixed place of residence, and no suitable or adequate accommodation alternatives <sup>79</sup> . People aged 15 and older enter homelessness at rates that are dependent on age, levels of psychological distress, unemployment and substance misuse. For people under the age of 15 years, rates of entering homelessness are dependent on age.	ABS <sup>80</sup>	The rate at which people enter homelessness was reduced by applying a multiplier of 0.95 from January 2024 onward.
Unemployment	Following the ABS, unemployment is defined as a state in which an individual who is of working age and ability, and who is actively seeking and available for work, is not employed in any paid work or self-employed work. People can transition between the states of being sufficiently employed, underemployed, unemployed or not-in-the-labor-force.	ABS <sup>81–83</sup>	The rates at which people enter unemployment from being sufficiently employed or underemployed was reduced by applying a multiplier of 0.95 from January 2024 onward.
Discontinue secondary education	Discontinuing secondary education refers to when a student enrolled in any high school grade between grade 7 and 12 withdraws their enrollment. Rates of discontinuation are dependent on levels of psychological distress in the population aged 12–17 years.	Australian Curriculum, Assessment and Reporting Authority (ACARA) <sup>84</sup>	The rate at which students discontinue secondary education was reduced by applying a multiplier of 0.95 from January 2024 onward.
Discontinue tertiary education	Discontinuing tertiary education refers to when an individual attending education for a qualification at certificate III or above withdraws their enrollment. Rates of discontinuation are influenced by psychological distress.	ABS <sup>83,85</sup>	The rate at which students discontinued tertiary education was reduced by applying a multiplier of 0.95 from January 2024 onward.
Substance misuse	We followed the ABS' definition of 12-month substance use disorder, and here, we refer to this as substance misuse. Substance misuse refers to the use of alcohol and other drugs in a way that produces harm to the consumer of the substance and others. Harm associated with substance misuse can include a deterioration of an individual's mental and physical health, financial state, social connectedness, education and employment. The rate at which people develop substance misuse is dependent on homelessness, levels of young people aged 15–24 not in education nor employment, levels of psychological distress and recovery from substance misuse treatment services.	ABS <sup>86,87</sup>	The rates at which people developed 12-month substance use disorder was reduced by applying a multiplier of 0.95 from January 2024 onward.
Underemployment	Based on the ABS' definition, underemployment refers to a state in which an individual is employed in paid work but has indicated that they would prefer to work more hours. People can transition between the states of being sufficiently employed, underemployed, unemployed or not-in-the-labor-force.	ABS <sup>81–83</sup>	The rates at which people transition from being sufficiently employed into underemployment were reduced by applying a multiplier of 0.95 from January 2024 onward.



**Fig. 1 | Incremental QALYs, 5% improvement in social determinants versus business as usual.** Cumulative QALYs over 11 years, 2024–2034, discounted, versus business as usual.




**Fig. 2 | Incremental cost (health care perspective), 5% improvement in social determinants versus business as usual.** Cumulative costs over 11 years, 2024–2034, discounted, 2020–2021 Australian dollars versus business as usual. ed., education.

QALYs) (Fig. 1; Supplementary Table 3 in Supplementary Information part B). A 5% improvement in the other determinants was also associated with an increase in QALYs but to a lesser extent. Reducing substance misuse and improving social cohesion individually resulted in the greatest number of suicide deaths avoided (Supplementary Table 3 in Supplementary Information part B). Reducing substance misuse, improving social cohesion and reducing childhood difficulties individually realized the greatest number of avoided self-harm hospitalizations. Improving childhood difficulties and social cohesion avoided the greatest number of mental health-related emergency department presentations.

Improving the social determinants of mental health substantially reduced downstream costs. For the health care perspective, the greatest

reduction in costs was achieved by a 5% improvement in homelessness (\$16.4 million), followed by social cohesion (\$16.3 million) and childhood difficulties (\$11.7 million) (Fig. 2; Supplementary Table 1 in Supplementary Information part B). A 5% improvement in substance misuse resulted in \$1 million of downstream cost savings. Improvements in unemployment, underemployment and discontinuation of secondary or tertiary education resulted in cost reductions less than this.

For the societal perspective, the greatest reductions in costs were seen by 5% improvements in substance misuse (\$274.8 million), unemployment (\$97.3 million), social cohesion (\$38 million) and homelessness (\$38 million) (Fig. 3; Supplementary Table 1 in Supplementary Information part B). The downstream cost savings



**Fig. 3 | Incremental cost (societal perspective), 5% improvement in social determinants versus business as usual.** Cumulative costs over 11 years, 2024–2034, discounted, 2020–2021 Australian dollars versus business as usual. ed., education.

were attributed largely to a reduction in lost productivity. These productivity savings amounted to \$157.3 million for substance misuse, \$95 million for unemployment, and \$20.6 million for social cohesion, for the societal perspective.

When analyzing cost impacts by payer for the societal perspective, homelessness (\$13.6 million), substance misuse (\$10.4 million) and social cohesion (\$7.3 million) were the most important determinants for the Australian (National) Government. Five percent improvements in substance misuse (\$107.8 million), homelessness (\$19.1 million), social cohesion (\$16.3 million) and childhood difficulties (\$11.5 million) achieved the greatest reductions in costs for the Queensland (State) Government. Homelessness was the only determinant that reduced costs for nongovernment payers, which predominantly consists of individuals in the form of out-of-pocket costs. Improvements in all other determinants increased costs for nongovernment payers. (This was mainly due to a reduction in welfare payments received by this group. Changes in these transfer payments result in a decrease in costs to government and zero net impact on total costs. We have not included employment income, nor income tax, in this Analysis, which would be expected to more than offset the reduced welfare payments for individuals.) For reductions in the cost of lost productivity incurred by the general economy, improvements in substance misuse (\$157.3 million), unemployment (\$95 million), social cohesion (\$20.6 million) and underemployment (\$10.8 million) achieved the greatest changes.

For all social determinants except social cohesion, the estimated incremental total costs that flowed from a 10% improvement in social determinants were slightly greater than simply doubling the estimates for a 5% improvement (Supplementary Table 1 in Supplementary Information part B).

When costs and QALYs were combined with a willingness to pay per QALY of \$83,004 to derive incremental net monetary benefit (INMB) compared with business as usual (a full explanation of these terms and methodology is available in the Methods), a 5% improvement in substance misuse resulted in an INMB of \$281.67 million, childhood difficulties \$234.50 million, social cohesion \$146.64 million and homelessness \$38.75 million (Fig. 4; Supplementary Table 2 in Supplementary Information part B). A comparison between 3-year and 11-year time horizons reveals that taking a long-term view would

enable decision-makers to exponentially capitalize on the full extent of the net benefit anticipated from improving the social determinants of mental health. For example, for the societal perspective, the INMB for a 5% improvement in social cohesion was \$3.99 million at 3 years and \$146.64 million at the end of 11 years. Similar findings reinforcing the importance of a long-term view were found for health outcomes (Supplementary Table 3 in Supplementary Information part B). For example, a 5% improvement in social cohesion achieved a reduction of 648 mental health-related emergency department presentations over 11 years, but this was only 12 by the end of 3 years.


Univariate sensitivity analysis found that varying utility values, homelessness costs and substance misuse costs did not change the rank ordering of social determinants based on INMB for the societal perspective (Supplementary Information part B).

## Discussion

System dynamics modeling (SDM), including a participatory model-building process, was used to demonstrate the health and economic benefits of improving eight social determinants of mental health. We observe from the model that even modest improvements in determinants resulted in material increases in health outcomes and reductions in costs for both the health care and societal perspectives. This simulation model was developed in collaboration with a diverse group of local stakeholders for a region of 1.2 million people, a process that could be replicated and customized for other regions.

This Analysis provides quantitative support for ‘moving upstream’ to target the underlying causes of a large portion of distress and mental health challenges<sup>17,43,44</sup>. A decision-maker contemplating how to prioritize the allocation of scarce resources towards improving mental health via social determinants could utilize these estimates by first identifying what their most important objectives are. If their main objective is to maximize the improvement of mental health outcomes in terms of QALYs, then social cohesion and childhood difficulties would be the primary determinants to target. If their main objective is to reduce suicide deaths, targeting substance misuse and social cohesion would maximize this outcome. If their objective is reducing pressure on health care services, improving social cohesion and childhood difficulties achieves the greatest reductions in mental health-related





**Fig. 4 | INMB (societal perspective), 5% improvement in social determinants.** Cumulative costs over 11 years, 2024–2034, discounted, 2020–2021 Australian dollars versus business as usual. ed., education.

emergency department (ED) presentations, and reducing substance misuse attains the greatest reduction in self-harm hospitalizations.

The greatest reductions in costs for the health care perspective were attained by reductions in childhood difficulties and homelessness. When this perspective was expanded to account for costs incurred outside the health care sector and lost productivity for the societal perspective, reductions in substance misuse and unemployment realized the greatest downstream cost savings.

Ideally, a decision-maker would seek to maximize allocative efficiency through consideration of both health benefits (in this case, in terms of QALYs) and costs by using the INMB, which also accounts for the willingness to pay for a QALY. The ranking based on the highest INMB over 11 years for the societal perspective was (1) substance misuse, (2) childhood difficulties, (3) social cohesion and (4) unemployment. Caution should be applied when interpreting the INMB estimates in this Analysis, as these exclude the cost of interventions normally included in the calculation of INMB.

Overall, social cohesion, childhood difficulties, substance misuse and unemployment are a group of social determinants that could be prioritized on the basis that they were forecast to improve a variety of final (QALYs and suicide deaths) and intermediate (hospitalizations and ED presentations) health outcomes and/or reduce costs for the societal perspective. This adds to, and is consistent with, existing literature on the link between unemployment and suicide<sup>45,46</sup>, the influence of early life circumstances on mental disorders in later life<sup>47–49</sup>, the importance of social connection for mental health and well-being<sup>39,50–53</sup>, and the association between substance misuse and mental health challenges<sup>54,55</sup>. Given the limitations of universal prevention in mental health<sup>4</sup>, this Analysis provides quantitative support for targeting more specific, underlying causes. Box 3 provides a summary of systematic review-level evidence of interventions targeting the social determinants of mental health.

The findings of this study highlight the importance of taking a long-term view when considering whether the social determinants of mental health warrant the investment of scarce resources. When comparing the estimates produced here between 3 years and 11 years, much more value, in terms of both health benefits and costs, became apparent by

11 years. The choice of 11 years as the maximum time horizon is itself arbitrary, and the model timeframe could be extended to multiple decades. This would allow a more comprehensive capture of health benefits accrued by people who are younger at the start of the simulation as they age, as well as the potential for intergenerational prevention effects. The ranking of social determinants based on total cost reductions for both the health care and societal perspectives changed over the 11-year timeframe (Figs. 2 and 3) and may change beyond this time horizon.

Taking a societal perspective was useful for analyzing the economic impact of improving the social determinants of mental health, with cost estimates much larger for the societal perspective than for the health care perspective. This was due to the inclusion of additional cost categories, such as lost productivity, the criminal justice system, and non-health care social services for homelessness and substance misuse. There have been calls for greater use of the societal perspective<sup>56</sup>, or at least reporting both the health care and societal perspectives in economic evaluations<sup>57</sup>. Perspective can alter the conclusions in economic evaluations for mental health. For example, a systematic review of economic evaluations for depression found that the results in 24% of studies changed quadrant of the cost-effectiveness plane (representing positive or negative differences in costs or health outcomes) when the societal perspective was taken and 5% of studies changed conclusions of whether the intervention was cost-effective or not<sup>58</sup>. In the case of this Analysis, a societal perspective helps to highlight a more comprehensive picture of benefits available from targeting the social determinants of mental health.

There are a number of limitations to this Analysis. First, only a selection of the social determinants of mental health have been included in this Analysis. These were identified as priority determinants thought to be driving youth mental health outcomes in the region. Some important determinants were not modeled owing to a current lack of data. Examples of excluded determinants include level of income, measures of relative socioeconomic advantage/disadvantage, food security, loneliness, discrimination and inequalities experienced by marginalized groups (for example, First Nations peoples, LGBTQ+, migrants, and culturally and linguistically diverse people). Although all of these are captured to a degree by the composite measure of social cohesion included in

**BOX 1****Lived experience perspective**

Economic arguments are sometimes portrayed as being at odds with lived/living experience frameworks as ways to guide mental health investment and public health policy. Lived/living expertise is highly experiential and highly personal—it speaks to a first-person perspective rather than a population approach to understanding community need. However, lived/living experience also provides a clear rationale for why better economic evidence and a strong approach to reducing the burden of social determinants is so necessary. Most young people, when asked to reflect on how or why they came to experience a mental health or substance use challenge, will not say that they experience a neurochemical imbalance or that they inherited an epigenetic characteristic from their parents. They will say things like: “I had a rough childhood,” or “I can’t find a job, so I’m stressed about rent every week” or “I don’t know what the future looks like for me after I spent high school in lockdown.” Understanding the impact of social determinants on the widening mental health and well-being divide between younger generations and older ones is essential to begin to curb the downward trend in youth mental health. Similarly, young people with lived experience will also tell you that the current approach of investing in hospital-based and community service provider models, even at an individual level, doesn’t work well for most people. Better economic evidence is necessary to guide a social response to mental health and substance use challenges to ensure investment is placed where it will have the most impact. Tailoring this evidence to respond to the local context of a region is also in line with lived/living experience frameworks—this approach allows for the nuances, complexity and dynamism of a specific community to guide the knowledge production that sits behind the evidence. This study speaks to a particular region that has particular needs and constructs a clear rationale for evidence-based investment in the social determinants driving distress in that region. It is a fantastic blueprint for better decision-making, with young people with lived experience at the center of the process.

by Jordan van Rosmalen

this Analysis (Table 1), analyzing these individually would provide a more nuanced picture to a decision-maker to aid their prioritization of scarce resources to particular areas. Second, the choice of 5% and 10% improvements in the social determinants is arbitrary and hypothetical. Five percent was used as the focus for results as something that could be attainable and set as a target by a policymaker. Related to this, we cannot say whether these improvements in social determinants are cost-effective as the cost of the interventions or policy changes required to achieve them is unknown. The cost of implementation would need to be compared with the reported cost offsets and health benefits to determine cost-effectiveness through simulation modeling. Third, although we have estimated cost impacts by payer, the Australian funding system is complex, and no attempt has been made in this Analysis to account for the impacts on original funding sources. Fourth, the capacity for homelessness and substance misuse to have more of a dramatic impact on cost reductions than their increase in QALYs would suggest is because a large portion of their costs is linked directly with that determinant rather than being mediated through levels of distress (which determines QALYs). For example, criminal justice costs are incurred for both substance misuse and homelessness, and any proportional reduction in the prevalence of these determinants has an equivalent reduction in criminal justice costs, regardless of their association and

**BOX 2****Systems perspective**

The findings of this Analysis reinforce the need to take a wider perspective on mental illness in their context of the daily lives of young people—in their families and communities, interacting with systems as well as natural, built and digital environments. This ecological approach has been adopted by The Nest<sup>88</sup>, Australia’s well-being framework for children and young people up to 24 years old. It conceptualizes well-being as six interrelated domains: feeling valued, loved and safe; being healthy, learning; participating; having material basics; and possessing a strong sense of identity and culture. To have the best possible well-being, a young person needs to be adequately resourced in all six domains at an individual level as well as within their family, community and wider society including online. Enhancing social determinants as a means of preventing and/or alleviating youth mental illness (as well as fostering well-being in its own right) is therefore the responsibility of many, and requires a multifaceted systems approach<sup>89</sup>. The results of this study clearly demonstrate the importance and cumulative effect of such an approach, especially when a timeframe beyond the typical duration of funding or political cycles is applied. They provide a clear, compelling argument for policymakers and funders to adopt, articulate, appropriately resource and collaboratively enact a proactive, long-term, cross-sectoral vision to interrupt the prevalence and impact of youth mental illness. Doing so will also enhance the conditions that will enable our young people—and their social networks, including future generations—a better chance to thrive.

by Sophie Morson

effect on levels of distress or suicide deaths. The same consideration applies to unemployment and underemployment and the productivity impacts linked to employment status. Fifth, we have not included all costs that might be relevant for the societal perspective. One example of this is the increase in earnings that would be received by healthier or newly employed people (represented here by the payer category of ‘nongovernment including individuals’), offsetting the reduction in welfare payments to this group. This has been excluded due to our focus on productivity-related costs. Another example would be carbon emissions and other environmental impacts. These were excluded due to time and resource constraints on the model development process. Another limitation is that only limited univariate sensitivity analysis was carried out, including the two different levels of improvement to determinants (of 5% and 10%) and two different time horizons (3 years and 11 years) (Supplementary Information part B). The main reason for this was the exclusion of defined and costed interventions from the analysis (that is, cost-effectiveness analysis has not been performed). One-way and probabilistic sensitivity analyses are useful for cost-effectiveness analyses because they reveal the circumstances for which the conclusions about cost-effectiveness might change and deriving the uncertainty intervals around economic summary measures. For the present Analysis, decision rules based on the incremental cost-effectiveness ratio, for example, are irrelevant due to the exclusion of interventions. One-way sensitivity analysis of the discount rate, for example, would only serve to change the magnitude of estimates. The limited sensitivity analysis that was performed found that varying utility values, homelessness costs and substance misuse costs did not change the rank ordering of social determinants based on INMB for the societal perspective. Another limitation is that we have included some costs related to carers but not changes in quality of life of carers due to time and resource constraints. This will be included in a future iteration

## BOX 3

## Interventions targeting the social determinants of mental health

Although we have not included specific interventions in this Analysis, some evidence exists for various programs and services that could be implemented to move toward these hypothetical improvements. Here, we describe several systematic reviews that summarize this evidence and identify gaps that can be filled by future research. The objective of this section is to provide pointers to high-level evidence in the field for interested readers, rather than a thorough, systematic assessment of effectiveness for each individual type of intervention.

An umbrella review (systematic review of reviews) included 101 reviews of interventions targeting social determinants of mental disorders<sup>42</sup>, mapping them to the United Nations Sustainable Development Goals and grouping them across five domains following the framework set out by Lund et al.<sup>16</sup>: demographic, economic, environmental events, neighborhood and sociocultural. This umbrella review serves as a useful resource covering a vast range of interventions<sup>42</sup>. A small sample of interventions identified by this review that could be prioritized for investment based on existing evidence includes: digital and brief advocacy for female intimate partner violence survivors; cash transfer programs for low- and middle-income countries; and investment in psychosocial support for vulnerable individuals following environmental events. However, the authors found important gaps in current evidence, at least in terms of review-level evidence. No reviews were found that met their inclusion criteria on: interventions to address income inequality or employment precarity; increased access to or completion of education; or interventions targeting climate change, vulnerable ecosystems or disaster preparedness<sup>42</sup>. This review did not look for cost-effectiveness analyses.

Another umbrella review, of interventions for reducing loneliness specifically, included 211 studies and found that social support, social cognitive training and meditation or mindfulness interventions decreased loneliness<sup>90</sup>. This review did not look for economic evidence on loneliness interventions.

Kirkbride et al., in the context of a broad review of the social determinants of mental health, argue for the prioritization of three kinds of intervention for improving the social determinants of mental health: (1) those that target the early life course (prenatal, childhood and adolescence) to interrupt intergenerational transmission between social determinants and mental health problems; (2) those that impact multiple domains such as physical, mental and social outcomes; and (3) those that focus on alleviating poverty because this is often a root cause related to many social determinants and poor mental health<sup>37</sup>. Examples of interventions that target the early life course are online parenting programs and school-based suicide prevention programs. The Triple P online parenting program is supported by evidence on its effectiveness<sup>91</sup> and cost-effectiveness, using both conventional<sup>92</sup> and dynamic modeling techniques<sup>93</sup>. The Thrive by Five International Program empowers parents with co-designed and place-based psychoeducational information coupled with practical

activities that easily integrate into daily routines to support the health social, emotional and cognitive development in early childhood<sup>94,95</sup>. School-based suicide prevention programs<sup>96–98</sup> are more effective than school-based mindfulness, well-being, anxiety or depression programs<sup>99–102</sup>, but the few cost-effectiveness analyses that exist report mixed conclusions on their value for money<sup>25,27,93,103</sup>. This suggests that country of implementation, context of the school and population, design of the intervention, side effects<sup>104</sup> and accounting for broader systemic impacts (for example, productivity of teachers and demand for mental health services) are important determinants of the economic credentials of school-based interventions.

Examples of interventions that focus on alleviating poverty include the employment program Individual Placement and Support (IPS), as well as direct economic support, such as universal/guaranteed incomes or cash transfers<sup>38</sup>. Systematic reviews with meta-analysis have found that IPS is effective in terms of vocational outcomes but has only indirect effects on nonvocational outcomes, such as quality life and mental health<sup>105,106</sup>. The cost effectiveness of IPS depends on the country of implementation, intervention design and the underlying level of unemployment based on reviews of economic literature<sup>107,108</sup>. A systematic review and meta-analysis of the effects of changes in income on mental health found that lifting individuals out of poverty was associated with an improvement in mental health measures and well-being<sup>109</sup>. Although effect sizes were modest, they were also consistent and likely to have a large impact at the population health level, particularly when targeted at lifting people out of poverty. Subgroup analysis revealed no consistent evidence that effect size was influenced by income source. The authors compared these results with antidepressants and cognitive behavioral therapy (CBT) and concluded that “interventions that move people above the poverty line might be roughly half as effective in improving mental health as antidepressants and a quarter as effective as CBT”. However, because “these treatments are primarily studied in individuals at high-risk rather than general population samples, the potential impacts of anti-poverty interventions at a population mental health level could be substantial”<sup>109</sup>.

To summarize the review-level evidence on interventions for improving the social determinants of mental health, there is sufficient evidence to support greater implementation of certain interventions now, but this is an area where much more research could be conducted, particularly evaluations of real-world implementation and generation of economic evidence, both within-trial and modeled cost-effectiveness/cost-benefit analyses. One limitation of the studies mentioned here is that they are review-level evidence. If a review has not been conducted on a particular topic, that does not mean there is no evidence supporting its effectiveness. Similarly, studies may have been published since the review was conducted, or they did not meet the inclusion criteria of a particular review. Thorough analysis of each individual type of intervention is outside the scope of the present Analysis.

of the model. Adding QALYs to the model for carers would be expected to enhance the value of improving the social determinants of health.

Although specific interventions have not been included in this Analysis, the results reported here do provide some guidance for decision-makers considering investing in interventions that target the social determinants of mental health. For example, if an intervention is expected to achieve a 5% improvement in social cohesion (as defined in Table 1), it could cost up to approximately \$16 million over 11 years

(discounted) and still be considered cost saving for the health care perspective based on the figures provided in Supplementary Table 1 (Supplementary Information part B) as well as the health benefits specified in Supplementary Table 3 (Supplementary Information part B). For the societal perspective, as long as the intervention cost less than \$38 million over 11 years (discounted), it would be considered cost saving. Specific strategies are likely to be context dependent, and policymakers, academics and local communities could work together to both



identify contextually appropriate interventions and develop models that can accurately evaluate them to reduce implementation barriers.

Future research directions could include using a similar modeling approach to investigate the cost-effectiveness of interventions targeted at improving the social determinants of mental health based on those that are supported by evidence of effectiveness. This would enable more specific policy actions by estimating the cost of implementing the intervention in a particular region and using the best available evidence to inform its anticipated effectiveness. Future iterations of this modeling could also seek to include an expanded set of benefits and costs that fall in the societal perspective, such as measures of well-being or environmental impacts, as relevant to the intervention and decision context being analyzed. Future research could also investigate the value of improving several social determinants concurrently or combinations of interventions concurrently.

Planning to respond to mental health is often health focused, and typically bound by the realities of funding limitations. While these limitations persist, the modeling approach outlined here vastly expands our capacity to better identify opportunities to respond to community mental health priorities, transcending the health system to more fully account for other key areas of concern, such as employment and housing. This increases our understanding of the relative utility of different interventions and their material impact on people's lives and on the broader community.

## Methods

The Consolidated Health Economic Evaluation Reporting Standards (CHEERS) 2022 were followed when reporting the results and methods of analysis<sup>59</sup>. The completed CHEERS checklist is available as Supplementary Information. Protocols related to this project have been published on the SDM approach<sup>60</sup>, economic analysis<sup>61</sup> and participatory model-building process<sup>62</sup>. More detailed methods are available as Supplementary Information.

### Model development process

A participatory systems modeling process was undertaken to develop a system dynamics simulation model to estimate the health and economic consequences of three types of system change aimed at improving mental health outcomes for young people: (1) increasing or decreasing the growth rates of health services capacity; (2) implementing or upscaling evidence-based interventions; and (3) changing various social determinants of mental health. The present Analysis was focused on the third group of system changes. The participatory systems modeling component of the project included three in-person workshops with attendance from a range of stakeholders including mental health professionals, primary care physicians, allied health service providers, young people with a lived experience of mental health conditions and their carers, and health agency representatives.

- In workshop 1, participants identified outcomes of interest, mapped the youth mental health service system in the geographical catchment of Brisbane South Primary Health Network, and identified programs and initiatives of interest, as well as the social determinants that were most important to youth mental health in their region.
- In workshop 2, the research team presented the basic structure of the model based on the workshop 1 systems mapping exercise to participants for feedback. Participants also provided detailed advice on what new interventions would look like if implemented in their region.
- In workshop 3, a draft, interactive version of the model was provided to participants for user testing and exploration of key policy and planning insights. This process facilitated 'sense-checking' of the model output and identification of bugs or unanticipated findings.

Regular meetings were held with a model development advisory group between the workshops for more detailed components of the model development process. Where model inputs or development decisions were informed by expert advice, this could be informed through any of the workshops, model development group meetings or personal communication with subject matter experts from any of the following roles, noting that these categorizations are somewhat arbitrary and individual participants are likely to span multiple roles and identities:

- Young people with a lived experience of mental health conditions and their carers.
- Clinician subject experts such as psychiatrists, psychologists and general practitioners.
- Academic researchers including health economists, systems modelers, data scientists and research support personnel.
- Policy, health administration and management representatives, including those from the Primary Health Network and State Department of Health.
- Representatives from nonhealth sectors such as education and housing.

Research evidence and data were used to parameterize the model, and historical time series data were used to calibrate the model. With regard to searching the literature for mechanisms of effectiveness, a rapid review approach was used, relying where possible on systematic reviews. Parameter values that could not be derived directly from these sources were estimated or calibrated via constrained optimization, using historical time series data for a wide range of sociodemographic and health-related outcomes, including participation and unemployment rates, the prevalence of moderate to very high psychological distress, intentional self-harm hospitalization and suicide mortality rates and rates of mental health service usage. Powell's method was used to obtain the set of optimal parameter values minimizing the mean of the absolute differences between the observed time series values and the corresponding model outputs, where each difference was expressed as a percentage of the observed value<sup>63</sup>.

### Model structure overview

SDM was chosen for this Analysis owing to its potential for capturing complexity, interactions, dynamics and broader systemic effects relevant to mental health planning<sup>34</sup>. The model is a logically consistent mathematical framework that integrates best available data and evidence sources with expert and local knowledge (including the expert knowledge of those with a lived experience of mental health conditions and their carers). The model captures population and demographic dynamics, pathways to youth mental health care, service interactions and workforce capacity, and the potentially nonadditive effects of intervention combinations. The model was validated through face validity among stakeholders and by observing its ability to reproduce historic trends across observed data in the region from 2011 to 2022.

The sectors included in the model are:

- Population, which models the resident population divided into six age brackets (0–4-year-olds, 5–11, 12–14, 15–17, 18–24, and 25 and older),
- Education, which models students enrolled in primary, secondary and post-secondary education, and people with different levels of highest qualifications,
- Labor force, which models unemployment, underemployment and participation rates,
- Not in education, employment nor training (NEET), which models the youth population aged 15–24 years not in education, employment nor training,
- Homelessness, which models the population experiencing homelessness,

- Substance misuse, which models the prevalence of 12-month substance misuse disorder and substance misuse closed treatment episodes,
- Psychological distress/disorder, which models the prevalence of low psychological distress and the prevalence of moderate to very high psychological distress based on the Kessler Psychological Distress Scale (K10)<sup>64</sup>. The population with moderate to very high psychological distress is further dichotomized by whether or not they meet the criteria for any 12-month mental disorder,
- Strengths and difficulties (used as the measure of childhood difficulties), which models the prevalence of behavioral and emotional difficulties among children aged 0–11 years as measured by the Strengths and Difficulties Questionnaire (SDQ),
- Social cohesion, which models the population level of social cohesion according to the Scanlon–Monash Index of Social Cohesion,
- Suicidal behaviors, which models the rates of suicide attempts and suicide deaths, and
- Mental health services, which models mental health services delivered by health professionals and online mental health services.

Supplementary Fig. 1 in Supplementary Information part A provides an overview of the sectors and the causal connections between them. Supplementary Information part A also contains a detailed description of each sector, including its structure, sources of input data and calibration graphs.

Hypothetical improvements in each social determinant of 5% and 10% were chosen as attainable but meaningful levels of improvement that could be set by a policymaker as targets to strive toward. Table 1 contains further details of each determinant and what the levels of improvement mean for each scenario. Additional detail is provided in Supplementary Information part B, where the projection for each social determinant is provided over the 11-year time horizon along with the expected changes with 5% and 10% improvements.

## Costs

Supplementary Information part B contains a detailed itemization of each unit cost, its source and derivation methods (Supplementary Tables 4 and 5). In brief, the cost of health services was obtained from the Australian Institute of Health and Welfare and the Independent Health and Aged Care Pricing Authority. Costs associated with homelessness and substance misuse were obtained from gray literature reports estimating the associated economic burden. Employee earnings used in the calculation of productivity estimates were obtained from the Australian Bureau of Statistics.

## Health measures and QALYs

The detailed sector descriptions in Supplementary Information part A describe the calculation methods by which key health outcomes are estimated: suicide deaths, ED presentations and self-harm hospitalizations. The QALY is an additional measure of final health outcomes that is useful because it is a composite measure of health that includes both changes in survival and quality of life, reflecting the preference-based utilities of distress-related health states. The prevalence of people experiencing higher levels of distress is a key driver of accumulated QALYs over the timeframe of the model. Because the K10 measure is used to quantify levels of distress in the model, utilities for people aged 12 years and over were derived by using a mapping algorithm to convert low, moderate, high and very high distress levels based on the K10 to the EQ-5D multiattribute utility instrument<sup>65</sup>, applying mapped utility decrements to Australian age-based population norms<sup>66</sup> (Supplementary Table 6 in Supplementary Information part B). Distressed health states for children aged 11 years and younger were based on the SDQ<sup>67</sup>. A mapping algorithm developed by Sharma et al., with a study population of Australian children, was used to transform SDQ scores to

CHU9D utilities (Supplementary Table 7 in Supplementary Information part B)<sup>68</sup>. The CHU9D has strong psychometric performance in both mental health<sup>69</sup> and general pediatric<sup>70</sup> populations. A recent systematic review confirmed this was the most recent and relevant mapping algorithm with the SDQ as the starting measure<sup>71</sup>.

## Other economic methods

The time horizon was set to 11 years to allow sufficient time for cost and health consequences to play out after a change to a social determinant but short enough to minimize the uncertainty in future states of the world beyond this timeframe. Sensitivity analysis using a time horizon of 3 years was also conducted and reported here for health outcomes and INMB. A conventional lifetime time horizon is not relevant due to the dynamic nature of the population, where births and positive net migration ensure the population continues to grow. Costs and health consequences were discounted at 5%, based on the reference discount rate set by the Pharmaceutical Benefits Advisory Committee<sup>72</sup>. Both health care and societal perspectives were adopted due to the diverse stakeholders and decision-makers involved in the participatory model building process, and all results are presented for both perspectives. The societal perspective is broader than, and fully encapsulates, the health care perspective. One of the key differences between the two perspectives is the inclusion of productivity costs in the societal perspective. Costs incurred by nonhealth sectors are also included in the societal perspective. The costs that apply to each perspective are detailed in Supplementary Information part B. Neither univariate nor probabilistic sensitivity analyses were carried out for this Analysis (beyond alternative timeframes, degrees of improvement, utility values and costs related to homelessness and substance misuse). Our choice to exclude sensitivity analysis is predominantly due to the hypothetical nature of the improvements to social determinants. In conventional cost-effectiveness analysis where specific, defined interventions are included, the primary purpose of the sensitivity analysis is to establish the degree of uncertainty around summary economic measures based on the joint probability of all relevant uncertain parameters (probabilistic), or to what degree conclusions may change when individual parameters are varied within plausible ranges, that is, to establish the degree of confidence in whether the binary choice of implementing an intervention represents allocative efficiency. For the present Analysis, the degree of improvement in a social determinant is arbitrary, so the choice of distribution around this would also be arbitrary. Similarly, any univariate change in a parameter would only serve to change the magnitude in anticipated health or cost consequences.

One way sensitivity analysis was conducted on a select group of parameters expected to have the most influential impact on the rank order of social determinants: utility values for ages 12+ varied  $\pm 10\%$ ; utility values for ages 0–11 varied  $\pm 10\%$ ; costs related to homelessness varied  $\pm 20\%$ ; costs related to substance misuse varied  $\pm 20\%$ . Results for this sensitivity analysis are reported in Supplementary Information part B.

## Economic summary measures

Net monetary benefit (NMB) is a conversion of health into financial units to aid the comparability of strategies and presentation of results. NMB is given by multiplying QALYs by the willingness to pay for a QALY, AUD\$83,004, then subtracting costs. The willingness to pay per QALY was obtained from Shiroya et al.<sup>73</sup> and indexed to the 2021–2022 financial year. We adopted the social value of a QALY (demand for health) approach for valuing the threshold rather than an opportunity cost (supply side) approach<sup>74</sup>. This is relevant for the present context because the social determinants operate outside the health care sector, so any intervention or policy change would be happening at a broader level than the health sector that is not constrained by the health care budget. Using a willingness to pay for a QALY value from the literature gives us the added benefit of using a threshold based on empirical

evidence, rather than an arbitrary threshold (usually AUD\$50,000 per QALY in Australia, although this has never been publicly set by a government agency in Australia). INMB is the difference in NMB for a scenario of change compared with business as usual. Although INMBs reported here cannot be interpreted in the conventional manner because the costs of interventions have not been accounted for, they do provide useful estimates of the collective economic value of both health and cost consequences in a single figure.

### Reporting summary

Further information on research design is available in the Nature Portfolio Reporting Summary linked to this article.

### Data availability

This simulation model used data from a variety of sources, all of which are referenced in the Supplementary Information.

### Code availability

The model file is not available online; however, we welcome sharing of our models and methods with interested researchers who wish to collaborate with us. One of the reasons for a collaborative approach to sharing our models is the ability to provide training and support to users.

### References

- Meadows, G. N. et al. Resolving the paradox of increased mental health expenditure and stable prevalence. *Aust. N. Z. J. Psychiatry* **53**, 844–850 (2019).
- Budget 2021-22: Generational change and record investment in the health of Australians. *Department of Health and Aged Care* <https://www.health.gov.au/ministers/the-hon-greg-hunt-mp/media/budget-2021-22-generational-change-and-record-investment-in-the-health-of-australians> (2021).
- The National Children's Mental Health and Wellbeing Strategy* (National Mental Health Commission, 2021).
- Cuijpers, P. Preventing the onset of depressive disorders: state of the art and future directions. *Curr. Dir. Psychol. Sci.* **34**, 51–56 (2025).
- GBD 2021 Diseases and Injuries Collaborators. Global incidence, prevalence, years lived with disability (YLDs), disability-adjusted life-years (DALYs), and healthy life expectancy (HALE) for 371 diseases and injuries in 204 countries and territories and 811 subnational locations, 1990–2021: a systematic analysis for the Global Burden of Disease Study 2021. *Lancet* **403**, 2133–2161 (2024).
- Wilkins, R., Vera-Toscano, E. & Botha, F. *The Household, Income and Labour Dynamics in Australia Survey: Selected Findings from Waves 1 to 21* (Melbourne Institute: Applied Economic and Social Research, 2024).
- Causes of death, Australia. *Australian Bureau of Statistics* <https://www.abs.gov.au/statistics/health/causes-death/causes-death-australia/2023> (2024).
- Biddle, N., Edwards, B. & Rehill, P. *The Relationship between COVID-19 Policies and Subjective Wellbeing—August 2022* (The ANU Centre for Social Research and Methods, 2022).
- Skinner, A., Occhipinti, J. A., Song, Y. J. C. & Hickie, I. B. Population mental health improves with increasing access to treatment: evidence from a dynamic modelling analysis. *BMC Psychiatry* **22**, 692 (2022).
- Haidt, J. *The Anxious Generation: How the Great Rewiring of Childhood Is Causing an Epidemic of Mental Illness* (Random House, 2024).
- McGorry, P. D. et al. The Lancet Psychiatry Commission on youth mental health. *Lancet Psychiatry* **11**, 731–774 (2024).
- Patel, V. et al. The Lancet Commission on global mental health and sustainable development. *Lancet* **392**, 1553–1598 (2018).
- Scott, K. M. et al. Association of mental disorders with subsequent chronic physical conditions: world mental health surveys from 17 countries. *JAMA Psychiatry* **73**, 150–158 (2016).
- Clark, C. et al. Impact of childhood and adulthood psychological health on labour force participation and exit in later life. *Psychol. Med.* **47**, 1597–1608 (2017).
- Santomauro, D. F., Purcell, C., Whiteford, H. A., Ferrari, A. J. & Vos, T. Grading disorder severity and averted burden by access to treatment within the GBD framework: a case study with anxiety disorders. *Lancet Psychiatry* **10**, 272–281 (2023).
- Lund, C. et al. Social determinants of mental disorders and the Sustainable Development Goals: a systematic review of reviews. *Lancet Psychiatry* **5**, 357–369 (2018).
- Occhipinti, J.-A. et al. The influence of economic policies on social environments and mental health. *Bull. World Health Org.* **102**, 323–329 (2024).
- Braveman, P. & Gottlieb, L. The social determinants of health: it's time to consider the causes of the causes. *Publ. Health Rep.* **129**, 19–31 (2014).
- Occhipinti, J.-A. et al. Estimating the Mental Wealth of nations: valuing social production and investment. *Nat. Ment. Health* **1**, 247–253 (2023).
- Occhipinti, J. A. et al. Measuring, modeling, and forecasting the mental wealth of nations. *Front. Publ. Health* **10**, 879183 (2022).
- Bloom D. E. et al. *The Global Economic Burden of Noncommunicable Diseases* (World Economic Forum, 2011).
- Roehrig, C. Mental disorders top the list of the most costly conditions in the United States: \$201 billion. *Health Aff.* **35**, 1130–1135 (2016).
- Knapp, M. et al. Investing in recovery: making the business case for effective interventions for people with schizophrenia and psychosis. *LSE Research Online Documents on Economics* 56773 (London School of Economics and Political Science, LSE Library, 2014).
- Mental Health* (Productivity Commission, 2020); <https://www.pc.gov.au/inquiries/completed/mental-health/report>
- Kularatna, S. et al. Cost-effectiveness analysis of paediatric mental health interventions: a systematic review of model-based economic evaluations. *BMC Health Serv. Res.* **22**, 542 (2022).
- McDaid, D., Park, A. L. & Wahlbeck, K. The economic case for the prevention of mental illness. *Annu. Rev. Publ. Health* **40**, 373–389 (2019).
- Le, L. K. et al. Cost-effectiveness evidence of mental health prevention and promotion interventions: a systematic review of economic evaluations. *PLoS Med.* **18**, e1003606 (2021).
- Feldman, I., Gebreslassie, M., Sampaio, F., Nystrand, C. & Ssegonja, R. Economic evaluations of public health interventions to improve mental health and prevent suicidal thoughts and behaviours: a systematic literature review. *Adm. Policy Ment. Health* **48**, 299–315 (2021).
- Knapp, M. & Wong, G. Economics and mental health: the current scenario. *World Psychiatry* **19**, 3–14 (2020).
- Lathe, J., Silverwood, R. J., Hughes, A. D. & Patalay, P. Examining how well economic evaluations capture the value of mental health. *Lancet Psychiatry* **11**, 221–230 (2024).
- Ha, N. T., Huong, N. T., Anh, V. N. & Anh, N. Q. Modelling in economic evaluation of mental health prevention: current status and quality of studies. *BMC Health Serv. Res.* **22**, 906 (2022).
- National Report Card 2023: Monitoring the Performance of Australia's Mental Health System* (National Mental Health Commission, 2024).
- Rosenberg, S. & Salvador-Carulla, L. PERSPECTIVES: accountability for mental health: the Australian experience. *J. Ment. Health Policy Econ.* **20**, 37–54 (2017).



34. Crosland, P. et al. Incorporating complexity and system dynamics into economic modelling for mental health policy and planning. *Pharmacoeconomics* **42**, 1301–1315 (2024).
35. Allen, J., Balfour, R., Bell, R. & Marmot, M. Social determinants of mental health. *Int. Rev. Psychiatry* **26**, 392–407 (2014).
36. Lund, C. Global mental health and its social determinants: how should we intervene? *Behav. Res. Ther.* **169**, 104402 (2023).
37. Kirkbride, J. B. et al. The social determinants of mental health and disorder: evidence, prevention and recommendations. *World Psychiatry* **23**, 58 (2024).
38. Ridley, M., Rao, G., Schilbach, F. & Patel, V. Poverty, depression, and anxiety: causal evidence and mechanisms. *Science* **370**, eaay0214 (2020).
39. Na, P. J. et al. Social determinants of health and suicide-related outcomes: a review of meta-analyses. *JAMA Psychiatry* **82**, 337–346 (2025).
40. *Guidance on Mental Health Policy and Strategic Action Plans* (World Health Organization, 2025).
41. Littleton, C. & Reader, C. To what extent do Australian child and youth health, and education wellbeing policies, address the social determinants of health and health equity?: a policy analysis study. *BMC Public Health* **22**, 2290 (2022).
42. Oswald, T. K. et al. Interventions targeting social determinants of mental disorders and the Sustainable Development Goals: a systematic review of reviews. *Psychol. Med.* **54**, 1475–1499 (2024).
43. Occhipinti, J. A. et al. Mental health: build predictive models to steer policy. *Nature* **597**, 633–636 (2021).
44. Occhipinti, J. A. et al. Reducing youth suicide: systems modelling and simulation to guide targeted investments across the determinants. *BMC Med.* **19**, 61 (2021).
45. Skinner, A., Osgood, N. D., Occhipinti, J.-A., Song, Y. J. C. & Hickie, I. B. Unemployment and underemployment are causes of suicide. *Sci. Adv.* **9**, eadg3758 (2023).
46. Nordt, C., Warnke, I., Seifritz, E. & Kawohl, W. Modelling suicide and unemployment: a longitudinal analysis covering 63 countries, 2000–11. *Lancet Psychiatry* **2**, 239–245 (2015).
47. Davies, C. et al. Prenatal and perinatal risk and protective factors for psychosis: a systematic review and meta-analysis. *Lancet Psychiatry* **7**, 399–410 (2020).
48. Grummitt, L., Baldwin, J. R., Lafoa'i, J., Keyes, K. M. & Barrett, E. L. Burden of mental disorders and suicide attributable to childhood maltreatment. *JAMA Psychiatry* **81**, 782–788 (2024).
49. Danese, A. & Widom, C. S. Associations between objective and subjective experiences of childhood maltreatment and the course of emotional disorders in adulthood. *JAMA Psychiatry* **80**, 1009–1016 (2023).
50. Holt-Lunstad, J. Social connection as a critical factor for mental and physical health: evidence, trends, challenges, and future implications. *World Psychiatry* **23**, 312–332 (2024).
51. Breedvelt, J. J. F. et al. The effects of neighbourhood social cohesion on preventing depression and anxiety among adolescents and young adults: rapid review. *BJPsych Open* **8**, e97 (2022).
52. Fone, D. et al. Does social cohesion modify the association between area income deprivation and mental health? A multilevel analysis. *Int. J. Epidemiol.* **36**, 338–345 (2007).
53. Birrell, L., Werner-Seidler, A., Davidson, L., Andrews, J. L. & Slade, T. Social connection as a key target for youth mental health. *Ment. Health Prev.* **37**, 200395 (2025).
54. Iorfino, F. et al. Multidimensional outcomes in youth mental health care: what matters and why? *Med. J. Aust.* **211**, S4–S11 (2019).
55. Iorfino, F. et al. Patterns of emergency department presentations for a youth mental health cohort: data-linkage cohort study. *BJPsych Open* **9**, e170 (2023).
56. Jonsson, B. Ten arguments for a societal perspective in the economic evaluation of medical innovations. *Eur. J. Health Econ.* **10**, 357–359 (2009).
57. Neumann, P. J., Sanders, G. D., Russell, L. B., Siegel, J. E. & Ganiats, T. G. *Cost-Effectiveness in Health and Medicine* 2nd edn (Oxford Univ. Press, 2017).
58. Duevel, J. A. et al. Considering the societal perspective in economic evaluations: a systematic review in the case of depression. *Health Econ. Rev.* **10**, 32 (2020).
59. Husereau, D. et al. Consolidated Health Economic Evaluation Reporting Standards 2022 (CHEERS 2022) statement: updated reporting guidance for health economic evaluations. *Int. J. Technol. Assess. Health Care* **38**, e13 (2022).
60. Occhipinti, J. A. et al. Which social, economic, and health sector strategies will deliver the greatest impacts for youth mental health and suicide prevention? protocol for an advanced, systems modelling approach. *Front. Psychiatry* **12**, 759343 (2021).
61. Lawson, K. D. et al. A dynamic approach to economic priority setting to invest in youth mental health and guide local implementation: economic protocol for eight system dynamics policy models. *Front. Psychiatry* **13**, 835201 (2022).
62. Freebairn, L. et al. Participatory methods for systems modeling of youth mental health: implementation protocol. *JMIR Res. Protoc.* **11**, e32988 (2022).
63. Powell, M. *The BOBYQA Algorithm for Bound Constrained Optimization Without Derivatives: Technical Report* (Department of Applied Mathematics and Theoretical Physics, Univ. Cambridge, 2009).
64. Kessler, R. C. et al. Short screening scales to monitor population prevalences and trends in non-specific psychological distress. *Psychol. Med.* **32**, 959–976 (2002).
65. Gamst-Klaussen, T., Lamu, A. N., Chen, G. & Olsen, J. A. Assessment of outcome measures for cost-utility analysis in depression: mapping depression scales onto the EQ-5D-5L. *BJPsych Open* **4**, 160–166 (2018).
66. McCaffrey, N., Kaambwa, B., Currow, D. C. & Ratcliffe, J. Health-related quality of life measured using the EQ-5D-5L: South Australian population norms. *Health Qual. Life Outcomes* **14**, 133 (2016).
67. Goodman, A. & Goodman, R. Strengths and Difficulties Questionnaire as a dimensional measure of child mental health. *J. Am. Acad. Child Adolesc. Psychiatry* **48**, 400–403 (2009).
68. Sharma, R., Gu, Y., Sinha, K., Aghdaee, M. & Parkinson, B. Mapping the Strengths and Difficulties questionnaire onto the child health utility 9D in a large study of children. *Qual. Life Res.* **28**, 2429–2441 (2019).
69. O'Loughlin, R. et al. Comparing the psychometric performance of generic paediatric health-related quality of life instruments in children and adolescents with ADHD, anxiety and/or depression. *Pharmacoeconomics* **42**, 57–77 (2024).
70. Jones, R. et al. Comparative psychometric performance of common generic paediatric health-related quality of life instrument descriptive systems: results from the Australian paediatric multi-instrument comparison study. *Pharmacoeconomics* **42**, 39–55 (2024).
71. Oliveira Goncalves, A. S., Werdin, S., Kurth, T. & Panteli, D. Mapping studies to estimate health-state utilities from nonpreference-based outcome measures: a systematic review on how repeated measurements are taken into account. *Value Health* **26**, 589–597 (2023).
72. *Guidelines for Preparing a Submission to the Pharmaceutical Benefits Advisory Committee Version 5.0* (Pharmaceutical Benefits Advisory Committee, 2016).



73. Shirowa, T. et al. International survey on willingness-to-pay (WTP) for one additional QALY gained: what is the threshold of cost effectiveness? *Health Econ.* **19**, 422–437 (2010).
74. Vallejo-Torres, L. et al. On the estimation of the cost-effectiveness threshold: why, what, how? *Value Health* **19**, 558–566 (2016).
75. O'Donnell, J. *Mapping Social Cohesion 2023* (Scanlon Foundation Research Institute, 2023).
76. Markus, A. & Arnup, J. *Mapping Social Cohesion 2009—The Scan Foundation Surveys* (Monash Univ., 2010).
77. Growing up in Australia—The Longitudinal Study of Australian Children—data and documentation. *The Australian Institute of Family Studies* <https://growingupinaustralia.gov.au/data-and-documentation> (2018).
78. Growing up in Australia: Longitudinal Study of Australian Children (LSAC) Release 9.1 C2 (Waves 1–9C) [ADA Dataverse]. *The Australian Institute of Family Studies* <https://growingupinaustralia.gov.au/> (2022).
79. Homelessness operational groups (OPGP). *Australian Bureau of Statistics* <https://www.abs.gov.au/census/guide-census-data/census-dictionary/2021/variables-topic/housing/homelessness-operational-groups-opgp> (2021).
80. Estimating homelessness: census. *Australian Bureau of Statistics* <https://www.abs.gov.au/statistics/people/housing/estimating-homelessness-census> (2021).
81. Labour force, Australia, detailed. *Australian Bureau of Statistics* <https://www.abs.gov.au/statistics/labour/employment-and-unemployment/labour-force-australia-detailed/> (2023).
82. Labour force, Australia. *Australian Bureau of Statistics* <https://www.abs.gov.au/statistics/labour/employment-and-unemployment/labour-force-australia/> (2023).
83. Education and work [Tablebuilder]. *Australian Bureau of Statistics* <https://www.abs.gov.au/statistics/microdata-tablebuilder/tablebuilder> (2014–2022).
84. *Enrolments by Grade 2008 to 2021* (Australian Curriculum Assessment and Reporting Authority, 2021).
85. *Education and Work, Australia (Reference Period: May 2021)* (Australian Bureau of Statistics, 2021).
86. National study of mental health and wellbeing. *Australian Bureau of Statistics* <https://www.abs.gov.au/statistics/health/mental-health/national-study-mental-health-and-wellbeing/2020-2022> (2022).
87. National Survey of Mental Health and Wellbeing: summary of results, 2007. *Australian Bureau of Statistics* <https://www.abs.gov.au/statistics/health/mental-health/national-study-mental-health-and-wellbeing/2007> (2008).
88. Goodhue, R., Dakin, P. & Noble, K. What's in the nest? Exploring Australia's wellbeing framework for children and young people. *Canberra ARACY* [https://ihcsupportagency.org.au/wp-content/uploads/2021/09/WhatsInTheNest2021\\_EVersion.pdf](https://ihcsupportagency.org.au/wp-content/uploads/2021/09/WhatsInTheNest2021_EVersion.pdf) (2021).
89. Hogan, M., Hatfield-Dodds, L., Barnes, L. & Struthers, K. *Systems Leadership for Child Wellbeing Project: Stage 1 Synthesis Report* (Every Child and Australia and New Zealand School of Government (ANZSOG), Australia, 2021).
90. Veronese, N. et al. Interventions for reducing loneliness: an umbrella review of intervention studies. *Health Social Care Commun.* **29**, e89–e96 (2021).
91. Li, N., Peng, J. & Li, Y. Effects and moderators of Triple P on the social, emotional, and behavioral problems of children: systematic review and meta-analysis. *Front. Psychol.* **12**, 709851 (2021).
92. Sampaio, F. et al. Population cost-effectiveness of the Triple P parenting programme for the treatment of conduct disorder: an economic modelling study. *Eur. Child Adolesc. Psychiatry* **27**, 933–944 (2018).
93. Crosland, P. et al. Cost-effectiveness of system-level mental health strategies for young people in the Australian Capital Territory: a dynamic simulation modelling study. *Lancet Psychiatry* **11**, 123–133 (2024).
94. Loblay, V. et al. Enhancing equitable engagement for digital health promotion: Lessons from evaluating a childrearing app in Indonesia. *Digit. Health* **9**, 20552076231222112 (2023).
95. LaMonica, H. M. et al. Developing a parenting app to support young children's socioemotional and cognitive development in culturally diverse low- and middle-income countries: protocol for a co-design study. *JMIR Res. Protoc.* **11**, e39225 (2022).
96. Walsh, E. H., McMahon, J. & Herring, M. P. Research review: the effect of school-based suicide prevention on suicidal ideation and suicide attempts and the role of intervention and contextual factors among adolescents: a meta-analysis and meta-regression. *J. Child Psychol. Psychiatry* **63**, 836–845 (2022).
97. Walsh, E. H., Herring, M. P. & McMahon, J. A systematic review of school-based suicide prevention interventions for adolescents, and intervention and contextual factors in prevention. *Prev. Sci.* **24**, 365–381 (2023).
98. Breet, E., Matooane, M., Tomlinson, M. & Bantjes, J. Systematic review and narrative synthesis of suicide prevention in high-schools and universities: a research agenda for evidence-based practice. *BMC Public Health* **21**, 1116 (2021).
99. Dunning, D. et al. Do mindfulness-based programmes improve the cognitive skills, behaviour and mental health of children and adolescents? An updated meta-analysis of randomised controlled trials. *Evid. Based Ment. Health* **25**, 135–142 (2022).
100. Caldwell, D. M. et al. School-based interventions to prevent anxiety and depression in children and young people: a systematic review and network meta-analysis. *Lancet Psychiatry* **6**, 1011–1020 (2019).
101. Werner-Seidler, A. et al. School-based depression and anxiety prevention programs: an updated systematic review and meta-analysis. *Clin. Psychol. Rev.* **89**, 102079 (2021).
102. Gunawardena, H., Voukelatos, A., Nair, S., Cross, S. & Hickie, I. B. Efficacy and effectiveness of universal school-based wellbeing interventions in australia: a systematic review. *Int. J. Environ. Res. Public Health* **20**, 6508 (2023).
103. Schmidt, M. et al. Universal mental health interventions for children and adolescents: a systematic review of health economic evaluations. *Appl. Health Econ. Health Policy* **18**, 155–175 (2020).
104. Foulkes, L. & Stringaris, A. Do no harm: can school mental health interventions cause iatrogenic harm? *BJPsych Bull* **47**, 267–269 (2023).
105. Frederick, D. E. & VanderWeele, T. J. Supported employment: meta-analysis and review of randomized controlled trials of individual placement and support. *PLoS ONE* **14**, e0212208 (2019).
106. Brinchmann, B. et al. A meta-regression of the impact of policy on the efficacy of individual placement and support. *Acta Psychiatr. Scand.* **141**, 206–220 (2020).
107. Bond, G. R. *Cost-Effectiveness of Individual Placement and Support* (Advancing State Policy Integration for Recovery and Employment, 2023).
108. Zheng, K., Stern, B. Z., Wafford, Q. E. & Kohli-Lynch, C. N. Trial-based economic evaluations of supported employment for adults with severe mental illness: a systematic review. *Adm. Policy Ment. Health* **49**, 440–452 (2022).
109. Thomson, R. M. et al. How do income changes impact on mental health and wellbeing for working-age adults? A systematic review and meta-analysis. *Lancet Public Health* **7**, e515–e528 (2022).

## Acknowledgements

We acknowledge the attendees of the participatory systems modeling workshops who provided vital skills and expertise, particularly young people with a lived experience of mental health conditions and their carers. We acknowledge the staff of the Statistical Services Branch, Queensland Health, the Healthcare Improvement Unit, Queensland Health and the Mental Health Alcohol and Other Drugs Branch, Queensland Health for the datasets used for this project: the Emergency Data Collection, the Queensland Hospital Admitted Patient Data Collection and the Consumer Integrated Mental Health Application. We acknowledge the staff of the Australian Curriculum, Assessment and Reporting Authority for providing Student Enrolment and Senior Secondary Outcomes data, and the staff of Brisbane South Primary Health Network for providing headspace occasions of service data. This Analysis uses unit record data from Growing Up in Australia, the Longitudinal Study of Australian Children. The study is conducted in partnership between the Department of Social Services (DSS), the Australian Institute of Family Studies (AIFS) and the Australian Bureau of Statistics (ABS). The findings and views reported in this Analysis are those of the authors and should not be attributed to DSS, AIFS or the ABS. I.B.H. discloses support for the research of this work from NHMRC (APP2016346). D.A.M. discloses partial salary support for the research of this work from the Svare Chair in Health Economics, Value and Impact.

## Author contributions

P.C.: conceptualization, methodology, software, formal analysis, investigation, writing—original draft, writing for Box 3, visualization. N.H.: methodology, software, formal analysis, investigation, data curation, writing—review and editing. K.-H.N.: validation, writing—review and editing, investigation. K.T.: validation, writing—review and editing, investigation, data curation. S.H.H.: methodology, software, data curation, writing—review and editing. C.V.: methodology, software, investigation, writing—review and editing. AS: writing—review and editing, supervision. J.v.R.: writing for Box 1, writing—review and editing. S.R.: writing—review and editing. F.I.: writing—review and editing. V.L.: writing—review and editing. O.I.: investigation, writing—review and editing. S.P.: writing—review and editing. Y.J.C.S.: conceptualization, supervision, project administration, funding acquisition and resources. S.M.: writing for Box 2, writing—review and editing. J.M.G.P.: writing—review and editing. A.C.: writing—review and editing. D.A.M.: writing—review and editing, supervision. I.B.H.: conceptualization, writing—review and editing, supervision, funding acquisition. J.-A.O.: conceptualization, resources, supervision, project administration, funding acquisition, writing—review and editing.

## Funding

Open access funding provided by the University of Sydney

## Competing interests

This study was conducted under the Right care, first time, where you live Program (Brain and Mind Centre, University of

Sydney), supported by a AUD\$12.8 million grant provided by the BHP Foundation. The program is developing decision support infrastructure to guide investments and actions to foster the mental health and well-being of young people in their communities. J.-A.O. is both Head of Systems Modelling, Simulation & Data Science at the University of Sydney's Brain and Mind Centre and Managing Director of Computer Simulation & Advanced Research Technologies (CSART). I.H. is the Co-Director, Health and Policy at the Brain and Mind Centre (BMC) University of Sydney. The BMC operates an early-intervention youth services at Camperdown under contract to headspace. He is the Chief Scientific Advisor to, and a 3.2% equity shareholder in, InnoWell Pty Ltd, which aims to transform mental health services through the use of innovative technologies. V.L. is a board member for Matana Foundation, a philanthropic organization that provides funding to programs for disadvantaged young people. She does not receive any financial benefit for this role.

## Additional information

**Supplementary information** The online version contains supplementary material available at <https://doi.org/10.1038/s44220-025-00459-7>.

**Correspondence and requests for materials** should be addressed to Paul Crosland.

**Peer review information** *Nature Mental Health* thanks Laura Shields-Zeeman and the other, anonymous reviewer(s) for their contribution to the peer review of this work.

**Reprints and permissions information** is available at [www.nature.com/reprints](http://www.nature.com/reprints).

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2025

<sup>1</sup>Youth Mental Health and Technology, Brain and Mind Centre, Faculty of Medicine and Health, Central Clinical School, University of Sydney, Sydney, New South Wales, Australia. <sup>2</sup>Faculty of Medicine, The University of Queensland, Brisbane, Queensland, Australia. <sup>3</sup>Global Brain Health Institute, Trinity College, Dublin, Republic of Ireland. <sup>4</sup>Health Research Institute, University of Canberra, Canberra, Australian Capital Territory, Australia. <sup>5</sup>Thriving Queensland Kids Partnership, Brisbane, Queensland, Australia. <sup>6</sup>Children's Health Queensland Hospital and Health Service, Brisbane, Queensland, Australia. <sup>7</sup>Intelligence Unit, Better Health North Queensland, Mackay Hospital and Health Service, Queensland Government, Mackay, Queensland, Australia. <sup>8</sup>Cumming School of Medicine, University of Calgary and Alberta Children's Hospital Research Institute, Calgary, Alberta, Canada. <sup>9</sup>Computer Simulation and Advanced Research Technologies, Camperdown, New South Wales, Australia. <sup>10</sup>These authors contributed equally: Ian B. Hickie, Jo-An Occhipinti. ✉ e-mail: [paul.crosland@sydney.edu.au](mailto:paul.crosland@sydney.edu.au)

Reporting Summary

Nature Portfolio wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Portfolio policies, see our [Editorial Policies](#) and the [Editorial Policy Checklist](#).

Statistics

For all statistical analyses, confirm that the following items are present in the figure legend, table legend, main text, or Methods section.

- |                                     |   |
|-------------------------------------|---|
| n/a                                 | Confirmed   |
| <input checked="" type="checkbox"/> | <input type="checkbox"/> The exact sample size ( <i>n</i> ) for each experimental group/condition, given as a discrete number and unit of measurement   |
| <input checked="" type="checkbox"/> | <input type="checkbox"/> A statement on whether measurements were taken from distinct samples or whether the same sample was measured repeatedly  |
| <input checked="" type="checkbox"/> | <input type="checkbox"/> The statistical test(s) used AND whether they are one- or two-sided<br><i>Only common tests should be described solely by name; describe more complex techniques in the Methods section.</i>   |
| <input checked="" type="checkbox"/> | <input type="checkbox"/> A description of all covariates tested   |
| <input type="checkbox"/>            | <input checked="" type="checkbox"/> A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons   |
| <input checked="" type="checkbox"/> | <input type="checkbox"/> A full description of the statistical parameters including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient) AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals) |
| <input checked="" type="checkbox"/> | <input type="checkbox"/> For null hypothesis testing, the test statistic (e.g. <i>F</i> , <i>t</i> , <i>r</i> ) with confidence intervals, effect sizes, degrees of freedom and <i>P</i> value noted<br><i>Give P values as exact values whenever suitable.</i>                     |
| <input checked="" type="checkbox"/> | <input type="checkbox"/> For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings   |
| <input checked="" type="checkbox"/> | <input type="checkbox"/> For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes   |
| <input checked="" type="checkbox"/> | <input type="checkbox"/> Estimates of effect sizes (e.g. Cohen's <i>d</i> , Pearson's <i>r</i> ), indicating how they were calculated   |

Our web collection on [statistics for biologists](#) contains articles on many of the points above.

Software and code

Policy information about [availability of computer code](#)

- |                 |   |
|-----------------|---|
| Data collection | This simulation model used data from a variety of sources, all of which are referenced in the supplementary material.   |
| Data analysis   | The model file is not available online however we welcome sharing of our models and methods with interested researchers who wish to collaborate with us. One of the reasons for a collaborative approach to sharing our models is the ability to provide training and support to users. |

For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors and reviewers. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Portfolio [guidelines for submitting code & software](#) for further information.

Data

Policy information about [availability of data](#)

All manuscripts must include a [data availability statement](#). This statement should provide the following information, where applicable:

- Accession codes, unique identifiers, or web links for publicly available datasets
- A description of any restrictions on data availability
- For clinical datasets or third party data, please ensure that the statement adheres to our [policy](#)

This simulation model used data from a variety of sources, all of which are referenced in the supplementary material.

## Research involving human participants, their data, or biological material

Policy information about studies with [human participants or human data](#). See also policy information about [sex, gender \(identity/presentation\), and sexual orientation](#) and [race, ethnicity and racism](#).

Reporting on sex and gender	This simulation model does not distinguish between sex or gender.
Reporting on race, ethnicity, or other socially relevant groupings	Not applicable - the present analysis does not distinguish between these subpopulations.
Population characteristics	The simulation model structure disaggregates the modelled population by age, level of distress and mental health diagnostic category as described in the supplementary material.
Recruitment	Not applicable. The modelled population is a simulated group of people based on the population characteristics of the region.
Ethics oversight	Sydney Local Health District / Royal Prince Alfred Hospital - X21-0151 & 2021/ETH00553

Note that full information on the approval of the study protocol must also be provided in the manuscript.

## Field-specific reporting

Please select the one below that is the best fit for your research. If you are not sure, read the appropriate sections before making your selection.

☐ Life sciences ☒ Behavioural & social sciences ☐ Ecological, evolutionary & environmental sciences

For a reference copy of the document with all sections, see [nature.com/documents/nr-reporting-summary-flat.pdf](https://nature.com/documents/nr-reporting-summary-flat.pdf)

## Behavioural & social sciences study design

All studies must disclose on these points even when the disclosure is negative.

Study description	System dynamics simulation modelling in Stella Architect informed by participatory model-building processes
Research sample	Simulated population of Brisbane South Primary Health Network region
Sampling strategy	Not applicable
Data collection	Variety of sources were used to inform the quantitative aspects of the simulation model as described in supplementary material.
Timing	Historical data based on 2011 to 2023 and modelled forecasts from 2024 onwards
Data exclusions	Not applicable. Model sectors included fully described in supplementary material.
Non-participation	Not applicable
Randomization	Not applicable

## Reporting for specific materials, systems and methods

We require information from authors about some types of materials, experimental systems and methods used in many studies. Here, indicate whether each material, system or method listed is relevant to your study. If you are not sure if a list item applies to your research, read the appropriate section before selecting a response.

### Materials & experimental systems

n/a	Involved in the study
<input checked="" type="checkbox"/>	<input type="checkbox"/> Antibodies
<input checked="" type="checkbox"/>	<input type="checkbox"/> Eukaryotic cell lines
<input checked="" type="checkbox"/>	<input type="checkbox"/> Palaeontology and archaeology
<input checked="" type="checkbox"/>	<input type="checkbox"/> Animals and other organisms
<input checked="" type="checkbox"/>	<input type="checkbox"/> Clinical data
<input checked="" type="checkbox"/>	<input type="checkbox"/> Dual use research of concern
<input checked="" type="checkbox"/>	<input type="checkbox"/> Plants

### Methods

n/a	Involved in the study
<input checked="" type="checkbox"/>	<input type="checkbox"/> ChIP-seq
<input checked="" type="checkbox"/>	<input type="checkbox"/> Flow cytometry
<input checked="" type="checkbox"/>	<input type="checkbox"/> MRI-based neuroimaging