


A network outcome analysis of psychological risk factors driving suicide risk in emergency department patients

Received: 27 September 2023

Accepted: 15 January 2025

Published online: 20 February 2025

 Check for updatesRené Freichel¹✉, Matthew K. Nock^{2,3,4} & Brian A. O'Shea⁵

Different theories of suicide propose somewhat different psychological factors that lead to suicidal thoughts and behaviors. For example, Beck's theory highlights hopelessness, while the interpersonal–psychological theory of suicide emphasizes burdensomeness, lack of belonging and fearlessness about death. Surprisingly, few studies have tested which theoretically proposed psychological factors are most predictive of suicidal thoughts and behaviors. We used network outcome analysis to disentangle the effects of these constructs in predicting suicidal ideation, suicide plans and attempts. Participants were 1,412 patients presenting to an emergency department with psychiatric complaints, with follow-up assessments one month and six months ($n = 938$) later. Here we showed that different psychological factors predicted different parts of the continuum of suicidal thoughts and behaviors. Lack of belongingness was most predictive of suicidal ideation (partial correlation (pcor) = 0.14), acquired capability for death (that is, fearlessness of death) was most predictive of suicide planning (pcor = 0.08), and hopelessness was most predictive of suicide attempts (pcor = 0.12). Individuals' explicit associations with death (that is, death = me) prospectively predicted all three outcomes (pcor = 0.13–0.23). The occurrence of suicidal thoughts and behaviors is best predicted using constructs from several different theories of suicide. Future theoretical and empirical work should integrate components of existing theories.

Suicide is among the leading causes of death¹, with rates in the United States increasing from 1999 through 2018 (ref. 2). Different theoretical frameworks have been proposed to explain the role of psychological factors (that is, cognitions, emotions) in the transition from suicidal ideation to suicide plans and suicide attempts. Initially, simple univariate theories, such as the hopelessness theory by Beck³, pose that negative expectations for oneself in the future lead to suicidal behavior. Slightly more complex theories have emerged that contextualize a range of co-occurring psychological affective states in the prediction of suicide-related

outcomes. Joiner's interpersonal theory of suicide^{4,5} proposes that three key factors explain the emergence of suicidal behavior, hypothesizing that perceived burdensomeness only in combination with thwarted belongingness will lead to suicidal ideation, whereas fearlessness about injury/death predicts the transition from suicidal thought to behavior. The integrated motivational–volitional model of suicidal behavior⁶ hypothesizes that specific psychological states, such as entrapment, predict suicidal ideation, and different volitional moderators, including fearlessness about death and impulsivity, predict suicidal behavior.

¹Department of Psychology, University of Amsterdam, Amsterdam, the Netherlands. ²Department of Psychology, Harvard University, Cambridge, MA, USA. ³Department of Psychiatry, Massachusetts General Hospital, Boston, MA, USA. ⁴Mental Health Research, Franciscan Children's, Brighton, MA, USA.

⁵School of Psychology, University of Nottingham, Nottingham, England. ✉e-mail: r.freichel@uva.nl

Table 1 | Comparison of ideation and no-ideation groups at 6-month follow-up (n=938)

Measure	No ideation at 6-month follow-up	Ideation at 6-month follow-up	t	P	95% confidence interval	
	M (s.d.)	M (s.d.)			Lower	Upper
n	456	482				
Adverse psychological state						
IntEmo	3.93 (1.18)	4.23 (0.99)	−4.09	<0.01	−0.43	−0.15
Hopeless	2.98 (1.40)	3.8 (1.18)	−9.59	<0.01	−0.98	−0.65
TimeRunOut	3.07 (1.43)	3.76 (1.32)	−7.66	<0.01	−0.87	−0.51
TrappedProb	3.74 (1.32)	4.3 (1.00)	−7.37	<0.01	−0.72	−0.42
Anxious	3.52 (1.30)	4.1 (1.05)	−7.41	<0.01	−0.72	−0.42
LifImpossible	3.09 (1.46)	3.76 (1.25)	−7.47	<0.01	−0.84	−0.49
Impulsive	3.08 (1.37)	3.54 (1.28)	−5.25	<0.01	−0.63	−0.29
Burden	2.92 (1.46)	3.77 (1.30)	−9.39	<0.01	−1.03	−0.67
NotBelong	3.10 (1.41)	4.04 (1.18)	−11.09	<0.01	−1.11	−0.78
NotAfraidDeath	2.84 (1.47)	3.33 (1.48)	−5.08	<0.01	−0.68	−0.30
NoSocSupport	2.32 (1.35)	2.67 (1.38)	−4.03	<0.01	−0.53	−0.18
NoPurpose	3.00 (1.33)	3.55 (1.32)	−6.29	<0.01	−0.71	−0.37
Explicit–implicit association						
ExplAssoDeath	34.56 (31.29)	56.33 (30.05)	−10.86	<0.01	−25.71	−17.84
IatDScore	−0.42 (0.32)	−0.35 (0.34)	−3.39	0.01	−0.12	−0.03

The 95% confidence interval is for the difference in means between the two groups. Two-sided t tests were conducted without corrections for multiple comparisons. Different adverse psychological states are abbreviated as follows (Supplementary Table 1): IntEmo, intensity of emotions/emotional reactivity (I feel emotions very intensely); Hopeless, hopelessness (I feel hopeless about the future); TimeRunOut, despair/entrapment (time is running out for me); TrappedProb, feeling trapped (I feel trapped by my problems); Anxious, anxiety (I am constantly anxious and on edge); LifImpossible, desperation (life is impossible if things don't change); Impulsive, impulsivity (I often act without thinking when I am upset); Burden, burdensomeness (I am a burden to others); NotBelong, lack of belongingness (I feel like I don't belong); NotAfraidDeath, acquired capability for death (I am not afraid of death or injury); No SocSupport, social support (I have no social support); NoPurpose, purposefulness (I do not have a sense of direction and purpose in life). All group differences were significant at the ≤ 0.01 level. Standard deviations are reported in parentheses. All psychological state items range from 1 to 5. The explicit association with death score (ExplAssoDeath) ranges on a scale from 0 to 100, and the IAT D score (IatDScore) ranges from −2 to 2.

Most current theories of suicide do not specifically articulate why these factors (for example, hopelessness, lack of belongingness) lead to suicide rather than other behaviors that might serve the same function as suicide such as escaping from aversive psychological states, through alcohol or substance use or other forms of self-injury. One theoretical model proposes that one's identification with specific behaviors (for example, suicide, alcohol use) may predispose people toward engaging in these specific behaviors when in distress⁷. Such identification can be measured via implicit association tests⁸ or explicit self-reports of such associations (for example, single-item measure: 'To what extent do you think of death as being like you versus not like you?'). Implicit death association tests measure the extent to which individuals associate the concept 'death' (as opposed to 'life') with the self. Such computerized reaction-time-based tests (detailed description in Methods) were designed to assess implicit biases, which some have described as more automatic/less controllable than explicit biases^{9,10}. Implicit biases represent markers of vulnerability for suicide attempts¹¹ that may overcome challenges with self-reported suicide risk assessment. Such challenges include deliberate faking, reluctance to disclose suicidal ideation, unawareness of the true extent of suicide risk and the transient nature of suicidal thoughts, where an individual might accurately report no suicidal thoughts at one moment, only to experience them again soon after¹². Although no formal theory has been developed or specifically added implicit biases into their models of suicide, Nock¹³ has proposed that alternative methods such as the Death/Suicide Implicit Association Test (IAT)^{11,14} might provide insight into markers of vulnerability for suicide risk¹¹.

Previous studies have demonstrated that implicit and explicit associations with the concepts of suicide and non-suicidal self-injury are specifically predictive of engagement in these behaviors both concurrently and prospectively (that is, over 6–12 months)^{15,16}.

However, surprisingly few studies have examined psychological factors from different theories together to examine which are most predictive of suicidal outcomes. Contemporary theories of suicide integrate many intercorrelated and overlapping factors representing adverse psychological states¹⁷. Meehl¹⁸ further described this problem of intercorrelation among many variables of interest ('crud factor') when using standard statistical approaches for theory testing. Thus, there is a need to better understand the unique contribution of different cognitive, social and affective predictors across theories.

In the past decade, network analysis has been proposed as a statistical toolbox^{19,20} for identifying conditional (that is, partial) associations between a range of variables. Network models consist of nodes (that is, variables) that are connected through edges (that is, pairwise conditional associations). The present study uses a novel network analysis approach, namely network outcome analysis²¹, to examine predictive associations between a range of different psychological factors and the development of suicidal ideation, suicide plans, and suicide attempts. A crucial benefit of network analysis concerns the separation of direct effects (for example, edge between impulsivity and suicidal ideation) and indirect effects (for example, association between impulsivity and suicidal ideation through other cognitions)²¹. Based on prior research, we predicted that hopelessness, burdensomeness, lack of belongingness, entrapment, and identification with death would be the strongest proximate predictors of suicide-related outcomes.

Results

Sample characteristics and group comparison

Participants were patients presenting to the Massachusetts General Hospital Emergency Department (ED) between February 2015 and March 2017 due to severe psychiatric distress. There were two follow-up surveys conducted over the telephone or email, 1 month and 6 months




Fig. 1 | Symptom network with suicidal ideation at 6-month follow-up (hyperparameter = 0.25). The connections (positive edges shown in blue) represent conditional associations. The thickness of each line describes the strength of the association. All edge weights (strengths of associations) can be

found in Supplementary Tables 7–9. The red square indicates the suicide-related outcome (suicidal ideation) assessed at the 6-month follow-up; all other variables have been assessed at baseline.

after the baseline visit. All psychological factors (description in Methods) were evaluated at baseline, while all suicide-related outcomes (that is, ideation, suicide plan, attempt) were assessed during both follow-up surveys. The sample at baseline ($n = 1,412$) was predominantly male (54.32%; female, 43.96%; transgender, 1.72%), young ($M = 34.95$, $s.d. = 13.57$) and relatively diverse with respect to ethnicity and race (non-Hispanic white, 68.0%; non-Hispanic African American, 7.22%; Hispanic other, 7.15%; non-Hispanic other, 6.22%; Hispanic white, 5.79%; non-Hispanic Asian, 3.86%; and other race or ethnicity, 1.76%). At the 1-month follow-up survey, there were complete data for reports of suicidal ideation for 900 participants (63.74%). By the 6-month follow-up, 938 participants (66.43%) completed reports on measures of suicidal ideation. A supplemental analysis (Supplementary Table 3) showed that older age, female gender, lower income and a lifetime history of a suicide attempt were significantly associated with slight increases in the likelihood of dropout.

Table 1 describes participants' responses to assessments about psychological factors as well as their explicit and implicit association with death. We examined group differences in the mean level on the basis of whether individuals reported suicidal ideation (Table 1), suicide plan (Supplementary Table 5) and a suicide attempt (Supplementary Table 6) at the 6-month follow-up. Participants with suicidal ideation, suicide plans and suicide attempts at 6-month follow-up reported generally more adverse psychological states and a stronger explicit association with death compared with individuals who did not report any of these suicide-related outcomes at 6-month follow-up. Participants' implicit association with death is stronger in individuals experiencing suicidal ideation; however, there were no significant group differences in this implicit score for the comparison by suicide plans ($P = 0.29$) and suicide attempts ($P = 0.52$).

Network outcome analysis

The network outcome analysis consistently identified direct predictive associations between explicit association with death at baseline and

all suicide-related outcomes (suicidal ideation 100% retrieved, suicide plan 99% retrieved, suicide attempt 65% retrieved) at the 6-month follow-up. We present all network models with the 1-month follow-up outcomes in Supplementary Figs. 1–3. These direct associations with explicit association with death also hold at the 1-month follow-up, except for suicide attempt.

Figure 1 displays the network model with suicidal ideation at the 6-month follow-up. Importantly, lack of belongingness constitutes the single adverse psychological state that shows predictive associations (retrieved in 95% of bootstraps) with suicidal ideation at the 6-month follow-up visit. Lack of belongingness is strongly associated with feelings of burdensomeness (100% retrieved). Implicit associations with death and several adverse psychological states, including the acquired capability for death, lack of belongingness ('I feel like I don't belong') and purposefulness, are most strongly associated with individuals' explicit association with death, which in turn predicts suicidal ideation (100% retrieved).

The network outcome analyses for suicide plan and suicide attempt are shown in Fig. 2. Interestingly, in addition to the explicit association with death, the acquired capability for death was the only adverse psychological state that showed direct predictive associations with suicide plans (48% retrieved) at the 6-month follow-up. For the prediction of suicide attempts, hopelessness at baseline was the single key predictor (47% retrieved) in addition to individuals' explicit association with death (65% retrieved).

The estimated network structures and all-important edges with the prospective suicide outcomes presented in the preceding showed sufficient stability in our bootstrapping analysis (Supplementary Figs. 7–9). The network outcome models that were estimated with a conservative hyperparameter ($\gamma = 0.5$) are shown in Supplementary Figs. 4–6. This conservative estimation process showed consistent associations between all suicide-related outcomes and individuals' explicit association with death. Networks that included sex and age as covariates in the network largely replicated the results (Supplementary Figs. 10–12).




Fig. 2 | Symptom network with suicidal plan and suicide attempt at 6-month follow-up. a, Suicide plans (SuiPlan) at the 6-month follow-up. **b**, Suicide attempts (SuiAttempt) at the 6-month follow-up. All other nodes have been assessed at baseline.

Discussion

The present study used an innovative methodological approach, prospective network outcome analysis, to identify proximal risk factors of suicide-related outcomes, specifically suicidal ideation, suicide plans and suicide attempts. The crucial finding from our study concerns the specificity of associations found between theory-derived psychological states and suicide-related outcomes.

Individuals' explicit association with death was consistently associated with all suicide-related outcomes, and it emerged as an important bridge node connecting a range of psychological states (that is, lack of purpose, hopelessness) with suicidal ideation. Thus, individuals' explicit association with death may represent a rather general or more abstract cognition that may be the result of various more-specific negative psychological states, such as lack of purpose or hopelessness. Similarly, our findings suggest that implicit measures (suicide IAT D score) represent a convergent marker of explicit suicide cognitions that, in turn, predict suicide-related outcomes. This is in line with recent studies showing no evidence for direct associations between suicide IAT scores and suicide attempts in high-risk samples²².

A crucial finding from our study is the specificity of associations between suicide-related outcomes and adverse psychological states. In line with Joiner's interpersonal theory of suicide^{4,5}, suicidal ideation closely relates to feelings of thwarted belongingness that are strongly associated with perceived burdensomeness. This lack of social support and feelings of loneliness may lead to a sense of helplessness that ultimately contributes to suicidal ideation. During the progression from suicidal ideation to suicide attempt, individuals' feelings of not being afraid of death may represent one proximal factor that contributes to individuals' capability for suicide. This finding is in line with previous studies showing associations between acquired capability for death and past suicide attempts²³. Finally, our results show that hopelessness is the key adverse psychological state that is uniquely associated with suicide attempts. This is consistent with Beck's theory of hopelessness³; however, our study represents one of the first to show this specificity with attempts while controlling for (that is, partialing out) a range of other related factors (for example, feelings of not belonging).

Our findings suggest that thwarted belongingness along with the closely related concept of perceived burdensomeness are two critical factors closely connected to suicidal ideation. By contrast, the acquired capability for death appears to be more closely linked to suicidal intent (that is, suicide plans). Although our analysis does not directly examine moderation effects, these key components align with the integrated motivational–volitional model of suicidal behavior⁶, where they serve as motivational (thwarted belongingness, perceived burdensomeness) and volitional (acquired capability for death) moderators.

We believe that our study offers valuable insights for advancing future suicide research and clinical implications concerning the assessment of suicide risk¹. The present study is one of the first to simultaneously test the associations between psychological factors from across different theories and suicidal outcomes to determine which are more predictive of such outcomes. Our findings indicate that the occurrence of suicidal thoughts and behaviors is best predicted using constructs from several different theories of suicide. This underscores the importance of adopting an integrative approach in future studies aimed at predicting suicidal outcomes. We believe that the multimodal nature of our study represents a key strength as we included self-reported psychosocial variables (for example, lack of social support), behavioral measures (that is, implicit self-death associations) and prospectively assessed suicide-related cognitions and behaviors (that is, ideation, plan, attempt). In line with previous work¹⁶, the behavioral indicator of implicit suicide cognition did not provide additional predictive value for suicide-related outcomes beyond what was already captured by the explicit self-reported measure. Future research should explore whether additional behavioral measures, such as attentional biases toward suicide assessed through the Suicide Stroop Task²⁴ or behavioral indicators of executive functioning, including cognitive inflexibility²⁵, enhance the prediction of suicidal thoughts and behaviors beyond key self-reported risk factors.

Regarding clinical implications, our findings underscore the importance of addressing three conceptually modifiable and outcome-specific cognitions: hopelessness, lack of belongingness and the acquired capability for death. Clinical intervention studies are needed

to investigate two key questions: (1) the extent to which these cognitions are malleable through targeted interventions and (2) whether addressing these factors at specific risk stages of the process (that is, ideation → lack of belongingness; planning → acquired capability for death; attempt → hopelessness) can effectively prevent the progression from suicidal ideation to suicide attempts.

Several important limitations should be noted. First, our assessment of adverse psychological states and explicit association with death was based on individuals' self-report, which may naturally be biased, particularly in the high-stress context of an emergency room. The various theory-derived factors were either adapted from existing scales (Supplementary Table 1) or developed by clinical experts. As a result, some items lack empirical validation, and future work should aim to replicate our results using validated instruments. Second, there was a substantial rate of dropout of participants, with only 66.43% of individuals from the baseline providing valid responses to the 6-month follow-up survey. We used complete case analysis, and thus, our results may be biased by this high rate of attrition. Moreover, a subgroup of individuals chose not to participate in this study due to impairments related to their psychiatric symptoms²⁶, possibly including strong suicidal urges, and thus, there is a risk of non-participation bias. Third, our analyses integrated a prospective marker of suicide-related outcomes in the network to investigate unique associations with measures assessed at baseline. As a result, our findings should be viewed as temporal associations rather than directed causal effects. Future longitudinal analyses with a continuous assessment of different adverse psychological states and suicide-related outcomes at different measurement waves (that is, panel studies) are necessary to establish temporal within-person pathways²⁷. Fifth, while the present study assessed constructs from several prominent theories of suicide, it did not cover all relevant theoretical models. For example, the three-step theory of suicide²⁸ emphasizes the role of psychological pain, connectedness and acquired, dispositional and practical capability for suicide. Future studies should aim to replicate our findings while incorporating a broader range of theoretical models.

Our study was one of the first to simultaneously investigate the predictive utility of different domains of risk factors (that is, explicit associations, adverse psychological states, behavioral measures of implicit suicide cognition) in a high-risk sample for suicidal ideation, suicide plan and suicide attempts. Our results highlight the importance of attending to individuals' explicit association with death and key adverse states, namely, hopelessness, lack of belongingness and the acquired capability for death to prevent a transition from suicidal ideation to suicide attempts.

Methods

Data source and procedure

The study complies with all ethical regulations and received ethical approval (number 2014P001620/MGH) from Harvard University and Massachusetts General Hospital. Patients presenting due to severe psychiatric distress to the Massachusetts General Hospital Emergency Department (ED) (between February 2015 and March 2017) were approached by study research assistants. Reasons for not obtaining informed consent included impairment due to psychiatric complaints, discharge, refusal to participate, language barriers and various other reasons²⁶. To be included in the study, participants had to be adults (18 years or older; no upper age limit), have presented at the Acute Psychiatry Service, and have had access to a phone or email over the 6-month follow-up period. Exclusion criteria encompassed any condition that hinders an individual's capacity to provide informed consent and participate in the study, including (1) the inability to speak or read English, (2) severe cognitive impairments due to conditions, such as florid psychosis, intellectual disability, dementia, and acute intoxication, or (3) the presence of extremely agitated or violent behavior.

The attending psychiatrist on duty determined the applicability of these criteria, with the primary focus being the safety of both participants and research staff. Participants could receive a total compensation of \$30 for completing the in-person session and two follow-up surveys (\$10 for each), along with additional benefits such as parking and cafeteria vouchers.

After obtaining written informed consent, the baseline visit while in the emergency room included an evaluation from the treating clinician, brief clinical self-report measures (administered through a tablet), and the completion of a brief implicit association test²⁹. There were two follow-up surveys conducted over the telephone or email, one month and six months after the baseline visit. The study, sample, and enrollment procedures are described in more detail elsewhere²⁶. The data supporting this study are not publicly available due to their highly sensitive nature.

Measures

Suicide-related outcomes. The presence of suicidal ideation ('Did you have thoughts of killing yourself?') and suicide plan ('Did you ever make a plan to kill yourself?') were assessed through self-report at 1 month and 6 months post-ED visit. Presence of a suicide attempt at the 1-month and 6-month follow-up visits was assessed through both self-report ('Did you make a suicide attempt (that is, purposefully hurt yourself with at least some intent to die)?') and the review of medical records (indication of suicide attempt via either self-report or medical record was considered evidence of a suicide attempt). These three items assessing suicidal thoughts and behaviors were adapted from the Self-Injurious Thoughts and Behaviors Interview³⁰.

Psychological constructs. We assessed several theory-derived psychological factors that have been shown to correlate with suicidal thoughts and behaviors. Each was assessed on a five-point Likert scale (1 = exactly like me; 5 = not at all like me). We recoded the reversely scored items such that a higher score on all items indicates more negative/adverse psychological states/cognitions. All 12 items can be found in Supplementary Table 1. These included emotion reactivity ('when I experience emotions, I feel them very intensely'³¹), hopelessness ('I am hopeless about the future'; 'life is impossible if things don't change right away'³²), entrapment ('I feel trapped by my problems'³³), anxiety/agitation ('I am constantly anxious and on edge'³⁴), impulsiveness ('when I am upset, I often act without thinking'³⁵), burdensomeness ('I am a burden to others'³⁶), lack of belongingness ('I feel like I don't belong'), acquired capability for death ('I am not afraid of injury or death'³⁷) as well as the protective factors of belongingness and meaning in life³⁸ ('I have people who care about me and are there for me whenever I need them'; 'I have a sense of direction and purpose in life').

Explicit association with death. We assessed participants' explicit association with death through a single-item measure ('To what extent do you think of death as being 'like you' versus 'not like you?') on a scale from 0 to 100. The use of such semantic differential items that match the respective categories of the IAT is standard practice in implicit cognition research¹⁰, and these explicit measures have been used in previous work^{16,39}.

Death IAT. A brief version (5 minutes) of the Death IAT was used to assess participants' implicit association with death^{8,29}. In this computerized reaction-time-based task, participants are instructed to classify target words (for example, 'funeral') using a left and right keyboard button press. The target words are presented on the center of the screen, and the category labels ('life' or 'death'/'me' or 'not me') appear on the top left and right of the screen. The order of these categories changes across the two critical blocks, such that on one block, death- and me-related words are assigned the same key press, while on the other block, 'life' and 'me' share the same key press. A commonly used difference

score (D score⁴⁰) is computed, which indicates the relative strength of associations between the self and the concepts of death versus life. Higher D scores represent stronger death–me associations indicated through faster responses on the death/me compared with the life/me blocks³⁹. The split-half reliability of the IAT used in the current measure was 0.37. This is lower than previous reliability estimates for the Death IAT ($r = 0.61$, $n = 1,979$ (ref. 39)). A Spearman's rank correlation was computed to assess the relationship between the D scores and the explicit association with death. There was a positive correlation between the explicit and implicit measures, $\rho = 0.15$, $P < 0.01$.

Data analysis

Network outcome analysis. We estimated the networks using mixed graphical models⁴¹ that allow for both continuous (for example, psychological states) and binary (for example, suicide attempt) variables. The network consists of both nodes (variables) that are connected through edges (conditional associations, similar to partial correlations) between two pairs of variables that control for all other variables in the network. Following the ‘network outcome analysis’ approach by Blanken et al.²¹, we integrated the prospective outcomes (for example, suicide attempt at 6-month follow-up) as a node in the network. We estimated separate networks for each prospective outcome (that is, ideation, plan, attempt) to prevent multicollinearity issues that are common in network analysis. This network outcome approach allowed us to separate direct predictive pathways (that is, direct connection with the outcome) and indirect predictive associations (that is, connecting with the outcome through other nodes; predictive mediation).

Network estimation. Mixed graphical models are based on a node-wise regression approach in which a variable is predicted by all other variables in the model using a regularized generalized linear model framework⁴². Depending on the type and distribution of the variable (continuous with Gaussian distribution or binary with Bernoulli distribution), an appropriate link function was used. We used the least absolute shrinkage and selection operator regularization method to avoid spurious false-positive edges in the network. The extended Bayesian information criterion (EBIC) was used as the method to select the optimal tuning parameter (lambda). This EBIC model selection method tends to be a more conservative approach than model selection using cross-validation. An EBIC hyperparameter gamma determines the extent of regularization and level of sparsity. Hyperparameters close to 0.5 may be too conservative and miss true positive edges, whereas parameters close to 0 may lead to false-positive edges and err on the side of discovery⁴³. A moderately conservative default hyperparameter of 0.25 was chosen to increase specificity and facilitate interpretability⁴⁴.

Network stability. To estimate the robustness and stability of the estimated networks, we used a nonparametric bootstrapping stability analysis with 1,000 bootstraps⁴⁵. This involves re-estimating the model on sampled data (with replacement) across 1,000 iterations. The bootstrapped confidence interval around the edge weights (Supplementary Figs. 7–9) and the proportion of edge presence in all bootstraps describe the precision of these estimates. We report the retrieved percentage of edge presence in ‘Results’. All analyses were conducted in RStudio (R version 4.4.0)⁴⁶. Networks were estimated using the packages *bootnet*⁴⁷ and *mgm*⁴¹ and visualized using *qgraph*⁴⁸. We used the layout of the 6-month follow-up suicidal ideation network for all visualizations to facilitate visual comparisons. No additional thresholds or scaling parameters were used for the network visualization.

Reporting summary

Further information on research design is available in the Nature Portfolio Reporting Summary linked to this article.

Data availability

Data available upon reasonable request due to the sensitive nature of the data.

Code availability

All preprocessing and analyses were performed using RStudio (R version 4.4.0). Network estimation was conducted using the *bootnet* (version 1.6) and *mgm* (version 1.2-14) R packages. Network visualizations were created using *qgraph* (version 1.9.8). Supplemental analyses utilized the *stats* package (version 4.4.0). The Death IAT was implemented using JavaScript.

References

- Nock, M. K. et al. Suicide and suicidal behavior. *Epidemiol. Rev.* **30**, 133–154 (2008).
- Hedegaard H., Curtin S. C. & Warner M. *Increase in Suicide Mortality in the United States, 1999–2018* (NCHS, 2020).
- Beck, A. T. Hopelessness as a predictor of eventual suicide. *Ann. N. Y. Acad. Sci.* **487**, 90–96 (1986).
- Joiner T. E. *Why People Die by Suicide* (Harvard Univ. Press, 2005).
- Van Orden, K. A. et al. The interpersonal theory of suicide. *Psychol. Rev.* **117**, 575–600 (2010).
- O'Connor, R. C. & Kirtley, O. J. The integrated motivational–volitional model of suicidal behaviour. *Phil. Trans. R. Soc. Lond. B* **373**, 20170268 (2018).
- Nock, M. K. Why do people hurt themselves?: New insights into the nature and functions of self-injury. *Curr. Dir. Psychol. Sci.* **18**, 78–83 (2009).
- Nock, M. K. et al. Measuring the suicidal mind: implicit cognition predicts suicidal behavior. *Psychol. Sci.* **21**, 511–517 (2010).
- Freichel, R. & O'Shea, B. A. Suicidality and mood: the impact of trends, seasons, day of the week, and time of day on explicit and implicit cognitions among an online community sample. *Transl. Psychiatry* **13**, 157 (2023).
- Werntz, A. J., Steinman, S. A., Glenn, J. J., Nock, M. K. & Teachman, B. A. Characterizing implicit mental health associations across clinical domains. *J. Behav. Ther. Exp. Psychiatry* **52**, 17–28 (2016).
- Sohn, M. N., McMorris, C. A., Bray, S. & McGirr, A. The death–implicit association test and suicide attempts: a systematic review and meta-analysis of discriminative and prospective utility. *Psychol. Med.* **51**, 1789–1798 (2021).
- Glenn, C. R. et al. Implicit identification with death predicts change in suicide ideation during psychiatric treatment in adolescents. *J. Child Psychol. Psychiatry* **58**, 1319–1329 (2017).
- Nock, M. K. Recent and needed advances in the understanding, prediction, and prevention of suicidal behavior. *Depress. Anxiety* **33**, 460–463 (2016).
- Harrison, D. P., Stritzke, W. G. K., Fay, N., Ellison, T. M. & Hudaib, A. R. Probing the implicit suicidal mind: does the Death/Suicide Implicit Association Test reveal a desire to die, or a diminished desire to live? *Psychol. Assess.* **26**, 831–840 (2014).
- Glenn, C. R., Millner, A. J., Esposito, E. C., Porter, A. C. & Nock, M. K. Implicit identification with death predicts suicidal thoughts and behaviors in adolescents. *J. Clin. Child Adolesc. Psychol.* **48**, 263–272 (2019).
- Freichel, R., Kahveci, S. & O'Shea, B. How do explicit, implicit, and sociodemographic measures relate to concurrent suicidal ideation? A comparative machine learning approach. *Suicide Life Threat. Behav.* **54**, 49–60 (2024).

17. Millner, A. J., Robinaugh, D. J. & Nock, M. K. Advancing the understanding of suicide: the need for formal theory and rigorous descriptive research. *Trends Cogn. Sci.* **24**, 704–716 (2020).
18. Meehl, P. E. Appraising and amending theories: the strategy of Lakatosian defense and two principles that warrant it. *Psychol. Inq.* **1**, 108–141 (1990).
19. Freichel R. Symptom network analysis tools for applied researchers with cross-sectional and panel data—a brief overview and multiverse analysis. *Psychol. Rep.* <https://doi.org/10.1177/00332941231213649> (2023).
20. McNally, R. J. Network analysis of psychopathology: controversies and challenges. *Annu. Rev. Clin. Psychol.* **17**, 31–53 (2021).
21. Blanken, T. F., Borsboom, D., Penninx, B. W. & Van Someren, E. J. Network outcome analysis identifies difficulty initiating sleep as a primary target for prevention of depression: a 6 year prospective study. *Sleep* **43**, zsz288 (2020).
22. Rath, D. et al. Predicting suicidal behavior by implicit associations with death? Examination of the death IAT in two inpatient samples of differing suicide risk. *Psychol. Assess.* **33**, 287–299 (2021).
23. Malone, K. M. et al. Protective factors against suicidal acts in major depression: reasons for living. *Am. J. Psychiatry* **157**, 1084–1088 (2000).
24. Cha, C. B., Najmi, S., Park, J. M., Finn, C. T. & Nock, M. K. Attentional bias toward suicide-related stimuli predicts suicidal behavior. *J. Abnorm. Psychol.* **119**, 616–622 (2010).
25. Miranda, R., Gallagher, M., Bauchner, B., Vaysman, R. & Marroquin, B. Cognitive inflexibility as a prospective predictor of suicidal ideation among young adults with a suicide attempt history. *Depress. Anxiety* **29**, 180–186 (2012).
26. Nock, M. K. et al. Prediction of suicide attempts using clinician assessment, patient self-report, and electronic health records. *JAMA Netw. Open* **5**, e2144373 (2022).
27. Freichel, R., Wiers, R., O'Shea, B., McNally, R. J. & de Beurs, D. Between the group and the individual: the need for within-person panel study approaches in suicide research. *Psychiatry Res.* **330**, 115549 (2023).
28. Klonsky, E. D. & May, A. M. The three-step theory (3ST): a new theory of suicide rooted in the 'ideation-to-action' framework. *Int. J. Cogn. Ther.* **8**, 114–129 (2015).
29. Millner, A. J., Coppersmith, D. D. L., Teachman, B. A. & Nock, M. K. The brief death implicit association test: scoring recommendations, reliability, validity, and comparisons with the death implicit association test. *Psychol. Assess.* **30**, 1356–1366 (2018).
30. Nock, M. K., Holmberg, E. B., Photos, V. I. & Michel, B. D. Self-injurious thoughts and behaviors interview: development, reliability, and validity in an adolescent sample. *Psychol. Assess.* **19**, 309–317 (2007).
31. Nock, M. K., Wedig, M. M., Holmberg, E. B. & Hooley, J. M. The emotion reactivity scale: development, evaluation, and relation to self-injurious thoughts and behaviors. *Behav. Ther.* **39**, 107–116 (2008).
32. Beck, A. T., Weissman, A., Lester, D. & Trexler, L. The measurement of pessimism: the hopelessness scale. *J. Consult. Clin. Psychol.* **42**, 861–865 (1974).
33. Gilbert, P. & Allan, S. The role of defeat and entrapment (arrested flight) in depression: an exploration of an evolutionary view. *Psychol. Med.* **28**, 585–598 (1998).
34. Spitzer, R. L., Kroenke, K., Williams, J. B. W. & Löwe, B. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch. Intern. Med.* **166**, 1092–1097 (2006).
35. Shaw, T. C. Suicide and sanity. *Lancet* **169**, 1067–1069 (1907).
36. Van Orden, K. A., Cukrowicz, K. C., Witte, T. K. & Joiner, T. E. Thwarted belongingness and perceived burdensomeness: construct validity and psychometric properties of the Interpersonal Needs Questionnaire. *Psychol. Assess.* **24**, 197–215 (2012).
37. Van Orden, K. A., Witte, T. K., Gordon, K. H., Bender, T. W. & Joiner, T. E. Jr Suicidal desire and the capability for suicide: tests of the interpersonal-psychological theory of suicidal behavior among adults. *J. Consult. Clin. Psychol.* **76**, 72–83 (2008).
38. Kleiman, E. M. & Beaver, J. K. A meaningful life is worth living: meaning in life as a suicide resiliency factor. *Psychiatry Res.* **210**, 934–939 (2013).
39. O'Shea, B. A., Glenn, J. J., Millner, A. J., Teachman, B. A. & Nock, M. K. Decomposing implicit associations about life and death improves our understanding of suicidal behavior. *Suicide Life Threat. Behav.* **50**, 1065–1074 (2020).
40. Greenwald, A. G., Nosek, B. A. & Banaji, M. R. Understanding and using the implicit association test: i. an improved scoring algorithm. *J. Pers. Soc. Psychol.* **85**, 197–216 (2003).
41. Haslbeck, J. M. B. & Waldorp, L. J. mgm: estimating time-varying mixed graphical models in high-dimensional data. *J. Stat. Softw.* **93**, 1–46 (2020).
42. Betz, L. T., Penzel, N. & Kambeitz, J. A network approach to relationships between cannabis use characteristics and psychopathology in the general population. *Sci. Rep.* **12**, 7163 (2022).
43. Epskamp, S. & Fried, E. I. A tutorial on regularized partial correlation networks. *Psychol. Methods* **23**, 617–634 (2018).
44. Hevey, D. Network analysis: a brief overview and tutorial. *Health Psychol. Behav. Med.* **6**, 301–328 (2018).
45. Epskamp, S., Borsboom, D. & Fried, E. I. Estimating psychological networks and their accuracy: a tutorial paper. *Behav. Res. Methods* **50**, 195–212 (2018).
46. R Core Team R: *A Language and Environment for Statistical Computing* (R Foundation for Statistical Computing, 2021); <https://www.R-project.org/>
47. Epskamp, S. & Fried, E. I. *bootnet: Bootstrap Methods for Various Network Estimation Routines* (CRAN R-Proj Org., 2015).
48. Epskamp, S., Cramer, A. O. J., Waldorp, L. J., Schmittmann, V. D. & Borsboom, D. qgraph: network visualizations of relationships in psychometric data. *J. Stat. Softw.* **48**, 1–18 (2012).

Acknowledgments

This study was supported by the American Foundation for Suicide Prevention (grant number RFA-O-020-13; M.K.N.). The funders had no role in study design, data collection and analysis, decision to publish or preparation of the manuscript.

Author contributions

All of the authors made a substantial contribution to this study. R.F. drafted the paper. R.F. and B.A.O. analyzed the data. M.K.N. designed the study and was responsible for study conception, funding acquisition and supervision of all activities. All authors contributed to revising the paper, and all authors approved the final version of the paper for submission.

Competing interests

M.K.N. receives publication royalties from Macmillan, Pearson and UpToDate. He has been a paid consultant in the past 3 years for Cambridge Health Alliance and for legal cases regarding a death by suicide. He has stock options in Cerebral Inc. He is an unpaid scientific adviser for Empatica, Koko and TalkLife. R.F. and B.A.O. declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s44220-025-00389-4>.

Correspondence and requests for materials should be addressed to René Freichel.

Peer review information *Nature Mental Health* thanks Raymond Chan, Emma H. Moscardini and Yuan Pang Wang for their contribution to the peer review of this work.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2025

Reporting Summary

Nature Portfolio wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Portfolio policies, see our [Editorial Policies](#) and the [Editorial Policy Checklist](#).

Statistics

For all statistical analyses, confirm that the following items are present in the figure legend, table legend, main text, or Methods section.

n/a	Confirmed
<input type="checkbox"/>	<input checked="" type="checkbox"/> The exact sample size (<i>n</i>) for each experimental group/condition, given as a discrete number and unit of measurement
<input type="checkbox"/>	<input checked="" type="checkbox"/> A statement on whether measurements were taken from distinct samples or whether the same sample was measured repeatedly
<input type="checkbox"/>	<input checked="" type="checkbox"/> The statistical test(s) used AND whether they are one- or two-sided <i>Only common tests should be described solely by name; describe more complex techniques in the Methods section.</i>
<input type="checkbox"/>	<input checked="" type="checkbox"/> A description of all covariates tested
<input type="checkbox"/>	<input checked="" type="checkbox"/> A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons
<input type="checkbox"/>	<input checked="" type="checkbox"/> A full description of the statistical parameters including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient) AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals)
<input type="checkbox"/>	<input checked="" type="checkbox"/> For null hypothesis testing, the test statistic (e.g. <i>F</i> , <i>t</i> , <i>r</i>) with confidence intervals, effect sizes, degrees of freedom and <i>P</i> value noted <i>Give P values as exact values whenever suitable.</i>
<input checked="" type="checkbox"/>	<input type="checkbox"/> For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings
<input checked="" type="checkbox"/>	<input type="checkbox"/> For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes
<input checked="" type="checkbox"/>	<input type="checkbox"/> Estimates of effect sizes (e.g. Cohen's <i>d</i> , Pearson's <i>r</i>), indicating how they were calculated

Our web collection on [statistics for biologists](#) contains articles on many of the points above.

Software and code

Policy information about [availability of computer code](#)

Data collection	The death-IAT was implemented using JavaScript and run through the Project Implicit servers based at Harvard University.
Data analysis	All preprocessing and analyses were performed using RStudio (R version 4.4.0). Network estimation was conducted using the bootnet (version 1.6) and mgm (version 1.2-14) R packages. Network visualizations were created using qgraph (version 1.9.8). Supplemental analyses utilized the stats package (version 4.4.0). Relevant code for this method is already widely available online and no custom scripts have been used.

For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors and reviewers. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Portfolio [guidelines for submitting code & software](#) for further information.

Data

Policy information about [availability of data](#)

All manuscripts must include a [data availability statement](#). This statement should provide the following information, where applicable:

- Accession codes, unique identifiers, or web links for publicly available datasets
- A description of any restrictions on data availability
- For clinical datasets or third party data, please ensure that the statement adheres to our [policy](#)

Data available upon reasonable request due to the sensitive nature of the data. Data was collected from patients presenting due to severe psychiatric distress to the Massachusetts General Hospital Emergency Department (ED) (between February 2015 and March 2017).

Research involving human participants, their data, or biological material

Policy information about studies with [human participants or human data](#). See also policy information about [sex, gender \(identity/presentation\), and sexual orientation](#) and [race, ethnicity and racism](#).

Reporting on sex and gender	The sample at baseline (n =1412) was predominantly male (54.32%; female: 43.96%, transgender: 1.72%). See results section.
Reporting on race, ethnicity, or other socially relevant groupings	The sample was relatively diverse with respect to ethnicity and race (non-Hispanic white: 68.0%, non-Hispanic African American: 7.22%, Hispanic Other: 7.15%, non-Hispanic other: 6.22%, Hispanic White: 5.79%, non-Hispanic Asian: 3.86%, and other race or ethnicity: 1.76%).
Population characteristics	See Table 1 and results section for detailed population characteristics. The study sample consisted of participants presenting to the emergency room. Study inclusion criteria include adult status (≥18 years-old) and presentation at the ER.
Recruitment	Patients presenting due to severe psychiatric distress to the Massachusetts General Hospital Emergency Department (ED) (between February 2015 and March 2017) were approached by study research assistants.
Ethics oversight	The study received ethical approval from Harvard University and Massachusetts General Hospital.

Note that full information on the approval of the study protocol must also be provided in the manuscript.

Field-specific reporting

Please select the one below that is the best fit for your research. If you are not sure, read the appropriate sections before making your selection.

☐ Life sciences ☒ Behavioural & social sciences ☐ Ecological, evolutionary & environmental sciences

For a reference copy of the document with all sections, see nature.com/documents/nr-reporting-summary-flat.pdf

Behavioural & social sciences study design

All studies must disclose on these points even when the disclosure is negative.

Study description	Patients presenting due to severe psychiatric distress to the Massachusetts General Hospital Emergency Department (ED) (between February 2015 and March 2017) were approached by study research assistants. There were two follow-up surveys conducted over the telephone or email, one month and six months after the baseline visit. The data collected was quantitative - see methods section.
Research sample	Patients presenting to the ER. The sample was predominantly male (54.32%; female: 43.96%, transgender: 1.72%), young (M = 34.95, SD = 13.57), and relatively diverse with respect to ethnicity and race (non-Hispanic white: 68.0%, non-Hispanic African American: 7.22%, Hispanic Other: 7.15%, non-Hispanic other: 6.22%, Hispanic White: 5.79%, non-Hispanic Asian: 3.86%, and other race or ethnicity: 1.76%).
Sampling strategy	Convenience sampling. Patients presenting due to severe psychiatric distress to the Massachusetts General Hospital Emergency Department (ED) (between February 2015 and March 2017) were approached by study research assistants. A target sample size of 2,000 participants was determined based on a power analysis for sufficient statistical power (.80) with alpha set at .05.
Data collection	Data was collected in-person at the emergency room. There was also additional data collection through email or telephone 1 and 6 months after the baseline visit. Study research assistants were present during data collection. Blinding not applicable. Behavioral tasks were administered through iPads or laptops.
Timing	Baseline visit between February 2015 and March 2017
Data exclusions	Exclusion criteria encompassed 1) the inability to speak or read English, 2) severe cognitive impairments due to conditions, such as florid psychosis, intellectual disability, dementia, acute intoxication, or 3) the presence of extremely agitated or violent behavior. The main reasons for failure to obtain consent were psychiatric impairment (n = 193), lack of contact information for follow-up (n = 137), participant refusal (n = 132), language or physical barriers (n = 29), discharge prior to completing the consent process (n = 16), refusal by family or friends (n = 11), and various other factors (n = 14). See http://doi.org/10.1001/jamanetworkopen.2021.44373 for all details.
Non-participation	See results section. At the 1-month follow-up survey, there were complete data for reports of suicidal ideation for 900 participants (63.74%). By the 6-month follow-up, 938 participants (66.43%) completed reports on measures of suicidal ideation. A supplemental analysis (see Table S3) showed that older age, female gender, lower income, and a lifetime history of a suicide attempt were significantly associated with slight increases in the likelihood of dropout.
Randomization	Not applicable as there were no conditions to be randomized to.

Reporting for specific materials, systems and methods

We require information from authors about some types of materials, experimental systems and methods used in many studies. Here, indicate whether each material, system or method listed is relevant to your study. If you are not sure if a list item applies to your research, read the appropriate section before selecting a response.

Materials & experimental systems

n/a	Involved in the study
<input checked="" type="checkbox"/>	<input type="checkbox"/> Antibodies
<input checked="" type="checkbox"/>	<input type="checkbox"/> Eukaryotic cell lines
<input checked="" type="checkbox"/>	<input type="checkbox"/> Palaeontology and archaeology
<input checked="" type="checkbox"/>	<input type="checkbox"/> Animals and other organisms
<input checked="" type="checkbox"/>	<input type="checkbox"/> Clinical data
<input checked="" type="checkbox"/>	<input type="checkbox"/> Dual use research of concern
<input checked="" type="checkbox"/>	<input type="checkbox"/> Plants

Methods

n/a	Involved in the study
<input checked="" type="checkbox"/>	<input type="checkbox"/> ChIP-seq
<input checked="" type="checkbox"/>	<input type="checkbox"/> Flow cytometry
<input checked="" type="checkbox"/>	<input type="checkbox"/> MRI-based neuroimaging

Plants

Seed stocks

Report on the source of all seed stocks or other plant material used. If applicable, state the seed stock centre and catalogue number. If plant specimens were collected from the field, describe the collection location, date and sampling procedures.

Novel plant genotypes

Describe the methods by which all novel plant genotypes were produced. This includes those generated by transgenic approaches, gene editing, chemical/radiation-based mutagenesis and hybridization. For transgenic lines, describe the transformation method, the number of independent lines analyzed and the generation upon which experiments were performed. For gene-edited lines, describe the editor used, the endogenous sequence targeted for editing, the targeting guide RNA sequence (if applicable) and how the editor was applied.

Authentication

Describe any authentication procedures for each seed stock used or novel genotype generated. Describe any experiments used to assess the effect of a mutation and, where applicable, how potential secondary effects (e.g. second site T-DNA insertions, mosaicism, off-target gene editing) were examined.