

## SYSTEMATIC REVIEW OPEN



## Shared cortical characteristics in major depressive disorder, anxiety disorder, and chronic pain: a structural MRI meta-analysis study

Wei Yu<sup>1,2,3,4</sup>, Bo Tao<sup>1,2,3,4</sup>, Fei Zhu<sup>1,2,3</sup>, Ziyang Gao<sup>1,2,3</sup>, Yuan Xiao<sup>1,2,3</sup> , Qiyong Gong<sup>1,2,3</sup> and Su Lui<sup>1,2,3</sup> ✉

© The Author(s) 2025

Chronic pain (CP) is closely related with major depressive disorder (MDD) and anxiety disorders (ANX), with high comorbidity and shared risk factors. Prior studies have demonstrated common neural correlates across the three disorders, but their neuroanatomic basis is not fully clear. Hence, the preregistered meta-analysis (CRD42019119709) intended to explore common alterations in cortical thickness among CP, MDD, and ANX, a widely used parameter for quantitatively assessing various cerebral conditions with high sensitivity to pathology in neuropsychology. A total of 68 studies comprising 3072 patients and 3427 healthy controls were finally included. Across the disorders, four common clusters with a significant reduction in cortical thickness were identified, including right insula, left anterior cingulate (AC), triangular part of the left inferior gyrus (IFG), and left middle temporal gyrus (MTG). Our findings suggested the shared cortical deficits involving ACC-insula/IFG circuit and left MTG in CP, MDD and ANX, revealing common neural correlates for cognitive and emotional processing in these highly comorbid disorders.

*Translational Psychiatry* (2025)15:430; <https://doi.org/10.1038/s41398-025-03424-1>

## INTRODUCTION

Chronic pain (CP), a leading cause of disability globally, is a complex disorder characterized by pain that persists beyond the normal tissue healing time (3 months) in the absence of an evident organic lesion, as defined by the International Association for the Study of Pain (IASP) [1]. Prevalence rates of CP vary within the range of 11% to 40%, affecting one in ten adults annually. Long-term pain significantly impacts individual's health and well-being, healthcare services, and society as a whole [1, 2].

Affective disorders, such as major depressive disorder (MDD) and anxiety disorders (ANX), are also major contributors to the overall burden of disability [3, 4]. Increasing evidence supports a strong relationship between CP, MDD, and ANX [5]. High comorbidity and shared risk factors, including life stress and adverse childhood experiences, are prevalent among these disorders [6–8]. Depression and anxiety can heighten stress responses, alter pain processing, and increase sensitivity to pain [9, 10]. Conversely, individuals with CP are more vulnerable to developing MDD and/or ANX compared to those without persistent pain [9, 10]. Consequently, recognizing and addressing this bidirectional association is essential for enhancing therapeutic adherence and patient outcomes, given the negative effects of comorbidity.

Pre-clinical studies over the past two decades have provided valuable insights into the transdiagnostic neural procedures underlying in CP, MDD and ANX [9, 11, 12]. Similar findings have also been observed in neuroimaging studies conducted on human

beings. Shared structural and functional brain abnormalities are identified in the cortico- and meso-limbic circuit, which consist of the insula, amygdala, hippocampus, anterior cingulate cortex (ACC), and prefrontal cortex (PFC), in all three disorders [11, 13, 14]. A recent meta-analysis of multimodal magnetic resonance imaging (MRI) has further supported these findings, revealing common reductions in gray matter (GM) volume in the insula and medial PFC across MDD, ANX, and CP [15]. However, none of MRI studies and meta-analyses so far have focused on whether changes in surface-based brain morphometry (eg., cortical thickness, CT) consistently overlap in MDD, ANX, and CP.

CT is one of the most widely used biomarker in the field of neuropsychology, known for its ability to quantitatively assess various cerebral conditions with high sensitivity to pathology compared to GM volume [16, 17]. Individual meta-analyses have been conducted for CP, MDD, and ANX, respectively, revealing significant cortical changes for each disorder, and some of these deficits partially overlapped across the three disorders [13, 18, 19]. Moreover, the systematic review conducted by Sindermann et al. summarized that a thinner cortex in the ACC and the orbital frontal region was found in both MDD and ANX compared to healthy controls (HCs) [20]. However, the aforementioned cortical abnormalities cannot fully reflect the common neural mechanisms underlying CP, MDD, and ANX, due to the lack of direct statistical analysis among the three disorders.

Therefore, the present study aimed to identify evidence of transdiagnostic neural alterations through meta-analyses of MRI

<sup>1</sup>Department of Radiology, Functional and Molecular Imaging Key Laboratory of Sichuan Province, West China Hospital of Sichuan University, No. 37 Guoxue Xiang, Chengdu 610041, China. <sup>2</sup>Huaxi MR Research Center (HMRRC), Department of Radiology, West China Hospital of Sichuan University, No. 37 Guoxue Xiang, Chengdu 610041, China.

<sup>3</sup>Research Unit of Psychoradiology, Chinese Academy of Medical Sciences, No. 37 Guoxue Xiang, Chengdu 610041, China. <sup>4</sup>These authors contributed equally: Wei Yu, Bo Tao.

✉email: [lusuwcmcs@tom.com](mailto:lusuwcmcs@tom.com)

Received: 8 April 2024 Revised: 30 April 2025 Accepted: 9 June 2025

Published online: 24 October 2025

studies on CT comparing patients with CP, MDD, and ANX with HCs. We first tested (i) cortical deficits in CP, MDD, and ANX, respectively, and then (ii) for common cortical changes across the three disorders. Also, we explored the disorder-specific cortical alterations for each disorder. This in-depth investigation of cortical alterations related to CP and affective disorders (MDD & ANX) may disentangle transdiagnostic neural underpinnings.

## METHODS

### Search and study selection

Meta-analysis procedure was performed using the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) reporting guideline [21] (Supplementary Table S1). The protocol was registered at PROSPERO (CRD42022361809). English-language literatures were searched until October 30, 2022 in Cochrane Library, Embase, PubMed and Web of Science. The relevant keywords and search strategy were described in detail in Supplementary methods. The studies comparing patients with CP, MDD, or ANX with HCs were selected, and the inclusion criteria were as follows: i) studies published in a peer-reviewed journal rather than theoretical papers, case reports, reviews, and meta-analyses; (ii) analyzed whole-brain CT; (iii) provided the coordinates of significant clusters in Talairach or Montreal Neurological Institute (MNI) space; (iv) individuals without any neurological or severe medical comorbidity in the term of the exclusion of CP and psychiatric comorbidity; and (v) only collected baseline findings in longitudinal or intervention/challenge studies. When the samples overlapped between studies, the study with the larger sample size was included. Furthermore, we made no restrictions on age, duration of illness, medication status and symptom severity of a patient to maximally cover any useful records.

### Data extraction

Demographic and clinical characteristics of each study, along with methodological information, were independently obtained by two authors (W.Y. and B.T), consisting of sample size, age, gender, illness duration, medication status, symptom severity, strength of MRI scanner, the method used to preprocess imaging data, and peak coordinates and effect sizes of the main brain regions that showed significant between-group differences. Any inconsistencies during the data extraction were discussed by the third person to reach a consensus decision.

### Meta-analysis

Seed-based mapping (SDM) software version 5.15 (<https://www.sdmproject.com/>) was used for meta-analyses of CT studies between patients with CP or affective disorders and control subjects [22]. Here, we briefly described this method as follow. First, peak coordinates of brain clusters with significant between-group differences and their statistical  $t$  values were extracted from each included study to create an effect-size signed map, and negative findings were also enrolled to get a null effect size. The  $Z$  and  $P$  values were converted into  $t$ -statistic values online once  $t$  values were not reported ([www.sdmproject.com/utilities/show-Statistics](https://www.sdmproject.com/utilities/show-Statistics)). Studies included that used Talairach coordinates were first converted into MNI space to standardize the coordinate space. Next, a random-effects analysis for the study maps was performed to obtain the mean meta-analytic signed map, accounting for both positive and negative effects. Finally, a threshold of  $P = 0.005$  with peak  $Z = 1$  and cluster extent = 10 was used.

We identified common and disorder-specific deficits of cortical morphology between CP, MDD and ANX via a three-step meta-analytic approach that has been used in other studies [23, 24]. (i) We initially conducted separate meta-analyses for the patient group of each disorder with their respective HCs to characterize robust CT changes for each disorder. (iii) For common CT

abnormalities across the three disorders (relative to their respective HCs), a conjunction analysis was performed by computing errors for the estimated  $P$  value of each vertex from the separate meta-analytic maps. (ii) Disorder-specific CT abnormalities (i.e., CP vs MDD, CP vs ANX, ANX vs MDD, relative to their respective HCs), were determined using a quantitative contrast method, which calculated the difference in each vertex covarying for age and sex, and used standard randomization tests to establish statistical significance. Additionally, we carefully controlled for comorbidity effects when identifying shared and distinct structural alterations in the comparative and conjunctive meta-analysis procedures.

To explore the potential impacts of confounding factors (e.g. age, sex, duration of illness and percentage of comorbidity) on cortical deficits from the separate meta-analyses, a linear random-effects model was used for meta-regression analyses in each patient group and a threshold of  $P$  value less than 0.0005 was adopted [25]. And at least 9 studies should be required for each meta-regression analysis, as recommended by Radua et al. [26].

### Jackknife sensitivity analysis

A whole-brain jackknife sensitivity analysis was performed to examine the reliability of meta-analysis results. It needed to iteratively repeat the analysis each time when one dataset was discarded. The results were considered highly replicable if a given cluster remained significant in 75% or more of the combinations of studies [27]. We conducted jackknife analysis for each pooled meta-analysis.

### Analysis of heterogeneity and publication bias

$I^2$  statistics was used to evaluate the inter-study heterogeneity of individual clusters obtained from the meta-analyses. Publication bias was evaluated by funnel plots. Egger's test and any results with  $P < 0.05$  were considered to be significant for publication bias [28].

## RESULTS


### Included studies and sample characteristics

The PRISMA flowchart of the literature search and selection was shown in Fig. 1. 35 studies comprising 1508 CP patients and 1549 HCs, 22 studies comprising 1133 MDD patients and 1515 HCs, and 11 studies comprising 431 patients with ANX and 363 HCs were included in the present meta-analysis. Demographic and clinical characteristics of these included studies were provided in Supplementary Table S2–S4. One-way ANOVA revealed significant differences in mean age ( $F = 16.948$ ,  $P < 0.001$ ,  $\eta^2 = 0.3475$ ) and female ratio ( $F = 4.531$ ,  $P = 0.014$ ,  $\eta^2 = 0.1224$ ) in the three patient groups. Post-hoc tests revealed that mean age and female ratio of the CP patient group were higher than those of MDD and ANX ( $P < 0.001$  &  $P < 0.001$ ).

### Meta-analysis

*CP, MDD and ANX patients versus HCs, respectively.* Compared with HCs, CP patients showed a significant reduction of CT only in the right insula (Table 1, Fig. 2A). MDD patients showed significantly reduced CT in four clusters in comparison with HCs: right middle temporal gyrus (MTG), right supplementary motor area (SMA), left superior temporal gyrus (STG) and left superior frontal gyrus (SFG), dorsolateral (Table 1, Fig. 2B). For ANX, thinner cortex was observed in four clusters: bilateral insula, triangular part of the right inferior frontal gyrus (IFG), and right median cingulate/paracingulate gyri (MG/PC) (Table 1, Fig. 2C). No clusters in each disorder had a greater cortex than that in control subjects. Controlling for comorbidity in the analysis, the cortical results of CP and MDD remained stable in comparison with HCs, with the exception of ANX (Supplementary Table S5).

Given that the complex etiology and clinical presentations of CP, subgroup analysis may be helpful in clarifying distinct



**Fig. 1 PRISMA flow diagram of study selection in this meta-analysis.** ANX, anxiety disorder; MDD, major depressive disorder; CP, chronic pain; HC, healthy control.

neurological mechanisms involved in different illness conditions. Therefore, CP patients were divided into neuropathic- and non-neurogenic pain groups, the former comprising 21 studies, with 1139 patients and 1040 HCs while the latter including 14 studies with 408 patients and 498 HCs. Subgroup meta-analyses revealed increased CT in a wide range of brain regions including left striatum, right paracentral lobule, and reduced CT in right superior parietal gyrus, left precentral gyrus in neurogenic pain patients (Supplementary Table S6). Patients with non-neurogenic pain showed increased CT in the left precentral gyrus (PreCG), orbital part of the bilateral middle frontal gyrus (MFG), and thinner CT in the right insula, median




cingulum, MTG and triangular part of the left IFG compared to HCs (Supplementary Table S7).

**Comparative and conjunction analyses of CT between CP, MDD and ANX.** Studies with comorbidity were excluded to avoid the confounding impacts of comorbid conditions, further identifying the common and specific CT changes for each disorder. Decreased and increased CT in ANX relative to CP were found in the right angular gyrus (ANG) and the right insula, respectively (Table 2, Fig. 3A). And greater CT was observed in four clusters in CP compared to MDD: right SMA, right MTG, orbital part of the left MFG, and left PreCG (Table 2, Fig. 3B), while ANX patients showed thinner cortex

**Table 1.** Whole-brain meta-analysis results for CT studies comparing CP, ANX, or MDD with their respective HCs.

MNI coordinates	SDM Z	Voxels	Region	P value, uncorrected	BA	Egger's bias	Egger's p
<b>CP &lt; HCs</b>							
36, 10, 4	-1.189	1504	Right insula	0.000283837	48	-0.46	0.243
<b>MDD &lt; HCs</b>							
60, -42, -2	-1.778	596	Right middle temporal gyrus	0.000051618	21, 22	0.93	0.091
-52, -24, 4	-1.374	203	Left superior temporal gyrus	0.000500619	48	1.45	0.004
8, 14, 52	-1.305	180	Right supplementary motor area	0.000898004	6	0.04	0.945
-24, 34, 36	-1.117	45	Left superior frontal gyrus	0.002755880	9	0.13	0.859
<b>ANX &lt; HCs</b>							
-36, 2, 0	-1.230	307	Left insula	0.000278711	48	1.16	0.268
42, 38, 0	-1.263	207	Right inferior frontal gyrus, triangular part	0.000233330	45, 47	1.12	0.286
6, 0, 38	-1.015	186	Right median cingulate/paracingulate gyri	0.001924992	24	-0.31	0.833
32, 26, 0	-1.230	161	Right insula	0.000278711	24	1.16	0.268

BA Brodmann area; MNI montreal neurological institute; SDM seed-based d mapping; CP chronic pain; MDD major depressive disorder; ANX anxiety disorder; HCs healthy controls.

**(A) CP < HCs****(B) MDD < HCs****(C) ANX < HCs**

**Fig. 2** Cortical changes of CP, MDD and ANX in comparison with HCs. A–C Reduced cortical deficits of CP, MDD and ANX in comparison with HCs, respectively. CP chronic pain, MDD major depressive disorder, ANX anxiety disorders, HCs healthy controls, STG superior temporal gyrus, SFG superior frontal gyrus, SMA supplementary motor area, IFG inferior Frontal Gyrus, MC/PG median cingulate/paracingulate gyri, L left, R right.

**Table 2.** Comparative meta-analysis results across CP, ANX, and MDD.

MNI coordinates	SDM Z	Voxels	Region	P value, uncorrected	BA	Egger's bias	Egger's p
<b>CP &gt; ANX</b>							
32, 26, 0	1.359	71	Right insula	0.000753462	47	−0.12	0.678
<b>CP &lt; ANX</b>							
34, −64, 46	−1.040	836	Right angular gyrus	0.000036120	7	0.12	0.725
<b>CP &gt; MDD</b>							
60, −42, −2	1.210	361	Right middle temporal gyrus	0.000361264	21	0.22	0.319
−30, 52, −6	1.165	218	Left middle frontal gyrus, orbital part	0.000526428	11, 47	0.42	0.058
−32, −28, 58	1.066	224	Left precentral gyrus	0.001032174	3, 4	0.31	0.081
8, 12, 54	1.081	195	Right supplementary motor area	0.000923812	6, 32	0.04	0.875
<b>ANX &lt; MDD</b>							
32, 24, 0	−1.545	258	Right insula	0.000134170	47	0.21	0.617
−8, 56, 6	−1.205	80	Left superior frontal gyrus, medial	0.001966298	10	−0.40	0.404
−34, 2, 0	−1.469	112	Left insula	0.000227094	48	−0.01	0.990
42, 38, 0	−1.515	91	Right inferior frontal gyrus, triangular part	0.000113547	45	0.05	0.895

BA Brodmann area; MNI montreal neurological institute; SDM seed-based d mapping; CP chronic pain; MDD major depression disorder; ANX anxiety disorder; HCs, healthy controls.

than MDD patients in 4 clusters: bilateral insula, left SFG and triangular part of the right IFG (Table 2, Fig. 3C).

Four common clusters with a reduction in CT were identified in conjunction analyses (CP ∩ MDD ∩ ANX): right insula, left anterior cingulate (AC), triangular part of the left IFG, and left MTG (Table 3, Fig. 3D). In addition, conjunction meta-analyses were also performed between any two disorders, and the relevant results were provided in Supplementary Results.

**Meta-regression.** In MDD, age was positively associated with thinner CT in the right MTG ( $r = 0.476$ ,  $p = 0.025$ ). There were no significant associations between CT deficits and any other confounding factors in CP, MDD and ANX.

**Analyses of jackknife, heterogeneity and publication bias.** The results of the jackknife sensitivity analysis were highly replicable (Supplementary Table S8–S10). Heterogeneity statistics  $I^2$  extracted from the significant clusters were low heterogeneity ( $I^2 < 50\%$ ). And there was significant publication bias in the left superior temporal gyrus in MDD (Table 1). No significant publication bias was identified by Egger's test for CP and ANX (Table 1).

## DISCUSSIONS


This meta-analysis is the first to quantitatively evaluate common cortical changes across CP, MDD, and ANX, in comparison to HCs. Shared decreases in CT were observed in the ACC-insula/IFG circuit and left MTG, shedding light on neural correlates of comorbidity and shared neural mechanisms for cognitive and emotional processing. Also, specific cortical changes for each disorder replicated and extended prior findings, suggesting additional disorder-specific mechanisms.

A recent publication by Brandl et al. found common GM volume reduction in insular and medial-prefrontal cortices across these three disorders, providing powerful evidence of neural correlates for comorbidity and a shared neural basis [15]. According to the 'perception-action' model for pain, the posterior insula, parietal operculum, and the primary and secondary somatosensory cortices form the primary pain unit responsible for pain perception and localization [29, 30]. The anterior insula and ACC,

as key components of the second cortical pain unit, are associated with attention to pain [31]. Both pain matrices collaborate to shape the affective experience of pain, without directly initiating behavioral responses. Series of studies have demonstrated abnormalities in the insular-ACC circuit in patients with chronic pain [13, 14, 32]. Also, this neurocircuit is engaged in the perception and attention aspects of negative emotion [33]. Several MRI studies identified disrupted connectivity between the insula cortex and ACC in depression, social anxiety disorder, and panic disorder [34, 35]. Thus, it suggested that the insula-ACC circuit is a pivotal node in the brain governing cognitive and emotional processing and behavior. Similar roles can be also observed in the insula-IFG circuit. Tops and Boksem speculated that the insula and IFG areas within the ventral corticolimbic control pathway first integrate limbic emotion-motivational and sensory inputs, and then produce different responses based on the required level of precision for a stimulus or task [33]. Findings from functional MRI (fMRI), electroencephalogram (EEG) and event-related potential (ERP) studies have confirmed this point [33]. However, the above two common neurocircuits are partially separable due to their respective adaptations to various conditions involving pain and emotion, even though they interact [33]. For example, a PET study identified correlations between an activity shift from the insula-IFG region to the dorsolateral cortex and antidepressant effects of medication in patients with depression [36]. Regarding the left MTG, both CP and affective disorder patients exhibited functional deficits in this region [37–39]. The MTG is responsible for emotional processing, selective attention, working memory, and other cognitive functions [40–42]. Abnormal activity in this area may be associated with negative emotions and cognitive impairment [43]. In conclusion, our research has identified common cortical changes in the insula-ACC/IFG circuit and left MTG among CP, MDD and ANX, offering insights into the development of potential treatment biomarkers for shared neural mechanisms, especially Transcranial Magnetic Stimulation (TMS).

Comparative meta-analyses in our study showed that specific cortical changes for CP, MDD and ANX partially overlapped with not only common CT changes across disorders but also with CT deficits in each disorder when compared to HCs, including the insula, IFG, MTG, and SMA. This implies the need for further





**Fig. 3 Disorder-specific and common cortical changes across CP, ANX and MDD.** Disorder-specific (A–C) and common (D) cortical deficits across these disorders. The brain regions marked in blue and red indicate decreased and increased cortical thickness, respectively, in patients compared to HCs. CP chronic pain, MDD major depressive disorder, ANX anxiety disorders, HCs healthy controls, ANG angular gyrus, SMA supplementary motor area, MTG middle temporal gyrus, PreCG precentral gyrus, MFG middle frontal gyrus, SFG superior frontal gyrus, IFG inferior Frontal Gyrus, ACC anterior cingulate, L left, R right.

research to uncover the complex neurological processes underlying this finding and to understand its implications for clinical treatment. Thus, here we only discuss CT changes in the angular gyrus specific to CP vs. ANX, as well as in the PreCG and MFG for CP vs. MDD, and the SFG for ANX vs. MDD. The angular gyrus, situated posteriorly in the inferior parietal lobule, serves as a multimodal integrative hub that converges multisensory inputs from multiple brain regions [44]. This area is associated with several cognitive domains, including language processing, memory retrieval, attention, and social cognition [44, 45]. Prior fMRI studies have reported dysregulated connectivity between the angular gyrus and other brain regions in patients with migraine,

fibromyalgia, and anxiety, respectively [46–50]. Here, we showed a significant reduction of CT in the angular gyrus in individuals with CP compared to those with ANX, indicating a potential vulnerability of this region to CP. Similar psychopathological mechanisms may underlie cortical deficits between CP and MDD in our study. Taken together, our observation of disorder-specific cortical changes enhances our comprehension of unique neural mechanisms associated with each disorder. Beyond the regional morphological changes associated with clinical conditions, mapping intrinsic brain connectivity networks can provide a potential mechanistic framework for understanding various aspects of human behavior, as reported in numerous studies [51–53]. These

**Table 3.** Conjunction meta-analysis results across CP, ANX, and MDD.

MNI coordinates	Voxels	P value, uncorrected	Region	BA
<b>Common CT-increased: (CP &gt; HCs) ∩ (ANX &gt; HCs) ∩ (MDD &gt; HCs)</b>				
None				
<b>Common CT-decreased: (CP &lt; HCs) ∩ (ANX &lt; HCs) ∩ (MDD &lt; HCs)</b>				
36, 10, 4	899	<0.00005	Right insula	48
−2, 40, 8	533	<0.00005	Left anterior cingulate	32
−52, 18, 8	227	<0.00005	Left inferior frontal gyrus, triangular part	48
−58, −16, −22	229	<0.00005	Left middle temporal gyrus	20

CT cortical thickness; BA Brodmann area; CP chronic pain; MDD major depression disorder; ANX anxiety disorder; HCs, healthy controls.

networks, which include functional and structural connections within the brain, can provide deeper insights into the neural mechanisms underlying cognitive processes, emotional regulation, and social interactions. In anticipation of future endeavors, we aim to examine the functional networks of these three diseases, thereby delving deeper into the shared and distinct functional network alterations among them.

There were some limitations that are worth highlighting. First, this study was based on summarized data (i.e. reported coordinates) rather than raw original data, and a few results with small or moderate effect sizes may have been neglected. Second, only 11 MRI studies related to ANX was included in this meta-analysis, which is smaller in comparison to those involving CP (35 studies) and MDD (22 studies), although the number of included studies for ANX met the recommended number of studies for performing SDM [26]. Third, some uncontrollable influence of medication on cortical deficits across these three disorders might still exist, as we did not exclude patients using drugs in our study. It was difficult to consistently and accurately record medication usage for all the included studies. The potential effect existed between the different field strength, as we did not limit the parameters. The reconstruction of cortices will be influenced by field strength and sequence parameters. This factor might subtly affect the contrast between gray and white matter and the extraction of white matter surface and the pial surfaces. Thus, the differences of MRI parameters across studies represent an important consideration when interpreting our findings. Finally, the included disorders were quite heterogeneous; for example, CP encompasses different conditions like chronic back pain or fibromyalgia (see detailed information in Supplementary Materials). This inherent heterogeneity, common in many psychiatric diagnostic categories, could potentially diminish statistical power and elevate the risk of obtaining false-negative results.

As the first meta-analysis quantitatively evaluating cortical characteristics among CP, MDD, and ANX, shared deficits involving ACC-insula/IFG circuit and left MTG suggested common neural correlates for cognitive and emotional processing in these highly comorbid disorders.

## REFERENCES

- Cohen SP, Vase L, Hooten WM. Chronic pain: an update on burden, best practices, and new advances. *Lancet*. 2021;397:2082–97.
- Raffaelli W, Arnaudo E. Pain as a disease: an overview. *J Pain Res*. 2017;10:2003–8.
- Chisholm D, Sweeny K, Sheehan P, Rasmussen B, Smit F, Cuijpers P, et al. Scaling-up treatment of depression and anxiety: a global return on investment analysis. *Lancet Psychiatry*. 2016;3:415–24.
- Trautmann S, Rehm J, Wittchen HU. The economic costs of mental disorders: Do our societies react appropriately to the burden of mental disorders? *EMBO Rep*. 2016;17:1245–9.
- GBD 2017 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet*. 2018;392:1789–858.
- Slavich GM. Life stress and health: a review of conceptual issues and recent findings. *Teach Psychol*. 2016;43:346–55.
- Antoniu G, Lambourg E, Steele JD, Colvin LA. The effect of adverse childhood experiences on chronic pain and major depression in adulthood: a systematic review and meta-analysis. *Br J Anaesth*. 2023;130:729–46.
- Chopra K, Arora V. An intricate relationship between pain and depression: clinical correlates, coactivation factors and therapeutic targets. *Expert Opin Ther Targets*. 2014;18:159–76.
- Kimura LF, Novaes LS, Picolo G, Munhoz CD, Cheung CW, Camarini R. How environmental enrichment balances out neuroinflammation in chronic pain and comorbid depression and anxiety disorders. *Br J Pharmacol*. 2022;179:1640–60.
- Chang MH, Hsu JW, Huang KL, Su TP, Bai YM, Li CT, et al. Bidirectional association between depression and fibromyalgia syndrome: a nationwide longitudinal study. *J Pain*. 2015;16:895–902.
- Serafini RA, Pryce KD, Zachariou V. The mesolimbic dopamine system in chronic pain and associated affective comorbidities. *Biol Psychiatry*. 2020;87:64–73.
- Humo M, Lu H, Yalcin I. The molecular neurobiology of chronic pain-induced depression. *Cell Tissue Res*. 2019;377:21–43.
- Henn AT, Larsen B, Frahm L, Xu A, Adebimpe A, Scott JC, et al. Structural imaging studies of patients with chronic pain: an anatomical likelihood estimate meta-analysis. *Pain*. 2023;164:e10–e24.
- Wang Z, Yuan M, Xiao J, Chen L, Guo X, Dou Y, et al. Gray matter abnormalities in patients with chronic primary pain: a coordinate-based meta-analysis. *Pain Physician*. 2022;25:1–13.
- Brandl F, Weise B, Mulej Bratec S, Jassim N, Hoffmann Ayala D, Bertram T, et al. Common and specific large-scale brain changes in major depressive disorder, anxiety disorders, and chronic pain: a transdiagnostic multimodal meta-analysis of structural and functional MRI studies. *Neuropsychopharmacology*. 2022;47:1071–80.
- Fjell AM, Westlye LT, Grydeland H, Amlie I, Espeseth T, Reinvang I, et al. Accelerating cortical thinning: unique to dementia or universal in aging? *Cereb Cortex*. 2014;24:919–34.
- Winkler AM, Kochunov P, Blangero J, Almasy L, Zilles K, Fox PT, et al. Cortical thickness or grey matter volume? The importance of selecting the phenotype for imaging genetics studies. *Neuroimage*. 2010;53:1135–46.
- Sobral M, Morgado S, Moreira H, Guiomar R, Ganho-Ávila A. Association between cortical thickness and anxiety measures: a scoping review. *Psychiatry Res Neuroimaging*. 2022;319:111423.
- Li Q, Zhao Y, Chen Z, Long J, Dai J, Huang X, et al. Meta-analysis of cortical thickness abnormalities in medication-free patients with major depressive disorder. *Neuropsychopharmacology*. 2020;45:703–12.
- Sindermann L, Redlich R, Opel N, Böhnlein J, Dannlowski U, Leehr EJ. Systematic transdiagnostic review of magnetic-resonance imaging results: depression, anxiety disorders and their co-occurrence. *J Psychiatr Res*. 2021;142:226–39.
- Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*. 2009;6:e1000097.
- Albajes-Eizaguirre A, Solanes A, Vieta E, Radua J. Voxel-based meta-analysis via permutation of subject images (PSI): theory and implementation for SDM. *Neuroimage*. 2019;186:174–84.
- Liu X, Klugah-Brown B, Zhang R, Chen H, Zhang J, Becker B. Pathological fear, anxiety and negative affect exhibit distinct neurostructural signatures: evidence from psychiatric neuroimaging meta-analysis. *Transl Psychiatry*. 2022;12:405.
- Klugah-Brown B, Zhou X, Pradhan BK, Zweerings J, Mathiak K, Biswal B, et al. Common neurofunctional dysregulations characterize obsessive-compulsive,

- substance use, and gaming disorders-An activation likelihood meta-analysis of functional imaging studies. *Addict Biol.* 2021;26:e12997.
25. Radua J, Rubia K, Canales-Rodríguez EJ. Anisotropic kernels for coordinate-based meta-analyses of neuroimaging studies. *Front Psychiatry.* 2014;5:13.
  26. Radua J, Mataix-Cols D. Voxel-wise meta-analysis of grey matter changes in obsessive-compulsive disorder. *Br J Psychiatry.* 2009;195:393–402.
  27. Radua J, Mataix-Cols D, Phillips ML, El-Hage W, Kronhaus DM, Cardoner N, et al. A new meta-analytic method for neuroimaging studies that combines reported peak coordinates and statistical parametric maps. *Eur Psychiatry.* 2012;27:605–11.
  28. Radua J, Grau M, van den Heuvel OA, Thiebaut de Schotten M, Stein DJ, Canales-Rodríguez EJ, et al. Multimodal voxel-based meta-analysis of white matter abnormalities in obsessive-compulsive disorder. *Neuropsychopharmacology.* 2014;39:1547–57.
  29. Fombergstein K, Qadri S, Ramani R. Functional MRI and pain. *Curr Opin Anaesthesiol.* 2013;26:588–93.
  30. Iannetti GD, Mouraux A. From the neuromatrix to the pain matrix (and back). *Exp Brain Res.* 2010;205:1–12.
  31. Walter C, Oertel BG, Felden L, Kell CA, Nöth U, Vermehren J, et al. Brain mapping-based model of  $\Delta(9)$ -Tetrahydrocannabinol effects on connectivity in the pain matrix. *Neuropsychopharmacology.* 2016;41:1659–69.
  32. Dehghan M, Schmidt-Wilcke T, Pfleiderer B, Eickhoff SB, Petzke F, Harris RE, et al. Coordinate-based (ALE) meta-analysis of brain activation in patients with fibromyalgia. *Hum Brain Mapp.* 2016;37:1749–58.
  33. Tops M, Boksem MA. A potential role of the inferior frontal gyrus and anterior insula in cognitive control, brain rhythms, and event-related potentials. *Front Psychol.* 2011;2:330.
  34. Mulders PC, van Eijndhoven PF, Schene AH, Beckmann CF, Tendolcar I. Resting-state functional connectivity in major depressive disorder: a review. *Neurosci Biobehav Rev.* 2015;56:330–44.
  35. Peterson A, Thome J, Frewen P, Lanius RA. Resting-state neuroimaging studies: a new way of identifying differences and similarities among the anxiety disorders? *Can J Psychiatry.* 2014;59:294–300.
  36. Wu JC, Gillin JC, Buchsbaum MS, Schachar C, Darnall LA, Keator DB, et al. Sleep deprivation PET correlations of Hamilton symptom improvement ratings with changes in relative glucose metabolism in patients with depression. *J Affect Disord.* 2008;107:181–6.
  37. Liu L, Lyu TL, Fu MY, Wang LP, Chen Y, Hong JH, et al. Changes in brain connectivity linked to multisensory processing of pain modulation in migraine with acupuncture treatment. *Neuroimage Clin.* 2022;36:103168.
  38. Zhu Z, Wang Y, Lau WKW, Wei X, Liu Y, Huang R, et al. Hyperconnectivity between the posterior cingulate and middle frontal and temporal gyrus in depression: Based on functional connectivity meta-analyses. *Brain Imaging Behav.* 2022;16:1538–51.
  39. Yun JY, Kim JC, Ku J, Shin JE, Kim JJ, Choi SH, et al. The left middle temporal gyrus in the middle of an impaired social-affective communication network in social anxiety disorder. *J Affect Disord.* 2017;214:53–59.
  40. Fox MD, Corbetta M, Snyder AZ, Vincent JL, Raichle ME, et al. Spontaneous neuronal activity distinguishes human dorsal and ventral attention systems. *Proc Natl Acad Sci USA.* 2006;103:10046–51.
  41. Gaffrey MS, Luby JL, Botteron K, Repovš G, Barch DM. Default mode network connectivity in children with a history of preschool onset depression. *J Child Psychol Psychiatry.* 2012;53:964–72.
  42. Corbetta M, Shulman GL. Control of goal-directed and stimulus-driven attention in the brain. *Nat Rev Neurosci.* 2002;3:201–15.
  43. Zhang Q, Li X, Yan H, Wang Y, Ou Y, Yu Y, et al. Associations between abnormal spontaneous neural activity and clinical variables, eye movements, and event-related potential indicators in major depressive disorder. *Front Neurosci.* 2022;16:1056868.
  44. Seghier ML. The angular gyrus: multiple functions and multiple subdivisions. *Neuroscientist.* 2013;19:43–61.
  45. Andrews-Hanna JR. The brain's default network and its adaptive role in internal mentation. *Neuroscientist.* 2012;18:251–70.
  46. Yuan M, Liu B, Yang B, Dang W, Xie H, Lui S, et al. Dysfunction of default mode network characterizes generalized anxiety disorder relative to social anxiety disorder and post-traumatic stress disorder. *J Affect Disord.* 2023;334:35–42.
  47. Fan S, Yu Y, Wu Y, Kai Y, Wang H, Chen Y, et al. Altered brain entropy and functional connectivity patterns in generalized anxiety disorder patients. *J Affect Disord.* 2023;332:168–75.
  48. Tian Z, Yin T, Xiao Q, Dong X, Yang Y, Wang M, et al. The altered functional connectivity with pain features integration and interaction in migraine without aura. *Front Neurosci.* 2021;15:646538.
  49. Pando-Naude V, Barrios FA, Alcauter S, Pasaye EH, Vase L, Brattico E, et al. Functional connectivity of music-induced analgesia in fibromyalgia. *Sci Rep.* 2019;9:15486.
  50. Garza-Villarreal EA, Jiang Z, Vuust P, Alcauter S, Vase L, Pasaye EH, et al. Music reduces pain and increases resting state fMRI BOLD signal amplitude in the left angular gyrus in fibromyalgia patients. *Front Psychol.* 2015;6:1051.
  51. Zhang X, Xu R, Ma H, Qian Y, Zhu J. Brain structural and functional damage network localization of suicide. *Biol Psychiatry.* 2024;95:1091–9.
  52. Mo F, Zhao H, Li Y, Cai H, Song Y, Wang R, et al. Network localization of state and trait of auditory verbal hallucinations in schizophrenia. *Schizophr Bull.* 2024;50:1326–36.
  53. Cheng Y, Cai H, Liu S, Yang Y, Pan S, Zhang Y, et al. Brain network localization of gray matter atrophy and neurocognitive and social cognitive dysfunction in schizophrenia. *Biol Psychiatry.* 2025;97:148–56.

## ACKNOWLEDGEMENTS

We thank Rebekka Lencer for helpful comments.

## AUTHOR CONTRIBUTIONS

Conceptualization: Wei Yu, Bo Tao; Data curation, and validation: Wei Yu, Bo Tao, Fei Zhu, Ziyang Gao, Yuan Xiao; Formal analysis: Wei Yu, Bo Tao, Fei Zhu, Yuan Xiao; Supervision: Qiyong Gong; Rebekka Lencer; Su Lui; Funding acquisition: Su Lui; Writing-review & editing: All authors.

## FUNDING

This study was supported by the National Natural Science Foundation of China (Project Nos. 82120108014, 82441007), National Key R&D Program of China (Project Nos. 2022YFC2009901, 2022YFC2009900), Chengdu Science and Technology Office, major technology application demonstration project (Project Nos. 2022-YF09-00062-SN, 2022-GH03-00017-HZ), 1.3.5 project for disciplines of excellence, West China Hospital, Sichuan University (Project Nos. ZYGD23003, ZYAI24010), the Fundamental Research Funds for the Central Universities (Project No. ZYGX2022YGRH008).

## COMPETING INTERESTS

The authors declare no competing interests.

## ADDITIONAL INFORMATION

**Supplementary information** The online version contains supplementary material available at <https://doi.org/10.1038/s41398-025-03424-1>.

**Correspondence** and requests for materials should be addressed to Su Lui.

**Reprints and permission information** is available at <http://www.nature.com/reprints>

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Open Access** This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

© The Author(s) 2025