



ORIGINAL ARTICLE

Can socioeconomic status moderate the effect of a conflictive family environment on brain structure and externalizing/internalizing behavior in children and adolescents?

Elizabeth Trejos-Castillo,¹ Ann Mastergeorge,¹ Hugo Cogo-Moreira,² Vinicius Oliveira Santana,³ André Zugman,³ Célia Maria de Araújo,³ Pedro Mario Pan,^{3,4} Felipe Picon,^{4,5,6} Luis Augusto Paim Rohde,^{4,5,6} Andrea P. Jackowski³

¹Human Development & Family Sciences, Texas Tech University, Lubbock, Texas, USA. ²Department of Education, Information and Communications Technology and Learning, Østfold University College, Halden, Norway. ³Laboratório Interdisciplinar de Neurociências Clínicas, Departamento de Psiquiatria, Universidade Federal de São Paulo, São Paulo, SP, Brazil. ⁴Instituto Nacional de Ciência e Tecnologia de Psiquiatria do Desenvolvimento para a Infância e Adolescência, São Paulo, SP, Brazil. ⁵Programa de Déficit de Atenção/Hiperatividade, Divisão Adulto, Hospital de Clínicas de Porto Alegre, Porto Alegre, RS, Brazil. ⁶Departamento de Psiquiatria, Faculdade de Medicina, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, Brazil.

Objective: This study examined the effects of socioeconomic status as a possible moderator of the effects of family conflict on externalizing/internalizing behavior and hippocampal and amygdala volume.

Methods: A longitudinal complete-case analysis of 714 children and adolescents (mean age: 11.2 years; 46.2% female) was conducted using data from the Brazilian High-Risk Cohort Study for Psychiatric Disorders in Childhood. At baseline, parents/guardians completed the Family Environment Scale and a socioeconomic status scale. Three years after baseline assessment, the same participants underwent brain magnetic resonance imaging, and the Child Behavior Checklist was administered. Automated segmentation of the amygdala and hippocampus was performed in FreeSurfer 5.1.

Results: Although family conflict at baseline predicted externalizing/internalizing behavior at follow-up, we found no evidence that family conflict and socioeconomic status affected brain structure or that family conflict had a moderating effect on psychopathology and brain outcomes conditioned on socioeconomic status.

Conclusion: These results are consistent with emerging evidence that family conflict is a risk factor for externalizing/internalizing behavior in youth. These findings warrant further attention, focusing on prevention and intervention efforts and social policy development.

Keywords: Child behavior; socioeconomic status; family conflict; brain structure

Introduction

The family environment significantly influences child behavioral and neurodevelopmental outcomes, encompassing various factors, such as family composition, home environment, parenting behaviors, interaction styles, and parental mental health and functioning. For instance, positive parenting practices – such as warmth, support, and effective communication – foster resilience and healthy emotional development in children.¹ Conversely, a dysfunctional family environment marked by low parental acceptance and family conflict is associated with an increased risk of psychopathology in children.

Specifically, higher levels of family conflict can lead to both internalizing behaviors, such as anxiety and depression, and externalizing behaviors, such as aggression and defiance.² Moreover, parental mental health issues, including depression and anxiety, can negatively affect parenting quality, further exacerbating behavioral challenges in children.³

During brain development, windows of vulnerability from the fetal period to adolescence render the brain particularly sensitive to environmental influences. Key factors include family dynamics, parental attitudes, maternal depression, social deprivation, and low socioeconomic status (SES).⁴ Early childhood⁵ and puberty^{6,7}

Correspondence: Andrea P. Jackowski, Laboratório Interdisciplinar de Neurociências Clínicas, Departamento de Psiquiatria, Universidade Federal de São Paulo, Rua Pedro de Toledo 669, 3º andar, CEP 04039-032, São Paulo, SP, Brazil.

E-mail: andrea.jackowski@unifesp.br

Submitted Feb 24 2025, accepted May 07 2025.

How to cite this article: Trejos-Castillo E, Mastergeorge A, Cogo-Moreira H, Santana VO, Zugman A, de Araújo CM, et al. Can socioeconomic status moderate the effect of a conflictive family environment on brain structure and externalizing/internalizing behavior in children and adolescents? Braz J Psychiatry. 2025;47:e20254195. <http://doi.org/10.47626/1516-4446-2025-4195>

are especially critical periods, characterized by rapid neurobiological and hormonal changes that can profoundly impact neurodevelopmental outcomes. The interplay of these environmental factors can adversely affect cognitive and emotional development, underscoring the detrimental impact of a negative or unresponsive environment during these formative years.

In developmental psychopathology, the relationship between brain structure and behavior is complex, particularly concerning whether reductions in cerebral grey matter volume are more closely associated with comorbid dimensions like internalizing or externalizing factors than specific neurodevelopmental disorders.⁸ In a large community sample of children ($n=254$), it was found that grey matter volume reductions in the prefrontal regions are related to general psychopathology and that volumetric reductions in the limbic regions (amygdala, hippocampus, and insula) are associated with internalizing but not externalizing factors.⁹ However, a longitudinal study found no alterations in the hippocampal or prefrontal regions in relation to internalizing or externalizing behaviors, although it did find that smaller amygdala volumes in adolescent males were related to greater early childhood externalizing behavior.¹⁰ A recent study reported an association between higher parent-reported aggression scores and reduced hippocampal volume in a follow-up analysis but not at baseline, suggesting that this relationship is driven by the trajectory of hippocampal development during adolescence.⁷

Studies indicate that SES is associated with health, cognitive abilities, academic performance, brain development, and socioemotional outcomes in children and adolescents.¹¹⁻¹⁶ Adolescents with low SES have less access to the resources and experiences of those with high SES, putting them at risk for developmental and behavioral problems.¹⁷ A number of studies have demonstrated an association between SES and behavioral and brain development.¹⁸⁻²¹ SES plays a crucial role in the growth and maturation of brain structure,²² with lower total grey matter,²³ frontal lobe (orbitofrontal cortex and cingulate),²⁴ and parietal lobe volumes,²⁵ as well as reduced cortical thickness.²⁶ Volumetric changes in the hippocampus^{27,28} and amygdala have been observed to vary according to SES status.^{29,30} Studies on typically developing children have found that lower SES predicts a smaller hippocampus^{26,27,31,32} and amygdala,³¹ and that low SES is associated with a reduction in grey matter volumes in the frontal and temporal regions.⁹ Furthermore, associations have been demonstrated between family conflict and hippocampal functional connectivity³³ and volume, with parental rejection being associated with lower hippocampal volume in adulthood.³⁴

Studies on Brazilian youth have examined similar constructs, including SES, parenting practices, child behavior,³⁵ psychopathology,³⁶ and externalizing/internalizing behaviors.³⁷ Altafim et al.³⁵ investigated the relationship between SES, parenting practices, and child behavior problems in a sample of 204 Brazilian mothers with children aged 3 to 8 years, finding that lower SES was associated with more negative parenting practices and a higher incidence of internalizing problems in

children.³⁵ Gallo et al.³⁶ analyzed data from 3,715 participants in a birth cohort study in Pelotas, Brazil. Perinatal SES data were collected at birth, and depressive symptoms were assessed when participants were 18 years old.³⁶ The study identified several SES-related risk factors for depression in both males and females, including maternal smoking during pregnancy, low maternal education, and low family income.³⁶ For girls, additional risk factors included having a young mother, while for boys, depression was associated with non-White maternal race and maternal mental health problems.³⁶ These findings highlight the role of SES in increasing vulnerability to depression, particularly among girls exposed to early-life adversity. Biazoli et al.³⁷ used data from the Brazilian High-Risk Cohort Study for Psychiatric Disorders in Childhood (BHRC) to examine the relationship between SES, brain function, and emotional and behavioral outcomes in school-age children. The study included 655 children aged 7 to 12 years from public schools in São Paulo. Resting-state functional magnetic resonance imaging (MRI) data revealed a negative correlation between spontaneous brain activity in the right superior temporal gyrus and internalizing behaviors.³⁷ Additionally, higher SES was associated with increased activity in this brain region, which is involved in emotional regulation, suggesting that children from higher SES backgrounds may be at lower risk for internalizing problems such as anxiety and depression.³⁷ However, these studies have not examined how these variables interact to influence brain structure or how they may moderate the relationship between socioeconomic factors and child behavior.

Families with lower SES often face challenges, such as financial instability, limited resources, and social isolation, which can increase family conflict and negatively impact children's emotional and behavioral outcomes.³⁸ However, despite these hardships, many low-income families build strong social networks that provide emotional support and a sense of community. These networks can foster resilience, improve mental health, and offer coping strategies, highlighting the complex nature of low SES.³⁹

Given these considerations, we hypothesized that baseline family conflict (focal predictor) would be associated with externalizing /internalizing behaviors and neuroimaging outcomes in follow-up (i.e., an *a priori* hypothesis of hippocampal and amygdala volume change). Additionally, we expected SES to act as a potential moderator, with stronger effects observed in families with lower SES. This suggests that the negative impact of an adverse family environment would be more pronounced among those with lower SES. Therefore, we hypothesized that there would be an interaction between family conflict and SES concerning hippocampus and amygdala volumes, as well as externalizing/internalizing behaviors.

Methods

Participants

We included 714 participants (5-14 years old; mean age: 9.92 years; 57.6% males) who underwent brain MRI and

had complete data for the 11 variables required for the covariance model. These participants were drawn from a larger cohort (the BHRC) focused on school-age children.⁴⁰ The study involved students from 57 schools in the cities of São Paulo and Porto Alegre. On each school's registration day, we invited the biological parents of 6- to 12-year-old children to participate in the study.

Biological parents, predominantly mothers (87.3% of cases), from 8,012 families (representing 9,937 children) completed the family history screening test.⁴¹ The BHRC was created by combining the two strata ($n=2,511$). The first stratum, "the random group" ($n=958$), comprised randomly selected individuals. The second stratum, "the high-risk group" ($n=1,553$), included children at risk for psychopathology, selected using a validated prioritization algorithm.⁴⁰ Only one child per family was included. Details regarding the cohort recruitment, data collection procedures, and the methodological approach have been described previously.^{37,40,42,43}

Materials

The parents completed the Child Behavior Checklist (CBCL)⁴⁴ and the Family Environment Scale (FES).⁴⁵ Behavioral and emotional problems were assessed using the validated Brazilian version of the CBCL,⁴⁶ which allows and assessment of dimensions of psychopathology that have shown to be valid in several cultures. Two summary scores were computed based on the CBCL scores: internalizing (summarizing the anxious/depressed, withdrawn/depressed, and somatic complaint scores; range 0-55) and externalizing (summarizing the rule-breaking and aggressive behavior scores; range 0-51). Wagner et al.⁴⁷ provided psychometric evidence for the validity of this measure.

Family functioning was measured using the complete Brazilian version of the FES,⁴⁸ which consists of six subscales across three domains: 1) interpersonal relationships (cohesion, expressivity, and conflict subscales); 2) personal growth (achievement-oriented subscale); and 3) system maintenance (control and organization subscales). Our analysis focused specifically on the conflict subscale.

The Brazilian SES scale, developed by the Associação Brasileira de Empresas de Pesquisa (Brazilian Association of Research Enterprises),⁴⁹ was used to define SES (the scores were directly and positively correlated with

SES). This scale is the conventional and official method for SES stratification in Brazil, assessing estimated income, access to goods and services (regularly reviewed to reflect the national experience), and parental educational status. Scores range from 0-46. For reference in terms of SES groups and their relation to the continuous scores and respective income, we present the mean estimated median family income by SES group (SD) based on the currency rate as of January 2, 2012: 1) USD 1,087.5 (SD, 797.3) for those with low SES (classes E and D; scores 0-13); 2) USD 1,423.9 (SD, 772.3) for those with mid-level SES (classes C and B; scores 14-34); and 3) USD 2,450.6 (SD, 1,725.0) for those with high SES (class A; scores 35-46). Our sample's scores ranged from 6-36 (average 18.27; SD, 4.23) (Table 1).

MRI was performed using a 1.5T MRI scanner (GE Sigma HDX and GE Sigma HD, GE Healthcare, Chicago, IL, USA) at two sites (São Paulo and Porto Alegre). Both systems followed the same protocols, and MRIs were performed with the following T1-weighted image parameters: TR = 10.916 ms, TE = 4.2 ms, thickness = 1.2 mm, flip angle = 15°, NEX = 1, matrix size = 256 × 192, FOV = 245 mm, and bandwidth = 122.109, yielding 156 axial slices. Image processing and analyses were performed using an automated, unbiased, atlas-based Bayesian segmentation method in FreeSurfer 5.1.⁵⁰ Volumes of the four subcortical structures (amygdala and hippocampus) were extracted using FreeSurfer. Due to the sensitivity of brain MRI signals and segmentation algorithms to artifacts such as head movement and tilt,^{51,52} we conducted a visual quality control inspection of the extracted volumes and hippocampal segmentation for all outliers. As a result, we excluded four subjects from the analysis. We controlled for intracranial volume, age, and sex in all analyses.

The SES and the FES scales were applied at baseline, whereas the CBCL and the MRI scan were performed at the follow-up evaluation 3 years after baseline.

Statistical analysis

Based on the hypotheses, multiple analyses of variance were used to evaluate the effects of the family conflict score (the focal predictor) and SES (the potential moderator) on the follow-up outcomes of CBCL internalizing and externalizing factor scores and on imaging results in four brain areas: the right/left hippocampus and

Table 1 Complete-case descriptive statistics for the study variables ($n=714$; male = 411 [57.6%])

Variable	Minimum	Maximum	Mean \pm SD
Age at baseline (years)	5.83	14.34	9.92 \pm 1.85
SES	6.00	36.00	18.21 \pm 4.37
FES – Conflict Score	0.00	10.00	3.44 \pm 2.23
CBCL – Internalizing Score	0.00	55.00	13.79 \pm 9.78
CBCL – Externalizing Score	0.00	51.00	12.93 \pm 10.14
Left hippocampus (mm ³)	2,501.00	5,090.00	3,818.76 \pm 366.11
Right hippocampus (mm ³)	2,536.00	5,159.20	3,918.59 \pm 374.71
Left amygdala (mm ³)	832.90	2,114.30	1,459.95 \pm 196.14
Right amygdala (mm ³)	879.80	2,464.80	1,574.46 \pm 214.04
Intracranial volume (mm ³)	919,993.00	1,864,950.00	1,394,864.48 \pm 131,792.09

CBCL = Child Behavior Checklist; FES = Family Environment Scale; SES = socioeconomic status.

right/left amygdala, adjusting for sex, age, and intracranial volume in all analyses. We opted for multivariate analysis of variance over multiple univariate approaches because multivariate tests more accurately capture the complex interrelated nature of our research questions, which involve modelling a combination of neuroimaging and behavioral scales that share variance and covary.^{53,54} We constructed a model incorporating interaction terms between SES and family conflict for the four outcomes to examine the potential moderating effect of SES. We hypothesized that more detrimental family contexts might exacerbate the direct impact of poverty on brain and behavior, leading to more deleterious effects. Conversely, a healthy family environment could act as a protective factor, mitigating the deleterious effects of poverty on neurodevelopmental outcomes. This moderating approach does not assume causal directionality⁵⁵ among SES, family environment, the brain, and behavioral outcomes because both the main predictor and outcomes were not collected at the two time points, which would at least have allowed us to apply the Granger causality test. Moreover, our framework is based on an ordinary regression approach, not the average moderated treatment effect under the potential outcomes' framework and its recent developments.⁵⁶

Modelling CBCL and neuroimaging measures simultaneously as outcomes mitigated multiple testing, reducing the false discovery rate. One strength of testing this hypothesis using two time points from the BHRC data was that data on the focal predictor (family conflict) and the moderator (SES) were collected at baseline. However, behavioral and neuroimaging outcomes were assessed at different time points.

A sensitivity analysis determined the robustness of the estimates under "missingness," which was expected in the longitudinal design. We assumed that missing mechanisms were missing at random and used full-information maximum likelihood to deal with missingness among the variables in the follow-up (outcomes) and baseline (predictor and moderator) measures.

Complete-case analyses using IBM SPSS Statistics determined effect sizes based on the partial eta square. Missing data analysis under full-information maximum likelihood was conducted using Mplus.⁵⁷

Ethics statement

The ethics committees of the participating institutions approved all procedures. Written informed consent was obtained from all participants and their parents or guardians.

Results

Table 1 presents the descriptive statistics of the continuous variables. In terms of SES, 16.7% ($n=753$) were classified as strata A/B and 72.4% ($n=3,273$) and 10.9% ($n=495$) were classified as strata D and E, respectively. Table 2 shows the Pearson correlations between the continuous variables included in the statistical models.

Pillai's trace showed that of the two main predictors (SES and family conflict scores), only family conflict was a significant predictor ($F_{\text{conflict}} [6,703] = 15.664$, $p < 0.001$, $\eta^2 = 0.118$). Sex, age, and intracranial volume (covariates) were also significant: $F_{\text{sex}} (6,703) = 7.698$, $p < 0.001$, $\eta^2 = 0.062$; $F_{\text{age}} (6,703) = 3.446$, $p = 0.002$, $\eta^2 = 0.029$; and $F_{\text{icv}} (6,703) = 79.787$, $p < 0.001$, $\eta^2 = 0.405$, respectively.

Between-subject analyses revealed significant associations between externalizing ($\beta = 1.511$ standard error = 0.160, $p < 0.001$, $\eta^2 = 0.112$) and internalizing ($\beta = 0.989$, standard error = 0.160, $p < 0.001$, $\eta^2 = 0.051$) behavior and family conflict. This indicates that greater family conflict predicted higher externalizing and internalizing behavior scores during follow-up.

There was no evidence of an association between family conflict and SES with the four neuroimaging outcomes. Information regarding between-subject effects is presented in Table 3.

Furthermore, Pillai's trace showed no evidence that the association of FES conflict score and SES had a moderating effect on neuroimaging outcomes ($F_{\text{conflict} * \text{SES}} [6,702] = 1.004$, $p = 0.422$, $\eta^2 = 0.009$). Thus, there is no evidence that FES conflict scores have a conditional effect on neuroimaging outcomes according to different SES values.

Sensitivity analysis under full-information maximum likelihood, in which the number of analyzed cases was 2,501, also indicated no evidence that there was an interaction between SES and FES conflict scores. Moreover, the association between FES conflict, externalization, and internalization was preserved. Table 4 presents the standardized coefficients for the conditional and unconditional models using 2,501 participants.

Discussion

We examined the associations between SES, child-family conflict, externalizing/internalizing behaviors, and bilateral hippocampus and amygdala volumes. Additionally, we examined the moderating effects of SES on the associations between family conflict, externalizing/internalizing behaviors, and brain structures. Family conflict emerged as a predictor of externalizing/internalizing behaviors in children during follow-up. Our findings corroborate those of previous studies conducted with Brazilian children regarding the connections between family conflict and externalizing/internalizing behaviors in children.^{35,36}

Our results showed that family conflict had a substantial effect on internalizing and externalizing behavior (the effect being greater on externalizing behavior) but not on subcortical brain structure volumes. Studies investigating parenting from early toddlerhood to adolescence have typically emphasized the quality of parenting behavior, concentrating on parental warmth, hostility or conflict, and control. Among these dimensions, parental hostility is most associated with adverse neurodevelopmental outcomes in children, including externalizing/internalizing behaviors and poorer executive function.⁵⁸ Two meta-analyses reported a robust association between conflictive and hostile/aggressive parent-child

Table 2 Pearson correlations between the continuous variables included in the statistical models

	1	2	3	4	5	6	7	8	9	10
Age (years) (1)										
Pearson correlation	1									
Sig. (2-tailed)										
SES (2)										
Pearson correlation	0.031	1								
Sig. (2-tailed)	0.41									
FES: conflict score (3)										
Pearson correlation	0.037	-0.152**	1							
Sig. (2-tailed)	0.33	< 0.01								
CBCL Neuro: internalizing (4)										
Pearson correlation	0.062	-0.135**	0.242**	1						
Sig. (2-tailed)	0.10	< 0.01	< 0.01							
CBCL Neuro: externalizing (5)										
Pearson correlation	0.061	-0.134**	0.355**	0.591**	1					
Sig. (2-tailed)	0.11	< 0.01	< 0.01	< 0.01						
Left hippocampus (6)										
Pearson correlation	0.181**	0.081*	-0.074*	-0.006	-0.044	1				
Sig. (2-tailed)	< 0.01	0.03	0.05	0.87	0.24					
Right hippocampus (7)										
Pearson correlation	0.177**	0.068	-0.060	0.001	-0.047	0.833**	1			
Sig. (2-tailed)	< 0.01	0.07	0.11	0.97	0.21	< 0.01				
Left amygdala (8)										
Pearson correlation	0.182**	0.067	-0.070	-0.118**	-0.124**	0.567**	0.571**	1		
Sig. (2-tailed)	< 0.01	0.07	0.06	< 0.01	< 0.01	< 0.01	< 0.01			
Right amygdala (9)										
Pearson correlation	0.206**	0.046	-0.036	-0.066	-0.065	0.565**	0.581**	0.776**	1	
Sig. (2-tailed)	< 0.01	0.21	0.34	0.08	0.08	< 0.01	< 0.01	< 0.01		
Intracranial volume (10)										
Pearson correlation	0.210**	0.049	-0.030	-0.036	-0.064	0.624**	0.620**	0.588**	0.588**	1
Sig. (2-tailed)	< 0.01	0.19	0.42	0.33	0.09	< 0.01	< 0.01	< 0.01	< 0.01	

CBCL = Child Behavior Checklist; FES = Family Environment Scale; SES = socioeconomic status.

* Correlation is significant at the 0.05 level (2-tailed).

** Correlation is significant at the 0.01 level (2-tailed).

relationships and externalizing behaviors in children and adolescents.⁵⁹ Moreover, a parent effects model, which is described as conflictual or negative parenting that results in and/or exacerbates child or adolescent externalizing behavior, has been reported as responsible for this association.⁶⁰ In this context, our results on the association between family conflict and externalizing/internalizing behaviors support these findings.

The neural correlates of SES have been consistently addressed by studies investigating brain structure and mental health in adults and children. In contrast to previous studies,⁶¹ we did not find sufficient evidence that SES affects youth developmental outcomes. Ursache et al.⁶² pointed out that although understanding of the relationship between SES and brain and developmental outcomes has improved, there is still no consensus on the definition and conceptualization of the SES construct. Although we used a conventional Brazilian SES assessment, which was appropriate for our cultural construct, it limits comparison with other countries.

For instance, a large study of 2,043 children aged 11-13 years found that although low SES was associated with childhood mental health problems, this association was modified by parental emotional well-being and parenting practices.⁶³ A systematic review¹⁷ showed that the impact of low SES is more pronounced in early childhood than in adolescence, and lower SES was more strongly associated with externalizing disorders than internalizing disorders. We found a minimal association between SES and externalizing/internalizing behaviors ($r = -0.134$). This weak relationship might be attributable to the instrument we used to measure SES, which, while appropriate for the culture in which the data were collected, limits the generalizability of the findings to broader populations and cultures. Additionally, SES was assessed only at baseline and not during the 3-year follow-up, meaning we cannot assume that SES remained stable throughout this period. The fact that this measure was not assessed longitudinally could further weaken the observed relationship between SES and behavioral

Table 3 Between-subject effects, unstandardized regression coefficients with 95%CI and effect size

Outcome/covariates	B	SE	t	p-value	95% CI		Partial eta squared
					Lower bound	Upper bound	
Left hippocampus							
Intercept	1.336.414	131.372	10.173	< 0.001	1.078.488	1.594.340	0.128
Male	13.753	23.826	0.577	0.564	-33.024	60.531	0.000
Age	11.647	5.943	1.960	0.050	-0.022	23.316	0.005
FES conflict	-8.778	4.850	-1.810	0.071	-18.300	0.744	0.005
ICV	0.002	9.151E-05	18.238	0.000	0.001	0.002	0.320
SES	3.347	2.477	1.352	0.177	-1.515	8.210	0.003
Right hippocampus							
Intercept	1.413.081	135.183	10.453	< 0.001	1.147.673	1.678.489	0.134
Male	24.399	24.517	0.995	0.320	-23.735	72.534	0.001
Age	12.687	6.116	2.074	0.038	0.679	24.694	0.006
FES conflict	-7.034	4.991	-1.409	0.159	-16.832	2.764	0.003
ICV	0.002	9.417E-05	17.850	< 0.001	0.001	0.002	0.310
SES	2.479	2.549	0.973	0.331	-2.524	7.483	0.001
Left amygdala							
Intercept	290.432	71.822	4.044	< 0.001	149.423	431.441	0.023
Male	62.210	13.026	4.776	< 0.001	36.637	87.784	0.031
Age	8.279	3.249	2.548	0.011	1.899	14.658	0.009
FES conflict	-5.168	2.651	-1.949	0.052	-10.374	0.037	0.005
ICV	0.001	5.003E-05	15.004	< 0.001	0.001	0.001	0.241
SES	1.221	1.354	0.902	0.368	-1.438	3.879	0.001
Right amygdala							
Intercept	274.930	78.598	3.498	< 0.001	120.616	429.243	0.017
Male	61.697	14.255	4.328	< 0.001	33.711	89.683	0.026
Age	11.703	3.556	3.291	0.001	4.722	18.685	0.015
FES conflict	-2.723	2.902	-0.938	0.348	-8.420	2.974	0.001
ICV	0.001	5.475E-05	15.030	< 0.001	0.001	0.001	0.242
SES	0.516	1.482	0.348	0.728	-2.393	3.425	0.000
Internalizing							
Intercept	11.343	4.325	2.622	0.009	2.851	19.834	0.010
Male	-1.987	0.784	-2.534	0.012	-3.528	-0.447	0.009
Age	0.461	0.196	2.355	0.019	0.077	0.845	0.008
FES conflict	0.989	0.160	6.196	< 0.001	0.676	1.303	0.051
ICV	-1.159E-07	3.013E-06	-0.038	0.969	-6.031E-06	5.800E-06	0.000
SES	-0.232	0.082	-2.843	0.005	-0.392	-0.072	0.011
Externalizing							
Intercept	15.766	4.339	3.634	< 0.001	7.247	24.285	0.018
Male	1.276	0.787	1.622	0.105	-0.269	2.821	0.004
Age	0.489	0.196	2.490	0.013	0.103	0.874	0.009
FES conflict	1.511	0.160	9.435	< 0.001	1.197	1.826	0.112
ICV	-7.229E-06	3.023E-06	-2.392	0.017	-1.316E-05	-1.295E-06	0.008
SES	-0.194	0.082	-2.377	0.018	-0.355	-0.034	0.008

FES = Family Environment Scale; ICV = intracranial volume; SE = standard error; SES = socioeconomic status.

outcomes. Despite these limitations, our study contributes to the global understanding of child and youth neurodevelopment by supporting previous findings in a Brazilian sample and offering new insights for future cross-cultural and cross-national research.

Internalizing and externalizing behaviors have been implicated in the development of numerous psychopathologies, such as disruptive behavior disorders,⁶⁴ substance abuse,^{65,66} ADHD,⁶⁷ autism spectrum disorder,⁶⁸ etc. As mentioned earlier, the windows of vulnerability observed in adolescence are critical for brain development and, although the results of the present study showed no association with brain measures, the fact that our analyses were conducted on a very young population (mean age: 11.2 years) must be considered. We cannot

disregard the possibility that brain alterations may appear later in life, since hippocampal reductions and other structural brain alterations associated with mental illness have been commonly observed in adult populations.⁶⁹

Furthermore, our study showed the influence of family conflict on the behavior of children, shedding some light on the mechanisms involved in the association between psychopathology and behavioral symptoms, highlighting the need for interventions that target not only the children or adolescents affected by family conflict but provide parental training and education as well. For example, a systematic review⁷⁰ showed that three different parental behavior training programs were quite efficacious for externalizing and internalizing symptoms in children and adolescents with disruptive behavior disorders,

Table 4 Sensitivity analysis for moderation modeling using full information maximum likelihood

Outcome/covariates	Conditional model			Unconditional model		
	Standardized beta	SE	p-value	Standardized beta	SE	p-value
Left hippocampus						
Sex	-0.017	0.031	0.583	0.006	0.025	0.818
Age	0.059	0.030	0.047	0.038	0.025	0.132
SES	-0.010	0.052	0.844	0.032	0.024	0.181
FES conflict	-0.193	0.113	0.087	-0.055	0.030	0.064
ICV	0.589	0.035	< 0.001	0.625	0.020	< 0.001
Interaction (SES*FES)	0.150	0.120	0.210	0.006	0.025	0.818
Right hippocampus						
Sex	-0.028	0.031	0.363	-0.012	0.025	0.626
Age	0.061	0.030	0.039	0.042	0.025	0.089
SES	-0.042	0.050	0.400	0.024	0.024	0.305
FES conflict	-0.233	0.107	0.030	-0.046	0.029	0.111
ICV	0.571	0.039	< 0.001	0.627	0.020	< 0.001
Interaction (SES*FES)	0.206	0.113	0.069	-	-	-
Left amygdala						
Sex	-0.163	0.035	0.000	-0.179	0.027	< 0.001
Age	0.082	0.032	0.011	0.071	0.026	0.006
SES	0.044	0.056	0.434	0.029	0.025	0.247
FES conflict	-0.023	0.131	0.858	-0.052	0.030	0.076
ICV	0.522	0.030	< 0.001	-0.179	0.024	< 0.001
Interaction (SES*FES)	-0.039	0.137	0.778	0.545	0.027	< 0.001
Right amygdala						
Sex	-0.148	0.034	< 0.001	-0.167	0.026	< 0.001
Age	0.108	0.032	0.001	0.086	0.025	0.001
SES	0.058	0.054	0.287	0.021	0.024	0.396
FES conflict	0.100	0.126	0.425	-0.023	0.029	0.425
ICV	0.515	0.035	< 0.001	0.563	0.023	< 0.001
Interaction (SES*FES)	-0.140	0.130	0.281	-0.167	0.026	< 0.001
Internalizing						
Sex	0.108	0.038	0.005	0.102	0.030	0.001
Age	0.087	0.037	0.017	0.139	0.030	< 0.001
SES	-0.138	0.067	0.040	-0.111	0.029	< 0.001
FES conflict	0.178	0.147	0.227	0.221	0.034	< 0.001
ICV	0.001	0.040	0.982	0.010	0.032	0.747
Interaction (SES*FES)	0.047	0.158	0.764	0.102		
Externalizing						
Sex	-0.069	0.036	0.054	-0.060	0.030	0.043
Age	0.083	0.035	0.017	0.112	0.030	< 0.001
SES	-0.065	0.065	0.314	-0.111	0.028	< 0.001
FES conflict	0.389	0.116	0.001	0.327	0.033	< 0.001
ICV	-0.096	0.038	0.011	-0.055	0.033	0.093
Interaction (SES*FES)	-0.088	0.142	0.536	-0.060	-	-

SFES = Family Environment Scale; ICV = intracranial volume; SE = standard error; SES = socioeconomic status.

reinforcing the relevance of studies that seek a better understanding of the environmental factors involved in neurodevelopmental disorders.

Future directions in the field could involve innovative testing and robust models to determine whether behaviors change over time for a continuous outcome, such as externalizing scores (via latent change score), or a categorical approach in which latent homogenous groups of behaviors (via latent transition analysis) could be predicted by external factors, such as family conflict. Moreover, the synchronicity between changes in behavior and neuroimaging indicators could be tested for both outcomes through repeated measures over time.⁷¹ One limitation is that the scanner sites (one each in São Paulo

and Porto Alegre) were not considered to reduce model complexity in multilevel data analysis. Moreover, it is important to note that SES will likely be underestimated if observations within clusters are correlated (i.e., if there is an intra-cluster correlation for neuroimaging and CBCL scores). However, the point estimates (coefficients) are usually unaffected – clustering mainly affects standard error, not the estimated effects (the reported effect sizes). Therefore, the effect sizes are still trustworthy. However, if the model is misspecified (e.g., omitting cluster-level fixed/random effects), the coefficients could also be biased.

In conclusion, brain developmental trajectories are affected by biological and environmental factors throughout

childhood and adolescence. The impact of the family environment and SES on brain structure and mental health appears undeniable. Our findings are consistent with emerging evidence on an association between family conflict and externalizing/internalizing behavior in youth. It is important to consider that neurodevelopment is influenced not only by these windows of vulnerability but also by windows of opportunity that could buffer and ameliorate negative developmental outcomes. Understanding both developmental hazards and buffering effects on developmental trajectories warrants further attention in future studies regarding both prevention and intervention efforts in psychopathology, as well as determining the effects of low SES for social policy.

Acknowledgements

This study was mainly funded by Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP; process 15/50469-0) and in part by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES; Finance Code 001).

AZ has received funding from Coordenação de Aperfeiçoamento de Pessoal de Nível Superior/Conselho Nacional de Desenvolvimento Científico e Tecnológico and the National Institute of Mental Health /National Institutes of Health outside the scope of this work.

Disclosure

PMP has received payments or honoraria for lectures and presentations at educational events for Sandoz, Daiichi Sankyo, Eurofarma, Abbot, Libbs, Instituto Israelita de Pesquisa e Ensino Albert Einstein, and Instituto D'Or de Pesquisa e Ensino. LAPR has received a grant or research support from, served as a consultant to, and served on the speakers' bureau of Aché, Bial, Medice, Novartis/Sandoz, Pfizer/Upjohn, and Shire/Takeda over the last 3 years; and has received authorship royalties from Oxford Press and ArtMed. The Programa de Déficit de Atenção/Hiperatividade and Programa Ambulatorial para Transtorno Bipolar Juvenil, chaired by LAPR, have received unrestricted educational and research support from the following pharmaceutical companies over the last 3 years: Novartis/Sandoz and Shire/Takeda. The other authors report no conflicts of interest.

Data availability statement

Data dictionary is available at <https://osf.io/ktz5h/wiki/Data%20Dictionaries/> and <https://osf.io/w3jr4> to direct download. Individual-level data are available upon request to the Brazilian High-Risk Cohort Study research committee, by following the instructions and filling the research form available at <https://osf.io/ktz5h/wiki/home/>.

Author contributions

ETC: Conceptualization, Data curation, Methodology, Formal analysis, Writing – original draft, Writing – review & editing.

AM: Conceptualization, Data curation, Methodology, Formal analysis, Writing – original draft, Writing – review & editing.

HCM: Conceptualization, Data curation, Methodology, Formal analysis, Writing – original draft, Writing – review & editing.

VOS: Data curation, Formal analysis, Writing – original draft, Writing – review & editing.

AZ: Data curation, Formal analysis, Writing – original draft, Writing – review & editing.

CMA: Data curation, Formal analysis, Writing – original draft, Writing – review & editing.

PMP: Data curation, Formal analysis, Writing – original draft, Writing – review & editing.

FP: Data curation, Formal analysis, Writing – original draft, Writing – review & editing.

LAPR: Data curation, Formal analysis, Writing – original draft, Writing – review & editing.

APJ: Conceptualization, Supervision, Data curation, Methodology, Formal analysis, Writing – original draft, Writing – review & editing.

All authors have read and approved of the final version to be published.

Handling Editor: Rodolfo Damiano

References

- 1 Bush NR, Wakschlag LS, LeWinn KZ, Hertz-Picciotto I, Nozadi SS, Pieper S, et al. Family environment, neurodevelopmental risk, and the Environmental Influences on Child Health Outcomes (ECHO) initiative: looking back and moving forward. *Front Psychiatry*. 2020;11:547.
- 2 Flores SM, Salum GA, Manfro GG. Dysfunctional family environments and childhood psychopathology: the role of psychiatric comorbidity. *Trends Psychiatry Psychother*. 2014;36:147-51.
- 3 Michals S, Reeder K. Parental mental health impacts a child's well-being: parent-focused interventions can help. *J Stud Res*. 2023;12.
- 4 Luby J, Belden A, Botteron K, Marrus N, Harms MP, Babb C, et al. The effects of poverty on childhood brain development: the mediating effect of caregiving and stressful life events. *JAMA Pediatr*. 2013;167:1135-42.
- 5 Bush NR, Boyce WT. The contributions of early experience to biological development and sensitivity to context. In: Lewis M, Rudolph KD, editors *Handbook of developmental psychopathology* Boston, MA: Springer US; 2014. p. 287-309.
- 6 Byrne ML, Whittle S, Vijayakumar N, Dennison M, Simmons JG, Allen NB. A systematic review of adrenarche as a sensitive period in neurobiological development and mental health. *Dev Cogn Neurosci*. 2017;25:12-28.
- 7 Holder MK, Blaustein JD. Puberty and adolescence as a time of vulnerability to stressors that alter neurobehavioral processes. *Front Neuroendocrinol*. 2014;35:89-110.
- 8 Snyder HR, Hankin BL, Sandman CA, Head K, Davis EP. Distinct patterns of reduced prefrontal and limbic gray matter volume in childhood general and internalizing psychopathology. *Clin Psychol Sci*. 2017;5:1001-13.
- 9 Caldwell JZK, Armstrong JM, Hanson JL, Sutterer MJ, Stodola DE, Koenigs M, et al. Preschool externalizing behavior predicts gender-specific variation in adolescent neural structure. *PLoS One*. 2015;10:e0117453.
- 10 Bos MGN, Wierenga LM, Blankenstein NE, Schreuders E, Tamnes CK, Crone EA. Longitudinal structural brain development and externalizing behavior in adolescence. *J Child Psychol Psychiatry*. 2018;59:1061-72.
- 11 Bradley RH, Corwyn RF. Socioeconomic status and child development. *Annu Rev Psychol*. 2002;53:371-99.

- 12 Hackman DA, Farah MJ. Socioeconomic status and the developing brain. *Trends Cogn Sci*. 2009;13:65-73.
- 13 Hackman DA, Farah MJ, Meaney MJ. Socioeconomic status and the brain: mechanistic insights from human and animal research. *Nat Rev Neurosci*. 2010;11:651-9.
- 14 Judd N, Sauce B, Wiedenhoeft J, Tromp J, Chaarani B, Schliep A, et al. Cognitive and brain development is independently influenced by socioeconomic status and polygenic scores for educational attainment. *Proc Natl Acad Sci USA*. 2020;117:12411-8.
- 15 Raizada RDS, Kishiyama MM. Effects of socioeconomic status on brain development, and how cognitive neuroscience may contribute to leveling the playing field. *Front Hum Neurosci*. 2010;4:3.
- 16 Rakesh D, Zalesky A, Whittle S. Similar but distinct – effects of different socioeconomic indicators on resting state functional connectivity: findings from the Adolescent Brain Cognitive Development (ABCD) Study®. *Dev Cogn Neurosci*. 2021;51:101005.
- 17 Reiss F. Socioeconomic inequalities and mental health problems in children and adolescents: a systematic review. *Soc Sci Med*. 2013;90:24-31.
- 18 Dufford AJ, Evans GW, Dmitrieva J, Swain JE, Liberzon I, Kim P. Prospective associations, longitudinal patterns of childhood socioeconomic status, and white matter organization in adulthood. *Hum Brain Mapp*. 2020;41:3580-93.
- 19 McEwen CA, McEwen BS. Social structure, adversity, toxic stress, and intergenerational poverty: an early childhood model. *Annu Rev Sociol*. 2017;43:445-72.
- 20 Leijser LM, Siddiqi A, Miller SP. Imaging evidence of the effect of socio-economic status on brain structure and development. *Semin Pediatr Neurol*. 2018;27:26-34.
- 21 Brito NH, Noble KG. Socioeconomic status and structural brain development. *Front Neurosci*. 2014;8:276.
- 22 Jednoróg K, Altarelli I, Monzalvo K, Fluss J, Dubois J, Billard C, et al. The influence of socioeconomic status on children's brain structure. *PLoS One*. 2012;7:e42486.
- 23 Holz NE, Boecker R, Hohm E, Zohsel K, Buchmann AF, Blomeyer D, et al. The long-term impact of early life poverty on orbitofrontal cortex volume in adulthood: results from a prospective study over 25 years. *Neuropsychopharmacology*. 2015;40:996-1004.
- 24 Hanson JL, Hair N, Shen DG, Shi F, Gilmore JH, Wolfe BL, et al. Family poverty affects the rate of human infant brain growth. *PLoS One*. 2013;8:e80954.
- 25 Lawson GM, Duda JT, Avants BB, Wu J, Farah MJ. Associations between children's socioeconomic status and prefrontal cortical thickness. *Dev Sci*. 2013;16:641-52.
- 26 Noble KG, Houston SM, Kan E, Sowell ER. Neural correlates of socioeconomic status in the developing human brain. *Dev Sci*. 2012;15:516-27.
- 27 Hair NL, Hanson JL, Wolfe BL, Pollak SD. Association of child poverty, brain development, and academic achievement. *JAMA Pediatr*. 2015;169:822-9.
- 28 Hanson JL, Nacewicz BM, Sutterer MJ, Cayo AA, Schaefer SM, Rudolph KD, et al. Behavioral problems after early life stress: contributions of the hippocampus and amygdala. *Biol Psychiatry*. 2015;77:314-23.
- 29 Noble KG, Korgaonkar MS, Grieve SM, Brickman AM. Higher education is an age-independent predictor of white matter integrity and cognitive control in late adolescence. *Dev Sci*. 2013;16:653-64.
- 30 Shaked D, Katzel LI, Seliger SL, Gullapalli RP, Davatzikos C, Erus G, et al. Dorsolateral prefrontal cortex volume as a mediator between socioeconomic status and executive function. *Neuropsychology*. 2018;32:985-95.
- 31 Hanson JL, Chandra A, Wolfe BL, Pollak SD. Association between income and the hippocampus. *PLoS One*. 2011;6:e18712.
- 32 Hanson JL, Adkins DJ, Nacewicz BM, Barry KR. Impact of socioeconomic status on amygdala and hippocampus subdivisions in children and adolescents. *bioRxiv* [Preprint]. 2023 Mar 13:2023.03.10.532071. doi: 10.1101/2023.03.10.532071
- 33 Feng R, Bao W, Zhuo L, Gao Y, Yao H, Li Y, et al. Family conflict associated with intrinsic hippocampal-ofc connectivity in adolescent depressive disorder. *Front Psychiatry*. 2022;12:797898.
- 34 Kawamoto M, Takagishi H, Ishihara T, Takagi S, Kanai R, Sugihara G, et al. Hippocampal volume mediates the relationship of parental rejection in childhood with social cognition in healthy adults. *Sci Rep*. 2023;13:19167.
- 35 Altafim ERP, McCoy DC, Linhares MBM. Relations between parenting practices, socioeconomic status, and child behavior in Brazil. *Child Youth Serv Rev*. 2018;89:93-102.
- 36 Gallo EAG, De Mola CL, Wehrmeister F, Gonçalves H, Kieling C, Murray J. Childhood maltreatment preceding depressive disorder at age 18 years: a prospective Brazilian birth cohort study. *J Affect Disord*. 2017;217:218-24.
- 37 Biazoli CE Jr, Salum GA, Gadelha A, Rebello K, Moura LM, Pan PM, et al. Socioeconomic status in children is associated with spontaneous activity in right superior temporal gyrus. *Brain Imaging Behav*. 2020;14:961-70.
- 38 Repetti RL, Taylor SE, Seeman TE. Risky families: family social environments and the mental and physical health of offspring. *Psychol Bull*. 2002;128:330-66.
- 39 Plesko CM, Yu Z, Tobin K, Gross D. Social connectedness among parents raising children in low-income communities: an integrative review. *Res Nurs Health*. 2021;44:957-69.
- 40 Salum GA, Gadelha A, Pan PM, Moriyma TS, Graeff-Martins AS, Tamanaha AC, et al. High risk cohort study for psychiatric disorders in childhood: rationale, design, methods and preliminary results. *Int J Methods Psych Res*. 2015;24:58-73.
- 41 Weissman MM, Wickramaratne P, Adams P, Wolk S, Verdelli H, Olfson M. Brief screening for family psychiatric history: the family history screen. *Arch Gen Psychiatry*. 2000;57:675-82.
- 42 Sato JR, Salum GA, Gadelha A, Crossley N, Vieira G, Manfro GG, et al. Default mode network maturation and psychopathology in children and adolescents. *J Child Psychol Psychiatry*. 2016;57:55-64.
- 43 Sato JR, Biazoli CE Jr, Salum GA, Gadelha A, Crossley N, Vieira G, et al. Associations between children's family environment, spontaneous brain oscillations, and emotional and behavioral problems. *Eur Child Adolesc Psychiatry*. 2019;28:835-45.
- 44 Achenbach TM, Rescorla LA. Child behavior checklist for ages 6-18 (CBCL/6-18, ASEBA CBCL/6-18) [Database record]. 2001. APA PsycTests.
- 45 Moos RH, Moos BS. Family environment scale manual: development, applications, research Palo Alto, Ca: Consulting Psychologists Press; 1994.
- 46 Bordin IA, Rocha MM, Paula CS, Teixeira MCTV, Achenbach TM, Rescorla LA, et al. Child Behavior Checklist (CBCL), Youth Self-Report (YSR) and Teacher's Report Form (TRF): an overview of the development of the original and Brazilian versions. *Cad Saude Publica*. 2013;29:13-28.
- 47 Wagner F, Martel MM, Cogo-Moreira H, Maia CRM, Pan PM, Rohde LA, et al. Attention-deficit/hyperactivity disorder dimensionality: the reliable 'g' and the elusive 's' dimensions. *Eur Child Adolesc Psychiatry*. 2016;25:83-90.
- 48 Vianna VPT, Silva EA, Souza-Formigoni MLO. [Portuguese version of the Family Environment Scale: application and validation]. *Rev Saude Publica*. 2007;41:419-26.
- 49 Associação Brasileira de Empresas de Pesquisa (ABEP). Critério Brasil: critério de classificação econômica Brasil2019. abep.org/critério-brasil/
- 50 FreeSurfer software suite. surfer.nmr.mgh.harvard.edu/
- 51 Ducharme S, Albaugh MD, Nguyen T-V, Hudziak JJ, Mateos-Pérez JM, Labbe A, et al. Trajectories of cortical thickness maturation in normal brain development – the importance of quality control procedures. *Neuroimage*. 2016;125:267-79.
- 52 Hedges EP, Dimitrov M, Zahid U, Brito Vega B, Si S, Dickson H, et al. Reliability of structural MRI measurements: the effects of scan session, head tilt, inter-scan interval, acquisition sequence, Free-Surfer version and processing stream. *Neuroimage*. 2022;246:118751.
- 53 Fish LJ. Why multivariate methods are usually vital. *Meas Eval Couns Dev*. 1988;21:130-7.
- 54 Thompson B. Why multivariate methods are usually vital in research: some basic concepts. 1994 [cited 2025 Jun 10]. <https://eric.ed.gov/?id=ED367687>
- 55 Holz NE, Laucht M, Meyer-Lindenberg A. Recent advances in understanding the neurobiology of childhood socioeconomic disadvantage. *Curr Opin Psychiatry*. 2015;28:365-70.
- 56 Dong N, Kelcey B, Spybrook J. Identifying and estimating causal moderation for treated and targeted subgroups. *Multivariate Behav Res*. 2023;58:221-40.
- 57 Muthén L, Muthén B. Mplus user's guide. 2020.

- 58 Khaleque A. Perceived parental hostility and aggression, and children's psychological maladjustment, and negative personality dispositions: a meta-analysis. *J Child Fam Stud*. 2017;26:977-88.
- 59 Weymouth BB, Buehler C, Zhou N, Henson RA. A meta-analysis of parent-adolescent conflict: disagreement, hostility, and youth maladjustment. *J Fam Theory Rev*. 2016;8:95-112.
- 60 Collins WA, Maccoby EE, Steinberg L, Hetherington EM, Bornstein MH. Contemporary research on parenting: the case for nature and nurture. *Am Psychol*. 2000;55:218-32.
- 61 Essex MJ, Kraemer HC, Armstrong JM, Boyce WT, Goldsmith HH, Klein MH, et al. Exploring risk factors for the emergence of children's mental health problems. *Arch Gen Psychiatry*. 2006;63:1246-56.
- 62 Ursache A, Noble KG; Pediatric Imaging, Neurocognition and Genetics Study. Socioeconomic status, white matter, and executive function in children. *Brain Behav*. 2016;6:e00531.
- 63 Bøe T, Sivertsen B, Heiervang E, Goodman R, Lundervold AJ, Hysing M. Socioeconomic status and child mental health: the role of parental emotional well-being and parenting practices. *J Abnorm Child Psychol*. 2014;42:705-15.
- 64 Katzmann J, Döpfner M, Götz-Dorten A. Child-based treatment of oppositional defiant disorder: mediating effects on parental depression, anxiety and stress. *Eur Child Adolesc Psychiatry*. 2018;27:1181-92.
- 65 Esposito-Smythers C, Spirito A. Adolescent substance use and suicidal behavior: a review with implications for treatment research. *Alcohol Clin Exp Res*. 2004;28:77S-88S.
- 66 Meque I, Dachew BA, Maravilla JC, Salom C, Alati R. Externalizing and internalizing symptoms in childhood and adolescence and the risk of alcohol use disorders in young adulthood: a meta-analysis of longitudinal studies. *Aust N Z J Psychiatry*. 2019;53:965-75.
- 67 Tung I, Li JJ, Meza JI, Jezior KL, Kianmahd JSV, Hentschel PG, et al. Patterns of comorbidity among girls with ADHD: a meta-analysis. *Pediatrics*. 2016;138:e20160430.
- 68 Ding N, Gao H, Jiang J, Zhai M, Shao H, Fu L, et al. The characteristics and factors of the internalizing and externalizing behaviours of children at high risk for autism spectrum disorder. *BMC Psychiatry*. 2021;21:523.
- 69 Cao P, Chen C, Si Q, Li Y, Ren F, Han C, et al. Volumes of hippocampal subfields suggest a continuum between schizophrenia, major depressive disorder and bipolar disorder. *Front Psychiatry*. 2023;14:1191170.
- 70 Zarakoviti E, Shafran R, Papadimitriou D, Bennett SD. The efficacy of parent training interventions for disruptive behavior disorders in treating untargeted comorbid internalizing symptoms in children and adolescents: a systematic review. *Clin Child Fam Psychol Rev*. 2021;24:542-52.
- 71 Kievit RA, Fuhrmann D, Borgeest GS, Simpson-Kent IL, Henson RNA. The neural determinants of age-related changes in fluid intelligence: a pre-registered, longitudinal analysis in UK Biobank. *Wellcome Open Res*. 2018;3:38.