



ORIGINAL ARTICLE

Parental style, childhood trauma, and impulsivity contribute to the severity of crack cocaine addiction: a cluster analysis

Daniela Benzano Bumaguin,^{1,2} Joana Corrêa de Magalhães Narváez,^{1,3} Jaqueline B. Schuch,^{1,2,4} Deborah Daitschman,¹ Sérgio Kakuta Kato,³ Lisia von Diemen,^{1,2} Felix H.P. Kessler^{1,2}

¹Centro de Pesquisa em Álcool e Drogas, Hospital de Clínicas de Porto Alegre, Universidade Federal do Rio Grande do Sul (UFRGS), Porto Alegre, RS, Brazil. ²Programa de Pós-Graduação em Psiquiatria e Ciências do Comportamento, UFRGS, Porto Alegre, RS, Brazil. ³Universidade Federal de Ciências da Saúde de Porto Alegre, Porto Alegre, RS, Brazil. ⁴Responsabilidade Social, Hospital Moinhos de Vento, Porto Alegre, RS, Brazil.

Objective: To determine clusters (groups) of substance use in crack cocaine users through severity scores on the sixth version of the Addiction Severity Index (ASI-6) and compare the groups in relation to risk factors, such as parental style, childhood maltreatment, and impulsivity.

Methods: This cross-sectional study included 531 men with substance use disorder who were admitted to an inpatient addiction treatment unit in southern Brazil. To detect more homogeneous groups of individuals, the K-means clustering based on ASI-6 scores was used to create groups of individuals with similar severity scores in different areas. Parenting styles were assessed using the Measure of Parental Style, childhood trauma was assessed using the Childhood Trauma Questionnaire, and impulsivity was measured using the Barratt Impulsivity Scale 11. Poisson regression was used for association analysis.

Results: Two distinct clusters were identified, which differed significantly across all composite scores ($p < 0.001$). These associations were further confirmed through Poisson regression analysis. The more severe cluster showed significantly higher scores for maternal abuse ($p = 0.026$), sexual abuse ($p = 0.003$), motor impulsivity ($p = 0.014$), and unplanned impulsivity above the 75th percentile ($p = 0.032$) than the less severe group. Other parenting styles, trauma types, and impulsivity did not differ significantly between the groups.

Conclusion: Dividing patients into severity clusters can contribute to more targeted treatments. Further research on outpatients would reinforce the importance of early life factors and impulsivity treatment.

Keywords: Childhood; behavior; crack cocaine; addiction; cluster analysis

Introduction

The worldwide estimated prevalence of crack cocaine use is between 0.3 and 0.5%,¹ with an estimated annual usage prevalence of 1.03% in South America.¹ Although snorted cocaine consumption has declined in the European Union, crack cocaine consumption has not.² In Brazil in 2015, the lifetime estimated use of snorted cocaine and crack cocaine was 3.1% and 0.9%, respectively.³ However, this estimated use of cocaine/crack reached 12.9% in a recent study of students from metropolitan São Paulo.⁴ Crack cocaine use disorder ranked first as a motive for detoxification treatment in Argentina, Uruguay, and Chile, and it ranked second in

Brazil.² This concerning scenario reinforces the need for further studies in the area.

Crack cocaine use has serious repercussions on the lives of users and on society. Including a major impact on the physical and mental health of users.⁵ Crack cocaine users have historically suffered social marginalization, a history of imprisonment, violence, and sex work, including exposure to sexually transmitted diseases.⁶⁻⁸ Different factors can affect the severity of addiction, including inherent sex differences, such as prevalence, metabolism, and hormonal responses, in addition to vulnerability and sociocultural factors.⁹⁻¹¹ Overall, clinical presentation and biological effects distinctly affect the course of addiction according to sex. Hence, sex-specific

Correspondence: Daniela Benzano Bumaguin, Universidade Federal do Rio Grande do Sul, Hospital de Clínicas de Porto Alegre, Centro de Pesquisa em Álcool e Drogas, Rua Ramiro Barcelos, 2350, CEP 90035-903, Porto Alegre, RS, Brazil.
E-mail: danielabenzano@gmail.com
Submitted Mar 27 2024, accepted Jan 19 2025.

How to cite this article: Bumaguin DB, Narváez JCM, Schuch JB, Daitschman D, Kato SK, von Diemen L, et al. Parental style, childhood trauma, and impulsivity contribute to the severity of crack cocaine addiction: a cluster analysis. Braz J Psychiatry. 2025;47:e20243645. <http://doi.org/10.47626/1516-4446-2024-3645>

analyses are necessary to better understand the role of different factors in addiction severity and reduce sex-related bias. Parenting styles and childhood maltreatment have also been associated with substance use. For instance, authoritarian, permissive, and neglectful parenting styles have been more associated with drug use and addiction tendency than indulgent parenting styles (warmth without strictness).^{12,13} Furthermore, in a meta-analysis Rakovski et al.¹⁴ found evidence of a relationship between different types of abuse and the use of licit and illicit substances. In particular, there was a relationship between physical and sexual abuse and marijuana use, and between sexual abuse and cocaine, hallucinogen, and other unspecified substance use.¹⁴ Impulsive behavior has also been associated with substance use through different mechanisms. One such mechanism is reduced cognitive and response inhibition. Another factor is damaged brain structure and function due to acute or chronic substance use. Finally, genetic and environmental factors can also influence distinct behaviors related to substance use disorder.¹⁵

Individuals with addiction have difficulties accessing health systems and a high prevalence of medical and psychiatric illness. Social inclusion is fundamental but still insufficient and should also be linked to addiction treatment.¹⁶ Recent studies have highlighted factors associated with a lack of adherence after hospitalization or outpatient treatment, such as the lack of family support and the fact that some patients have been homeless in the last 6 months.¹⁷

Over the last few decades, psychiatric studies have classified substance users to better understand use and severity profiles, mainly due to the widely heterogeneous characteristics mentioned above. Grouping users to better understand them can lead to more personalized care and more effective treatment strategies. To find behavioral differences among polydrug users, a British study identified three patterns of use: extensive, moderate, and none.¹⁸ Another recent study described four classes among crack cocaine users: light use, moderate use, daily use, and very high quantity use.¹⁹ In this context, it is extremely important to find a more reliable way, based on the clinical context of users, to understand consumption patterns and repercussions, considering socioemotional aspects and behavior.

Dividing users into groups based on severity, as well as affective and behavioral expression, would allow a better response to their care needs, especially in Brazil's sociocultural context. Since substance users face problems in social, medical, and legal areas, a comprehensive assessment can help create more homogeneous groups. The Addiction Severity Index (ASI) provides a global assessment of addiction severity. Using this instrument to group users into clusters would be of great importance due to its scope.

Therefore, the aims of this study were to determine severity clusters of substance use in crack cocaine users according to severity scores on the sixth version of the ASI (ASI-6) and to compare these groups regarding risk factors cited in the literature, such as parental style, childhood maltreatment, and impulsivity.

Methods

Sample

The sample consisted of 531 men diagnosed with substance use disorder according to DSM-5 criteria who were admitted to an inpatient addiction treatment unit in Porto Alegre, Brazil. Patients aged at least 18 years who reported that crack cocaine was their preferred substance and provided written informed consent were included in the study. We excluded patients with psychotic disorders or cognitive deficits – assessed through interviews with specialists – that prevented them from responding to the questionnaires. An expected difference between the clusters for all variables, with an effect size $d = 1.5$, considering a separation between the centroids $\Delta > 4$ would result in a power of 100%.

Procedures

Trained health researchers, supervised by senior psychologists and psychiatrists, assessed all participants using the Brazilian versions of clinical and semi-structured interviews. Evaluator training guaranteed the correct application of the instruments, as well as the veracity and accuracy of the measurements. Patients were invited to participate in the study on the first or second day after hospitalization, provided the effects of intoxication did not interfere with their ability to provide informed consent. The questionnaires and scales were applied during the first week of detoxification, although this was delayed if it was determined that acute symptoms could interfere in the assessment.

To assess substance use severity, sociodemographic and clinical data were collected, in addition to ASI-6²⁰ results. The Measure of Parental Style is a self-report instrument that assesses the relationship between respondents and their parents during childhood and adolescence.²¹ It measures three parenting styles: indifference, abuse, and overcontrol. The Childhood Trauma Questionnaire, which is also self-reported,²² assesses the occurrence of maltreatment, abuse, and neglect during childhood, but does not necessarily involve parents. It consists of five subscales: emotional abuse, physical abuse, sexual abuse, emotional neglect, and physical neglect. Childhood trauma is categorized as absent or present based on scores for each subtype. Childhood trauma is classified as scores ≥ 10 in the physical neglect and physical abuse subscales, ≥ 15 in the emotional neglect subscale, ≥ 13 in the emotional abuse subscale, and ≥ 8 in the physical abuse subscale. Patient impulsivity was measured using the Barratt Impulsivity Scale 11,²³ which consists of three subscales: attentional, motor, and unplanned impulsivity. All questionnaires had been previously translated and validated for Brazilian Portuguese.

Statistical analyses

The REDCap platform was used to include and extract data. Statistical analyses were performed in R 4.2.1 and

SPSS 20.0. Data normality was assessed using the Kolmogorov-Smirnov test.

Clusters were identified among crack cocaine users using K-means cluster analysis. This type of analysis creates sets of very similar individuals who differ from other groups. Severity clusters were defined using the K-means technique from the composite scores of the ASI-6 medical, employment/support, alcohol, drugs, family/social, legal, and psychiatric dimensions. The composite score for family and children was removed because many users did not have children. Participants with complete ASI-6 composite scores were included in the analysis. The optimal number of clusters was determined using the NbClust package in R. This package uses 30 indexes to determine the number of clusters and proposes the best clustering scheme. After identifying the clusters by composite ASI severity scores, the standardized mean difference (Cohen's *d*) was calculated to assess the magnitude of the difference within user groups. The variables were quantitatively compared using Student's *t*-test for independent samples or the Mann-Whitney test according to data distribution. Associations among categorical variables were determined using the chi-square test with Yates's continuity correction. Poisson regression model with robust variance was used to evaluate the relationship between the different factors and the more severe usage cluster, with adjustment variables selected according to their association in the bivariate analysis. The blocks for the hierarchical analysis were 1) parenting style, 2) childhood trauma, and 3) impulsivity (dichotomized as below or above the 75th percentile to highlight the relationship between high

impulsivity and more serious substance use). A 5% significance level was used in all analyses.

Ethics statement

This study was approved by the Hospital de Clínicas de Porto Alegre ethics committee (protocol no. 2014-0249). All patients provided written informed consent before inclusion.

Results

Data from 531 crack cocaine users were analyzed. Through statistical analysis, the optimal number of clusters was defined as two, one with 306 participants, and the other with 225 (Figure 1).

The clusters differed significantly for all composite scores (Supplementary Table S1). Cluster 1 had lower ASI-6 composite scores for all scales than cluster 2, except for the family/social support scale. To understand the magnitude of the difference between the two clusters, Cohen's standardized effect size was calculated. The effect sizes and their confidence intervals are shown in Figure 2. The composite scores with the greatest magnitude of standardized difference were drugs, alcohol, psychiatric, medical, and family and social problems. Although there were significant differences among the other composite scores, the effect magnitudes did not differ significantly.

Sociodemographic characteristics were compared between more and less severe clusters, and there were no significant differences regarding age, days of




Figure 1 More severe (cluster 2) and less severe (cluster 1) clusters of crack cocaine users admitted to a specialized addiction treatment unit (n=531).




Figure 2 Comparison of Cohen's standardized effect size between more and less severe clusters of crack cocaine users admitted to a specialized addiction treatment unit (n = 531). White circles represent the point estimate of Cohen's *d*, while the lines represent the 95%CI of the estimate. FS = family/social. † Between the dotted lines, Cohen's *d* is 0.5-0.8 (medium effect size). ‡ Between the dotted line and the right end of the graph, Cohen's *d* is > 0.8 (large effect size or more).

Table 1 Comparison of characteristics between the more and less severe clusters of crack cocaine users admitted to a specialized addiction treatment unit (n=531)

Characteristics	Least severe cluster (n=306)	Most severe cluster (n=225)	p-value
Age (years), mean (SD) [†]	35.3 (9.7)	35.8 (9.7)	0.084
Days of hospitalization, mean (SD) [†]	21.5 (16.0)	20.7 (15.3)	0.566
Race (White) [†]	216 (70.6)	148 (65.8)	0.278
Income for the last 6 months (BRL), median (IQR) [§]	7,000 (4,260-12,000)	5,400 (1,800-10,501)	0.014
Marital status (with partner) [†]	58 (19.0)	37 (16.6)	0.559
Education level [†]			0.941
Primary education	177 (59.0)	129 (57.6)	
High school	91 (30.3)	71 (31.7)	
Higher education	32 (10.7)	24 (10.7)	
Age of first crack cocaine use (years), median (IQR) [§]	18 (16-22)	18 (16-24)	0.611

Data presented as n (%), unless otherwise specified.
Bold type denotes statistical significance.
IQR = interquartile range.
[†] Compared using Student's *t*-test for independent samples.
[‡] Associated using the chi-square test or the chi-square test with Yates's continuity correction.
[§] Compared using the Mann-Whitney test.

hospitalization, race, marital status, education level, or age at first use of crack cocaine. Users in the more severe cluster had significantly lower income than those in the less severe cluster (Table 1). Moreover, in the more severe cluster, 39.6% used cannabis, while in the less severe cluster only 29.1% did so (*p* = 0.015). Alcohol use also differed significantly between the groups (69.3% in the more severe cluster vs. 44.8% in the less severe cluster, *p* < 0.001). No participants reported the use of other illicit substances.

Factors associated with more severe use, such as parenting style, childhood trauma, and impulsivity were also compared between the clusters. The more severe cluster had significantly higher scores for maternal

indifference (median 0.50 vs 0.33, *p* = 0.006), maternal abuse (median 0.60 vs 0.40, *p* = 0.010), and paternal indifference (median 1.17 vs 0.67, *p* = 0.010) than the less severe cluster. In addition, the more severe cluster had a higher childhood trauma score (median 49 vs 42, *p* < 0.001) and a higher frequency of physical abuse (46.8 vs 34.8%, *p* = 0.011), emotional neglect (26.6 vs 16.8%, *p* = 0.013), emotional abuse (48.8 vs 33.9%, *p* = 0.002), and sexual abuse (27.9 vs 14.7%, *p* = 0.001). The more severe group also had higher total impulsivity scores (median 79 vs 73, *p* < 0.001) and a higher frequency of participants with motor (29.6 and 16.4%, *p* = 0.001) and unplanned impulsivity scores above the 75th percentile (29.4 and 14.8%, *p* < 0.001).

Table 2 Factors associated with the more severe cluster of crack cocaine users admitted to a specialized addiction treatment unit (n=531)

Blocks	Crack cocaine	
	Adjusted PR	p-value
Parenting styles		
Maternal indifference	1.00 (0.83-1.20)	0.968
Maternal abuse	1.24 (1.03-1.51)	0.026
Paternal indifference	1.12 (0.93-1.34)	0.232
Paternal abuse	1.03 (0.84-1.25)	0.797
Moderate-to-severe trauma		
Physical neglect	1.06 (0.84-1.35)	0.632
Physical abuse	1.11 (0.85-1.45)	0.434
Emotional neglect	1.15 (0.89-1.48)	0.285
Emotional abuse	1.16 (0.89-1.51)	0.276
Sexual abuse	1.42 (1.13-1.79)	0.003
Impulsivity > 75th percentile		
Attentional	1.03 (0.78-1.36)	0.822
Motor	1.36 (1.06-1.73)	0.014
Unplanned	1.33 (1.03-1.73)	0.032

Bold type denotes statistical significance.

The blocks were adjusted for factors with $p < 0.05$ in the preceding block.

PR = prevalence ratio estimated using Poisson regression with robust variance.

In a Poisson regression model with robust variance estimation (Table 2) that was hierarchically divided into three blocks (parenting style, trauma, and impulsivity), maternal abuse, sexual abuse, motor impulsivity, and unplanned impulsivity scores above the 75th percentile were associated with the more severe cluster.

Discussion

Through robust statistical analysis, this study identified two severity clusters of male crack cocaine users among a sample of heavy users. The patients were divided into more and less severe groups, which had important differences in parental style, childhood trauma, and impulsivity scores, but not in most sociodemographic variables. This is the first Brazilian study to establish a grouping among illicit substance users through multi-dimensional assessment. One hypothesis was that a large number of clusters would be identified, given that in other studies a high heterogeneity of substances used usually translated into more subgroups.¹⁹ However, we detected only two distinct groups, probably due to a more homogeneous sample (i.e., all individuals were inpatients, and the main substance was crack cocaine).

Dividing individuals into clusters or groups that are relatively homogenous in terms of sex, use severity, need for hospitalization, and choice of drug can reveal important differences, especially regarding early trauma. Abusive and traumatic situations can impair neuropsychological and self-preservation functions, leading to greater impulsivity.²⁴ This can facilitate the development of more personalized therapeutic and preventive interventions to enable more efficient referrals to psychiatric and health care services.

The difference between the clusters in composite score for psychiatric, social, and family problems corroborates the current treatment guidelines. Thus, according

to our results, psychiatric treatment, psychosocial techniques, and family support should be supported.²⁵ Our results also align with other studies in which users who have been homeless have worse scores due to their high vulnerability.⁷ Moreover, individuals with substance use disorder have a high frequency of clinical comorbidities,²⁶ and the large magnitude of difference in composite scores for medical problems also corroborates the literature. In these patients, there is a higher frequency of sexually transmitted infections, such as HIV and hepatitis,²⁷⁻²⁹ as well as other medical conditions, such as hypertension, respiratory, and digestive tract diseases.²⁶

We found significant differences in some important risk factors between severity clusters. The results of the adjusted models showed that maternal abuse, sexual abuse, and motor and unplanned impulsivity were associated with the more severe cluster. A recent study by our research team based on the same clinical sample found higher childhood trauma scores in crack cocaine users than alcohol users and individuals without addiction.³⁰ Dubowitz et al.³¹ also found different effects between trauma types and users of different substances, although sexual abuse had no effects, regardless substance type. In general, childhood maltreatment has differed significantly between substance users (such as crack cocaine) and controls.³² Although physical and emotional abuse tends to have a major impact, the potential harmful impact of sexual abuse must be highlighted. Previous studies have more commonly associated sexual abuse with female crack cocaine users. In a sample of pregnant adolescents, the chance of substance use was 13 times higher among those who suffered sexual abuse and a 17 times higher among those who suffered both sexual and physical abuse, suggesting a synergistic effect.³³ Although this large effect was not found in our sample of men, there was a significant association in the same direction. Moreover, early

experimentation and more severe drug use may be linked to the need to anesthetize early trauma in non-protective family contexts.²⁷ The relationship between sexual abuse and more serious substance use could lead to the conclusion that children who suffer from sexual abuse are more likely to become substance users due to the consequences of both the trauma itself and the underlying family conditions.

It is known that many users face early vulnerabilities due to their parental environment.^{24,27} Maternal abuse is linked to greater psychological disruption, especially in countries where single-mother families predominate. However, studies have shown that maternal and paternal care is a protective effect against substance use and abuse,³⁴ as well as that there is a relationship between parental style and multiple drug use.³⁵ Studies of adolescents have shown that those with higher levels of affection and communication with their parents had fewer criteria for substance dependency.³⁶ Other authors have found that poor parenting styles were risk factors for substance abuse in adolescents, a relationship that was mediated by social support.³⁷

In terms of behavioral expression, higher impulsivity scores have been found in crack cocaine users^{24,38} and have been associated with crack cocaine severity.^{15,39} Higher impulsivity scores have been strongly related to parameters of long-term crack cocaine use.⁴⁰ Several studies have found a difference between substance abusers and controls regarding a preference for short-term rewards over long-term rewards. There is evidence that more immediate and hedonistic behavior, such as eschewing delayed rewards, which is related to impulsivity, is higher among individuals with addictive behavior. Therefore, differences between severity clusters are to be expected.⁴¹ Previous studies have also found an association between higher impulsivity scores, childhood trauma, and more severe substance use among crack cocaine users. The relationship between childhood trauma and impulsivity, as well as between the latter and substance use severity, suggests that impulsivity plays a mediating role. Cognitive and decision-making processes are influenced by structural and historical modifications due to trauma, which, in turn, influence impulsivity and substance use severity. The relationship between childhood trauma and substance use severity may be partially mediated by impulsivity, and further studies would be extremely relevant.²⁴ Impulsivity can accelerate substance seeking behavior and substance use can increase impulsivity, a vicious circle that can spiral into uncontrolled behavior. Therefore, one approach to addiction treatment would be to reduce impulsive behavior and prevent the development of uncontrolled use or relapse.⁴²

This study has made important contributions to the Brazilian and international literature, using ASI-6 composite scores to investigate different aspects of substance use severity. However, it presents some limitations. Since our sample only consisted of male patients, the findings and analyses can only be generalized to men. Although the participants' responses could have been affected by memory bias, the questionnaire was applied by trained evaluators to guarantee higher quality. Being a cross-

sectional study, the patients were evaluated at a single point in time, and variations in substance use could not be investigated, only lifetime and recent use. Furthermore, the patients varied little in terms of severity since all of them were indicated for hospitalization at the time of assessment. Nevertheless, severity clusters were identified. Longitudinal studies are needed to clarify the relationship between use severity and association with a particular cluster. However, it may be challenging to monitor these patients and conduct this type of study. Moreover, addiction severity may vary more widely in non-hospitalized individuals. Further studies with non-hospitalized substance users are needed to determine whether the same factors observed in our study can also distinguish specific groups among them.

In conclusion, we were able to identify two clusters of patients with different severity levels, and these clusters differed regarding more distal aspects, such as parental style and childhood trauma, as well as more proximal aspects, such as impulsivity. This brings new perspective to this difficult-to-treat population, allowing a more thorough treatment approach aimed at resolving factors associated with psychiatric and clinical comorbidities, as well as current and early aspects of the user's life. Although its subject is highly complex, the present study presents a new approach that can guide preventive and therapeutic psychoeducational techniques aimed at this vulnerable population.

Disclosure

The authors report no conflicts of interest.

Data availability statement

The data that support this study are available in the body of the paper and/or supplementary materials.

Author contributions

DBB: Conceptualization, Data curation, Formal analysis, Funding acquisition, Investigation, Methodology, Project administration, Resources, Software, Validation, Visualization, Writing – original draft, Writing – review & editing. JCMN: Conceptualization, Data curation, Investigation, Validation, Visualization, Writing – original draft, Writing – review & editing.

JBS: Methodology, Supervision, Writing – original draft, Writing – review & editing.

DD: Conceptualization, Investigation, Resources, Software, Visualization, Writing – review & editing.

SKK: Methodology, Software, Validation, Writing – review & editing.

LD: Conceptualization, Funding acquisition, Investigation, Methodology, Project administration, Supervision, Writing – original draft, Writing – review & editing.

FHPK: Conceptualization, Funding acquisition, Investigation, Methodology, Project administration, Resources, Supervision, Writing – original draft, Writing – review & editing.

All authors have read and approved of the final version to be published.

Handling Editor: João Castaldelli-Maia

References

- United Nations (UN). Office on drugs and crime [Internet]. 2021 [cited 2024 Mar 18]. www.unodc.org/unodc/en/data-and-analysis/wdr2021.html
- United Nation (UN). World Drug Report 2022 [Internet]. [cited 2024 Mar 18]. www.unodc.org/unodc/en/data-and-analysis/world-drug-report-2022.html
- Bastos FIPM, de Vasconcellos MTL, De Boni RB, dos Reis NB, Coutinho CF de S. III Levantamento Nacional sobre o uso de drogas pela população brasileira. Rio de Janeiro: Ministério da Saúde, Fiocruz; 2017.
- Vellozo EP, Vitale MSDS, Passos MAZ, Niskier SR, Schoen TH, Hall PR, et al. Prevalence of psychoactive substance use by adolescents in public schools in a municipality in the São Paulo Metropolitan Area, Brazil. *Cad Saude Publica*. 2023;39:e00169722.
- Butler AJ, Rehm J, Fischer B. Health outcomes associated with crack-cocaine use: systematic review and meta-analyses. *Drug Alcohol Depend*. 2017;180:401-16.
- de Carvalho HB, Seibel SD. Crack cocaine use and its relationship with violence and HIV. *Clinics (São Paulo)*. 2009;64:857-66.
- Halpern SC, Scherer JN, Roglio V, Faller S, Sordi A, Ornell F, et al. [Clinical and social vulnerabilities in crack users according to housing status: a multicenter study in six Brazilian state capitals]. *Cad Saude Publica*. 2017;33:e00037517.
- Guimarães RA, de Oliveira Landgraf de Castro V, do Valle Leone de Oliveira SM, Stabile AC, Motta-Castro ARC, Dos Santos Carneiro MA, et al. Gender differences in patterns of drug use and sexual risky behaviour among crack cocaine users in Central Brazil. *BMC Psychiatry*. 2017;17:412.
- Lejuez CW, Bornovalova MA, Reynolds EK, Daughters SB, Curtin JJ. Risk factors in the relationship between gender and crack/cocaine. *Exp Clin Psychopharmacol*. 2007;15:165-75.
- Pope SK, Falck RS, Carlson RG, Leukefeld C, Booth BM. Characteristics of rural crack and powder cocaine use: gender and other correlates. *Am J Drug Alcohol Abuse*. 2011;37:491-6.
- Abdalla RR, Madruga CS, Ribeiro M, Pinsky I, Caetano R, Laranjeira R. Prevalence of cocaine use in Brazil: data from the II Brazilian National Alcohol and Drugs Survey (BNADS). *Addict Behav*. 2014;39:297-301.
- Villarejo S, Garcia OF, Alcaide M, Villarreal ME, Garcia F. Early family experiences, drug use, and psychosocial adjustment across the life span: is parental strictness always a protective factor? *Psychosoc Interv*. 2024;33:15-27.
- Fatemisough H, Moradi A, Dastgerdi MV, Fatemi R. Relationship between parenting styles and tendency to addiction in Iran: a systematic review and meta-analysis. *Iran J Public Health*. 2024;53:70-80.
- Rakovski C, Lalli M, Gu J, Hobson M, Wollenhaupt-Aguier B, Minuzzi L, et al. Childhood maltreatment as a predictor of substance use/misuse among youth: a systematic review and meta-analysis. *Neurosci Biobehav Rev*. 2024;166:105873.
- Kozak K, Lucatch AM, Lowe DJE, Balodis IM, MacKillop J, George TP. The neurobiology of impulsivity and substance use disorders: implications for treatment. *Ann N Y Acad Sci*. 2019;1451:71-91.
- Ribeiro M, Duailibi S, Frazinger R, Alonso ALS, Marchetti L, Williams AV, et al. The Brazilian 'Cracolândia' open drug scene and the challenge of implementing a comprehensive and effective drug policy. *Addiction*. 2016;111:571-3.
- Klein E, Ornell F, Roglio VS, Scherer JN, Sordi AO, Schuch JB, et al. Early discharge predictors among inpatient crack cocaine users. *Trends Psychiatry Psychother*. 2023;45:e20210401.
- Smith GW, Farrell M, Bunting BP, Houston JE, Shevlin M. Patterns of polydrug use in Great Britain: findings from a national household population survey. *Drug Alcohol Depend*. 2011;113:222-8.
- Liu Y, Vaddiparti K, Cheong J, Cottler LB. Identification of typologies of cocaine use based on quantity, frequency, and duration of use: a latent profile analysis. *J Addict Med*. 2021;15:211-8.
- Kessler F, Cacciola J, Alterman A, Faller S, Souza-Formigoni ML, Cruz MS, et al. Psychometric properties of the sixth version of the Addiction Severity Index (ASI-6) in Brazil. *Braz J Psychiatry*. 2012;34:24-33.
- Baeza FLC, Caldieraro MAK, Pinheiro DO, Fleck MP. Translation and cross-cultural adaptation into Brazilian Portuguese of the Measure of Parental Style (MOPS) - a self-reported scale - according to the International Society for Pharmacoeconomics and Outcomes Research (ISPOR) recommendations. *Braz J Psychiatry*. 2010;32:159-63.
- Grassi-Oliveira R, Stein LM, Pezzi JC. [Translation and content validation of the Childhood Trauma Questionnaire into Portuguese language]. *Rev Saude Publica*. 2006;40:249-55.
- Diemen LV, Szobot CM, Kessler F, Pechansky F. Adaptation and construct validation of the Barratt Impulsiveness Scale (BIS 11) to Brazilian Portuguese for use in adolescents. *Braz J Psychiatry*. 2007;29:153-6.
- Narvaez JCM, Magalhães PVS, Trindade EK, Vieira DC, Kauer-Sant'Anna M, Gama CS, et al. Childhood trauma, impulsivity, and executive functioning in crack cocaine users. *Compr Psychiatry*. 2012;53:238-44.
- Zaleski M, Laranjeira RR, Marques ACP, Ratto L, Romano M, Alves HNP, et al. Guidelines of the Brazilian Association of Studies on Alcohol and Other Drugs (ABEAD) for diagnosis and treatment of psychiatric comorbidity with alcohol and other substance and dependence. *Int Rev Psychiatry*. 2017;29:254-62.
- Lagisetty PA, Maust D, Heisler M, Bohnert A. Physical and mental health comorbidities associated with primary care visits for substance use disorders. *J Addict Med*. 2017;11:161-2.
- Narvaez JCM, Roglio VS, Di Tommaso B, Pechansky F. Transgenerational cycle of traumatization and HIV risk exposure among crack users. *Int J Environ Res Public Health*. 2023;20:5285.
- Barcellos NT, Fuchs SC, Fuchs FD. Prevalence of and risk factors for HIV infection in individuals testing for HIV at counseling centers in Brazil. *Sex Transm Dis*. 2003;30:166-73.
- de Azevedo RC, Botega NJ, Guimarães LA. Crack users, sexual behavior and risk of HIV infection. *Braz J Psychiatry*. 2007;29:26-30.
- Hoffmann A, Benzano D, Ornell F, Kessler FHP, von Diemen L, Schuch JB. Childhood trauma subtypes may influence the pattern of substance use and preferential substance in men with alcohol and/or crack-cocaine addiction. *Braz J Psychiatry*. 2022;44:416-9.
- Dubowitz H, Roesch S, Lewis T. Child maltreatment, early adult substance use, and mediation by adolescent behavior problems. *Child Maltreat*. 2021;26:238-48.
- Banducci AN, Hoffman E, Lejuez CW, Koenen KC. The relationship between child abuse and negative outcomes among substance users: psychopathology, health, and comorbidities. *Addict Behav*. 2014;39:1522-7.
- Berenson AB, San Miguel VV, Wilkinson GS. Violence and its relationship to substance use in adolescent pregnancy. *J Adolesc Health*. 1992;13:470-4.
- Eun JD, Paksarian D, He JP, Merikangas KR. Parenting style and mental disorders in a nationally representative sample of US adolescents. *Soc Psychiatry Psychiatr Epidemiol*. 2018;53:11-20.
- Valente JY, Cogo-Moreira H, Sanchez ZM. Predicting latent classes of drug use among adolescents through parental alcohol use and parental style: a longitudinal study. *Soc Psychiatry Psychiatr Epidemiol*. 2019;54:455-67.
- Escamilla I, Juan N, Benito A, Castellano-García F, Rodríguez-Ruiz F, Haro G. Substance addiction in adolescents: influence of parenting and personality traits. *Brain Sci*. 2024;14:449.
- Liu L, Meng W, Liu B. The mediating role of social support in the relationship between parenting styles and adolescent drug abuse identification. *Front Psychol*. 2021;12:802408.
- Miguel AQC, Madruga CS, Cogo-Moreira H, Yamauchi R, Simões V, Da Silva CJ, et al. Sociodemographic characteristics, patterns of crack use, concomitant substance use disorders, and psychiatric

- symptomatology in treatment-seeking crack-dependent individuals in Brazil. *J Psychoactive Drugs*. 2018;50:367-72.
- 39 von Diemen L, Bassani DG, Fuchs SC, Szobot CM, Pechansky F. Impulsivity, age of first alcohol use and substance use disorders among male adolescents: a population based case-control study. *Addiction*. 2008;103:1198-205.
 - 40 Vonmoos M, Hulka LM, Preller KH, Jenni D, Schulz C, Baumgartner MR, et al. Differences in self-reported and behavioral measures of impulsivity in recreational and dependent cocaine users. *Drug Alcohol Depend*. 2013;133:61-70.
 - 41 MacKillop J, Amlung MT, Few LR, Ray LA, Sweet LH, Munafò MR. Delayed reward discounting and addictive behavior: a meta-analysis. *Psychopharmacology (Berl)*. 2011;216:305-21.
 - 42 Perry JL, Carroll ME. The role of impulsive behavior in drug abuse. *Psychopharmacology (Berl)*. 2008;200:1-26.