

# Math Refresher for DS

Lecture 2



# Last Time

- Vector spaces
- Euclidian spaces (= vector spaces + dot product)
- Length of a vector
- Distances and angles between the vectors
- Orthogonality
- Orthogonal projections

# Today

- Back to vector spaces
  - Linear independence
  - Basis
- Basic operations with matrices.



# **Back to Vector Spaces**



# (Reminder) Vector Space: Definition

- A real-valued vector space  $(V, +, \cdot)$  is a set of vectors  $V$  with two operations
$$(1) +: V \times V \rightarrow V, \quad (2) \cdot: \mathbb{R} \times V \rightarrow V$$

that satisfy the following properties (axioms):

	Property	Meaning
1.	<b>Associativity</b> of addition	$x + (y + z) = (x + y) + z$
2.	<b>Commutativity</b> of addition	$x + y = y + x$
3.	<b>Identity element</b> of addition	$\exists 0 \in V: \forall x \in V \quad 0 + x = x$
4.	<b>Identity element</b> of scalar multiplication	$\forall x \in V \quad 1 \cdot x = x$
5.	<b>Inverse element</b> of addition	$\forall x \in V \quad \exists -x \in V: \quad x + (-x) = 0$
6.	<b>Compatibility</b> of scalar multiplication	$\alpha(\beta x) = (\alpha\beta)x$
7.	<b>Distributivity</b>	$(\alpha + \beta)x = \alpha x + \beta x$
8.		$\alpha(x + y) = \alpha x + \alpha y$

# (Reminder) Examples of Vector Spaces

- $\mathbb{R}^n$  - a set of vectors with  $n$  real entries.  
 $(\mathbb{R}^n, +, \cdot)$  is a vector space.

# (Reminder) Examples of Vector Spaces

- $\mathbb{R}^n$  - a set of vectors with  $n$  real entries.  
 $(\mathbb{R}^n, +, \cdot)$  is a vector space.
- $\mathbb{P}^n$  - a set of polynomials of degree  $\leq n$  with real coefficients  
 $(\mathbb{P}^n, +, \cdot)$  is also a vector space!  
“Vectors” here are polynomials.

# **Vector Subspaces**



# Vector Subspace

- $V = (\mathbb{V}, +, \cdot)$  - a vector space.

# Vector Subspace

- $V = (\mathbb{V}, +, \cdot)$  - a vector space.
- Consider  $\mathbb{U} \neq \emptyset$  – a subset of  $\mathbb{V}$  ( $\mathbb{U} \subseteq \mathbb{V}$ ).

# Vector Subspace

- $V = (\mathbb{V}, +, \cdot)$  - a vector space.
- Consider  $\mathbb{U} \neq \emptyset$  – a subset of  $\mathbb{V}$  ( $\mathbb{U} \subseteq \mathbb{V}$ ).
- $U = (\mathbb{U}, +, \cdot)$  - a *vector subspace* ( $U \subseteq V$ ) if  $U$  is a vector space with operations
  - $+ : \mathbb{U} \times \mathbb{U} \rightarrow \mathbb{U}$
  - $\cdot : \mathbb{R} \times \mathbb{U} \rightarrow \mathbb{U}$

# Vector Subspace

- How do we check if  $U = (\mathbb{U}, +, \cdot)$  is a vector space?

# Vector Subspace

- How do we check if  $U = (\mathbb{U}, +, \cdot)$  is a vector space?
- Since  $\mathbb{U} \subseteq \mathbb{V}$  and  $V = (\mathbb{V}, +, \cdot)$  is a vector space, many properties of  $+$  and  $\cdot$  hold automatically.

# Vector Subspace

- How do we check if  $U = (\mathbb{U}, +, \cdot)$  is a vector space?
- Since  $\mathbb{U} \subseteq \mathbb{V}$  and  $V = (\mathbb{V}, +, \cdot)$  is a vector space, many properties of  $+$  and  $\cdot$  hold automatically.
- In fact, we only need to check:
  1. that  $0 \in \mathbb{U}$


# Vector Subspace

- How do we check if  $U = (\mathbb{U}, +, \cdot)$  is a vector space?
- Since  $\mathbb{U} \subseteq \mathbb{V}$  and  $V = (\mathbb{V}, +, \cdot)$  is a vector space, many properties of  $+$  and  $\cdot$  hold automatically.
- In fact, we only need to check:
  1. that  $0 \in \mathbb{U}$
  2. closure of  $+$  and  $\cdot$ :
    - $\forall x, y \in \mathbb{U} \ x + y \in \mathbb{U}$
    - $\forall x \in \mathbb{U}, \lambda \in \mathbb{R} \ \lambda x \in \mathbb{U}$

# Vector Subspace: Examples




- Consider  $\mathbb{R}^2$ .
- Is  $A$  a vector subspace?



# Vector Subspace: Examples




- Consider  $\mathbb{R}^2$ .
- Is  $A$  a vector subspace?



# Vector Subspace: Examples




- Consider  $\mathbb{R}^2$ .
- Is  $A$  a vector subspace?



# Vector Subspace: Examples




- Consider  $\mathbb{R}^2$ .
- Is  $A$  a vector subspace?
- No!
  - $x \in A$  but  $2x \notin A \rightarrow$ 
    - operation isn't closed.



# Vector Subspace: Examples



- Consider  $\mathbb{R}^2$ .
- Is  $B$  a vector subspace?




# Vector Subspace: Examples



- Consider  $\mathbb{R}^2$ .
- Is  $B$  a vector subspace?
- No!


$0 \notin B$



# Vector Subspace: Examples




- Consider  $\mathbb{R}^2$ .
- Is  $D$  a vector subspace?



# Vector Subspace: Examples



- Consider  $\mathbb{R}^2$ .
- Is  $D$  a vector subspace?
- Yes!




# Vector Subspace: Examples



- Consider  $\mathbb{R}^2$ .
- Is  $D$  a vector subspace?
- Yes!

$\{0, +, \cdot\}$  is a trivial vector  
subspace of any vector space



# Vector Subspace: Examples

- Consider  $P^n = (\mathbb{P}^n, +, \cdot)$  - a vector space.

# Vector Subspace: Examples

- Consider  $P^n = (\mathbb{P}^n, +, \cdot)$  - a vector space.
- $\forall m \leq n \ P^m = (\mathbb{P}^m, +, \cdot) \subseteq P^n$  - a vector subspace:

# Vector Subspace: Examples

- Consider  $P^n = (\mathbb{P}^n, +, \cdot)$  - a vector space.
- $\forall m \leq n \ P^m = (\mathbb{P}^m, +, \cdot) \subseteq P^n$  - a vector subspace:
  - $0 \in \mathbb{P}^m$

# Vector Subspace: Examples

- Consider  $P^n = (\mathbb{P}^n, +, \cdot)$  - a vector space.
- $\forall m \leq n \ P^m = (\mathbb{P}^m, +, \cdot) \subseteq P^n$  - a vector subspace:
  - $0 \in \mathbb{P}^m$
  - Closure: when we add up polynomials of degree  $m \leq n$  or multiply them by a scalar, we always get a polynomial of of degree  $m \leq n$ .

# **Linear Combinations**



# Linear Combinations

- We can add vectors together and multiply them by scalars.  
Let's combine these two operations!

# Linear Combinations

- We can add vectors together and multiply them by scalars.  
Let's combine these two operations!
- $V$  – a vector space.

# Linear Combinations

- We can add vectors together and multiply them by scalars.  
Let's combine these two operations!
- $V$  – a vector space.
- $x_1, x_2, \dots, x_k \in V$  – some vectors.

# Linear Combinations

- We can add vectors together and multiply them by scalars.  
Let's combine these two operations!
- $V$  – a vector space.
- $x_1, x_2, \dots, x_k \in V$  – some vectors.
- $\lambda_1, \lambda_2, \dots, \lambda_k \in \mathbb{R}$  - some scalars.

# Linear Combinations

- We can add vectors together and multiply them by scalars.  
Let's combine these two operations!
- $V$  – a vector space.
- $x_1, x_2, \dots, x_k \in V$  – some vectors.
- $\lambda_1, \lambda_2, \dots, \lambda_k \in \mathbb{R}$  - some scalars.

$$v = \lambda_1 x_1 + \lambda_2 x_2 + \cdots + \lambda_k x_k = \sum_{i=1}^k \lambda_i x_i \in V -$$

a *linear combination* of  $x_1, x_2, \dots, x_k$ .

# Linear Combinations: Examples

- In  $(\mathbb{R}^2, +, \cdot)$ , consider vectors  $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$  and  $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ .

# Linear Combinations: Examples

- In  $(\mathbb{R}^2, +, \cdot)$ , consider vectors  $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$  and  $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ .

$u = 2e_1 - e_2 = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$  is a linear combination of  $e_1$  and  $e_2$ .

# Linear Combinations: Examples

- In  $(\mathbb{R}^2, +, \cdot)$ , consider vectors  $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$  and  $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ .  
 $u = 2e_1 - e_2 = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$  is a linear combination of  $e_1$  and  $e_2$ .
- In  $(\mathbb{P}^2, +, \cdot)$ , consider vectors  $e_0 = 1, e_1 = t, e_2 = t^2$ .

# Linear Combinations: Examples

- In  $(\mathbb{R}^2, +, \cdot)$ , consider vectors  $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$  and  $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ .

$u = 2e_1 - e_2 = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$  is a linear combination of  $e_1$  and  $e_2$ .

- In  $(\mathbb{P}^2, +, \cdot)$ , consider vectors  $e_0 = 1, e_1 = t, e_2 = t^2$ .

$u = 2t^2 - t + 1 = 2e_2 - e_1 + e_0$  is a linear combination of  $e_0, e_1$  and  $e_2$ .

# Linear Combinations: Examples

- In  $(\mathbb{R}^2, +, \cdot)$ , consider vectors  $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$  and  $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ .

$u = 2e_1 - e_2 = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$  is a linear combination of  $e_1$  and  $e_2$ .

- In  $(\mathbb{P}^2, +, \cdot)$ , consider vectors  $e_0 = 1, e_1 = t, e_2 = t^2$ .

$u = 2t^2 - t + 1 = 2e_2 - e_1 + e_0$  is a linear combination of  $e_0, e_1$  and  $e_2$ .

$v = 3t + 3 = 3e_1 + 3e_0$  is a linear combination of  $e_0$  and  $e_1$ .

# Span



# Span

- $V$  – a vector space,  $\mathbb{A} = \{x_1, x_2, \dots, x_k\} \subseteq V$  – a set of vectors.

# Span

- $V$  – a vector space,  $\mathbb{A} = \{x_1, x_2, \dots, x_k\} \subseteq V$  – a set of vectors.
- A set of all linear combinations of vectors in  $\mathbb{A}$  is called a *span* of  $\mathbb{A}$ :

# Span

- $V$  – a vector space,  $\mathbb{A} = \{x_1, x_2, \dots, x_k\} \subseteq V$  – a set of vectors.
- A set of all linear combinations of vectors in  $\mathbb{A}$  is called a *span* of  $\mathbb{A}$ :

$$span[x_1, \dots, x_n] = \{\lambda_1 x_1 + \dots + \lambda_n x_n | \lambda_1, \dots, \lambda_n \in \mathbb{R}\}$$

# Span

- $V$  – a vector space,  $\mathbb{A} = \{x_1, x_2, \dots, x_k\} \subseteq V$  – a set of vectors.
- A set of all linear combinations of vectors in  $\mathbb{A}$  is called a *span* of  $\mathbb{A}$ :


$$span[x_1, \dots, x_n] = \{\lambda_1 x_1 + \dots + \lambda_n x_n | \lambda_1, \dots, \lambda_n \in \mathbb{R}\}$$

- If  $\mathbb{A}$  spans vector space  $V$ , we write

$$V = span[\mathbb{A}] \text{ or } V = span[x_1, \dots, x_n].$$


# Span: Example 1

- Consider  $\mathbb{R}^2$ .




# Span: Example 1

- Consider  $\mathbb{R}^2$ .
- $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$




# Span: Example 1

- Consider  $\mathbb{R}^2$ .
- $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$   
 $span[e_1] = \left\{ \begin{bmatrix} \lambda \\ 0 \end{bmatrix} \mid \lambda \in \mathbb{R} \right\}$  - only vectors parallel to the first axis.




# Span: Example 1

- Consider  $\mathbb{R}^2$ .
- $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$   
 $span[e_1] = \left\{ \begin{bmatrix} \lambda \\ 0 \end{bmatrix} \mid \lambda \in \mathbb{R} \right\}$  - only vectors parallel to the first axis.
- $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$



# Span: Example 1

- Consider  $\mathbb{R}^2$ .
- $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$   
 $span[e_1] = \left\{ \begin{bmatrix} \lambda \\ 0 \end{bmatrix} \mid \lambda \in \mathbb{R} \right\}$  - only vectors parallel to the first axis.
- $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$   
 $span[e_1] = \left\{ \begin{bmatrix} \lambda_1 \\ \lambda_2 \end{bmatrix} \mid \lambda_{1,2} \in \mathbb{R} \right\} = \mathbb{R}^2$ .



# Span: Example 2

- Consider  $P^3 = (\mathbb{P}^3, +, \cdot)$ .

# Span: Example 2

- Consider  $P^3 = (\mathbb{P}^3, +, \cdot)$ .
- $e_0 = 1, e_1 = t, e_2 = t^2$

$$span[e_0, \dots, e_2] =$$

# Span: Example 2

- Consider  $P^3 = (\mathbb{P}^3, +, \cdot)$ .
- $e_0 = 1, e_1 = t, e_2 = t^2$

$$\text{span}[e_0, \dots, e_2] = \{at^2 + bt + c \mid a, b, c \in \mathbb{R}\} = P^2$$

# Span: Example 2

- Consider  $P^3 = (\mathbb{P}^3, +, \cdot)$ .

- $e_0 = 1, e_1 = t, e_2 = t^2$

$$span[e_0, \dots, e_2] = \{at^2 + bt + c \mid a, b, c \in \mathbb{R}\} = P^2 \subseteq P^3$$

# Span: Example 2

- Consider  $P^3 = (\mathbb{P}^3, +, \cdot)$ .

- $e_0 = 1, e_1 = t, e_2 = t^2$

$$\text{span}[e_0, \dots, e_2] = \{at^2 + bt + c \mid a, b, c \in \mathbb{R}\} = P^2 \subseteq P^3$$

- $e_0 = 1, e_1 = t, e_2 = t^2, e_3 = t^3$

$$\text{span}[e_0, \dots, e_3] =$$

# Span: Example 2

- Consider  $P^3 = (\mathbb{P}^3, +, \cdot)$ .

- $e_0 = 1, e_1 = t, e_2 = t^2$

$$\text{span}[e_0, \dots, e_2] = \{at^2 + bt + c \mid a, b, c \in \mathbb{R}\} = P^2 \subseteq P^3$$

- $e_0 = 1, e_1 = t, e_2 = t^2, e_3 = t^3$

$$\text{span}[e_0, \dots, e_3] = \{at^3 + bt^2 + ct + d \mid a, b, c, d \in \mathbb{R}\} = P^3$$

# Generating Set

- $V$  – a vector space.
- $\mathbb{A} = \{x_1, x_2, \dots, x_k\} \subseteq V$  – a set of vectors.

# Generating Set

- $V$  – a vector space.
- $\mathbb{A} = \{x_1, x_2, \dots, x_k\} \subseteq V$  – a set of vectors.
- If every vector  $v \in V$  can be expressed as a linear combination of  $x_1, x_2, \dots, x_k$ ,  $\mathbb{A}$  is called a *generating set* for  $V$ .

# Linear independence



# Linear Combinations

- A zero vector can always be represented as a trivial linear combination of  $x_1, x_2, \dots, x_k$ :

$$0 = \sum_{i=1}^k 0 \cdot x_i$$

# Linear Combinations

- A zero vector can always be represented as a trivial linear combination of  $x_1, x_2, \dots, x_k$ :

$$0 = \sum_{i=1}^k 0 \cdot x_i$$

- We are mostly interested in *non-trivial linear combinations* of  $x_1, x_2, \dots, x_k$  where not all  $\lambda_i$  are 0.

# Linear (In)dependence

- Consider a vector space  $V$ .
- $x_1, x_2, \dots, x_k \in V$  – some vectors.

# Linear (In)dependence

- Consider a vector space  $V$ .
- $x_1, x_2, \dots, x_k \in V$  – some vectors.
- If there is a non-trivial linear combination of  $x_1, x_2, \dots, x_k$  such that  $\sum_{i=1}^k \lambda_i x_i = 0$  with at least one  $\lambda_i \neq 0$ , vectors  $x_1, x_2, \dots, x_k$  are *linearly dependent*.

# Linear (In)dependence

- Consider a vector space  $V$ .
- $x_1, x_2, \dots, x_k \in V$  – some vectors.
- If there is a non-trivial linear combination of  $x_1, x_2, \dots, x_k$  such that  $\sum_{i=1}^k \lambda_i x_i = 0$  with at least one  $\lambda_i \neq 0$ , vectors  $x_1, x_2, \dots, x_k$  are *linearly dependent*.
- If only trivial solution exists, vectors  $x_1, x_2, \dots, x_k$  are *linearly independent*.

# Linear (In)dependence

- If there is a non-trivial linear combination of  $x_1, x_2, \dots, x_k$  such that  $\sum_{i=1}^k \lambda_i x_i = 0$  with at least one  $\lambda_i \neq 0$ , vectors  $x_1, x_2, \dots, x_k$  are *linearly dependent*.

# Linear (In)dependence

- If there is a non-trivial linear combination of  $x_1, x_2, \dots, x_k$  such that  $\sum_{i=1}^k \lambda_i x_i = 0$  with at least one  $\lambda_i \neq 0$ , vectors  $x_1, x_2, \dots, x_k$  are *linearly dependent*.

↔

- A set of vectors  $x_1, x_2, \dots, x_k$  is linearly dependent if and only if (at least) one of the vectors is a linear combination of the others

$$x_i = \alpha_1 x_1 + \cdots + \alpha_k x_k$$

# Linear (In)dependence

- If there is a non-trivial linear combination of  $x_1, x_2, \dots, x_k$  such that  $\sum_{i=1}^k \lambda_i x_i = 0$  with at least one  $\lambda_i \neq 0$ , vectors  $x_1, x_2, \dots, x_k$  are *linearly dependent*.

↔

- A set of vectors  $x_1, x_2, \dots, x_k$  is linearly dependent if and only if (at least) one of the vectors is a linear combination of the others

$$x_i = \alpha_1 x_1 + \cdots + \alpha_k x_k$$

**Why is this so? Try to prove this yourself.**

# Linear (In)dependence: Example 1

- Consider  $\mathbb{R}^2$ .

# Linear (In)dependence: Example 1

- Consider  $\mathbb{R}^2$ .
- Vectors  $u = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$  and  $v = \begin{bmatrix} -1 \\ -2 \end{bmatrix}$  are not linearly independent:  $u = -2v$ .

# Linear (In)dependence: Example 1

- Consider  $\mathbb{R}^2$ .
- Vectors  $u = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$  and  $v = \begin{bmatrix} -1 \\ -2 \end{bmatrix}$  are not linearly independent:  $u = -2v$ .
- Vectors  $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$  and  $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$  are linearly independent:  
there are no  $\lambda_1, \lambda_2 \in \mathbb{R}$  with at least one  $\lambda_i \neq 0$  such that

$$\lambda_1 e_1 + \lambda_2 e_2 = \begin{bmatrix} \lambda_1 \\ \lambda_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$

# Linear (In)dependence: Example 1

- Consider  $\mathbb{R}^2$ .
- Vectors  $u = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$  and  $v = \begin{bmatrix} -1 \\ -2 \end{bmatrix}$  are not linearly independent:  $u = -2v$ .

- Vectors  $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$  and  $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$  are linearly independent:  
there are no  $\lambda_1, \lambda_2 \in \mathbb{R}$  with at least one  $\lambda_i \neq 0$  such that

$$\lambda_1 e_1 + \lambda_2 e_2 = \begin{bmatrix} \lambda_1 \\ \lambda_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$

(Or: you cannot represent  $e_1$  as  $\lambda e_2$  or vice versa).

# Linear (In)dependence: Example 2

- Consider  $P = (\mathbb{P}^3, +, \cdot)$ .
- $1, t, t^2 \in P$  – vectors. Are they linearly independent?

# Linear (In)dependence: Example 2

- Consider  $P = (\mathbb{P}^3, +, \cdot)$ .
- $1, t, t^2 \in P$  – vectors. Are they linearly independent?
- Yes!

There is no way we can represent one of those vectors as a linear combination of the others.

# Linear (In)dependence: Example 3

- Are the following vectors in  $\mathbb{R}^4$  linearly independent?

$$x_1 = \begin{bmatrix} 1 \\ 2 \\ -3 \\ 4 \end{bmatrix}, \quad x_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \\ 2 \end{bmatrix}, \quad x_3 = \begin{bmatrix} -1 \\ -2 \\ 1 \\ 1 \end{bmatrix}$$

# Linear (In)dependence: Example 3

- Are the following vectors in  $\mathbb{R}^4$  linearly independent?

$$x_1 = \begin{bmatrix} 1 \\ 2 \\ -3 \\ 4 \end{bmatrix}, \quad x_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \\ 2 \end{bmatrix}, \quad x_3 = \begin{bmatrix} -1 \\ -2 \\ 1 \\ 1 \end{bmatrix}$$

- Are there  $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$  with at least one  $\lambda_i \neq 0$  such that

$$\lambda_1 x_1 + \lambda_2 x_2 + \lambda_3 x_3 = 0$$

# Linear (In)dependence: Example 3

- Are the following vectors in  $\mathbb{R}^4$  linearly independent?

$$x_1 = \begin{bmatrix} 1 \\ 2 \\ -3 \\ 4 \end{bmatrix}, \quad x_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \\ 2 \end{bmatrix}, \quad x_3 = \begin{bmatrix} -1 \\ -2 \\ 1 \\ 1 \end{bmatrix}$$

- Are there  $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$  with at least one  $\lambda_i \neq 0$  such that

$$\lambda_1 x_1 + \lambda_2 x_2 + \lambda_3 x_3 = 0 \leftrightarrow$$

$$\lambda_1 \begin{bmatrix} 1 \\ 2 \\ -3 \\ 4 \end{bmatrix} + \lambda_2 \begin{bmatrix} 1 \\ 1 \\ 0 \\ 2 \end{bmatrix} + \lambda_3 \begin{bmatrix} -1 \\ -2 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} \lambda_1 + \lambda_2 - \lambda_3 \\ 2\lambda_1 + \lambda_2 - 2\lambda_3 \\ -3\lambda_1 + \lambda_3 \\ 4\lambda_1 + 2\lambda_2 + \lambda_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

# Linear (In)dependence: Example 3

- Are there  $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$  with at least one  $\lambda_i \neq 0$  such that

$$\begin{cases} \lambda_1 + \lambda_2 - \lambda_3 = 0 \\ 2\lambda_1 + \lambda_2 - 2\lambda_3 = 0 \\ -3\lambda_1 + \lambda_3 = 0 \\ 4\lambda_1 + 2\lambda_2 + \lambda_3 = 0 \end{cases}$$

# Linear (In)dependence: Example 3

- Are there  $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$  with at least one  $\lambda_i \neq 0$  such that

$$\begin{cases} \lambda_1 + \lambda_2 - \lambda_3 = 0 \\ 2\lambda_1 + \lambda_2 - 2\lambda_3 = 0 \\ -3\lambda_1 + \lambda_3 = 0 \\ 4\lambda_1 + 2\lambda_2 + \lambda_3 = 0 \end{cases} \Leftrightarrow \begin{cases} \lambda_3 = 3\lambda_1 \end{cases}$$

# Linear (In)dependence: Example 3

- Are there  $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$  with at least one  $\lambda_i \neq 0$  such that

$$\begin{cases} \lambda_1 + \lambda_2 - \lambda_3 = 0 \\ 2\lambda_1 + \lambda_2 - 2\lambda_3 = 0 \\ -3\lambda_1 + \lambda_3 = 0 \\ 4\lambda_1 + 2\lambda_2 + \lambda_3 = 0 \end{cases} \Leftrightarrow \begin{cases} -2\lambda_1 + \lambda_2 = 0 \\ -4\lambda_1 + \lambda_2 = 0 \\ \lambda_3 = 3\lambda_1 \\ 7\lambda_1 + 2\lambda_2 = 0 \end{cases}$$

# Linear (In)dependence: Example 3

- Are there  $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$  with at least one  $\lambda_i \neq 0$  such that

$$\begin{cases} \lambda_1 + \lambda_2 - \lambda_3 = 0 \\ 2\lambda_1 + \lambda_2 - 2\lambda_3 = 0 \\ -3\lambda_1 + \lambda_3 = 0 \\ 4\lambda_1 + 2\lambda_2 + \lambda_3 = 0 \end{cases} \Leftrightarrow \begin{cases} -2\lambda_1 + \lambda_2 = 0 \\ -4\lambda_1 + \lambda_2 = 0 \\ \lambda_3 = 3\lambda_1 \\ 7\lambda_1 + 2\lambda_2 = 0 \end{cases} \Leftrightarrow \begin{cases} \lambda_2 = 2\lambda_1 \\ \lambda_3 = 3\lambda_1 \end{cases}$$

# Linear (In)dependence: Example 3

- Are there  $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$  with at least one  $\lambda_i \neq 0$  such that

$$\begin{cases} \lambda_1 + \lambda_2 - \lambda_3 = 0 \\ 2\lambda_1 + \lambda_2 - 2\lambda_3 = 0 \\ -3\lambda_1 + \lambda_3 = 0 \\ 4\lambda_1 + 2\lambda_2 + \lambda_3 = 0 \end{cases} \Leftrightarrow \begin{cases} -2\lambda_1 + \lambda_2 = 0 \\ -4\lambda_1 + \lambda_2 = 0 \\ \lambda_3 = 3\lambda_1 \\ 7\lambda_1 + 2\lambda_2 = 0 \end{cases} \Leftrightarrow \begin{cases} \lambda_2 = 2\lambda_1 \\ -2\lambda_1 = 0 \\ \lambda_3 = 3\lambda_1 \\ 11\lambda_1 = 0 \end{cases}$$

# Linear (In)dependence: Example 3

- Are there  $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$  with at least one  $\lambda_i \neq 0$  such that

$$\begin{cases} \lambda_1 + \lambda_2 - \lambda_3 = 0 \\ 2\lambda_1 + \lambda_2 - 2\lambda_3 = 0 \\ -3\lambda_1 + \lambda_3 = 0 \\ 4\lambda_1 + 2\lambda_2 + \lambda_3 = 0 \end{cases} \Leftrightarrow \begin{cases} -2\lambda_1 + \lambda_2 = 0 \\ -4\lambda_1 + \lambda_2 = 0 \\ \lambda_3 = 3\lambda_1 \\ 7\lambda_1 + 2\lambda_2 = 0 \end{cases} \Leftrightarrow \begin{cases} \lambda_2 = 2\lambda_1 \\ -2\lambda_1 = 0 \\ \lambda_3 = 3\lambda_1 \\ 11\lambda_1 = 0 \end{cases} \Leftrightarrow \lambda_{1,2,3} = 0$$

# Linear (In)dependence: Example 3

- Are there  $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$  with at least one  $\lambda_i \neq 0$  such that

$$\begin{cases} \lambda_1 + \lambda_2 - \lambda_3 = 0 \\ 2\lambda_1 + \lambda_2 - 2\lambda_3 = 0 \\ -3\lambda_1 + \lambda_3 = 0 \\ 4\lambda_1 + 2\lambda_2 + \lambda_3 = 0 \end{cases} \Leftrightarrow \begin{cases} -2\lambda_1 + \lambda_2 = 0 \\ -4\lambda_1 + \lambda_2 = 0 \\ \lambda_3 = 3\lambda_1 \\ 7\lambda_1 + 2\lambda_2 = 0 \end{cases} \Leftrightarrow \begin{cases} \lambda_2 = 2\lambda_1 \\ -2\lambda_1 = 0 \\ \lambda_3 = 3\lambda_1 \\ 11\lambda_1 = 0 \end{cases} \Leftrightarrow \lambda_{1,2,3} = 0$$

- Only a trivial linear combination of  $x_1, x_2$  and  $x_3$  gives a zero vector.

# Linear (In)dependence: Example 3

- Are there  $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$  with at least one  $\lambda_i \neq 0$  such that

$$\begin{cases} \lambda_1 + \lambda_2 - \lambda_3 = 0 \\ 2\lambda_1 + \lambda_2 - 2\lambda_3 = 0 \\ -3\lambda_1 + \lambda_3 = 0 \\ 4\lambda_1 + 2\lambda_2 + \lambda_3 = 0 \end{cases} \Leftrightarrow \begin{cases} -2\lambda_1 + \lambda_2 = 0 \\ -4\lambda_1 + \lambda_2 = 0 \\ \lambda_3 = 3\lambda_1 \\ 7\lambda_1 + 2\lambda_2 = 0 \end{cases} \Leftrightarrow \begin{cases} \lambda_2 = 2\lambda_1 \\ -2\lambda_1 = 0 \\ \lambda_3 = 3\lambda_1 \\ 11\lambda_1 = 0 \end{cases} \Leftrightarrow \lambda_{1,2,3} = 0$$

- Only a trivial linear combination of  $x_1, x_2$  and  $x_3$  gives a zero vector.

Therefore,  $x_1, x_2$  and  $x_3$  are linearly independent.

# Linear (In)dependence: Example 3

- Are there  $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$  with at least one  $\lambda_i \neq 0$  such that

$$\begin{cases} \lambda_1 + \lambda_2 - \lambda_3 = 0 \\ 2\lambda_1 + \lambda_2 - 2\lambda_3 = 0 \\ -3\lambda_1 + \lambda_3 = 0 \\ 4\lambda_1 + 2\lambda_2 + \lambda_3 = 0 \end{cases} \Leftrightarrow \begin{cases} -2\lambda_1 + \lambda_2 = 0 \\ -4\lambda_1 + \lambda_2 = 0 \\ \lambda_3 = 3\lambda_1 \\ 7\lambda_1 + 2\lambda_2 = 0 \end{cases} \Leftrightarrow \begin{cases} \lambda_2 = 2\lambda_1 \\ -2\lambda_1 = 0 \\ \lambda_3 = 3\lambda_1 \\ 11\lambda_1 = 0 \end{cases} \Leftrightarrow \lambda_{1,2,3} = 0$$

- Only a trivial linear combination of  $x_1, x_2$  and  $x_3$  gives a zero vector.

Therefore,  $x_1, x_2$  and  $x_3$  are linearly independent.

**Next lecture: a better way to solve  
such systems of equations**

# Dimension of a Linear Space

- The linear space  $V$  is called *n-dimensional* if the maximal number of linearly independent vectors in it is equal to  $n$ .

# Dimension of a Linear Space

- The linear space  $V$  is called *n-dimensional* if the maximal number of linearly independent vectors in it is equal to  $n$ .

We denote this as  $\dim(V) = n$ .

# Dimension: Example

- $\mathbb{R}^n$  is a  $n$ -dimensional vector space. Why?
- Consider  $n$  vectors  $e_1, \dots, e_n$ :

$$e_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \quad \dots, \quad e_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix}$$

- $e_1, \dots, e_n$  are linearly independent  $\rightarrow \dim(\mathbb{R}^n) \geq n$ .
- Can there be more than  $n$  linearly independent vectors in  $\mathbb{R}^n$ ?

# Dimension: Example

- $\mathbb{R}^n$  is a  $n$ -dimensional vector space. Why?

# Dimension: Example

- $\mathbb{R}^n$  is a  $n$ -dimensional vector space. Why?
- Consider  $n$  vectors  $e_1, \dots, e_n$ :

$$e_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \quad \dots, \quad e_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix}$$

# Dimension: Example

- $\mathbb{R}^n$  is a  $n$ -dimensional vector space. Why?
- Consider  $n$  vectors  $e_1, \dots, e_n$ :

$$e_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \quad \dots, \quad e_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix}$$

- $e_1, \dots, e_n$  are linearly independent  $\rightarrow \dim(\mathbb{R}^n) \geq n$ .

# Dimension: Example

- Can there be more than  $n$  linearly independent vectors in  $\mathbb{R}^n$ ?

# Dimension: Example

- Can there be more than  $n$  linearly independent vectors in  $\mathbb{R}^n$ ?

No!

Explanation: next lecture.

# Dimension: Another Example

- What is the dimensionality of  $P^3 = (\mathbb{P}^3, +, \cdot)$ ?

# Dimension: Another Example

- What is the dimensionality of  $P^3 = (\mathbb{P}^3, +, \cdot)$ ?
- We can find 4 linearly independent vectors in  $P^3$ :

$$1, \quad t, \quad t^2, \quad t^3$$

# Dimension: Another Example

- What is the dimensionality of  $P^3 = (\mathbb{P}^3, +, \cdot)$ ?
- We can find 4 linearly independent vectors in  $P^3$ :

$$1, \quad t, \quad t^2, \quad t^3$$

- All the other vectors (polynomials) from  $P^3$  are linear combinations of those.

# Dimension: Another Example

- What is the dimensionality of  $P^3 = (\mathbb{P}^3, +, \cdot)$ ?
- We can find 4 linearly independent vectors in  $P^3$ :

$$1, \quad t, \quad t^2, \quad t^3$$

- All the other vectors (polynomials) from  $P^3$  are linear combinations of those.

$$\rightarrow \dim(P^3) = 4.$$

# Basis



# Basis

- A set of  $n$  linearly independent vectors  $e_1, e_2, \dots, e_n$  in an  $n$ -dimensional space  $V$  is called a *basis* for  $V$ .

# Basis

- A set of  $n$  linearly independent vectors  $e_1, e_2, \dots, e_n$  in an  $n$ -dimensional space  $V$  is called a *basis* for  $V$ .
- Basis is A set of vectors with which we can represent every vector in the vector space by adding them together and scaling them.

# Basis: Example



**girafe**  
ai

---

100

# Basis: Example

- $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$  is a basis for  $\mathbb{R}^2$ .

# Basis: Example

- $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$  is a basis for  $\mathbb{R}^2$ .
- $e_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \dots, e_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix}$  is a basis for  $\mathbb{R}^n$ .

# Basis: Example

- $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$  is a basis for  $\mathbb{R}^2$ .
- $e_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \dots, e_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix}$  is a basis for  $\mathbb{R}^n$ .
- $e_0 = 0, e_1 = t, \dots, e_n = t^n$  is a basis for  $P^n$ .

# Basis: Example

- Find the basis of a vector space spanned by vectors

$$x = \begin{bmatrix} 1 \\ -2 \\ 3 \end{bmatrix}, \quad y = \begin{bmatrix} 0 \\ -3 \\ 3 \end{bmatrix}, \quad z = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$$

# Basis: Example

- Find the basis of a vector space spanned by vectors

$$x = \begin{bmatrix} 1 \\ -2 \\ 3 \end{bmatrix}, \quad y = \begin{bmatrix} 0 \\ -3 \\ 3 \end{bmatrix}, \quad z = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$$

- Vectors  $x$  and  $y$  are linearly independent.

# Basis: Example

- Find the basis of a vector space spanned by vectors

$$x = \begin{bmatrix} 1 \\ -2 \\ 3 \end{bmatrix}, \quad y = \begin{bmatrix} 0 \\ -3 \\ 3 \end{bmatrix}, \quad z = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$$

- Vectors  $x$  and  $y$  are linearly independent.
- Vector  $z$  is a linear combination of  $x$  and  $y$ :  $z = x - y$ .

# Basis: Example

- Find the basis of a vector space spanned by vectors

$$x = \begin{bmatrix} 1 \\ -2 \\ 3 \end{bmatrix}, \quad y = \begin{bmatrix} 0 \\ -3 \\ 3 \end{bmatrix}, \quad z = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$$

- Vectors  $x$  and  $y$  are linearly independent.
- Vector  $z$  is a linear combination of  $x$  and  $y$ :  $z = x - y$ .
- Therefore,  $V = \text{span}[x, y, z] = \text{span}[x, y]$ .  
 $B = \{x, y\}$  - basis of  $V$ .

# Coordinates

- Let's fixe the order of the vectors in the basis:

$$e_1, e_2, \dots, e_n$$

# Coordinates

- Let's fixe the order of the vectors in the basis:

$$e_1, e_2, \dots, e_n$$

- Every vector  $v$  in a vector space can be represented as a linear combination of  $e_1, e_2, \dots, e_n$ :

$$v = a_1 e_1 + a_2 e_2 + \cdots + a_n e_n$$

# Coordinates

- Let's fixe the order of the vectors in the basis:

$$e_1, e_2, \dots, e_n$$

- Every vector  $v$  in a vector space can be represented as a linear combination of  $e_1, e_2, \dots, e_n$ :


$$v = a_1 e_1 + a_2 e_2 + \cdots + a_n e_n$$

- $a_1, a_2, \dots, a_n$  - *coordinates* of the vector  $v$  in the basis  $e_1, e_2, \dots, e_n$ .

# Coordinates: Example



- Consider  $\mathbb{R}^2$ .




# Coordinates: Example



- Consider  $\mathbb{R}^2$ .

- $x = \begin{bmatrix} 4 \\ 6 \end{bmatrix}$



# Coordinates: Example




- Consider  $\mathbb{R}^2$ .

- $x = \begin{bmatrix} 4 \\ 6 \end{bmatrix}$

- Canonical basis:

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$



# Coordinates: Example




- Consider  $\mathbb{R}^2$ .

- $x = \begin{bmatrix} 4 \\ 6 \end{bmatrix}$

- Canonical basis:

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$x = 4e_1 + 6e_2$$



# Orthogonal Basis

- A basis  $e_1, \dots, e_n$  where  $\forall i, j \ e_i \perp e_j$  is an *orthogonal basis*.

# Orthogonal Basis

- A basis  $e_1, \dots, e_n$  where  $\forall i, j \ e_i \perp e_j$  is an *orthogonal basis*.
- If, additionally,  $\forall i \ \|e_i\| = 1$ , then  $e_1, \dots, e_n$  is an *orthonormal basis*.

# Orthogonal Basis

- A basis  $e_1, \dots, e_n$  where  $\forall i, j \ e_i \perp e_j$  is an *orthogonal basis*.
- If, additionally,  $\forall i \ \|e_i\| = 1$ , then  $e_1, \dots, e_n$  is an *orthonormal basis*.

Example:

$$e_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \ e_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \ \dots, \ e_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix}$$

is an orthonormal basis of  $\mathbb{R}^n$ .

# Orthogonal Basis

- A basis  $e_1, \dots, e_n$  where  $\forall i, j \ e_i \perp e_j$  is an *orthogonal basis*.
- If, additionally,  $\forall i \ \|e_i\| = 1$ , then  $e_1, \dots, e_n$  is an *orthonormal basis*.

Example:

$$e_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \ e_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \ \dots, \ e_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix}$$
 is an orthonormal basis of  $\mathbb{R}^n$ .

- *Gram-Schmidt process*: a way to convert any basis to an orthogonal one. More details: practical session.

# **Change of Basis**



# Basis

- A vector space has more than one basis.

# Basis

- A vector space has more than one basis.
- Example:  $\mathbb{R}^2$ 
  - $e = \left\{ e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$  - canonical basis;

# Basis

- A vector space has more than one basis.
- Example:  $\mathbb{R}^2$ 
  - $e = \left\{ e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$  - canonical basis;
  - $a = \left\{ a_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, a_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right\}$  - another basis;

# Basis

- A vector space has more than one basis.
- Example:  $\mathbb{R}^2$ 
  - $e = \left\{ e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$  - canonical basis;
  - $a = \left\{ a_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, a_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right\}$  - another basis;
  - $b = \left\{ b_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, b_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix} \right\}$  - yet another one.

# Basis

- A vector space has more than one basis.
- Example:  $\mathbb{R}^2$ 
  - $e = \left\{ e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$  - canonical basis;
  - $a = \left\{ a_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, a_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right\}$  - another basis;
  - $b = \left\{ b_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, b_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix} \right\}$  - yet another one.
- Different basis = different coordinates.  
How exactly do they change?



# Coordinate Change: Example

- Consider  $\mathbb{R}^2$  with canonical basis

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

# Coordinate Change: Example

- Consider  $\mathbb{R}^2$  with canonical basis

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

- New basis:

$$e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \quad e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$$

# Coordinate Change: Example

- Consider  $\mathbb{R}^2$  with canonical basis

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

- New basis:

$$e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \quad e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$$

- $x_{old} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$

# Coordinate Change: Example

- Consider  $\mathbb{R}^2$  with canonical basis

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

- New basis:

$$e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \quad e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$$

- $x_{old} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$

- What are the coordinates in the new basis?

$$x_{new} = ?$$

# Coordinate Change: General Case

- Consider a vector space  $V$  with basis  $e_1, e_2, \dots e_n$ .

# Coordinate Change: General Case

- Consider a vector space  $V$  with basis  $e_1, e_2, \dots, e_n$ .
- Imagine vector  $x = [x_1, x_2, \dots, x_n] \in V$   
 $x_1, x_2, \dots, x_n$  - coordinates in basis  $e_1, e_2, \dots, e_n$ .

# Coordinate Change: General Case

- Consider a vector space  $V$  with basis  $e_1, e_2, \dots, e_n$ .
- Imagine vector  $x = [x_1, x_2, \dots, x_n] \in V$   
 $x_1, x_2, \dots, x_n$  - coordinates in basis  $e_1, e_2, \dots, e_n$ .
- New basis:  $e'_1, e'_2, \dots, e'_n$ .

# Coordinate Change: General Case

- Consider a vector space  $V$  with basis  $e_1, e_2, \dots, e_n$ .
- Imagine vector  $x = [x_1, x_2, \dots, x_n] \in V$   
 $x_1, x_2, \dots, x_n$  - coordinates in basis  $e_1, e_2, \dots, e_n$ .
- New basis:  $e'_1, e'_2, \dots, e'_n$ .
- What are the coordinates of  $x$  in this new basis?

$$x'_1, x'_2, \dots, x'_n = ?$$

# Coordinate Change

- Old basis:  $e_1, e_2, \dots, e_n$   
New basis:  $e'_1, e'_2, \dots, e'_n$
- $x_{old} = [x_1, x_2, \dots, x_n], \quad x_{new} = [x'_1, x'_2, \dots, x'_n] = ?$
- Coordinates of the new basis in the old one:

# Coordinate Change

- Old basis:  $e_1, e_2, \dots, e_n$   
New basis:  $e'_1, e'_2, \dots, e'_n$
- $x_{old} = [x_1, x_2, \dots, x_n], \quad x_{new} = [x'_1, x'_2, \dots, x'_n] = ?$
- Coordinates of the new basis in the old one:

$$e'_1 = \alpha_{11}e_1 + \alpha_{21}e_2 + \cdots + \alpha_{n1}e_n$$

$$e'_2 = \alpha_{12}e_1 + \alpha_{22}e_2 + \cdots + \alpha_{n2}e_n$$

⋮

$$e'_i = \alpha_{1i}e_1 + \alpha_{2i}e_2 + \cdots + \alpha_{ni}e_n$$

⋮

$$e'_n = \alpha_{1n}e_1 + \alpha_{2n}e_2 + \cdots + \alpha_{nn}e_n$$

# Coordinate Change

$$x = x_1 e_1 + x_2 e_2 + \cdots + x_n e_n = x'_1 e'_1 + x'_2 e'_2 + \cdots + x'_n e'_n =$$

# Coordinate Change

$$x = x_1 e_1 + x_2 e_2 + \cdots + x_n e_n = x'_1 e'_1 + x'_2 e'_2 + \cdots + x'_n e'_n =$$

Remember:  $e'_i = \alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n$

# Coordinate Change

$$x = x_1 e_1 + x_2 e_2 + \cdots + x_n e_n = x'_1 e'_1 + x'_2 e'_2 + \cdots + x'_n e'_n =$$

Remember:  $e'_i = \alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n$

$$\begin{aligned} &= x'_1 \cdot (\alpha_{11} e_1 + \alpha_{21} e_2 + \cdots + \alpha_{n1} e_n) + \cdots + x'_i \cdot (\alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n) + \cdots \\ &\quad + x'_n (\alpha_{1n} e_1 + \alpha_{2n} e_2 + \cdots + \alpha_{nn} e_n) = \end{aligned}$$

# Coordinate Change

$$x = x_1 \mathbf{e}_1 + x_2 \mathbf{e}_2 + \cdots + x_n \mathbf{e}_n = x'_1 \mathbf{e}'_1 + x'_2 \mathbf{e}'_2 + \cdots + x'_n \mathbf{e}'_n =$$

Remember:  $e'_i = \alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n$

$$\begin{aligned} &= x'_1 \cdot (\alpha_{11} \mathbf{e}_1 + \alpha_{21} \mathbf{e}_2 + \cdots + \alpha_{n1} \mathbf{e}_n) + \cdots + x'_i \cdot (\alpha_{1i} \mathbf{e}_1 + \alpha_{2i} \mathbf{e}_2 + \cdots + \alpha_{ni} \mathbf{e}_n) + \cdots \\ &\quad + x'_n (\alpha_{1n} \mathbf{e}_1 + \alpha_{2n} \mathbf{e}_2 + \cdots + \alpha_{nn} \mathbf{e}_n) = \end{aligned}$$

$e_1, \dots, e_n$  linearly independent  $\rightarrow$  coefficients in front of them should be the same on the both sides of the equality:

# Coordinate Change

$$x = x_1 \mathbf{e}_1 + x_2 \mathbf{e}_2 + \cdots + x_n \mathbf{e}_n = x'_1 \mathbf{e}'_1 + x'_2 \mathbf{e}'_2 + \cdots + x'_n \mathbf{e}'_n =$$

Remember:  $e'_i = \alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n$

$$\begin{aligned} &= x'_1 \cdot (\alpha_{11} \mathbf{e}_1 + \alpha_{21} \mathbf{e}_2 + \cdots + \alpha_{n1} \mathbf{e}_n) + \cdots + x'_i \cdot (\alpha_{1i} \mathbf{e}_1 + \alpha_{2i} \mathbf{e}_2 + \cdots + \alpha_{ni} \mathbf{e}_n) + \cdots \\ &\quad + x'_n (\alpha_{1n} \mathbf{e}_1 + \alpha_{2n} \mathbf{e}_2 + \cdots + \alpha_{nn} \mathbf{e}_n) = \end{aligned}$$

$e_1, \dots, e_n$  linearly independent  $\rightarrow$  coefficients in front of them should be the same on the both sides of the equality:

$$x_1 = x'_1 \alpha_{11} + \cdots + x'_i \alpha_{1i} + \cdots + x'_n \alpha_{1n}$$

$$x_2 = x'_1 \alpha_{21} + \cdots + x'_i \alpha_{2i} + \cdots + x'_n \alpha_{2n}$$

⋮

$$x_n = x'_1 \alpha_{n1} + \cdots + x'_i \alpha_{ni} + \cdots + x'_n \alpha_{nn}$$

# Coordinate Change

$$x = x_1 \mathbf{e}_1 + x_2 \mathbf{e}_2 + \cdots + x_n \mathbf{e}_n = x'_1 \mathbf{e}'_1 + x'_2 \mathbf{e}'_2 + \cdots + x'_n \mathbf{e}'_n =$$

Remember:  $e'_i = \alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n$

$$\begin{aligned} &= x'_1 \cdot (\alpha_{11} \mathbf{e}_1 + \alpha_{21} \mathbf{e}_2 + \cdots + \alpha_{n1} \mathbf{e}_n) + \cdots + x'_i \cdot (\alpha_{1i} \mathbf{e}_1 + \alpha_{2i} \mathbf{e}_2 + \cdots + \alpha_{ni} \mathbf{e}_n) + \cdots \\ &\quad + x'_n (\alpha_{1n} \mathbf{e}_1 + \alpha_{2n} \mathbf{e}_2 + \cdots + \alpha_{nn} \mathbf{e}_n) = \end{aligned}$$

$x_{old}$

$$\boxed{\begin{aligned} x_1 &= x'_1 \alpha_{11} + \cdots + x'_i \alpha_{1i} + \cdots + x'_n \alpha_{1n} \\ x_2 &= x'_1 \alpha_{21} + \cdots + x'_i \alpha_{2i} + \cdots + x'_n \alpha_{2n} \\ &\vdots \\ x_n &= x'_1 \alpha_{n1} + \cdots + x'_i \alpha_{ni} + \cdots + x'_n \alpha_{nn} \end{aligned}}$$

# Coordinate Change

$$x = x_1 \mathbf{e}_1 + x_2 \mathbf{e}_2 + \cdots + x_n \mathbf{e}_n = x'_1 \mathbf{e}'_1 + x'_2 \mathbf{e}'_2 + \cdots + x'_n \mathbf{e}'_n =$$

Remember:  $e'_i = \alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n$

$$\begin{aligned} &= x'_1 \cdot (\alpha_{11} \mathbf{e}_1 + \alpha_{21} \mathbf{e}_2 + \cdots + \alpha_{n1} \mathbf{e}_n) + \cdots + x'_i \cdot (\alpha_{1i} \mathbf{e}_1 + \alpha_{2i} \mathbf{e}_2 + \cdots + \alpha_{ni} \mathbf{e}_n) + \cdots \\ &\quad + x'_n (\alpha_{1n} \mathbf{e}_1 + \alpha_{2n} \mathbf{e}_2 + \cdots + \alpha_{nn} \mathbf{e}_n) = \end{aligned}$$

$x_{old}$

$$\begin{aligned} x_1 &= x'_1 \alpha_{11} + \cdots + x'_i \alpha_{1i} + \cdots + x'_n \alpha_{1n} \\ x_2 &= x'_1 \alpha_{21} + \cdots + x'_i \alpha_{2i} + \cdots + x'_n \alpha_{2n} \\ &\vdots \\ x_n &= x'_1 \alpha_{n1} + \cdots + x'_i \alpha_{ni} + \cdots + x'_n \alpha_{nn} \end{aligned}$$


$x_{new}$

# Coordinate Change

$$x = x_1 \mathbf{e}_1 + x_2 \mathbf{e}_2 + \cdots + x_n \mathbf{e}_n = x'_1 \mathbf{e}'_1 + x'_2 \mathbf{e}'_2 + \cdots + x'_n \mathbf{e}'_n =$$

Remember:  $e'_i = \alpha_{1i}e_1 + \alpha_{2i}e_2 + \cdots + \alpha_{ni}e_n$

$$\begin{aligned} &= x'_1 \cdot (\alpha_{11}e_1 + \alpha_{21}e_2 + \cdots + \alpha_{n1}e_n) + \cdots + x'_i \cdot (\alpha_{1i}e_1 + \alpha_{2i}e_2 + \cdots + \alpha_{ni}e_n) + \cdots \\ &\quad + x'_n (\alpha_{1n}e_1 + \alpha_{2n}e_2 + \cdots + \alpha_{nn}e_n) = \end{aligned}$$




# Coordinate Change: Example

- Consider  $\mathbb{R}^2$  with basis  $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ ,  $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$
- New basis:  $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ ,  $e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$
- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$ ,  $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix}$


# Coordinate Change: Example

- Consider  $\mathbb{R}^2$  with basis  $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ ,  $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$
- New basis:  $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ ,  $e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$
- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$ ,  $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix}$



# Coordinate Change: Example


- Consider  $\mathbb{R}^2$  with basis  $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ ,  $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$
- New basis:  $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ ,  $e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$
- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$ ,  $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix}$



$$\begin{aligned}2 &= 2x'_1 - 1x'_2 \\-1 &= 1x'_1 - 1x'_2\end{aligned}$$

# Coordinate Change: Example

- Consider  $\mathbb{R}^2$  with basis  $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ ,  $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$
- New basis:  $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ ,  $e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$
- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$ ,  $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix}$




$$\begin{aligned}2 &= 2x'_1 - 1x'_2 \\-1 &= 1x'_1 - 1x'_2\end{aligned} \Leftrightarrow \begin{aligned}x'_1 &= 3 \\x'_2 &= 4\end{aligned}$$

# Coordinate Change: Example

- Consider  $\mathbb{R}^2$  with basis  $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ ,  $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$

- New basis:  $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ ,  $e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$

- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$ ,  $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix}$



$$\begin{aligned} 2 &= 2x'_1 - 1x'_2 \iff x'_1 = 3 \\ -1 &= 1x'_1 - 1x'_2 \iff x'_2 = 4 \iff x_{new} = \begin{bmatrix} 3 \\ 4 \end{bmatrix} \end{aligned}$$

# Coordinate Change

- Going from one basis to the other:

$$\begin{aligned} \mathbf{x}_{old} &= \boxed{x_1 = x'_1 \alpha_{11} + \dots + x'_i \alpha_{1i} + \dots + x'_n \alpha_{1n}} \\ &\quad \vdots \\ &= \boxed{x_2 = x'_1 \alpha_{21} + \dots + x'_i \alpha_{2i} + \dots + x'_n \alpha_{2n}} \\ &= \boxed{x_n = x'_1 \alpha_{n1} + \dots + x'_i \alpha_{ni} + \dots + x'_n \alpha_{nn}} \end{aligned}$$

$e'_i$

$\mathbf{x}_{new}$

```
graph TD; subgraph Old_Basis [Old Basis]; x1[x1] --- C1[ ]; x2[x2] --- C2[ ]; xn[xn] --- Cn[ ]; end; subgraph New_Basis [New Basis]; x1_prime[x1'] --- C1_prime[ ]; x2_prime[x2'] --- C2_prime[ ]; xn_prime[xn'] --- Cn_prime[ ]; end; C1_prime --> ei_prime[e'i]; C2_prime --> xn_new[xnew]; Cn_prime --> xn_prime;
```

# Coordinate Change

- Going from one basis to the other:

$$\begin{aligned} \mathbf{x}_{\text{old}} &= \boxed{x_1 = x'_1 \alpha_{11} + \dots + x'_i \alpha_{1i} + \dots + x'_n \alpha_{1n}} \\ &\quad \vdots \\ &= \boxed{x_2 = x'_1 \alpha_{21} + \dots + x'_i \alpha_{2i} + \dots + x'_n \alpha_{2n}} \\ &= \boxed{x_n = x'_1 \alpha_{n1} + \dots + x'_i \alpha_{ni} + \dots + x'_n \alpha_{nn}} \end{aligned}$$

$e'_i$

$\mathbf{x}_{\text{new}}$

- There is a more compact way of writing this down using **matrices**.

# Matrices



# A Matrix

- $A \in \mathbb{R}^{m \times n}$  - a matrix with  $m$  rows and  $n$  columns:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

# A Matrix

- $A \in \mathbb{R}^{m \times n}$  - a matrix with  $m$  rows and  $n$  columns:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

- Examples:

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \in \mathbb{R}^{2 \times 2}, \quad \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \in \mathbb{R}^{2 \times 3}$$

# Special Matrices

- Diagonal matrix:  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$  ( $a_{ii} \neq 0, a_{ij} = 0 \forall i \neq j$ )

# Special Matrices

- Diagonal matrix:  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$  ( $a_{ii} \neq 0, a_{ij} = 0 \forall i \neq j$ )
- Identity matrix:  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$  ( $a_{ii} = 1, a_{ij} = 0 \forall i \neq j$ )

# Special Matrices

- Diagonal matrix:  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$  ( $a_{ii} \neq 0, a_{ij} = 0 \forall i \neq j$ )
- Identity matrix:  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$  ( $a_{ii} = 1, a_{ij} = 0 \forall i \neq j$ )
- Symmetric matrix:  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$  ( $a_{ij} = a_{ji}$ )

# Special Matrices

- Diagonal matrix:  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$  ( $a_{ii} \neq 0, a_{ij} = 0 \forall i \neq j$ )
- Identity matrix:  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$  ( $a_{ii} = 1, a_{ij} = 0 \forall i \neq j$ )
- Symmetric matrix:  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$  ( $a_{ij} = a_{ji}$ )
- Triangular matrix:  $\begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 5 \\ 0 & 0 & 6 \end{bmatrix}, \begin{bmatrix} 1 & 0 & 0 \\ 2 & 3 & 0 \\ 4 & 5 & 6 \end{bmatrix}$  ( $a_{ij} = 0 \forall i > j \text{ or } \forall i < j$ )

# Vectors vs Matrices

- An  $n$ -dimensional vector can be considered a  $n \times 1$  matrix:

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \in \mathbb{R}^{n \times 1}$$

# **Operations with Matrices**



# Transpose of a Matrix

- Consider a matrix

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

# Transpose of a Matrix

- Consider a matrix

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

- Transpose = writing columns as rows:

$$A^T = \begin{bmatrix} a_{11} & a_{21} & \cdots & a_{m1} \\ a_{12} & a_{22} & \cdots & a_{m2} \\ \cdots & \cdots & \cdots & \cdots \\ a_{1n} & a_{2n} & \cdots & a_{mn} \end{bmatrix}, \quad \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}^T = [x_1, \dots, x_n]$$

# Transpose of a Matrix: Example

$$\bullet \begin{bmatrix} 0 & 1 & 2 \\ 2 & 1 & 5 \end{bmatrix}^T = \begin{bmatrix} 0 & 2 \\ 1 & 1 \\ 2 & 5 \end{bmatrix}$$

# Transpose of a Matrix: Example

- $\begin{bmatrix} 0 & 1 & 2 \\ 2 & 1 & 5 \end{bmatrix}^T = \begin{bmatrix} 0 & 2 \\ 1 & 1 \\ 2 & 5 \end{bmatrix}$
- Transposing a symmetrical matrix = no changes:

$$\begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & 2 \\ 3 & 2 & 1 \end{bmatrix}^T = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & 2 \\ 3 & 2 & 1 \end{bmatrix}$$

# Multiplying by a Scalar

- We can multiply matrix by a scalar:

$$\lambda A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} = \begin{bmatrix} \lambda a_{11} & \lambda a_{12} & \cdots & \lambda a_{1n} \\ \lambda a_{21} & \lambda a_{22} & \cdots & \lambda a_{2n} \\ \dots & \dots & \dots & \dots \\ \lambda a_{m1} & \lambda a_{m2} & \cdots & \lambda a_{mn} \end{bmatrix}$$

# Multiplying by a Scalar

- We can multiply matrix by a scalar:

$$\lambda A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} = \begin{bmatrix} \lambda a_{11} & \lambda a_{12} & \cdots & \lambda a_{1n} \\ \lambda a_{21} & \lambda a_{22} & \cdots & \lambda a_{2n} \\ \dots & \dots & \dots & \dots \\ \lambda a_{m1} & \lambda a_{m2} & \cdots & \lambda a_{mn} \end{bmatrix}$$

- Example:

$$5 \cdot \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 5 & 0 \\ 0 & 5 \end{bmatrix}$$

# Sum of Two Matrices

- We can sum up matrices of the same size:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}, \quad B = \begin{bmatrix} b_{11} & b_{12} & \cdots & b_{1n} \\ b_{21} & b_{22} & \cdots & b_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ b_{m1} & b_{m2} & \cdots & b_{mn} \end{bmatrix},$$

$$A + B = \begin{bmatrix} a_{11} + b_{11} & a_{12} + b_{12} & \cdots & a_{1n} + b_{1n} \\ a_{21} + b_{21} & a_{22} + b_{22} & \cdots & a_{2n} + b_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} + b_{m1} & a_{m2} + b_{m2} & \cdots & a_{mn} + b_{mn} \end{bmatrix}$$

# Sum of Two Matrices

- We can sum up matrices of the same size:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}, \quad B = \begin{bmatrix} b_{11} & b_{12} & \cdots & b_{1n} \\ b_{21} & b_{22} & \cdots & b_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ b_{m1} & b_{m2} & \cdots & b_{mn} \end{bmatrix},$$

$$A + B = \begin{bmatrix} a_{11} + b_{11} & a_{12} + b_{12} & \cdots & a_{1n} + b_{1n} \\ a_{21} + b_{21} & a_{22} + b_{22} & \cdots & a_{2n} + b_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} + b_{m1} & a_{m2} + b_{m2} & \cdots & a_{mn} + b_{mn} \end{bmatrix}$$

- Example:

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} -1 & 2 \\ 0 & 3 \end{bmatrix} = \begin{bmatrix} 0 & 2 \\ 0 & 4 \end{bmatrix}$$

# Matrices Also Form a Vector Space!

- $(\mathbb{R}^{m \times n}, +, \cdot)$  - a vector space.  
“Vectors” = matrices.

# Matrices Also Form a Vector Space!

- $(\mathbb{R}^{m \times n}, +, \cdot)$  - a vector space.  
“Vectors” = matrices.
- You can check yourself that the necessary axioms hold.

# Matrix Multiplication

- Consider two matrices  $A = \{a_{ij}\}_{m \times n}$  and  $b = \{b_{ij}\}_{n \times p}$ .
- $C = AB$  – product of two matrices.

# Matrix Multiplication

- Consider two matrices  $A = \{a_{ij}\}_{m \times n}$  and  $b = \{b_{ij}\}_{n \times p}$ .
- $C = AB$  – product of two matrices.
- Each element  $c_{ij}$  of  $C$  is a dot product of  $i$ -th row of  $A$  and  $j$ -th column of  $B$ :  $C = \{c_{ij}\}_{m \times p} = \{(A_i, B^j)\}_{m \times p}$

# Matrix Multiplication

- Consider two matrices  $A = \{a_{ij}\}_{m \times n}$  and  $b = \{b_{ij}\}_{n \times p}$ .
- $C = AB$  – product of two matrices.
- Each element  $c_{ij}$  of  $C$  is a dot product of  $i$ -th row of  $A$  and  $j$ -th column of  $B$ :  $C = \{c_{ij}\}_{m \times p} = \{(A_i, B^j)\}_{m \times p}$
- Example  $\mathbb{R}^{2 \times 2}$ : 
$$\begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \cdot \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix} = \begin{bmatrix} a_{11}b_{11} + a_{12}b_{21} & a_{11}b_{12} + a_{12}b_{22} \\ a_{21}b_{11} + a_{22}b_{21} & a_{21}b_{12} + a_{22}b_{22} \end{bmatrix}$$

# Matrix Multiplication: Example

$$\begin{bmatrix} 0 & 1 & 2 \\ 2 & 1 & 5 \end{bmatrix} \begin{bmatrix} 3 & 6 & 4 \\ 2 & 5 & 8 \\ 7 & 1 & 9 \end{bmatrix} =$$

# Matrix Multiplication: Example

$$\begin{bmatrix} 0 & 1 & 2 \\ 2 & 1 & 5 \end{bmatrix} \begin{bmatrix} 3 & 6 & 4 \\ 2 & 5 & 8 \\ 7 & 1 & 9 \end{bmatrix} =$$

$$= \begin{bmatrix} 0 + 2 + 14 & 0 + 5 + 2 & 0 + 8 + 18 \\ 6 + 2 + 35 & 12 + 5 + 5 & 8 + 8 + 45 \end{bmatrix} =$$

# Matrix Multiplication: Example

$$\begin{bmatrix} 0 & 1 & 2 \\ 2 & 1 & 5 \end{bmatrix} \begin{bmatrix} 3 & 6 & 4 \\ 2 & 5 & 8 \\ 7 & 1 & 9 \end{bmatrix} =$$

$$= \begin{bmatrix} 0 + 2 + 14 & 0 + 5 + 2 & 0 + 8 + 18 \\ 6 + 2 + 35 & 12 + 5 + 5 & 8 + 8 + 45 \end{bmatrix} =$$

$$= \begin{bmatrix} 16 & 7 & 26 \\ 43 & 22 & 61 \end{bmatrix}$$

# Coordinate Change: Matrix Notation




- Result obtained before:

$e_1, \dots, e_n$  - old basis

$e'_1, \dots, e'_n$  - new basis

$$x_{old} = [x_1, \dots, x_n], \quad x_{new} = [x'_1, \dots, x'_n]$$



# Coordinate Change: Matrix Notation




- Result obtained before:

$e_1, \dots, e_n$  - old basis

$e'_1, \dots, e'_n$  - new basis

$$x_{\text{old}} = [x_1, \dots, x_n], \quad x_{\text{new}} = [x'_1, \dots, x'_n]$$



- Transition matrix: columns = coordinates of the new basis in the old one.

$$A = \begin{bmatrix} \alpha_{11} & \alpha_{12} & \cdots & \alpha_{1n} \\ \alpha_{21} & \alpha_{22} & \cdots & \alpha_{21} \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_{n1} & \alpha_{n2} & \cdots & \alpha_{nn} \end{bmatrix}$$

# Coordinate Change: Matrix Notation




- Result obtained before:

$e_1, \dots, e_n$  - old basis

$e'_1, \dots, e'_n$  - new basis

$$x_{old} = [x_1, \dots, x_n], \quad x_{new} = [x'_1, \dots, x'_n]$$



- Transition matrix: columns = coordinates of the new basis in the old one.

$$A = \begin{bmatrix} \alpha_{11} & \alpha_{12} & \dots & \alpha_{1n} \\ \alpha_{21} & \alpha_{22} & \dots & \alpha_{21} \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_{n1} & \alpha_{n2} & \dots & \alpha_{nn} \end{bmatrix}$$

$$x_{old} = A^T x_{new}$$

# Coordinate Change: Example (again)

- Consider  $\mathbb{R}^2$  with basis  $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ ,  $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$
- New basis:  $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ ,  $e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$
- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$ ,  $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = ?$

# Coordinate Change: Example (again)

- Consider  $\mathbb{R}^2$  with basis  $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ ,  $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$
- New basis:  $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ ,  $e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$
- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$ ,  $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = ?$

$$A = \begin{bmatrix} 2 & -1 \\ 1 & -1 \end{bmatrix},$$

# Coordinate Change: Example (again)

- Consider  $\mathbb{R}^2$  with basis  $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ ,  $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$
- New basis:  $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ ,  $e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$
- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$ ,  $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = ?$

$$A = \begin{bmatrix} 2 & -1 \\ 1 & -1 \end{bmatrix}, \quad \begin{bmatrix} 2 \\ -1 \end{bmatrix} = x_{old} = A^T x_{new} =$$

# Coordinate Change: Example (again)

- Consider  $\mathbb{R}^2$  with basis  $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ ,  $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$
- New basis:  $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ ,  $e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$
- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$ ,  $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = ?$

$$A = \begin{bmatrix} 2 & -1 \\ 1 & -1 \end{bmatrix}, \quad \begin{bmatrix} 2 \\ -1 \end{bmatrix} = x_{old} = A^T x_{new} = \begin{bmatrix} 2 & 1 \\ -1 & -1 \end{bmatrix} \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} =$$

# Coordinate Change: Example (again)

- Consider  $\mathbb{R}^2$  with basis  $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ ,  $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$
- New basis:  $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ ,  $e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$
- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$ ,  $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = ?$

$$A = \begin{bmatrix} 2 & -1 \\ 1 & -1 \end{bmatrix}, \quad \begin{bmatrix} 2 \\ -1 \end{bmatrix} = x_{old} = A^T x_{new} = \begin{bmatrix} 2 & 1 \\ -1 & -1 \end{bmatrix} \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = \begin{bmatrix} 2x'_1 + x'_2 \\ -x'_1 - x'_2 \end{bmatrix}$$

# Coordinate Change: Example (again)

- Consider  $\mathbb{R}^2$  with basis  $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ ,  $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$
- New basis:  $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ ,  $e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$
- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$ ,  $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = ?$

$$A = \begin{bmatrix} 2 & -1 \\ 1 & -1 \end{bmatrix}, \quad \begin{bmatrix} 2 \\ -1 \end{bmatrix} = x_{old} = A^T x_{new} = \begin{bmatrix} 2 & 1 \\ -1 & -1 \end{bmatrix} \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = \begin{bmatrix} 2x'_1 + x'_2 \\ -x'_1 - x'_2 \end{bmatrix}$$

$$x_{new} = \begin{bmatrix} 3 \\ 4 \end{bmatrix}.$$

# To Sum Up

- Vector spaces
  - Linear (in)dependence
  - Span
  - Basis
- Matrices
  - Matrix operations
  - Change of coordinates

# Next Time

- More on matrices
- Systems of linear equations