




# Math Refresher for DS

Lecture 3



# Last Time

- Vector Spaces
  - Linear combinations
  - Spans
  - Bases
  - Change of coordinates
- Matrices

# Today

- More on matrices
  - matrix operations;
  - rank;
  - determinant.
- Linear transformations
- Systems of linear equations

# **Matrices: a small review**



# A Matrix

- $A \in \mathbb{R}^{m \times n}$  - a matrix with  $m$  rows and  $n$  columns:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

- Examples:

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \in \mathbb{R}^{2 \times 2}, \quad \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \in \mathbb{R}^{2 \times 3}$$

# Special Matrices

- Diagonal matrix:  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$  ( $a_{ii} \neq 0, a_{ij} = 0 \forall i \neq j$ )
- Identity matrix:  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$  ( $a_{ii} = 1, a_{ij} = 0 \forall i \neq j$ )
- Symmetric matrix:  $\begin{bmatrix} 1 & 4 & 5 \\ 4 & 2 & 6 \\ 5 & 6 & 3 \end{bmatrix}$  ( $a_{ij} = a_{ji}$ )
- Triangular matrix:  $\begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 5 \\ 0 & 0 & 6 \end{bmatrix}, \begin{bmatrix} 1 & 0 & 0 \\ 2 & 3 & 0 \\ 4 & 5 & 6 \end{bmatrix}$  ( $a_{ij} = 0 \forall i > j \text{ or } \forall i < j$ )

# Basic Operations with Matrices

- Addition:

$$A = \{a_{ij}\}_{i=1,\dots,m, j=1,\dots,n}, \quad B = \{b_{ij}\}_{i=1,\dots,m, j=1,\dots,n}, \quad A + B = \{a_{ij} + b_{ij}\}_{i=1,\dots,m, j=1,\dots,n}$$

# Basic Operations with Matrices

- Addition:

$$A = \{a_{ij}\}_{i=1,\dots,m, j=1,\dots,n}, \quad B = \{b_{ij}\}_{i=1,\dots,m, j=1,\dots,n}, \quad A + B = \{a_{ij} + b_{ij}\}_{i=1,\dots,m, j=1,\dots,n}$$

- Multiplication by a scalar:

$$A = \{a_{ij}\}_{i=1,\dots,m, j=1,\dots,n}, \quad \lambda \in \mathbb{R}, \quad \lambda A = \{\lambda a_{ij}\}_{i=1,\dots,m, j=1,\dots,n}$$

# Matrix Multiplication

- Matrix multiplication:

$$A = \{a_{ij}\}_{i=1, \dots, m, j=1, \dots, n}, \quad B = \{b_{ij}\}_{i=1, \dots, n, j=1, \dots, k}$$

$$A \cdot B = \{(A_i, B^j)\}_{i=1, \dots, m, j=1, \dots, k} = \left\{ \sum_{l=1, \dots, n} a_{il} \cdot b_{lj} \right\}_{i=1, \dots, m, j=1, \dots, k}$$

# Matrix Multiplication

- Matrix multiplication:

$$A = \{a_{ij}\}_{i=1, \dots, m, j=1, \dots, n}, \quad B = \{b_{ij}\}_{i=1, \dots, n, j=1, \dots, k}$$

$$A \cdot B = \{(A_i, B^j)\}_{i=1, \dots, m, j=1, \dots, k} = \left\{ \sum_{l=1, \dots, n} a_{il} \cdot b_{lj} \right\}_{i=1, \dots, m, j=1, \dots, k}$$

- Example  $\mathbb{R}^{2 \times 2}$ :

$$\begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \cdot \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix} = \begin{bmatrix} a_{11}b_{11} + a_{12}b_{21} & a_{11}b_{12} + a_{12}b_{22} \\ a_{21}b_{11} + a_{22}b_{21} & a_{21}b_{12} + a_{22}b_{22} \end{bmatrix}$$

# Matrix Multiplication

- For numbers:  $2 \times 3 = 3 \times 2 = 6$ .

# Matrix Multiplication

- For numbers:  $2 \times 3 = 3 \times 2 = 6$ .
- Matrix multiplication is (*in general*) not commutative:

$$AB \neq BA$$

# Matrix Multiplication

- For numbers:  $2 \times 3 = 3 \times 2 = 6$ .
- Matrix multiplication is (*in general*) not commutative:

$$AB \neq BA$$

- Example:

$$A = \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \quad B = \begin{bmatrix} 6 & 2 \\ 3 & 2 \end{bmatrix}, \quad AB = \begin{bmatrix} 30 & 14 \\ 12 & 6 \end{bmatrix}, \quad BA = \begin{bmatrix} 20 & 28 \\ 11 & 16 \end{bmatrix}$$

# Matrix Multiplication

- Multiplication by identity matrix  $E$ :

$$AE = EA = A$$

# Matrix Multiplication

- Multiplication by identity matrix  $E$ :

$$AE = EA = A$$

- Multiplication by zero matrix  $O$ :

$$AO = OA = O$$

# Transposing a Matrix

- The transpose of a matrix results from “flipping” the rows and columns:

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}, \quad A^T = \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix}.$$

# Transposing a Matrix

- The following properties of transposes are easily verified:
  - $A$  – symmetric matrix  $\Rightarrow A^T = A$
  - $(A^T)^T = A$
  - $(A + B)^T = A^T + B^T$
  - $(AB)^T = B^T A^T$

# Linear Transforms



*A more interesting way of looking  
at matrices.*

# Linear Transformation



Linear Transformation

# Linear Transformation



Linear Transformation

# Linear Transformation




Linear Transformation

$$x_{input} \rightarrow A \rightarrow x_{output}$$


$A$  — transformation

$x_{input}, x_{output}$  — vectors





# Transformation



# Transformation



# Transformation: Examples



**girafe  
ai**

Source: [Linear Transformations and Matrices](#)

# Linear Transformation



Linear Transformation

$$x_{input} \rightarrow A \rightarrow x_{output}$$

$A$  — transformation

$x_{input}, x_{output}$  — vectors

# Linear Transformation



## Linear Transformation

A transformation that satisfies two properties:

1.  $A(x + y) = A(x) + A(y)$
2.  $A(\lambda x) = \lambda Ax$

$$x_{\text{input}} \rightarrow A \rightarrow x_{\text{output}}$$

$A$  — transformation

$x_{\text{input}}, x_{\text{output}}$  — vectors

# Linear Transformation

- How to describe a linear transformation numerically?
-

# Linear Transformation

- How to describe a linear transformation numerically?
- With matrices! How?

# Linear Transformation

- How to describe a linear transformation numerically?
- With matrices! How?

$$x_{input} = x_1 e_1 + x_2 e_2 + \cdots + x_n e_n, \quad e_1, \dots, e_n - \text{basis}, \quad x_1, \dots, x_n - \text{coordinates}$$

# Linear Transformation

- How to describe a linear transformation numerically?
- With matrices! How?

$$x_{input} = x_1 e_1 + x_2 e_2 + \cdots + x_n e_n, \quad e_1, \dots, e_n - \text{basis}, \quad x_1, \dots, x_n - \text{coordinates}$$

$$x_{output} = A(x_{input}) = A(x_1 e_1 + x_2 e_2 + \cdots + x_n e_n) =$$

# Linear Transformation

- How to describe a linear transformation numerically?
- With matrices! How?

$x_{input} = x_1e_1 + x_2e_2 + \dots + x_ne_n$ ,  $e_1, \dots, e_n$  – basis,  $x_1, \dots, x_n$  – coordinates

$$x_{output} = A(x_{input}) = A(x_1e_1 + x_2e_2 + \dots + x_ne_n) = x_1A(e_1) + x_2A(e_2) + \dots + x_nA(e_n)$$

# Linear Transformation

- How to describe a linear transformation numerically?
- With matrices! How?

$x_{input} = x_1e_1 + x_2e_2 + \dots + x_ne_n$ ,  $e_1, \dots, e_n$  – basis,  $x_1, \dots, x_n$  – coordinates

$$x_{output} = A(x_{input}) = A(x_1e_1 + x_2e_2 + \dots + x_ne_n) = x_1A(e_1) + x_2A(e_2) + \dots + x_nA(e_n)$$

$$A := [A(e_1) \mid A(e_2) \mid \dots \mid A(e_n)]$$

# Linear Transformation

- How to describe a linear transformation numerically?
- With matrices! How?

$x_{input} = x_1 e_1 + x_2 e_2 + \dots + x_n e_n$ ,  $e_1, \dots, e_n$  – basis,  $x_1, \dots, x_n$  – coordinates


$$x_{output} = A(x_{input}) = A(x_1 e_1 + x_2 e_2 + \dots + x_n e_n) = x_1 A(e_1) + x_2 A(e_2) + \dots + x_n A(e_n)$$

$$A := [A(e_1) \mid A(e_2) \mid \dots \mid A(e_n)]$$

$$\Rightarrow x_{output} = A(x_{input}) = A \cdot x_{input}$$

# Example: Rotation


- Imagine that we want to rotate vectors in  $\mathbb{R}^2$   $90^\circ$  anti-clockwise.



# Example: Rotation

- Imagine that we want to rotate vectors in  $\mathbb{R}^2$   $90^\circ$  anti-clockwise.
- What happens to the basis vectors?


$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \rightarrow , \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \rightarrow$$



# Example: Rotation

- Imagine that we want to rotate vectors in  $\mathbb{R}^2$   $90^\circ$  anti-clockwise.
- What happens to the basis vectors?


$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \rightarrow \quad , \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \rightarrow$$



# Example: Rotation

- Imagine that we want to rotate vectors in  $\mathbb{R}^2$   $90^\circ$  anti-clockwise.
- What happens to the basis vectors?

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \rightarrow \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \rightarrow \begin{bmatrix} -1 \\ 0 \end{bmatrix}$$




# Example: Rotation

- Imagine that we want to rotate vectors in  $\mathbb{R}^2$   $90^\circ$  anti-clockwise.
- What happens to the basis vectors?

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \rightarrow \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \rightarrow \begin{bmatrix} -1 \\ 0 \end{bmatrix}$$

- Therefore,

$$R = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$





# Example: Rotation


- Imagine that we want to rotate vectors in  $\mathbb{R}^2$   $90^\circ$  anti-clockwise.
- What happens to the basis vectors?

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \rightarrow \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \rightarrow \begin{bmatrix} -1 \\ 0 \end{bmatrix}$$

- Therefore,

$$R = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

- Consider  $x = [1, 1]^T$ .





# Example: Rotation

- Imagine that we want to rotate vectors in  $\mathbb{R}^2$   $90^\circ$  anti-clockwise.
- What happens to the basis vectors?


$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \rightarrow \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \rightarrow \begin{bmatrix} -1 \\ 0 \end{bmatrix}$$

- Therefore,

$$R = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

- Consider  $x = [1, 1]^T$ . After rotation:

$$x_{rotated} =$$





# Example: Rotation

- Imagine that we want to rotate vectors in  $\mathbb{R}^2$   $90^\circ$  anti-clockwise.
- What happens to the basis vectors?


$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \rightarrow \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \rightarrow \begin{bmatrix} -1 \\ 0 \end{bmatrix}$$

- Therefore,

$$R = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

- Consider  $x = [1, 1]^T$ . After rotation:

$$x_{rotated} =$$



# Example: Rotation

- Imagine that we want to rotate vectors in  $\mathbb{R}^2$   $90^\circ$  anti-clockwise.
- What happens to the basis vectors?


$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \rightarrow \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \rightarrow \begin{bmatrix} -1 \\ 0 \end{bmatrix}$$

- Therefore,

$$R = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

- Consider  $x = [1, 1]^T$ . After rotation:

$$x_{rotated} = Rx = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}.$$



# Linear Transformation

- Every linear transformation can be defined by its matrix.

*Columns = how this transformation changes the vectors in the selected basis.*

# Linear Transformation

- Every linear transformation can be defined by its matrix.

*Columns = how this transformation changes the vectors in the selected basis.*

- Vice versa: every square matrix defines some linear transformation.

# Common Transforms



# Identity Transformation

- Doesn't change anything.
- Transformation matrix  $E$ :

$$Ex = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

# Identity Transformation

- Doesn't change anything.
- Transformation matrix  $E$ :

$$Ex = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

- Example:

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

# Stretching / Squeezing

- Enlarge (compress) all distances in a particular direction by a constant factor.
- Transformation matrix:

$$Kx = \begin{bmatrix} k_1 & 0 \\ 0 & k_2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

# Stretching / Squeezing

- Enlarge (compress) all distances in a particular direction by a constant factor.
- Transformation matrix:

$$Kx = \begin{bmatrix} k_1 & 0 \\ 0 & k_2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

- Example: stretch  $x$ -axis ( $\times 3$ ) and squeeze  $y$ -axis ( $\times 0.5$ ):

$$\begin{bmatrix} 3 & 0 \\ 0 & 0.5 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \end{bmatrix}$$

# Projection on an Axis

- Consider  $\mathbb{R}^3$ . Project on the  $XY$  – plane.
- Transformation matrix:

$$Px = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

# Projection on an Axis

- Consider  $\mathbb{R}^3$ . Project on the  $XY$  – plane.
- Transformation matrix:

$$Px = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

- Example:

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}$$

# Rotation

- Rotating points anticlockwise by  $\theta$ .
- Rotation matrix  $R_\theta$ :

$$R_\theta \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

# Rotation

- Rotating points anticlockwise by  $\theta$ .
- Rotation matrix  $R_\theta$ :

$$R_\theta \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

- Example: rotate by  $45^\circ$  anticlockwise:

$$\begin{bmatrix} \sqrt{2}/2 & -\sqrt{2}/2 \\ \sqrt{2}/2 & \sqrt{2}/2 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ \sqrt{2} \end{bmatrix}$$

# Combining Transforms



# Composition

- Let  $A$  and  $B$  be two linear transforms.  
What if we first apply  $A$  and then  $B$ ?

# Composition

- Let  $A$  and  $B$  be two linear transforms.  
What if we first apply  $A$  and then  $B$ ?
- Example: first rotate by  $90^\circ$ , then squeeze.

# Composition

- Let  $A$  and  $B$  be two linear transforms.  
What if we first apply  $A$  and then  $B$ ?
- Example: first rotate by  $90^\circ$ , then squeeze.

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \text{— rotation, } B = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} \text{— squeezing}$$

# Composition

- Let  $A$  and  $B$  be two linear transforms.  
What if we first apply  $A$  and then  $B$ ?
- Example: first rotate by  $90^\circ$ , then squeeze.

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \text{--- rotation, } B = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} \text{--- squeezing}$$


$$B(Ax) =$$

# Composition

- Let  $A$  and  $B$  be two linear transforms.  
What if we first apply  $A$  and then  $B$ ?
- Example: first rotate by  $90^\circ$ , then squeeze.

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \text{--- rotation, } B = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} \text{--- squeezing}$$

$$B(Ax) =$$




# Composition

- Let  $A$  and  $B$  be two linear transforms.  
What if we first apply  $A$  and then  $B$ ?
- Example: first rotate by  $90^\circ$ , then squeeze.

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \text{--- rotation, } B = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} \text{--- squeezing}$$

$$B(Ax) =$$




# Composition

- Let  $A$  and  $B$  be two linear transforms.  
What if we first apply  $A$  and then  $B$ ?
- Example: first rotate by  $90^\circ$ , then squeeze.

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \text{--- rotation, } B = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} \text{--- squeezing}$$

$$B(Ax) =$$




# Composition

- Let  $A$  and  $B$  be two linear transforms.  
What if we first apply  $A$  and then  $B$ ?
- Example: first rotate by  $90^\circ$ , then squeeze.

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \text{--- rotation, } B = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} \text{--- squeezing}$$

$$B(Ax) = (BA)x = Cx$$




# Composition

- Let  $A$  and  $B$  be two linear transforms.  
What if we first apply  $A$  and then  $B$ ?
- Example: first rotate by  $90^\circ$ , then squeeze.

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \text{--- rotation, } B = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} \text{--- squeezing}$$

$$B(Ax) = (BA)x = Cx$$

$$C = BA = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 0 & -0.5 \\ 0.5 & 0 \end{bmatrix}$$



# Composition


- Let  $A$  and  $B$  be two linear transforms.  
What if we first apply  $A$  and then  $B$ ?
- Example: first rotate by  $90^\circ$ , then squeeze.

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \text{— rotation, } B = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} \text{— squeezing}$$

$$B(Ax) = (BA)x = Cx$$

$$C = BA = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 0 & -0.5 \\ 0.5 & 0 \end{bmatrix}$$

= “rotate by  $90^\circ$  and squeeze”



# Sum of Transforms

- Let  $A$  and  $B$  be two linear transforms.

# Sum of Transforms

- Let  $A$  and  $B$  be two linear transforms.
- $C = A + B$  – also a linear transform:

$$C = Ax + Bx$$

# Sum of Transforms

- Let  $A$  and  $B$  be two linear transforms.
- $C = A + B$  – also a linear transform:

$$C = Ax + Bx$$

- Example:

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \text{ – rotation, } B = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} \text{ – squeezing}$$

# Sum of Transforms


- Let  $A$  and  $B$  be two linear transforms.
- $C = A + B$  – also a linear transform:

$$C = Ax + Bx$$

- Example:

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \text{ – rotation}, \quad B = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} \text{ – squeezing}$$

$$x = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$



# Sum of Transforms


- Let  $A$  and  $B$  be two linear transforms.
- $C = A + B$  – also a linear transform:

$$C = Ax + Bx$$

- Example:

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \text{ – rotation}, \quad B = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} \text{ – squeezing}$$

$$\mathbf{x} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$



# Sum of Transforms


- Let  $A$  and  $B$  be two linear transforms.
- $C = A + B$  – also a linear transform:

$$C = Ax + Bx$$

- Example:

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \text{ – rotation}, \quad B = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} \text{ – squeezing}$$

$$\mathbf{x} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$



# Sum of Transforms


- Let  $A$  and  $B$  be two linear transforms.
- $C = A + B$  – also a linear transform:

$$C = Ax + Bx$$

- Example:

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \text{ – rotation}, \quad B = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} \text{ – squeezing}$$

$$\mathbf{x} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$



# Sum of Transforms


- Let  $A$  and  $B$  be two linear transforms.
- $C = A + B$  – also a linear transform:

$$C = Ax + Bx$$

- Example:

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \text{ – rotation, } B = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} \text{ – squeezing}$$

$$x = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \quad Ax + Bx = \begin{bmatrix} -1 \\ 1 \end{bmatrix} + \begin{bmatrix} 0.5 \\ 0.5 \end{bmatrix} = \begin{bmatrix} -0.5 \\ 1.5 \end{bmatrix}$$



# Inverse Transform



# Inverse transform



**girafe**  
ai

# Inverse transform

- Consider a transform  $A$ .

# Inverse transform

- Consider a transform  $A$ .
- Inverse transform  $A^{-1}$  “maps everything back to where it was”:

# Inverse transform

- Consider a transform  $A$ .
- Inverse transform  $A^{-1}$  “maps everything back to where it was”:

$$A^{-1}(Ax) = x \Leftrightarrow A^{-1}A = E - \text{identity transformation.}$$

# Inverse transform

- Consider a transform  $A$ .
- Inverse transform  $A^{-1}$  “maps everything back to where it was”:

$$A^{-1}(Ax) = x \Leftrightarrow A^{-1}A = E - \text{identity transformation.}$$

- Not every transform has an inverse!

# Inverse transform

- Consider a transform  $A$ .
- Inverse transform  $A^{-1}$  “maps everything back to where it was”:

$$A^{-1}(Ax) = x \Leftrightarrow A^{-1}A = E - \text{identity transformation.}$$

- Not every transform has an inverse!
  - Rotation: yes (rotate it back)

# Inverse transform

- Consider a transform  $A$ .
- Inverse transform  $A^{-1}$  “maps everything back to where it was”:
$$A^{-1}(Ax) = x \Leftrightarrow A^{-1}A = E \text{ -- identity transformation.}$$
- Not every transform has an inverse!
  - Rotation: yes (rotate it back)
  - Projection: no

# Inverse of a Matrix

- An  $n \times n$  matrix  $A$  has an inverse if there exists  $A^{-1}$  such that

$$AA^{-1} = A^{-1}A = E$$

# Inverse of a Matrix

- An  $n \times n$  matrix  $A$  has an inverse if there exists  $A^{-1}$  such that

$$AA^{-1} = A^{-1}A = E$$

- A matrix that doesn't have an inverse is called **singular** or **degenerate**.

# Inverse of a Matrix

- An  $n \times n$  matrix  $A$  has an inverse if there exists  $A^{-1}$  such that

$$AA^{-1} = A^{-1}A = E$$

- A matrix that doesn't have an inverse is called **singular** or **degenerate**.
- Which matrices have an inverse?

# Determinant



# Determinant

- A numerical way to characterize a linear transformation (and its matrix):
  - absolute value = how much area changes;
  - sign = change of orientation.
- More info on the interpretation: see [video](#).

# Determinant

- $A$  – linear transform.

# Determinant


- $A$  – linear transform.
- The determinant is nonzero  $\Leftrightarrow A$  is invertible and the linear map represented by matrix  $A$  is an **isomorphism**:

# Determinant

- $A$  – linear transform.
- The determinant is nonzero  $\Leftrightarrow A$  is invertible and the linear map represented by matrix  $A$  is an **isomorphism**:

- $\det A \neq 0$ :

- $A$ :




# Determinant

- $A$  – linear transform.
- The determinant is nonzero  $\Leftrightarrow A$  is invertible and the linear map represented by matrix  $A$  is an **isomorphism**:

- $\det A \neq 0$ :


- $A$ :



- $\det A = 0$ :

-

- Several vectors are mapped onto the same vector  $\Leftrightarrow A$  maps original vector space onto a lower-dimensional space.



# Computing Determinant

$$\begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$$

# Computing Determinant

$$\begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$$

- Example:

$$R_{90} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} = 0 - (-1) = 1 \Leftrightarrow$$

“there is a transform inverse to rotation by  $90^\circ$  anticlockwise”.

# Computing Determinant

$$\begin{vmatrix} a & b & c \\ d & e & f \\ g & h & i \end{vmatrix} = a \begin{vmatrix} e & f \\ h & i \end{vmatrix} - b \begin{vmatrix} d & f \\ g & i \end{vmatrix} + c \begin{vmatrix} d & e \\ g & h \end{vmatrix}$$

# Computing Determinant

$$\begin{vmatrix} a & b & c \\ d & e & f \\ g & h & i \end{vmatrix} = a \begin{vmatrix} e & f \\ h & i \end{vmatrix} - b \begin{vmatrix} d & f \\ g & i \end{vmatrix} + c \begin{vmatrix} d & e \\ g & h \end{vmatrix}$$

- Example:

$$\begin{vmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{vmatrix} = 1 \cdot \begin{vmatrix} 1 & 0 \\ 0 & 0 \end{vmatrix} - 0 + 0 = 0 \Leftrightarrow$$

“there is no transpose inverse to projection onto  $XY$ -plane”

# Computing Determinant

- $A = \{a_{ij}\}$  –  $n \times n$  matrix.

# Computing Determinant

- $A = \{a_{ij}\}$  –  $n \times n$  matrix.
- $M_{ij}$  – its minor  $\Leftrightarrow M_{ij}$  is an  $(n - 1) \times (n - 1)$  matrix resulting from removing  $i$ -th row and  $j$ -th column from  $A$ .

# Computing Determinant

- $A = \{a_{ij}\}$  –  $n \times n$  matrix.
- $M_{ij}$  – its minor  $\Leftrightarrow M_{ij}$  is an  $(n - 1) \times (n - 1)$  matrix resulting from removing  $i$ -th row and  $j$ -th column from  $A$ .

$$\det A = \sum_{i=1}^n (-1)^{i+j} a_{ij} \det M_{ij} = \sum_{j=1}^n (-1)^{i+j} a_{ij} \det M_{ij}$$

# Computing Determinant

- $A = \{a_{ij}\}$  –  $n \times n$  matrix.
- $M_{ij}$  – its minor  $\Leftrightarrow M_{ij}$  is an  $(n - 1) \times (n - 1)$  matrix resulting from removing  $i$ -th row and  $j$ -th column from  $A$ .

$$\det A = \sum_{i=1}^n (-1)^{i+j} a_{ij} \det M_{ij} = \sum_{j=1}^n (-1)^{i+j} a_{ij} \det M_{ij}$$

- Laplace extension.

# Some Properties of the Determinant

- $\det A^T = \det A$
- $\det AB = \det A \cdot \det B$
- $\det A^{-1} = \frac{1}{\det A}$

# **Finding Inverse of a Matrix**



# Gaussian Elimination

- $A = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 2 & 1 \\ 1 & 1 & 1 \end{pmatrix}$
- $\det A \neq 0 \Rightarrow$  there exists  $A^{-1}$ . Let's find it!



# Gaussian Elimination

- $A = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 2 & 1 \\ 1 & 1 & 1 \end{pmatrix}$
- $\det A \neq 0 \Rightarrow$  there exists  $A^{-1}$ . Let's find it!
- Augment the initial matrix:

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 & 1 & 0 \\ 1 & 1 & 1 & 0 & 0 & 1 \end{array} \right)$$



# Gaussian Elimination

- $A = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 2 & 1 \\ 1 & 1 & 1 \end{pmatrix}$

- $\det A \neq 0 \Rightarrow$  there exists  $A^{-1}$ . Let's find it!
- Augment the initial matrix:

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 & 1 & 0 \\ 1 & 1 & 1 & 0 & 0 & 1 \end{array} \right)$$

- Perform elementary row operations and obtain identity matrix on the left. The inverse will be on the right!



# Gaussian Elimination

- Elementary row operations:
  - swap rows;
  - multiply rows by some number;
  - add / subtract one row to / from another.

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 & 1 & 0 \\ 1 & 1 & 1 & 0 & 0 & 1 \end{array} \right)$$

# Gaussian Elimination

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 & 1 & 0 \\ 1 & 1 & 1 & 0 & 0 & 1 \end{array} \right) \rightarrow \{(3) - (1)\} \rightarrow \left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & -1 & 0 & 1 \end{array} \right) \rightarrow$$

# Gaussian Elimination

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 & 1 & 0 \\ 1 & 1 & 1 & 0 & 0 & 1 \end{array} \right) \rightarrow \{(3) - (1)\} \rightarrow \left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & -1 & 0 & 1 \end{array} \right) \rightarrow$$

$$\rightarrow \{(2) - 2 \cdot (3)\} \rightarrow \left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 2 & 1 & -2 \\ 0 & 1 & 0 & -1 & 0 & 1 \end{array} \right)$$

# Gaussian Elimination

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 & 1 & 0 \\ 1 & 1 & 1 & 0 & 0 & 1 \end{array} \right) \rightarrow \{(3) - (1)\} \rightarrow \left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & -1 & 0 & 1 \end{array} \right) \rightarrow$$

$$\rightarrow \{(2) - 2 \cdot (3)\} \rightarrow \left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 2 & 1 & -2 \\ 0 & 1 & 0 & -1 & 0 & 1 \end{array} \right) \rightarrow \{(1) - (2)\} \rightarrow \left( \begin{array}{ccc|ccc} 1 & 0 & 0 & -1 & -1 & 2 \\ 0 & 0 & 1 & 2 & 1 & -2 \\ 0 & 1 & 0 & -1 & 0 & 1 \end{array} \right)$$

# Gaussian Elimination

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 & 1 & 0 \\ 1 & 1 & 1 & 0 & 0 & 1 \end{array} \right) \rightarrow \{(3) - (1)\} \rightarrow \left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & -1 & 0 & 1 \end{array} \right) \rightarrow$$

$$\rightarrow \{(2) - 2 \cdot (3)\} \rightarrow \left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 2 & 1 & -2 \\ 0 & 1 & 0 & -1 & 0 & 1 \end{array} \right) \rightarrow \{(1) - (2)\} \rightarrow \left( \begin{array}{ccc|ccc} 1 & 0 & 0 & -1 & -1 & 2 \\ 0 & 0 & 1 & 2 & 1 & -2 \\ 0 & 1 & 0 & -1 & 0 & 1 \end{array} \right) \rightarrow$$

$$\rightarrow \{\text{swap (2) and (3)}\} \rightarrow \left( \begin{array}{ccc|ccc} 1 & 0 & 0 & -1 & -1 & 2 \\ 0 & 1 & 0 & -1 & 0 & 1 \\ 0 & 0 & 1 & 2 & 1 & -2 \end{array} \right)$$

# Gaussian Elimination

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 & 1 & 0 \\ 1 & 1 & 1 & 0 & 0 & 1 \end{array} \right) \rightarrow \dots \rightarrow \left( \begin{array}{ccc|cccc} 1 & 0 & 0 & -1 & -1 & 2 \\ 0 & 1 & 0 & -1 & 0 & 1 \\ 0 & 0 & 1 & 2 & 1 & -2 \end{array} \right)$$

$$A = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 2 & 1 \\ 1 & 1 & 1 \end{pmatrix}, \quad A^{-1} = \begin{pmatrix} -1 & -1 & 2 \\ -1 & 0 & 1 \\ 2 & 1 & -2 \end{pmatrix}, \quad AA^{-1} = A^{-1}A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

# Rank



# Column Space

- Consider a square matrix  $A$ .
- Its columns  $A^1, \dots, A^n$  can be seen as vectors.

# Column Space

- Consider a square matrix  $A$ .
- Its columns  $A^1, \dots, A^n$  can be seen as vectors.
- $U = \text{span}\{A^1, \dots, A^n\}$  – **column space** of  $A$ .
  - All vectors that can be obtain by linearly combining columns of  $A$ .
  - $\Leftrightarrow$  image of linear transformation  $A$  (= all the vectors we can get by applying  $A$ ).

# Rank

- Column space  $U = \text{span}\{A^1, \dots, A^n\}$  is the image of linear transformation  $A$ .
- **Rank** of a matrix is the number of dimensions in its column space.

# Rank

- Column space  $U = \text{span}\{A^1, \dots, A^n\}$  is the image of linear transformation  $A$ .
- **Rank** of a matrix is the number of dimensions in its column space.
  - **Full rank** matrix:  $n$  columns, all linearly independent.
  - Lower-rank matrices: linearly dependent columns present.

# Rank: Examples

- $A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$

- $B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}$

- $C = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

# Rank: Examples

- $A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \text{rank}(A) = 1$

- $B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}$

- $C = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

# Rank: Examples

- $A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \text{rank}(A) = 1$

- $B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}, \text{rank}(A) = 2$

- $C = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

# Rank: Examples

- $A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \text{rank}(A) = 1$
- $B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}, \text{rank}(A) = 2$
- $C = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \text{rank}(A) = 3$

# Column vs Row Rank

- Column space of  $A$  = span of  $A$ 's columns.  
Its dimensionality = (column) rank.

# Column vs Row Rank

- Column space of  $A$  = span of  $A$ 's columns.  
Its dimensionality = (column) rank.
- We can also define the row space of  $A$  as a span of  $A$ 's rows.  
Its dimensionality = row rank.

# Column vs Row Rank

- Column space of  $A$  = span of  $A$ 's columns.  
Its dimensionality = (column) rank.
- We can also define the row space of  $A$  as a span of  $A$ 's rows.  
Its dimensionality = row rank.
- Column rank vs. row rank?

# Column vs Row Rank

- Column space of  $A$  = span of  $A$ 's columns.  
Its dimensionality = (column) rank.
- We can also define the row space of  $A$  as a span of  $A$ 's rows.  
Its dimensionality = row rank.
- Column rank vs. row rank?
- Fundamental result: the column rank and the row rank are always equal.  
See [proofs](#).

# Rank

- This is why there cannot be more than  $n$  linearly independent vectors in  $\mathbb{R}^n$ !

# Rank

- This is why there cannot be more than  $n$  linearly independent vectors in  $\mathbb{R}^n$ !
- Imagine  $x_1, x_2, \dots, x_m \in \mathbb{R}^n$ :

# Rank

- This is why there cannot be more than  $n$  linearly independent vectors in  $\mathbb{R}^n$ !
- Imagine  $x_1, x_2, \dots, x_m \in \mathbb{R}^n$ :

$$X = [x_1 \mid x_2 \mid \dots \mid x_n] = \begin{bmatrix} x_{11} & x_{12} & \dots & x_{1m} \\ x_{21} & x_{22} & \dots & x_{2m} \\ \dots & \dots & \dots & \dots \\ x_{n1} & x_{n2} & \dots & x_{nm} \end{bmatrix}$$

# Rank

- This is why there cannot be more than  $n$  linearly independent vectors in  $\mathbb{R}^n$ !
- Imagine  $x_1, x_2, \dots, x_m \in \mathbb{R}^n$ :

$$X = [x_1 \mid x_2 \mid \dots \mid x_n] = \begin{bmatrix} x_{11} & x_{12} & \dots & x_{1m} \\ x_{21} & x_{22} & \dots & x_{2m} \\ \dots & \dots & \dots & \dots \\ x_{n1} & x_{n2} & \dots & x_{nm} \end{bmatrix}$$

$$\text{rank}(X) \leq \min\{n, m\}$$

# Rank: Examples

- $A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \text{rank}(A) = 1$
- $B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}, \text{rank}(A) = 2$
- $C = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \text{rank}(A) = 3$

# Rank: Examples

- $A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$ ,  $\text{rank}(A) = 1 \Leftrightarrow \mathbb{R}^3$  is mapped onto a line
- $B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}$ ,  $\text{rank}(A) = 2$
- $C = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ ,  $\text{rank}(A) = 3$

# Rank: Examples

- $A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$ ,  $\text{rank}(A) = 1 \Leftrightarrow \mathbb{R}^3$  is mapped onto a line
- $B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}$ ,  $\text{rank}(A) = 2 \Leftrightarrow \mathbb{R}^3$  is mapped onto a plane
- $C = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ ,  $\text{rank}(A) = 3$

# Rank: Examples

- $A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$ ,  $\text{rank}(A) = 1 \Leftrightarrow \mathbb{R}^3$  is mapped onto a line
- $B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}$ ,  $\text{rank}(A) = 2 \Leftrightarrow \mathbb{R}^3$  is mapped onto a plane
- $C = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ ,  $\text{rank}(A) = 3 \Leftrightarrow \mathbb{R}^3$  is mapped on itself (isomorphism)

# Rank: Examples

- $A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$ ,  $\text{rank}(A) = 1 \Leftrightarrow \mathbb{R}^3$  is mapped onto a line  
Ininitely many vectors are mapped into a zero vector.
- $B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}$ ,  $\text{rank}(A) = 2 \Leftrightarrow \mathbb{R}^3$  is mapped onto a plane  
Ininitely many vectors are mapped into a zero vector.
- $C = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ ,  $\text{rank}(A) = 3 \Leftrightarrow \mathbb{R}^3$  is mapped on itself (isomorphism)  
Only a zero vector is mapped into a zero vector.

# Null space

- A set of vectors that are mapped to  $\mathbf{0}$  by a linear transformation  $A$ .

# Null space

- A set of vectors that are mapped to  $\mathbf{0}$  by a linear transformation  $A$ .
- Example: projection onto  $XY$ -plane:

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad A \begin{bmatrix} 0 \\ 0 \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \Rightarrow$$

Null space:  $\left\{ v \in \mathbb{R}^3 \mid v = \begin{bmatrix} 0 \\ 0 \\ z \end{bmatrix}, z \in \mathbb{R} \right\}$

# **Systems of Linear Equations**



# what is a SLE?

$$\begin{cases} 2x_1 + 5x_2 + 3x_3 = -3 \\ 4x_1 + 0x_2 + 8x_3 = 0 \\ 1x_1 + 3x_2 + 0x_3 = 2 \end{cases}$$

# Solutions to SLE



1. 
$$\begin{cases} x + y = 1 \\ x + y = 2 \end{cases}$$

2. 
$$\begin{cases} x + y = 1 \\ 2x + y = 2 \end{cases}$$

3. 
$$\begin{cases} x + y = 1 \\ 2x + 2y = 2 \end{cases}$$

# Solutions to SLE



1.

$$\begin{cases} x + y = 1 \\ x + y = 2 \end{cases}$$

No solutions.

2.

$$\begin{cases} x + y = 1 \\ 2x + y = 2 \end{cases}$$

A single solution:  $x = 1, y = 0$

3.

$$\begin{cases} x + y = 1 \\ 2x + 2y = 2 \end{cases}$$

Infinitely many solutions.

# SLE: Matrix Notation

$$\begin{cases} 2x_1 + 5x_2 + 3x_3 = -3 \\ 4x_1 + 0x_2 + 8x_3 = 0 \\ 1x_1 + 3x_2 + 0x_3 = 2 \end{cases}$$

# SLE: Matrix Notation

$$\begin{cases} 2x_1 + 5x_2 + 3x_3 = -3 \\ 4x_1 + 0x_2 + 8x_3 = 0 \\ 1x_1 + 3x_2 + 0x_3 = 2 \end{cases}$$

$$A = \begin{bmatrix} 2 & 5 & 3 \\ 4 & 0 & 8 \\ 1 & 3 & 0 \end{bmatrix}, \quad b = \begin{bmatrix} -3 \\ 0 \\ 2 \end{bmatrix}, \quad x = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

$$Ax = b$$

# SLE: Matrix Notation

$$\begin{cases} 2x_1 + 5x_2 + 3x_3 = -3 \\ 4x_1 + 0x_2 + 8x_3 = 0 \\ 1x_1 + 3x_2 + 0x_3 = 2 \end{cases}$$

$$A = \begin{bmatrix} 2 & 5 & 3 \\ 4 & 0 & 8 \\ 1 & 3 & 0 \end{bmatrix}, \quad b = \begin{bmatrix} -3 \\ 0 \\ 2 \end{bmatrix}, \quad x = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

$$Ax = b$$

“Find vector(s)  $x$  that are mapped into  $b$  by transform  $A$ ”

# Number of Solutions

- $Ax = b$ : “Find vector(s)  $x$  that are mapped into  $b$  by transform  $A$ ”.

# Number of Solutions

- $Ax = b$ : “Find vector(s)  $x$  that are mapped into  $b$  by transform  $A$ ”.
- If  $\det A \neq 0$ , there exists  $A^{-1}$ :  $x = A^{-1}b$ .  
So, there is a single solution.

# Number of Solutions

- $Ax = b$ : “Find vector(s)  $x$  that are mapped into  $b$  by transform  $A$ ”.
- If  $\det A \neq 0$ , there exists  $A^{-1}$ :  $x = A^{-1}b$ .  
So, there is a single solution.
- If  $\det A = 0$ , it’s not that simple:

# Number of Solutions

- $Ax = b$ : “Find vector(s)  $x$  that are mapped into  $b$  by transform  $A$ ”.
- If  $\det A \neq 0$ , there exists  $A^{-1}$ :  $x = A^{-1}b$ .  
So, there is a single solution.
- If  $\det A = 0$ , it’s not that simple:  
 $A$  maps the original space into a lower dimensional subspace.

# Number of Solutions

- $Ax = b$ : “Find vector(s)  $x$  that are mapped into  $b$  by transform  $A$ ”.
- If  $\det A \neq 0$ , there exists  $A^{-1}$ :  $x = A^{-1}b$ .  
So, there is a single solution.
- If  $\det A = 0$ , it’s not that simple:  
 $A$  maps the original space into a lower dimensional subspace.  
If we are lucky,  $b$  is there, and we have infinitely many solutions.  
If not, there’re no solutions.

# Solutions to SLE



1.

$$\begin{cases} x + y = 1 \\ x + y = 2 \end{cases}$$

No solutions.

2.

$$\begin{cases} x + y = 1 \\ 2x + y = 2 \end{cases}$$

A single solution:  $x = 1, y = 0$

$$A = \begin{bmatrix} 1 & 1 \\ 2 & 1 \end{bmatrix}, \det A \neq 0.$$

3.

$$\begin{cases} x + y = 1 \\ 2x + 2y = 2 \end{cases}$$

Infinitely many solutions.

# Solutions to SLE



$$1. \begin{cases} x + y = 1 \\ x + y = 2 \end{cases}$$

No solutions.

$A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}, \det A = 0$ .  $A$  maps  $\mathbb{R}^2$  onto a line,  $b = [1, 2]^T$  isn't there.

$$2. \begin{cases} x + y = 1 \\ 2x + y = 2 \end{cases}$$

A single solution:  $x = 1, y = 0$

$$A = \begin{bmatrix} 1 & 1 \\ 2 & 1 \end{bmatrix}, \det A \neq 0.$$

$$3. \begin{cases} x + y = 1 \\ 2x + 2y = 2 \end{cases}$$

Infinitely many solutions.

# Solutions to SLE



$$1. \begin{cases} x + y = 1 \\ x + y = 2 \end{cases}$$

No solutions.

$A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}, \det A = 0$ .  $A$  maps  $\mathbb{R}^2$  onto a line,  $b = [1, 2]^T$  isn't there.

$$2. \begin{cases} x + y = 1 \\ 2x + y = 2 \end{cases}$$

A single solution:  $x = 1, y = 0$

$$A = \begin{bmatrix} 1 & 1 \\ 2 & 1 \end{bmatrix}, \det A \neq 0.$$

$$3. \begin{cases} x + y = 1 \\ 2x + 2y = 2 \end{cases}$$

Infinitely many solutions.

$A = \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}, \det A = 0$ .  $A$  maps  $\mathbb{R}^2$  onto a line,  $b = [1, 2]^T$  is there.

# Number of Solutions

- $Ax = b$ : “Find vector(s)  $x$  that are mapped into  $b$  by transform  $A$ ”.
- If  $\det A \neq 0$ , there exists  $A^{-1}$ :  $x = A^{-1}b$ .  
So, there is a single solution.
- If  $\det A = 0$ , it’s not that simple:  
 $A$  maps the original space into a lower dimensional subspace.  
If we are lucky,  $b$  is there, and we have infinitely many solutions.  
If not, there’re no solutions.

How do we check that?

# Number of Solutions

- $Ax = b$  – SLE.
- Consider matrix  $(A|b) = \begin{bmatrix} a_{11} & \dots & a_{1n} & b_1 \\ \dots & \dots & \dots & \dots \\ a_{m1} & \dots & a_{mn} & b_n \end{bmatrix}$ .

# Number of Solutions

- $Ax = b$  – SLE.
- Consider matrix  $(A|b) = \begin{bmatrix} a_{11} & \dots & a_{1n} & b_1 \\ \dots & \dots & \dots & \dots \\ a_{m1} & \dots & a_{mn} & b_n \end{bmatrix}$ .
  - $Ax = b$  has a unique solution  $\Leftrightarrow \text{rank}(A|b) = \text{rank}(A) = n$ .

# Number of Solutions

- $Ax = b$  – SLE.
- Consider matrix  $(A|b) = \begin{bmatrix} a_{11} & \dots & a_{1n} & b_1 \\ \dots & \dots & \dots & \dots \\ a_{m1} & \dots & a_{mn} & b_n \end{bmatrix}$ .
  - $Ax = b$  has a unique solution  $\Leftrightarrow \text{rank}(A|b) = \text{rank}(A) = n$ .
  - $Ax = b$  has infinitely many solutions  $\Leftrightarrow \text{rank}(A|b) = \text{rank}(A) < n$ .

# Number of Solutions

- $Ax = b$  – SLE.
- Consider matrix  $(A|b) = \begin{bmatrix} a_{11} & \dots & a_{1n} & b_1 \\ \dots & \dots & \dots & \dots \\ a_{m1} & \dots & a_{mn} & b_n \end{bmatrix}$ .
  - $Ax = b$  has a unique solution  $\Leftrightarrow \text{rank}(A|b) = \text{rank}(A) = n$ .
  - $Ax = b$  has infinitely many solutions  $\Leftrightarrow \text{rank}(A|b) = \text{rank}(A) < n$ .
  - $Ax = b$  has no solutions  $\Leftrightarrow \text{rank}(A|b) > \text{rank}(A)$ .

# Solutions to SLE



1.

$$\begin{cases} x + y = 1 \\ x + y = 2 \end{cases}$$

No solutions.

$$1 = \text{rank}(A) < \text{rank}(A|b) = 2$$

2.

$$\begin{cases} x + y = 1 \\ 2x + y = 2 \end{cases}$$

A single solution:  $x = 1, y = 0$

$$\text{rank}(A) = \text{rank}(A|b) = 2$$

3.

$$\begin{cases} x + y = 1 \\ 2x + 2y = 2 \end{cases}$$

Infinitely many solutions.

$$\text{rank}(A) = \text{rank}(A|b) = 1 < 2$$

# Gaussian Elimination



# Gaussian elimination

- $Ax = b$

$$A = \begin{bmatrix} 1 & 3 & -2 \\ 3 & 5 & 6 \\ 2 & 4 & 3 \end{bmatrix}, \quad b = \begin{bmatrix} 5 \\ 7 \\ 8 \end{bmatrix}$$

# Gaussian elimination

- $Ax = b$

$$A = \begin{bmatrix} 1 & 3 & -2 \\ 3 & 5 & 6 \\ 2 & 4 & 3 \end{bmatrix}, \quad b = \begin{bmatrix} 5 \\ 7 \\ 8 \end{bmatrix}$$

- Elementary row operations:

$$\begin{array}{c} \left[ \begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 3 & 5 & 6 & 7 \\ 2 & 4 & 3 & 8 \end{array} \right] \sim \left[ \begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 0 & -4 & 12 & -8 \\ 2 & 4 & 3 & 8 \end{array} \right] \sim \left[ \begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 0 & -4 & 12 & -8 \\ 0 & -2 & 7 & -2 \end{array} \right] \sim \left[ \begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 0 & 1 & -3 & 2 \\ 0 & -2 & 7 & -2 \end{array} \right] \\ \sim \left[ \begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 0 & 1 & -3 & 2 \\ 0 & 0 & 1 & 2 \end{array} \right] \sim \left[ \begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 0 & 1 & 0 & 8 \\ 0 & 0 & 1 & 2 \end{array} \right] \sim \left[ \begin{array}{ccc|c} 1 & 3 & 0 & 9 \\ 0 & 1 & 0 & 8 \\ 0 & 0 & 1 & 2 \end{array} \right] \sim \left[ \begin{array}{ccc|c} 1 & 0 & 0 & -15 \\ 0 & 1 & 0 & 8 \\ 0 & 0 & 1 & 2 \end{array} \right] \end{array}$$

# Gaussian elimination

- $Ax = b$

$$A = \begin{bmatrix} 1 & 3 & -2 \\ 3 & 5 & 6 \\ 2 & 4 & 3 \end{bmatrix}, \quad b = \begin{bmatrix} 5 \\ 7 \\ 8 \end{bmatrix}$$

- Elementary row operations:

$$\begin{array}{c} \left[ \begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 3 & 5 & 6 & 7 \\ 2 & 4 & 3 & 8 \end{array} \right] \sim \left[ \begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 0 & -4 & 12 & -8 \\ 2 & 4 & 3 & 8 \end{array} \right] \sim \left[ \begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 0 & -4 & 12 & -8 \\ 0 & -2 & 7 & -2 \end{array} \right] \sim \left[ \begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 0 & 1 & -3 & 2 \\ 0 & -2 & 7 & -2 \end{array} \right] \\ \sim \left[ \begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 0 & 1 & -3 & 2 \\ 0 & 0 & 1 & 2 \end{array} \right] \sim \left[ \begin{array}{ccc|c} 1 & 3 & -2 & 5 \\ 0 & 1 & 0 & 8 \\ 0 & 0 & 1 & 2 \end{array} \right] \sim \left[ \begin{array}{ccc|c} 1 & 3 & 0 & 9 \\ 0 & 1 & 0 & 8 \\ 0 & 0 & 1 & 2 \end{array} \right] \sim \left[ \begin{array}{ccc|c} 1 & 0 & 0 & -15 \\ 0 & 1 & 0 & 8 \\ 0 & 0 & 1 & 2 \end{array} \right] \end{array}$$

Unique solution.

# Gaussian Elimination

- $Ax = b$  – SLE.

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & -3 \\ 2 & 1 & 5 \end{bmatrix}, \quad b = \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}$$

# Gaussian Elimination

- $Ax = b$  – SLE.

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & -3 \\ 2 & 1 & 5 \end{bmatrix}, \quad b = \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}$$

- Elementary row operations:

$$\left( \begin{array}{ccc|c} 1 & 1 & 1 & 2 \\ 0 & 1 & -3 & 1 \\ 2 & 1 & 5 & 0 \end{array} \right) \rightarrow \dots \rightarrow \left( \begin{array}{ccc|c} 1 & 0 & 4 & 1 \\ 0 & 1 & -3 & 1 \\ 0 & 0 & 0 & -3 \end{array} \right)$$

# Gaussian Elimination

- $Ax = b$  – SLE.

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & -3 \\ 2 & 1 & 5 \end{bmatrix}, \quad b = \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}$$

- Elementary row operations:

$$\left( \begin{array}{ccc|c} 1 & 1 & 1 & 2 \\ 0 & 1 & -3 & 1 \\ 2 & 1 & 5 & 0 \end{array} \right) \rightarrow \dots \rightarrow \left( \begin{array}{ccc|c} 1 & 0 & 4 & 1 \\ 0 & 1 & -3 & 1 \\ 0 & 0 & 0 & -3 \end{array} \right)$$

No solutions.

# Gaussian Elimination

- $Ax = b$  – SLE.

$$A = \begin{bmatrix} -3 & -5 & 36 \\ -1 & 0 & 7 \\ 1 & 1 & -10 \end{bmatrix}, \quad b = \begin{bmatrix} 10 \\ 5 \\ -4 \end{bmatrix}$$

-

# Gaussian Elimination

- $Ax = b$  – SLE.

$$A = \begin{bmatrix} -3 & -5 & 36 \\ -1 & 0 & 7 \\ 1 & 1 & -10 \end{bmatrix}, \quad b = \begin{bmatrix} 10 \\ 5 \\ -4 \end{bmatrix}$$

- Elementary row operations:

$$\left( \begin{array}{ccc|c} -3 & -5 & 36 & 10 \\ -1 & 0 & 7 & 5 \\ 1 & 1 & -10 & -4 \end{array} \right) \rightarrow \dots \rightarrow \left( \begin{array}{ccc|c} 1 & 0 & -7 & -5 \\ 0 & 2 & -3 & -1 \\ 0 & 0 & 0 & 0 \end{array} \right)$$

# Gaussian Elimination

- $Ax = b$  – SLE.

$$A = \begin{bmatrix} -3 & -5 & 36 \\ -1 & 0 & 7 \\ 1 & 1 & -10 \end{bmatrix}, \quad b = \begin{bmatrix} 10 \\ 5 \\ -4 \end{bmatrix}$$

- Elementary row operations:

$$\left( \begin{array}{ccc|c} -3 & -5 & 36 & 10 \\ -1 & 0 & 7 & 5 \\ 1 & 1 & -10 & -4 \end{array} \right) \rightarrow \dots \rightarrow \left( \begin{array}{ccc|c} 1 & 0 & -7 & -5 \\ 0 & 2 & -3 & -1 \\ 0 & 0 & 0 & 0 \end{array} \right)$$

Infinitely many solutions.

# **Homogeneous SLE**



# Homogeneous SLE

$$\begin{cases} 2x_1 + 5x_2 + 3x_3 = 0 \\ 4x_1 + 0x_2 + 8x_3 = 0 \\ 1x_1 + 3x_2 + 0x_3 = 0 \end{cases}$$

# Homogeneous SLE

$$Ax = 0$$

# Homogeneous SLE

$$Ax = 0$$

Solutions = null space of  $A$ .

# Homogeneous SLE

$$Ax = 0$$

Solutions = null space of  $A$ .

$\text{rank } A = \#\text{variables} \rightarrow \text{unique solution (0)}$

$\text{rank } A < \#\text{variables} \rightarrow \text{infinitely many solutions.}$

# Homogeneous SLE

- $Ax = 0$
- Let  $V$  be a set of solutions:

$$\forall v \in V \quad Av = 0$$

# Homogeneous SLE

- $Ax = 0$
- Let  $V$  be a set of solutions:

$$\forall v \in V \quad Av = 0$$

- Note that
  - $0 \in V$  because  $A \cdot 0 = 0$ .

# Homogeneous SLE

- $Ax = 0$
- Let  $V$  be a set of solutions:

$$\forall v \in V \quad Av = 0$$

- Note that
  - $0 \in V$  because  $A \cdot 0 = 0$ .
  - $\forall v \in V \rightarrow \lambda v \in V$   
Indeed,  $A(\lambda v) = \lambda Av = 0$

# Homogeneous SLE

- $Ax = 0$
- Let  $V$  be a set of solutions:

$$\forall v \in V \quad Av = 0$$

- Note that
  - $0 \in V$  because  $A \cdot 0 = 0$ .
  - $\forall v \in V \rightarrow \lambda v \in V$   
Indeed,  $A(\lambda v) = \lambda Av = 0$
  - $\forall v_1, v_2 \in V \rightarrow (v_1 + v_2) \in V.$   
Indeed,  $A(v_1 + v_2) = Av_1 + Av_2 = 0$

# Homogeneous SLE

- $Ax = 0$
- Let  $V$  be a set of solutions:

$$\forall v \in V \quad Av = 0$$

- Note that
  - $0 \in V$  because  $A \cdot 0 = 0$ .
  - $\forall v \in V \rightarrow \lambda v \in V$   
Indeed,  $A(\lambda v) = \lambda Av = 0$
  - $\forall v_1, v_2 \in V \rightarrow (v_1 + v_2) \in V$ .  
Indeed,  $A(v_1 + v_2) = Av_1 + Av_2 = 0$

**V is a linear subspace!**

# To sum up

- Matrices as linear transforms
- Examples of common transforms
- Inverse
- Determinant
- Rank
- Solutions to SLE