




# Math Refresher for DS

Practical Session 11



# Today: Integrals

- Indefinite integrals

$$\int f(x)dx$$

- Definite integrals

$$\int_a^b f(x)dx$$

- Improper integrals

$$\int_{-\infty}^{+\infty} f(x)dx$$

# Indefinite Integral



# Antiderivatives

- Which function  $F(x)$  should we differentiate to get

$$f(x) = x^4 + 3x - 9$$

# Antiderivatives

- Which function  $F(x)$  should we differentiate to get

$$f(x) = x^4 + 3x - 9$$

$$F(x) = \frac{1}{5}x^5 + \frac{3}{2}x^2 - 9x$$

# Antiderivatives

- Which function  $F(x)$  should we differentiate to get

$$f(x) = x^4 + 3x - 9$$

$$F(x) = \frac{1}{5}x^5 + \frac{3}{2}x^2 - 9x$$

$$F(x) = \frac{1}{5}x^5 + \frac{3}{2}x^2 - 9x + 10$$

# Antiderivatives

- Which function  $F(x)$  should we differentiate to get

$$f(x) = x^4 + 3x - 9$$

$$F(x) = \frac{1}{5}x^5 + \frac{3}{2}x^2 - 9x$$

$$F(x) = \frac{1}{5}x^5 + \frac{3}{2}x^2 - 9x + 10$$

$$F(x) = \frac{1}{5}x^5 + \frac{3}{2}x^2 - 9x + C, \quad C \in \mathbb{R}$$

# Antiderivatives

Given a function,  $f(x)$ , an **anti-derivative** of  $f(x)$  is any function  $F(x)$  such that

$$F'(x) = f(x)$$

If  $F(x)$  is any anti-derivative of  $f(x)$  then the most general anti-derivative of  $f(x)$  is called an **indefinite integral** and denoted,

$$\int f(x) dx = F(x) + c, \quad c \text{ is any constant}$$

In this definition the  $\int$  is called the **integral symbol**,  $f(x)$  is called the **integrand**,  $x$  is called the **integration variable** and the “ $c$ ” is called the **constant of integration**.



# Indefinite Integral

$$\int f(x) dx$$

# Indefinite Integral: Example

$$\int x^n dx = \quad , \quad n \neq -1$$

$$\int \frac{1}{x} dx =$$

$$\int \sin x dx =$$

$$\int e^x dx =$$

# Indefinite Integral: Example

$$\int x^n dx = \frac{1}{n+1} x^{n+1} + C, \quad n \neq -1$$

$$\int \frac{1}{x} dx =$$

$$\int \sin x dx =$$

$$\int e^x dx =$$

# Indefinite Integral: Example

$$\int x^n dx = \frac{1}{n+1} x^{n+1} + C, \quad n \neq -1$$

$$\int \frac{1}{x} dx = \ln x + C$$

$$\int \sin x dx =$$

$$\int e^x dx =$$

# Indefinite Integral: Example

$$\int x^n dx = \frac{1}{n+1} x^{n+1} + C, \quad n \neq -1$$

$$\int \frac{1}{x} dx = \ln x + C$$

$$\int \sin x dx = -\cos x + C$$

$$\int e^x dx =$$

# Indefinite Integral: Example

$$\int x^n dx = \frac{1}{n+1} x^{n+1} + C, \quad n \neq -1$$

$$\int \frac{1}{x} dx = \ln x + C$$

$$\int \sin x dx = -\cos x + C$$

$$\int e^x dx = e^x + C$$

# Indefinite Integral: Example

- Determine the function  $f(x)$  if

$$f'(x) = 4x^3 - 9 + 2 \sin x + 7e^x, \quad f(0) = 15$$

# Indefinite Integral: Example

- Determine the function  $f(x)$  if

$$f'(x) = 4x^3 - 9 + 2 \sin x + 7e^x, \quad f(0) = 15$$

$$f(x) = \int 4x^3 - 9 + 2 \sin x + 7e^x dx =$$



# Indefinite Integral: Example

- Determine the function  $f(x)$  if

$$f'(x) = 4x^3 - 9 + 2 \sin x + 7e^x, \quad f(0) = 15$$

$$f(x) = \int 4x^3 - 9 + 2 \sin x + 7e^x dx = x^4 - 9x - 2 \cos x + 7e^x + C$$

# Indefinite Integral: Example

- Determine the function  $f(x)$  if

$$f'(x) = 4x^3 - 9 + 2 \sin x + 7e^x, \quad f(0) = 15$$

$$f(x) = \int 4x^3 - 9 + 2 \sin x + 7e^x dx = x^4 - 9x - 2 \cos x + 7e^x + C$$

$$f(0) = 15 =$$

# Indefinite Integral: Example

- Determine the function  $f(x)$  if

$$f'(x) = 4x^3 - 9 + 2 \sin x + 7e^x, \quad f(0) = 15$$

$$f(x) = \int 4x^3 - 9 + 2 \sin x + 7e^x dx = x^4 - 9x - 2 \cos x + 7e^x + C$$

$$f(0) = 15 = -2 \cos 0 + 7e^0 = 5$$

# Indefinite Integral: Example

- Determine the function  $f(x)$  if

$$f'(x) = 4x^3 - 9 + 2 \sin x + 7e^x, \quad f(0) = 15$$

$$f(x) = \int 4x^3 - 9 + 2 \sin x + 7e^x dx = x^4 - 9x - 2 \cos x + 7e^x + C$$

$$f(0) = 15 = -2 \cos 0 + 7e^0 = 5 \quad \rightarrow \quad C = 10$$

# Indefinite Integral: Example

- Determine the function  $f(x)$  if

$$f'(x) = 4x^3 - 9 + 2 \sin x + 7e^x, \quad f(0) = 15$$

$$f(x) = \int 4x^3 - 9 + 2 \sin x + 7e^x dx = x^4 - 9x - 2 \cos x + 7e^x + C$$

$$f(0) = 15 = -2 \cos 0 + 7e^0 = 5 \rightarrow C = 10$$

$$f(x) = x^4 - 9x - 2 \cos x + 7e^x + 10$$

# **Integration techniques**



# Substitution Rule

- Compute the following integral:

$$\int 18x^2 \sqrt[4]{6x^3 + 5} \, dx =$$

# Substitution Rule

- Compute the following integral:

$$\int 18x^2 \sqrt[4]{6x^3 + 5} \, dx =$$

- Substitution rule:

$$\int f(g(x)) g'(x) \, dx = \int f(u) \, du, \quad \text{where, } u = g(x)$$



# Substitution Rule

- Compute the following integral:

$$\int 18x^2 \sqrt[4]{6x^3 + 5} \, dx = \{u = 6x^3 + 5, \quad du = \quad\quad\quad\} =$$

- Substitution rule:

$$\int f(g(x)) g'(x) \, dx = \int f(u) \, du, \quad \text{where, } u = g(x)$$

# Substitution Rule

- Compute the following integral:

$$\int 18x^2 \sqrt[4]{6x^3 + 5} \, dx = \{u = 6x^3 + 5, \quad du = 18x^2 dx\} =$$

- Substitution rule:

$$\int f(g(x)) g'(x) \, dx = \int f(u) \, du, \quad \text{where, } u = g(x)$$

# Substitution Rule

- Compute the following integral:

$$\begin{aligned}\int 18x^2 \sqrt[4]{6x^3 + 5} \, dx &= \{u = 6x^3 + 5, \quad du = 18x^2 dx\} = \\ &= \int \sqrt[4]{u} \, du = \frac{4}{5} u^{5/4} + C\end{aligned}$$

- Substitution rule:

$$\int f(g(x)) g'(x) \, dx = \int f(u) \, du, \quad \text{where, } u = g(x)$$

# Substitution Rule

- Compute the following integral:

$$\begin{aligned}\int 18x^2 \sqrt[4]{6x^3 + 5} \, dx &= \{u = 6x^3 + 5, \quad du = 18x^2 dx\} = \\ &= \int \sqrt[4]{u} \, du = \frac{4}{5} u^{5/4} + C = \frac{4}{5} \sqrt[4]{(6x^3 + 5)^5} + C.\end{aligned}$$

- Substitution rule:

$$\int f(g(x)) g'(x) \, dx = \int f(u) \, du, \quad \text{where, } u = g(x)$$

# Substitution Rule - Example

$$\int 3(8y - 1)e^{4y^2 - y} dy =$$

# Substitution Rule - Example

$$\int 3(8y - 1)e^{4y^2 - y} dy = \int 3e^{4y^2 - y} d(4y^2 - y) =$$

# Substitution Rule - Example

$$\begin{aligned}\int 3(8y - 1)e^{4y^2 - y} dy &= \int 3e^{4y^2 - y} d(4y^2 - y) = \\ &= 3e^{4y^2 - y} + C\end{aligned}$$

# Integration by Parts

- Consider the following integrals:

$$\int e^x dx =$$

$$\int xe^{x^2} dx =$$

$$\int xe^{6x} dx =$$



# Integration by Parts

- Consider the following integrals:

$$\int e^x dx = e^x + C$$

$$\int xe^{x^2} dx =$$

$$\int xe^{6x} dx =$$

# Integration by Parts

- Consider the following integrals:

$$\int e^x dx = e^x + C$$

$$\int x e^{x^2} dx = \frac{1}{2} \int e^{x^2} dx^2 =$$

$$\int x e^{6x} dx =$$

# Integration by Parts

- Consider the following integrals:

$$\int e^x dx = e^x + C$$

$$\int x e^{x^2} dx = \frac{1}{2} \int e^{x^2} dx^2 = \frac{1}{2} e^{x^2} + C$$

$$\int x e^{6x} dx =$$

# Integration by Parts

- Consider the following integrals:

$$\int e^x dx = e^x + C$$

$$\int x e^{x^2} dx = \frac{1}{2} \int e^{x^2} dx^2 = \frac{1}{2} e^{x^2} + C$$

$$\int x e^{6x} dx = \dots ?$$

# Integration by Parts

- Chain rule:

$$(u \cdot v)' = u' \cdot v + u \cdot v'$$

# Integration by Parts

- Chain rule:

$$(u \cdot v)' = u' \cdot v + u \cdot v'$$

$$u \cdot v = \int (u \cdot v)' dx = \int u' \cdot v dx + \int u \cdot v' dx =$$

# Integration by Parts

- Chain rule:

$$(u \cdot v)' = u' \cdot v + u \cdot v'$$

$$u \cdot v = \int (u \cdot v)' dx = \int u' \cdot v dx + \int u \cdot v' dx = \int v du + \int u dv$$

# Integration by Parts

- Chain rule:

$$(u \cdot v)' = u' \cdot v + u \cdot v'$$

$$u \cdot v = \int (u \cdot v)' dx = \int u' \cdot v dx + \int u \cdot v' dx = \int v du + \int u dv$$

$$\int u dv = u \cdot v - \int v du$$



# Integration by Parts

- Consider the following integrals:

$$\int e^x dx = e^x + C$$

$$\int x e^{x^2} dx = \frac{1}{2} \int e^{x^2} dx^2 = \frac{1}{2} e^{x^2} + C$$

$$\int x e^{6x} dx =$$

# Integration by Parts

- Consider the following integrals:

$$\int e^x dx = e^x + C$$

$$\int x e^{x^2} dx = \frac{1}{2} \int e^{x^2} dx^2 = \frac{1}{2} e^{x^2} + C$$

$$\int x e^{6x} dx = \frac{1}{6} \int x d e^{6x} =$$

# Integration by Parts

- Consider the following integrals:

$$\int e^x dx = e^x + C$$

$$\int x e^{x^2} dx = \frac{1}{2} \int e^{x^2} dx^2 = \frac{1}{2} e^{x^2} + C$$

$$\int x e^{6x} dx = \frac{1}{6} \int x de^{6x} = \frac{1}{6} x e^{6x} - \frac{1}{6} \int e^{6x} dx =$$

# Integration by Parts - – Example 2

$$\int x^2 \sin 10x \, dx =$$

# Integration by Parts - – Example 2

$$\int x^2 \sin 10x \, dx = -0.1 \int x^2 d \cos 10x =$$

# Integration by Parts - – Example 2

$$\int x^2 \sin 10x \, dx = -0.1 \int x^2 d \cos 10x =$$

$$-0.1x^2 \cos 10x + 0.1 \int \cos 10x \, dx^2 =$$

# Integration by Parts - – Example 2

$$\int x^2 \sin 10x \, dx = -0.1 \int x^2 d \cos 10x =$$

$$-0.1x^2 \cos 10x + 0.1 \int \cos 10x \, dx^2 = -0.1x^2 \cos 10x + 0.2 \int x \cos 10x \, dx$$

# Integration by Parts - – Example 2

$$\int x^2 \sin 10x \, dx = -0.1 \int x^2 d \cos 10x =$$

$$-0.1x^2 \cos 10x + 0.1 \int \cos 10x \, dx^2 = -0.1x^2 \cos 10x + 0.2 \int x \cos 10x \, dx$$

$$= -0.1x^2 \cos 10x + 0.02 \sin 10x + 0.002 \cos 10x + C$$



# Integration by Parts - – Example 2

$$\int x^2 \sin 10x \, dx = -0.1 \int x^2 d \cos 10x =$$

$$-0.1x^2 \cos 10x + 0.1 \int \cos 10x \, dx^2 = -0.1x^2 \cos 10x + 0.2 \int x \cos 10x \, dx$$

$$= -0.1x^2 \cos 10x + 0.02 \sin 10x + 0.002 \cos 10x + C$$

$$\int x \cos 10x \, dx =$$

# Integration by Parts - – Example 2

$$\int x^2 \sin 10x \, dx = -0.1 \int x^2 d \cos 10x =$$

$$-0.1x^2 \cos 10x + 0.1 \int \cos 10x \, dx^2 = -0.1x^2 \cos 10x + 0.2 \int x \cos 10x \, dx$$

$$= -0.1x^2 \cos 10x + 0.02 \sin 10x + 0.002 \cos 10x + C$$

$$\int x \cos 10x \, dx = 0.1 \int x d \sin 10x = 0.1 \sin 10x - 0.1 \int \sin 10x \, dx$$

=

# Integration by Parts - – Example 2

$$\int x^2 \sin 10x \, dx = -0.1 \int x^2 d \cos 10x =$$

$$-0.1x^2 \cos 10x + 0.1 \int \cos 10x \, dx^2 = -0.1x^2 \cos 10x + 0.2 \int x \cos 10x \, dx$$

$$= -0.1x^2 \cos 10x + 0.02 \sin 10x + 0.002 \cos 10x + C$$

$$\int x \cos 10x \, dx = 0.1 \int x d \sin 10x = 0.1 \sin 10x - 0.1 \int \sin 10x \, dx$$

$$= 0.1 \sin 10x + 0.01 \cos 10x + C.$$

# Integration by Parts – Example 3

$$\int \ln x \, dx =$$

# Integration by Parts – Example 3

$$\int \ln x \, dx =$$

$$= x \ln x - \int x d \ln x =$$

# Integration by Parts – Example 3

$$\int \ln x \, dx =$$

$$= x \ln x - \int x d \ln x = x \ln x - \int x \cdot \frac{1}{x} dx =$$

# Integration by Parts – Example 3

$$\int \ln x \, dx =$$

$$= x \ln x - \int x d \ln x = x \ln x - \int x \cdot \frac{1}{x} dx =$$


$$= x \ln x - x + C$$

# Definite Integral








# Definite Integral






# Definite Integral



# Definite Integral

$$\int_a^b f(x)dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i)\Delta x$$



# Definite Integral

- The fundamental theorem of Calculus:

$$\int_a^b f(x)dx = F(b) - F(a)$$

- Example:

$$\int_0^1 x^2 dx = \frac{1}{3} x^3 \Big|_0^1 = \frac{1}{3} - 0 = \frac{1}{3}$$

# Definite Integral: Properties

1.  $\int_a^b f(x) dx = - \int_b^a f(x) dx$ . We can interchange the limits on any definite integral, all that we need to do is tack a minus sign onto the integral when we do.
2.  $\int_a^a f(x) dx = 0$ . If the upper and lower limits are the same then there is no work to do, the integral is zero.
3.  $\int_a^b cf(x) dx = c \int_a^b f(x) dx$ , where  $c$  is any number. So, as with limits, derivatives, and indefinite integrals we can factor out a constant.
4.  $\int_a^b f(x) \pm g(x) dx = \int_a^b f(x) dx \pm \int_a^b g(x) dx$ . We can break up definite integrals across a sum or difference.
5.  $\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$  where  $c$  is any number. This property is more important than we might realize at first. One of the main uses of this property is to tell us how we can integrate a function over the adjacent intervals,  $[a, c]$  and  $[c, b]$ . Note however that  $c$  doesn't need to be between  $a$  and  $b$ .

# Definite Integral – Example 2

$$\int_0^1 2e^{-2x} dx =$$

# Definite Integral – Example 2

$$\int_0^1 2e^{-2x} dx = -e^{-2x} \Big|_0^1 =$$

# Definite Integral – Example 2

$$\int_0^1 2e^{-2x} dx = -e^{-2x} \Big|_0^1 = -e^{-2} + 1 = 1 - \frac{1}{e^2}$$



# Definite Integral – Example 3

$$\int_{-1}^1 \frac{1}{x^2} dx =$$

# Definite Integral – Example 3


$$\int_{-1}^1 \frac{1}{x^2} dx =$$

$\frac{1}{x^2}$  isn't defined at 0  
Not a definite integral!

# Infinite interval



**girafe**  
**ai**




# Definition



- If  $f(x)$  is continuous on  $[a; +\infty)$ , then

$$\int_a^{+\infty} f(x)dx = \lim_{t \rightarrow +\infty} \int_a^t f(x)dx$$



# Definition




- If  $f(x)$  is continuous on  $[a; +\infty)$ , then

$$\int_a^{+\infty} f(x)dx = \lim_{t \rightarrow +\infty} \int_a^t f(x)dx$$

- If  $f(x)$  is continuous on  $(-\infty; b]$ , then

$$\int_{-\infty}^b f(x)dx = \lim_{t \rightarrow -\infty} \int_t^b f(x)dx$$



# Example

$$\int_1^{+\infty} \frac{1}{x^2} dx =$$

# Example

$$\int_1^{+\infty} \frac{1}{x^2} dx = \lim_{t \rightarrow +\infty} \int_1^t \frac{1}{x^2} dx =$$

# Example

$$\int_1^{+\infty} \frac{1}{x^2} dx = \lim_{t \rightarrow +\infty} \int_1^t \frac{1}{x^2} dx =$$

$$= \lim_{t \rightarrow +\infty} \left( -\frac{1}{x} \Big|_1^t \right) =$$



# Example

$$\begin{aligned}\int_1^{+\infty} \frac{1}{x^2} dx &= \lim_{t \rightarrow +\infty} \int_1^t \frac{1}{x^2} dx = \\ &= \lim_{t \rightarrow +\infty} \left( -\frac{1}{x} \Big|_1^t \right) = \lim_{t \rightarrow +\infty} \left( -\frac{1}{t} \right) + 1 =\end{aligned}$$

# Example

$$\begin{aligned}\int_1^{+\infty} \frac{1}{x^2} dx &= \lim_{t \rightarrow +\infty} \int_1^t \frac{1}{x^2} dx = \\ &= \lim_{t \rightarrow +\infty} \left( -\frac{1}{x} \Big|_1^t \right) = \lim_{t \rightarrow +\infty} \left( -\frac{1}{t} \right) + 1 = \\ &= 0 + 1 = 1\end{aligned}$$

# Divergent integrals

- We call integrals **convergent** if associated limits exist, and **divergent** otherwise.
- Example:

$$\int_1^{+\infty} \frac{1}{x} dx =$$

# Divergent integrals

- We call integrals **convergent** if associated limits exist, and **divergent** otherwise.
- Example:

$$\int_1^{+\infty} \frac{1}{x} dx = \lim_{t \rightarrow +\infty} \int_1^t \frac{1}{x} dx =$$

# Divergent integrals

- We call integrals **convergent** if associated limits exist, and **divergent** otherwise.
- Example:

$$\begin{aligned}\int_1^{+\infty} \frac{1}{x} dx &= \lim_{t \rightarrow +\infty} \int_1^t \frac{1}{x} dx = \\ &= \lim_{t \rightarrow +\infty} \left( \log x \Big|_1^t \right) =\end{aligned}$$

# Divergent integrals

- We call integrals **convergent** if associated limits exist, and **divergent** otherwise.
- Example:

$$\begin{aligned}\int_1^{+\infty} \frac{1}{x} dx &= \lim_{t \rightarrow +\infty} \int_1^t \frac{1}{x} dx = \\ &= \lim_{t \rightarrow +\infty} \left( \log x \Big|_1^t \right) = \\ &= \lim_{t \rightarrow +\infty} \log t + 0\end{aligned}$$

# Divergent integrals

- We call integrals **convergent** if associated limits exist, and **divergent** otherwise.
- Example: the following integral is divergent

$$\begin{aligned}\int_1^{+\infty} \frac{1}{x} dx &= \lim_{t \rightarrow +\infty} \int_1^t \frac{1}{x} dx = \\ &= \lim_{t \rightarrow +\infty} \left( \log x \Big|_1^t \right) = \\ &= \lim_{t \rightarrow +\infty} \log t + 0 \rightarrow +\infty\end{aligned}$$

# One more example

- For which  $p$  is the following integral convergent ( $a > 0$ )?

$$\int_a^{+\infty} \frac{1}{x^p} dx =$$



# One more example

- For which  $p$  is the following integral convergent ( $a > 0$ )?

$$\begin{aligned} \int_a^{+\infty} \frac{1}{x^p} dx &= \\ &= -\frac{1}{p-1} \cdot \lim_{t \rightarrow +\infty} \frac{1}{x^{p-1}} \Big|_a^t = \end{aligned}$$

# One more example

- For which  $p$  is the following integral convergent ( $a > 0$ )?

$$\begin{aligned}\int_a^{+\infty} \frac{1}{x^p} dx &= \\&= -\frac{1}{p-1} \cdot \lim_{t \rightarrow +\infty} \frac{1}{x^{p-1}} \Big|_a^t = \\&= -\frac{1}{p-1} \cdot \lim_{t \rightarrow +\infty} \left( \frac{1}{t^{p-1}} - \frac{1}{a^{p-1}} \right) =\end{aligned}$$

# One more example

- For which  $p$  is the following integral convergent ( $a > 0$ )?

$$\begin{aligned}\int_a^{+\infty} \frac{1}{x^p} dx &= \\&= -\frac{1}{p-1} \cdot \lim_{t \rightarrow +\infty} \frac{1}{x^{p-1}} \Big|_a^t = \\&= -\frac{1}{p-1} \cdot \lim_{t \rightarrow +\infty} \left( \frac{1}{t^{p-1}} - \frac{1}{a^{p-1}} \right) = \\&= \frac{1}{p-1} \cdot \frac{1}{a^{p-1}} \\&\text{when } p-1 > 0 \Leftrightarrow p > 1.\end{aligned}$$

# One more example

- For which  $p$  is the following integral convergent ( $a > 0$ )?

$$\begin{aligned}\int_a^{+\infty} \frac{1}{x^p} dx &= \\&= -\frac{1}{p-1} \cdot \lim_{t \rightarrow +\infty} \frac{1}{x^{p-1}} \Big|_a^t = \\&= -\frac{1}{p-1} \cdot \lim_{t \rightarrow +\infty} \left( \frac{1}{t^{p-1}} - \frac{1}{a^{p-1}} \right) = \\&= \frac{1}{p-1} \cdot \frac{1}{a^{p-1}}\end{aligned}$$

when  $p - 1 > 0 \Leftrightarrow p > 1$ .

If  $p \leq 1$ , the limit doesn't exist.

# Two infinite limits

- If both  $\int_{-\infty}^a f(x)dx$  and  $\int_a^{+\infty} f(x)dx$  are convergent, then the improper integral of  $f$  over  $(-\infty; +\infty)$  is

$$\int_{-\infty}^{+\infty} f(x)dx = \int_{-\infty}^a f(x)dx + \int_a^{+\infty} f(x)dx$$

# Example 2

- Is the following integral convergent or divergent?

$$\int_{-\infty}^{+\infty} x e^{-x^2} dx =$$

## Example 2

- Is the following integral convergent or divergent?

$$\int_{-\infty}^{+\infty} x e^{-x^2} dx = \int_{-\infty}^0 x e^{-x^2} dx + \int_0^{+\infty} x e^{-x^2} dx =$$

# Example 2

- Is the following integral convergent or divergent?

$$\int_{-\infty}^{+\infty} x e^{-x^2} dx = \int_{-\infty}^0 x e^{-x^2} dx + \int_0^{+\infty} x e^{-x^2} dx = \left\{ \begin{array}{l} y = x^2 \\ dy = 2x dx \end{array} \right\} =$$



# Example 2

- Is the following integral convergent or divergent?

$$\begin{aligned}\int_{-\infty}^{+\infty} x e^{-x^2} dx &= \int_{-\infty}^0 x e^{-x^2} dx + \int_0^{+\infty} x e^{-x^2} dx = \left\{ \begin{array}{l} y = x^2 \\ dy = 2x dx \end{array} \right\} = \\ &= \frac{1}{2} \int_{-\infty}^0 e^{-y} dy + \frac{1}{2} \int_0^{+\infty} e^{-y} dy =\end{aligned}$$

# Example 2

- Is the following integral convergent or divergent?

$$\begin{aligned}\int_{-\infty}^{+\infty} x e^{-x^2} dx &= \int_{-\infty}^0 x e^{-x^2} dx + \int_0^{+\infty} x e^{-x^2} dx = \left\{ \begin{array}{l} y = x^2 \\ dy = 2x dx \end{array} \right\} = \\ &= \frac{1}{2} \int_{-\infty}^0 e^{-y} dy + \frac{1}{2} \int_0^{+\infty} e^{-y} dy = \\ &= -\frac{1}{2} e^y \Big|_{-\infty}^0 - \frac{1}{2} e^y \Big|_0^{+\infty} =\end{aligned}$$

# Example 2

- Is the following integral convergent or divergent?

$$\begin{aligned}\int_{-\infty}^{+\infty} x e^{-x^2} dx &= \int_{-\infty}^0 x e^{-x^2} dx + \int_0^{+\infty} x e^{-x^2} dx = \left\{ \begin{array}{l} y = x^2 \\ dy = 2x dx \end{array} \right\} = \\ &= \frac{1}{2} \int_{-\infty}^0 e^{-y} dy + \frac{1}{2} \int_0^{+\infty} e^{-y} dy = \\ &= -\frac{1}{2} e^y \Big|_{-\infty}^0 - \frac{1}{2} e^y \Big|_0^{+\infty} = \\ &= -\frac{1}{2} e^0 + 0 - 0 + \frac{1}{2} e^0 =\end{aligned}$$

# Example 2

- Is the following integral convergent or divergent?



$$\begin{aligned}\int_{-\infty}^{+\infty} x e^{-x^2} dx &= \int_{-\infty}^0 x e^{-x^2} dx + \int_0^{+\infty} x e^{-x^2} dx = \left\{ \begin{array}{l} y = x^2 \\ dy = 2x dx \end{array} \right\} = \\ &= \frac{1}{2} \int_{-\infty}^0 e^{-y} dy + \frac{1}{2} \int_0^{+\infty} e^{-y} dy = \\ &= -\frac{1}{2} e^y \Big|_{-\infty}^0 - \frac{1}{2} e^y \Big|_0^{+\infty} = \\ &= -\frac{1}{2} e^0 + 0 - 0 + \frac{1}{2} e^0 = -\frac{1}{2} + \frac{1}{2} = 0.\end{aligned}$$

# Comparison test

- There are many techniques to check if an integral is convergent or not.
- *Example:* comparison test

Suppose that  $f(x) \geq g(x) \geq 0$  for  $x \geq a$ . Then

- if  $\int_a^{+\infty} f(x)dx$  converges,  
 $\int_a^{+\infty} g(x)dx$  also converges
- if  $\int_a^{+\infty} f(x)dx$  diverges,  
 $\int_a^{+\infty} g(x)dx$  also diverges



# Comparison test - Example

- Check if the following integral converges:

$$\int_2^{+\infty} \frac{\cos^2 x}{x^2} dx$$

# Comparison test - Example

- Check if the following integral converges:

$$\int_2^{+\infty} \frac{\cos^2 x}{x^2} dx$$

Let's use the comparison test:

# Comparison test - Example

- Check if the following integral converges:

$$\int_2^{+\infty} \frac{\cos^2 x}{x^2} dx$$

Let's use the comparison test:

$$0 \leq \frac{\cos^2 x}{x^2} \leq \frac{1}{x^2}$$



# Comparison test - Example

- Check if the following integral converges:

$$\int_2^{+\infty} \frac{\cos^2 x}{x^2} dx$$

Let's use the comparison test:

$$0 \leq \frac{\cos^2 x}{x^2} \leq \frac{1}{x^2}$$

$$\int_2^{+\infty} \frac{1}{x^2} dx \text{ converges} \rightarrow$$

# Comparison test - Example

- Check if the following integral converges:

$$\int_2^{+\infty} \frac{\cos^2 x}{x^2} dx$$

Let's use the comparison test:

$$0 \leq \frac{\cos^2 x}{x^2} \leq \frac{1}{x^2}$$


$$\int_2^{+\infty} \frac{1}{x^2} dx \text{ converges} \rightarrow$$

$$\int_2^{+\infty} \frac{\cos^2 x}{x^2} dx \text{ also converges!}$$

# Discontinuous integrand



**girafe**  
**ai**



# Definition - 1

- If  $f(x)$  is continuous on  $(a; b]$ , then the improper integral of  $f$  over  $[a; b]$  is

$$\int_a^b f(x)dx = \lim_{t \rightarrow a^+} \int_t^b f(x)dx$$

- If  $f(x)$  is continuous on  $[a; b)$ , then the improper integral of  $f$  over  $[a; b]$  is

$$\int_a^b f(x)dx = \lim_{t \rightarrow b^-} \int_a^t f(x)dx$$

# Example - 1

$$\int_0^1 \frac{1}{x^2} dx =$$

# Example - 1

$$\int_0^1 \frac{1}{x^2} dx =$$

$$= \lim_{t \rightarrow 0^+} \int_t^1 \frac{1}{x^2} dx =$$

# Example - 1

$$\int_0^1 \frac{1}{x^2} dx =$$

$$= \lim_{t \rightarrow 0^+} \int_t^1 \frac{1}{x^2} dx =$$

$$= \lim_{t \rightarrow 0^+} \left( -\frac{1}{x} \right) \Big|_t^1 =$$

# Example - 1

$$\int_0^1 \frac{1}{x^2} dx =$$

$$= \lim_{t \rightarrow 0^+} \int_t^1 \frac{1}{x^2} dx =$$

$$= \lim_{t \rightarrow 0^+} \left( -\frac{1}{x} \right) \Big|_t^1 =$$

$$= -1 + \lim_{t \rightarrow 0^+} \frac{1}{t}$$



# Example - 1

$$\int_0^1 \frac{1}{x^2} dx =$$

$$= \lim_{t \rightarrow 0^+} \int_t^1 \frac{1}{x^2} dx =$$

$$= \lim_{t \rightarrow 0^+} \left( -\frac{1}{x} \right) \Big|_t^1 =$$

$$= -1 + \lim_{t \rightarrow 0^+} \frac{1}{t} \rightarrow \infty$$

# Definition - 2

- If  $f(x)$  has a discontinuity at  $x = c \in [a; b]$ , and both  $\int_a^c f(x)dx$  and  $\int_c^b f(x)dx$  are convergent, then

$$\int_a^b f(x)dx = \int_a^c f(x)dx + \int_c^b f(x)dx$$

# Example - 2

$$\int_0^{+\infty} \frac{1}{x^2} dx =$$

## Example - 2

$$\begin{aligned} \int_0^{+\infty} \frac{1}{x^2} dx &= \\ &= \int_0^1 \frac{1}{x^2} dx + \int_1^{+\infty} \frac{1}{x^2} dx = \end{aligned}$$

# Example - 2

$$\begin{aligned}\int_0^{+\infty} \frac{1}{x^2} dx &= \\&= \int_0^1 \frac{1}{x^2} dx + \int_1^{+\infty} \frac{1}{x^2} dx = \\&= \int_0^1 \frac{1}{x^2} dx + 1\end{aligned}$$

## Example - 2

$$\begin{aligned}\int_0^{+\infty} \frac{1}{x^2} dx &= \\&= \int_0^1 \frac{1}{x^2} dx + \int_1^{+\infty} \frac{1}{x^2} dx = \\&= \int_0^1 \frac{1}{x^2} dx + 1\end{aligned}$$

$$\int_0^1 \frac{1}{x^2} dx =$$

## Example - 2

$$\begin{aligned}\int_0^{+\infty} \frac{1}{x^2} dx &= \\&= \int_0^1 \frac{1}{x^2} dx + \int_1^{+\infty} \frac{1}{x^2} dx = \\&= \int_0^1 \frac{1}{x^2} dx + 1\end{aligned}$$

$$\int_0^1 \frac{1}{x^2} dx = \lim_{t \rightarrow 0^+} \left( -\frac{1}{x} \Big|_t^1 \right) =$$

## Example - 2

$$\begin{aligned}\int_0^{+\infty} \frac{1}{x^2} dx &= \\&= \int_0^1 \frac{1}{x^2} dx + \int_1^{+\infty} \frac{1}{x^2} dx = \\&= \int_0^1 \frac{1}{x^2} dx + 1\end{aligned}$$

$$\begin{aligned}\int_0^1 \frac{1}{x^2} dx &= \lim_{t \rightarrow 0^+} \left( -\frac{1}{x} \Big|_t^1 \right) = \\&= \lim_{t \rightarrow 0^+} \left( -1 + \frac{1}{t} \right)\end{aligned}$$



## Example - 2

$$\begin{aligned}\int_0^{+\infty} \frac{1}{x^2} dx &= \\&= \int_0^1 \frac{1}{x^2} dx + \int_1^{+\infty} \frac{1}{x^2} dx = \\&= \int_0^1 \frac{1}{x^2} dx + 1\end{aligned}$$

$$\begin{aligned}\int_0^1 \frac{1}{x^2} dx &= \lim_{t \rightarrow 0^+} \left( -\frac{1}{x} \Big|_t^1 \right) = \\&= \lim_{t \rightarrow 0^+} \left( -1 + \frac{1}{t} \right) \rightarrow \infty\end{aligned}$$

## Example - 2

$$\begin{aligned}\int_0^{+\infty} \frac{1}{x^2} dx &= \\&= \int_0^1 \frac{1}{x^2} dx + \int_1^{+\infty} \frac{1}{x^2} dx = \\&= \int_0^1 \frac{1}{x^2} dx + 1 \rightarrow \infty\end{aligned}$$

$$\begin{aligned}\int_0^1 \frac{1}{x^2} dx &= \lim_{t \rightarrow 0^+} \left( -\frac{1}{x} \Big|_t^1 \right) = \\&= \lim_{t \rightarrow 0^+} \left( -1 + \frac{1}{t} \right) \rightarrow \infty\end{aligned}$$