

Optimization methods

Lecture 3: Optimality conditions

Alexandr Katrutsa

Modern State of Artificial Intelligence Masters Program
Moscow Institute of Physics and Technology

Brief reminder of the previous lecture

- ▶ Matrix calculus

Brief reminder of the previous lecture

- ▶ Matrix calculus
- ▶ Non-differentiable convex functions and subdifferential

Brief reminder of the previous lecture

- ▶ Matrix calculus
- ▶ Non-differentiable convex functions and subdifferential
- ▶ Automatic differentiation technique

Brief reminder of the previous lecture

- ▶ Matrix calculus
- ▶ Non-differentiable convex functions and subdifferential
- ▶ Automatic differentiation technique
- ▶ L -smooth convex functions

Brief reminder of the previous lecture

- ▶ Matrix calculus
- ▶ Non-differentiable convex functions and subdifferential
- ▶ Automatic differentiation technique
- ▶ L -smooth convex functions
- ▶ Strongly convex functions properties

Plan for today

- ▶ Optimality conditions: general concept

Plan for today

- ▶ Optimality conditions: general concept
- ▶ First order optimality condition

Plan for today

- ▶ Optimality conditions: general concept
- ▶ First order optimality condition
- ▶ Equality constrained optimization problems

Plan for today

- ▶ Optimality conditions: general concept
- ▶ First order optimality condition
- ▶ Equality constrained optimization problems
- ▶ KKT optimality conditions

Plan for today

- ▶ Optimality conditions: general concept
- ▶ First order optimality condition
- ▶ Equality constrained optimization problems
- ▶ KKT optimality conditions
- ▶ Slater regularity conditions

Motivation

Question 1

In what cases a solution of an optimization problem exists?

Motivation

Question 1

In what cases a solution of an optimization problem exists?

Question 2

How to verify that the point is not a solution?

Motivation

Question 1

In what cases a solution of an optimization problem exists?

Question 2

How to verify that the point is not a solution?

Question 3

How to find a solution of an optimization problem?

Existence

Let $\mathcal{X} \subset \mathbb{R}^n$ be a compact set and let $f : \mathcal{X} \rightarrow \mathbb{R}$ is continuous function in \mathcal{X} . Then the point of global minimum does exist in \mathcal{X} .

- To be sure that we find solution, we need compact set

Existence

Let $\mathcal{X} \subset \mathbb{R}^n$ be a compact set and let $f : \mathcal{X} \rightarrow \mathbb{R}$ is continuous function in \mathcal{X} . Then the point of global minimum does exist in \mathcal{X} .

- ▶ To be sure that we find solution, we need compact set
- ▶ Most problems are defined exactly in the compact sets

Existence

Let $\mathcal{X} \subset \mathbb{R}^n$ be a compact set and let $f : \mathcal{X} \rightarrow \mathbb{R}$ is continuous function in \mathcal{X} . Then the point of global minimum does exist in \mathcal{X} .

- ▶ To be sure that we find solution, we need compact set
- ▶ Most problems are defined exactly in the compact sets
- ▶ The source of open feasible sets is domain of some convex functions

Existence

Let $\mathcal{X} \subset \mathbb{R}^n$ be a compact set and let $f : \mathcal{X} \rightarrow \mathbb{R}$ is continuous function in \mathcal{X} . Then the point of global minimum does exist in \mathcal{X} .

- ▶ To be sure that we find solution, we need compact set
- ▶ Most problems are defined exactly in the compact sets
- ▶ The source of open feasible sets is domain of some convex functions
- ▶ This issue can be fixed by sequential approximation of the open feasible set by the compact one

General form of optimality conditions

Optimization problem in general form

$$f(\mathbf{x}) \rightarrow \min_{\mathbf{x} \in \mathcal{X}}$$

General form of optimality conditions

Optimization problem in general form

$$f(\mathbf{x}) \rightarrow \min_{\mathbf{x} \in \mathcal{X}}$$

Optimality criterion

Let f be defined in a set $\mathcal{X} \subset \mathbb{R}^n$. Then

General form of optimality conditions

Optimization problem in general form

$$f(\mathbf{x}) \rightarrow \min_{\mathbf{x} \in \mathcal{X}}$$

Optimality criterion

Let f be defined in a set $\mathcal{X} \subset \mathbb{R}^n$. Then

1. If \mathbf{x}^* is a minimizer of f in \mathcal{X} , then $0 \in \partial_{\mathcal{X}} f(\mathbf{x}^*)$

General form of optimality conditions

Optimization problem in general form

$$f(\mathbf{x}) \rightarrow \min_{\mathbf{x} \in \mathcal{X}}$$

Optimality criterion

Let f be defined in a set $\mathcal{X} \subset \mathbb{R}^n$. Then

1. If \mathbf{x}^* is a minimizer of f in \mathcal{X} , then $0 \in \partial_{\mathcal{X}} f(\mathbf{x}^*)$
2. If for some point $\mathbf{x}^* \in \mathcal{X}$ there exists subdifferential $\partial_{\mathcal{X}} f(\mathbf{x}^*)$ and $0 \in \partial_{\mathcal{X}} f(\mathbf{x}^*)$, then \mathbf{x}^* is a minimizer of f in \mathcal{X} .

General form of optimality conditions

Optimization problem in general form

$$f(\mathbf{x}) \rightarrow \min_{\mathbf{x} \in \mathcal{X}}$$

Optimality criterion

Let f be defined in a set $\mathcal{X} \subset \mathbb{R}^n$. Then

1. If \mathbf{x}^* is a minimizer of f in \mathcal{X} , then $0 \in \partial_{\mathcal{X}} f(\mathbf{x}^*)$
2. If for some point $\mathbf{x}^* \in \mathcal{X}$ there exists subdifferential $\partial_{\mathcal{X}} f(\mathbf{x}^*)$ and $0 \in \partial_{\mathcal{X}} f(\mathbf{x}^*)$, then \mathbf{x}^* is a minimizer of f in \mathcal{X} .

Q: what drawbacks this criterion has?

Proof

1. If \mathbf{x}^* is a minimizer

Proof

1. If \mathbf{x}^* is a minimizer

- For any $\mathbf{x} \in \mathcal{X}$: $f(\mathbf{x}) \geq f(\mathbf{x}^*)$

1. If \mathbf{x}^* is a minimizer

- ▶ For any $\mathbf{x} \in \mathcal{X}$: $f(\mathbf{x}) \geq f(\mathbf{x}^*)$
- ▶ This inequality can be re-written in the form:
$$f(\mathbf{x}) \geq f(\mathbf{x}^*) + \langle 0, \mathbf{x} - \mathbf{x}^* \rangle$$

1. If \mathbf{x}^* is a minimizer

- ▶ For any $\mathbf{x} \in \mathcal{X}$: $f(\mathbf{x}) \geq f(\mathbf{x}^*)$
- ▶ This inequality can be re-written in the form:
$$f(\mathbf{x}) \geq f(\mathbf{x}^*) + \langle 0, \mathbf{x} - \mathbf{x}^* \rangle$$
- ▶ It means that $0 \in \partial_{\mathcal{X}} f(\mathbf{x}^*)$ by definition

1. If \mathbf{x}^* is a minimizer
 - ▶ For any $\mathbf{x} \in \mathcal{X}$: $f(\mathbf{x}) \geq f(\mathbf{x}^*)$
 - ▶ This inequality can be re-written in the form:
$$f(\mathbf{x}) \geq f(\mathbf{x}^*) + \langle 0, \mathbf{x} - \mathbf{x}^* \rangle$$
 - ▶ It means that $0 \in \partial_{\mathcal{X}} f(\mathbf{x}^*)$ by definition
2. If $0 \in \partial_{\mathcal{X}} f(\mathbf{x}^*)$ for some point \mathbf{x}^*

Proof

1. If \mathbf{x}^* is a minimizer

- ▶ For any $\mathbf{x} \in \mathcal{X}$: $f(\mathbf{x}) \geq f(\mathbf{x}^*)$
- ▶ This inequality can be re-written in the form:
$$f(\mathbf{x}) \geq f(\mathbf{x}^*) + \langle 0, \mathbf{x} - \mathbf{x}^* \rangle$$
- ▶ It means that $0 \in \partial_{\mathcal{X}} f(\mathbf{x}^*)$ by definition

2. If $0 \in \partial_{\mathcal{X}} f(\mathbf{x}^*)$ for some point \mathbf{x}^*

- ▶ By definition of subgradient: $f(\mathbf{x}) \geq f(\mathbf{x}^*) + \langle \mathbf{a}, \mathbf{x} - \mathbf{x}^* \rangle$

1. If \mathbf{x}^* is a minimizer

- ▶ For any $\mathbf{x} \in \mathcal{X}$: $f(\mathbf{x}) \geq f(\mathbf{x}^*)$
- ▶ This inequality can be re-written in the form:
$$f(\mathbf{x}) \geq f(\mathbf{x}^*) + \langle 0, \mathbf{x} - \mathbf{x}^* \rangle$$
- ▶ It means that $0 \in \partial_{\mathcal{X}} f(\mathbf{x}^*)$ by definition

2. If $0 \in \partial_{\mathcal{X}} f(\mathbf{x}^*)$ for some point \mathbf{x}^*

- ▶ By definition of subgradient: $f(\mathbf{x}) \geq f(\mathbf{x}^*) + \langle \mathbf{a}, \mathbf{x} - \mathbf{x}^* \rangle$
- ▶ If $0 \in \partial_{\mathcal{X}} f(\mathbf{x}^*)$, then we have inequality $f(\mathbf{x}) \geq f(\mathbf{x}^*)$

1. If \mathbf{x}^* is a minimizer

- ▶ For any $\mathbf{x} \in \mathcal{X}$: $f(\mathbf{x}) \geq f(\mathbf{x}^*)$
- ▶ This inequality can be re-written in the form:
$$f(\mathbf{x}) \geq f(\mathbf{x}^*) + \langle 0, \mathbf{x} - \mathbf{x}^* \rangle$$
- ▶ It means that $0 \in \partial_{\mathcal{X}} f(\mathbf{x}^*)$ by definition

2. If $0 \in \partial_{\mathcal{X}} f(\mathbf{x}^*)$ for some point \mathbf{x}^*

- ▶ By definition of subgradient: $f(\mathbf{x}) \geq f(\mathbf{x}^*) + \langle \mathbf{a}, \mathbf{x} - \mathbf{x}^* \rangle$
- ▶ If $0 \in \partial_{\mathcal{X}} f(\mathbf{x}^*)$, then we have inequality $f(\mathbf{x}) \geq f(\mathbf{x}^*)$
- ▶ It means that \mathbf{x}^* is a minimizer

Geometric interpretation

Necessary condition for unconstrained problem

$$\min_{\mathbf{x} \in \mathbb{R}^n} f(\mathbf{x}) \quad (1)$$

Theorem

If \mathbf{x}^* is a solution of problem (1) and f is differentiable, then $f'(\mathbf{x}^*) = 0$.

Necessary condition for unconstrained problem

$$\min_{\mathbf{x} \in \mathbb{R}^n} f(\mathbf{x}) \quad (1)$$

Theorem

If \mathbf{x}^* is a solution of problem (1) and f is differentiable, then $f'(\mathbf{x}^*) = 0$.

Proof

Necessary condition for unconstrained problem

$$\min_{\mathbf{x} \in \mathbb{R}^n} f(\mathbf{x}) \quad (1)$$

Theorem

If \mathbf{x}^* is a solution of problem (1) and f is differentiable, then $f'(\mathbf{x}^*) = 0$.

Proof

$$\begin{aligned} \blacktriangleright \quad & f(\mathbf{y}) = f(\mathbf{x}^*) + \langle f'(\mathbf{x}^*), \mathbf{y} - \mathbf{x}^* \rangle + r(\mathbf{x}^*, \mathbf{y}) \text{ and} \\ & \lim_{\mathbf{y} \rightarrow \mathbf{x}^*} \frac{r(\mathbf{x}^*, \mathbf{y})}{\|\mathbf{x}^* - \mathbf{y}\|_2} = 0 \quad (*) \end{aligned}$$

Necessary condition for unconstrained problem

$$\min_{\mathbf{x} \in \mathbb{R}^n} f(\mathbf{x}) \quad (1)$$

Theorem

If \mathbf{x}^* is a solution of problem (1) and f is differentiable, then $f'(\mathbf{x}^*) = 0$.

Proof

- ▶ $f(\mathbf{y}) = f(\mathbf{x}^*) + \langle f'(\mathbf{x}^*), \mathbf{y} - \mathbf{x}^* \rangle + r(\mathbf{x}^*, \mathbf{y})$ and $\lim_{\mathbf{y} \rightarrow \mathbf{x}^*} \frac{r(\mathbf{x}^*, \mathbf{y})}{\|\mathbf{x}^* - \mathbf{y}\|_2} = 0$ (*)
- ▶ If $f'(\mathbf{x}^*) \neq 0$, then consider $\mathbf{y}(\tau) = \mathbf{x}^* - \tau f'(\mathbf{x}^*)$, $\tau > 0$

Necessary condition for unconstrained problem

$$\min_{\mathbf{x} \in \mathbb{R}^n} f(\mathbf{x}) \quad (1)$$

Theorem

If \mathbf{x}^* is a solution of problem (1) and f is differentiable, then $f'(\mathbf{x}^*) = 0$.

Proof

- ▶ $f(\mathbf{y}) = f(\mathbf{x}^*) + \langle f'(\mathbf{x}^*), \mathbf{y} - \mathbf{x}^* \rangle + r(\mathbf{x}^*, \mathbf{y})$ and $\lim_{\mathbf{y} \rightarrow \mathbf{x}^*} \frac{r(\mathbf{x}^*, \mathbf{y})}{\|\mathbf{x}^* - \mathbf{y}\|_2} = 0$ (*)
- ▶ If $f'(\mathbf{x}^*) \neq 0$, then consider $\mathbf{y}(\tau) = \mathbf{x}^* - \tau f'(\mathbf{x}^*)$, $\tau > 0$
- ▶ $f(\mathbf{y}(\tau)) = f(\mathbf{x}^*) + \langle f'(\mathbf{x}^*), \mathbf{y}(\tau) - \mathbf{x}^* \rangle + r(\mathbf{x}^*, \mathbf{y}(\tau)) = f(\mathbf{x}^*) - \tau \|f'(\mathbf{x}^*)\|_2^2 + r(\mathbf{x}^*, \mathbf{y}(\tau))$

Necessary condition for unconstrained problem

$$\min_{\mathbf{x} \in \mathbb{R}^n} f(\mathbf{x}) \quad (1)$$

Theorem

If \mathbf{x}^* is a solution of problem (1) and f is differentiable, then $f'(\mathbf{x}^*) = 0$.

Proof

- ▶ $f(\mathbf{y}) = f(\mathbf{x}^*) + \langle f'(\mathbf{x}^*), \mathbf{y} - \mathbf{x}^* \rangle + r(\mathbf{x}^*, \mathbf{y})$ and
$$\lim_{\mathbf{y} \rightarrow \mathbf{x}^*} \frac{r(\mathbf{x}^*, \mathbf{y})}{\|\mathbf{x}^* - \mathbf{y}\|_2} = 0 \quad (*)$$
- ▶ If $f'(\mathbf{x}^*) \neq 0$, then consider $\mathbf{y}(\tau) = \mathbf{x}^* - \tau f'(\mathbf{x}^*)$, $\tau > 0$
- ▶
$$f(\mathbf{y}(\tau)) = f(\mathbf{x}^*) + \langle f'(\mathbf{x}^*), \mathbf{y}(\tau) - \mathbf{x}^* \rangle + r(\mathbf{x}^*, \mathbf{y}(\tau)) = f(\mathbf{x}^*) - \tau \|f'(\mathbf{x}^*)\|_2^2 + r(\mathbf{x}^*, \mathbf{y}(\tau))$$
- ▶ From (*) follows that there exists $\bar{\tau}$ such that for all $\tau \in (0, \bar{\tau})$ holds $f(\mathbf{y}(\tau)) - f(\mathbf{x}^*) \leq -\frac{\tau}{2} \|f'(\mathbf{x}^*)\|_2^2 < 0$

Necessary condition for unconstrained problem

$$\min_{\mathbf{x} \in \mathbb{R}^n} f(\mathbf{x}) \quad (1)$$

Theorem

If \mathbf{x}^* is a solution of problem (1) and f is differentiable, then $f'(\mathbf{x}^*) = 0$.

Proof

- ▶ $f(\mathbf{y}) = f(\mathbf{x}^*) + \langle f'(\mathbf{x}^*), \mathbf{y} - \mathbf{x}^* \rangle + r(\mathbf{x}^*, \mathbf{y})$ and $\lim_{\mathbf{y} \rightarrow \mathbf{x}^*} \frac{r(\mathbf{x}^*, \mathbf{y})}{\|\mathbf{x}^* - \mathbf{y}\|_2} = 0$ (*)
- ▶ If $f'(\mathbf{x}^*) \neq 0$, then consider $\mathbf{y}(\tau) = \mathbf{x}^* - \tau f'(\mathbf{x}^*)$, $\tau > 0$
- ▶ $f(\mathbf{y}(\tau)) = f(\mathbf{x}^*) + \langle f'(\mathbf{x}^*), \mathbf{y}(\tau) - \mathbf{x}^* \rangle + r(\mathbf{x}^*, \mathbf{y}(\tau)) = f(\mathbf{x}^*) - \tau \|f'(\mathbf{x}^*)\|_2^2 + r(\mathbf{x}^*, \mathbf{y}(\tau))$
- ▶ From (*) follows that there exists $\bar{\tau}$ such that for all $\tau \in (0, \bar{\tau})$ holds $f(\mathbf{y}(\tau)) - f(\mathbf{x}^*) \leq -\frac{\tau}{2} \|f'(\mathbf{x}^*)\|_2^2 < 0$
- ▶ Thus, \mathbf{x}^* is not a minimizer, that is a contradiction.

Remarks

- ▶ This theorem is a special case of the general optimality condition

Remarks

- ▶ This theorem is a special case of the general optimality condition
- ▶ The additional proof is provided to introduce the fact on the descent property of the direction $-f'(\mathbf{x})$

If function is convex...

Theorem

If the objective function in problem (1) is convex, then \mathbf{x}^* is a global minimum iff $f'(\mathbf{x}^*) = 0$

If function is convex...

Theorem

If the objective function in problem (1) is convex, then \mathbf{x}^* is a global minimum iff $f'(\mathbf{x}^*) = 0$

Proof

If function is convex...

Theorem

If the objective function in problem (1) is convex, then \mathbf{x}^* is a global minimum iff $f'(\mathbf{x}^*) = 0$

Proof

- If \mathbf{x}^* is a global minimum, then \mathbf{x}^* is local minimum

If function is convex...

Theorem

If the objective function in problem (1) is convex, then \mathbf{x}^* is a global minimum iff $f'(\mathbf{x}^*) = 0$

Proof

- ▶ If \mathbf{x}^* is a global minimum, then \mathbf{x}^* is local minimum
- ▶ Thus, $f'(\mathbf{x}^*) = 0$ according to the previous theorem

If function is convex...

Theorem

If the objective function in problem (1) is convex, then \mathbf{x}^* is a global minimum iff $f'(\mathbf{x}^*) = 0$

Proof

- ▶ If \mathbf{x}^* is a global minimum, then \mathbf{x}^* is local minimum
- ▶ Thus, $f'(\mathbf{x}^*) = 0$ according to the previous theorem
- ▶ Let \mathbf{x}^* be a point such that $f'(\mathbf{x}^*) = 0$ and f is convex

If function is convex...

Theorem

If the objective function in problem (1) is convex, then \mathbf{x}^* is a global minimum iff $f'(\mathbf{x}^*) = 0$

Proof

- ▶ If \mathbf{x}^* is a global minimum, then \mathbf{x}^* is local minimum
- ▶ Thus, $f'(\mathbf{x}^*) = 0$ according to the previous theorem
- ▶ Let \mathbf{x}^* be a point such that $f'(\mathbf{x}^*) = 0$ and f is convex
- ▶ Then according to the FO criterion

$$f(\mathbf{y}) \geq f(\mathbf{x}^*) + \langle f'(\mathbf{x}^*), \mathbf{y} - \mathbf{x}^* \rangle = f(\mathbf{x}^*)$$

If function is convex...

Theorem

If the objective function in problem (1) is convex, then \mathbf{x}^* is a global minimum iff $f'(\mathbf{x}^*) = 0$

Proof

- ▶ If \mathbf{x}^* is a global minimum, then \mathbf{x}^* is local minimum
- ▶ Thus, $f'(\mathbf{x}^*) = 0$ according to the previous theorem
- ▶ Let \mathbf{x}^* be a point such that $f'(\mathbf{x}^*) = 0$ and f is convex
- ▶ Then according to the FO criterion

$$f(\mathbf{y}) \geq f(\mathbf{x}^*) + \langle f'(\mathbf{x}^*), \mathbf{y} - \mathbf{x}^* \rangle = f(\mathbf{x}^*)$$

- ▶ It means that \mathbf{x}^* is a global minimum

Second-order sufficient condition

Theorem

Let f be twice continuously differentiable function. A point \mathbf{x}^* satisfies equation $f'(\mathbf{x}^*) = 0$. If $\mathbf{s}^\top f''(\mathbf{x}^*)\mathbf{s} > 0$ for all $\mathbf{s} \neq 0$, then \mathbf{x}^* is a point of local minimum.

Proof by contradiction

Second-order sufficient condition

Theorem

Let f be twice continuously differentiable function. A point \mathbf{x}^* satisfies equation $f'(\mathbf{x}^*) = 0$. If $\mathbf{s}^\top f''(\mathbf{x}^*)\mathbf{s} > 0$ for all $\mathbf{s} \neq 0$, then \mathbf{x}^* is a point of local minimum.

Proof by contradiction

- Assume that there exists some point \mathbf{y} close to \mathbf{x}^* such that $f(\mathbf{y}) < f(\mathbf{x}^*)$

Second-order sufficient condition

Theorem

Let f be twice continuously differentiable function. A point \mathbf{x}^* satisfies equation $f'(\mathbf{x}^*) = 0$. If $\mathbf{s}^\top f''(\mathbf{x}^*)\mathbf{s} > 0$ for all $\mathbf{s} \neq 0$, then \mathbf{x}^* is a point of local minimum.

Proof by contradiction

- ▶ Assume that there exists some point \mathbf{y} close to \mathbf{x}^* such that $f(\mathbf{y}) < f(\mathbf{x}^*)$
- ▶ Then consider Taylor expansion $f(\mathbf{y}) = f(\mathbf{x}^*) + \langle f'(\mathbf{x}^*), \mathbf{y} - \mathbf{x}^* \rangle + \frac{1}{2} \langle \mathbf{y} - \mathbf{x}^*, f''(\mathbf{x}^*)(\mathbf{y} - \mathbf{x}^*) \rangle + o(\|\mathbf{y} - \mathbf{x}^*\|_2^2)$

Second-order sufficient condition

Theorem

Let f be twice continuously differentiable function. A point \mathbf{x}^* satisfies equation $f'(\mathbf{x}^*) = 0$. If $\mathbf{s}^\top f''(\mathbf{x}^*)\mathbf{s} > 0$ for all $\mathbf{s} \neq 0$, then \mathbf{x}^* is a point of local minimum.

Proof by contradiction

- ▶ Assume that there exists some point \mathbf{y} close to \mathbf{x}^* such that $f(\mathbf{y}) < f(\mathbf{x}^*)$
- ▶ Then consider Taylor expansion $f(\mathbf{y}) = f(\mathbf{x}^*) + \langle f'(\mathbf{x}^*), \mathbf{y} - \mathbf{x} \rangle + \frac{1}{2} \langle \mathbf{y} - \mathbf{x}, f''(\mathbf{x}^*)(\mathbf{y} - \mathbf{x}) \rangle + o(\|\mathbf{y} - \mathbf{x}\|_2^2)$
- ▶ From stationarity follows that $f(\mathbf{y}) - f(\mathbf{x}^*) = \frac{1}{2} \langle \mathbf{y} - \mathbf{x}, f''(\mathbf{x}^*)(\mathbf{y} - \mathbf{x}) \rangle + o(\|\mathbf{y} - \mathbf{x}\|_2^2)$

Second-order sufficient condition

Theorem

Let f be twice continuously differentiable function. A point \mathbf{x}^* satisfies equation $f'(\mathbf{x}^*) = 0$. If $\mathbf{s}^\top f''(\mathbf{x}^*)\mathbf{s} > 0$ for all $\mathbf{s} \neq 0$, then \mathbf{x}^* is a point of local minimum.

Proof by contradiction

- ▶ Assume that there exists some point \mathbf{y} close to \mathbf{x}^* such that $f(\mathbf{y}) < f(\mathbf{x}^*)$
- ▶ Then consider Taylor expansion $f(\mathbf{y}) = f(\mathbf{x}^*) + \langle f'(\mathbf{x}^*), \mathbf{y} - \mathbf{x} \rangle + \frac{1}{2} \langle \mathbf{y} - \mathbf{x}, f''(\mathbf{x}^*)(\mathbf{y} - \mathbf{x}) \rangle + o(\|\mathbf{y} - \mathbf{x}\|_2^2)$
- ▶ From stationarity follows that $f(\mathbf{y}) - f(\mathbf{x}^*) = \frac{1}{2} \langle \mathbf{y} - \mathbf{x}, f''(\mathbf{x}^*)(\mathbf{y} - \mathbf{x}) \rangle + o(\|\mathbf{y} - \mathbf{x}\|_2^2)$
- ▶ If $\mathbf{y} \rightarrow \mathbf{x}^*$, then we have a direction $\mathbf{z} \neq 0$ such that $\mathbf{z}^\top f''(\mathbf{x}^*)\mathbf{z} \leq 0$, that is contradiction

Saddle points

Definition

A point \mathbf{y} is called saddle point for a function f if there are directions \mathbf{z}_1 and \mathbf{z}_2 such that $f(\mathbf{y} + \mathbf{z}_1) > f(\mathbf{y})$, but $f(\mathbf{y} + \mathbf{z}_2) < f(\mathbf{y})$

Summary on unconstrained problems

- ▶ Use subdifferential for non-differentiable case

Summary on unconstrained problems

- ▶ Use subdifferential for non-differentiable case
- ▶ Use FOOC for convex differentiable function

Summary on unconstrained problems

- ▶ Use subdifferential for non-differentiable case
- ▶ Use FOOC for convex differentiable function
- ▶ Use second order sufficient condition for non-convex twice continuously differentiable

Summary on unconstrained problems

- ▶ Use subdifferential for non-differentiable case
- ▶ Use FOOC for convex differentiable function
- ▶ Use second order sufficient condition for non-convex twice continuously differentiable
- ▶ Saddle points are possible in the non-convex settings

Differential criterion for constrained problem

$$\min_{\mathbf{x} \in \mathcal{X}} f(\mathbf{x}) \quad (2)$$

Theorem

A point \mathbf{x}^* is a solution of the problem (2), where f is convex function, iff $\mathbf{x}^* \in \mathcal{X}$ and $\langle f'(\mathbf{x}^*), \mathbf{y} - \mathbf{x}^* \rangle \geq 0$ for all $\mathbf{y} \in \mathcal{X}$.

Differential criterion for constrained problem

$$\min_{\mathbf{x} \in \mathcal{X}} f(\mathbf{x}) \quad (2)$$

Theorem

A point \mathbf{x}^* is a solution of the problem (2), where f is convex function, iff $\mathbf{x}^* \in \mathcal{X}$ and $\langle f'(\mathbf{x}^*), \mathbf{y} - \mathbf{x}^* \rangle \geq 0$ for all $\mathbf{y} \in \mathcal{X}$.

Proof

Differential criterion for constrained problem

$$\min_{\mathbf{x} \in \mathcal{X}} f(\mathbf{x}) \quad (2)$$

Theorem

A point \mathbf{x}^* is a solution of the problem (2), where f is convex function, iff $\mathbf{x}^* \in \mathcal{X}$ and $\langle f'(\mathbf{x}^*), \mathbf{y} - \mathbf{x}^* \rangle \geq 0$ for all $\mathbf{y} \in \mathcal{X}$.

Proof

- ▶ Let $\mathbf{x}^* \in \mathcal{X}$ and the inequality holds. Then according to the first order criterion of convexity function f : $f(\mathbf{y}) \geq f(\mathbf{x}^*)$

Differential criterion for constrained problem

$$\min_{\mathbf{x} \in \mathcal{X}} f(\mathbf{x}) \quad (2)$$

Theorem

A point \mathbf{x}^* is a solution of the problem (2), where f is convex function, iff $\mathbf{x}^* \in \mathcal{X}$ and $\langle f'(\mathbf{x}^*), \mathbf{y} - \mathbf{x}^* \rangle \geq 0$ for all $\mathbf{y} \in \mathcal{X}$.

Proof

- ▶ Let $\mathbf{x}^* \in \mathcal{X}$ and the inequality holds. Then according to the first order criterion of convexity function f : $f(\mathbf{y}) \geq f(\mathbf{x}^*)$
- ▶ Let \mathbf{x}^* be a solution of the problem (2), but there exists $\tilde{\mathbf{y}}$ such that $\langle f'(\mathbf{x}^*), \tilde{\mathbf{y}} - \mathbf{x}^* \rangle < 0$

Differential criterion for constrained problem

$$\min_{\mathbf{x} \in \mathcal{X}} f(\mathbf{x}) \quad (2)$$

Theorem

A point \mathbf{x}^* is a solution of the problem (2), where f is convex function, iff $\mathbf{x}^* \in \mathcal{X}$ and $\langle f'(\mathbf{x}^*), \mathbf{y} - \mathbf{x}^* \rangle \geq 0$ for all $\mathbf{y} \in \mathcal{X}$.

Proof

- ▶ Let $\mathbf{x}^* \in \mathcal{X}$ and the inequality holds. Then according to the first order criterion of convexity function f : $f(\mathbf{y}) \geq f(\mathbf{x}^*)$
- ▶ Let \mathbf{x}^* be a solution of the problem (2), but there exists $\tilde{\mathbf{y}}$ such that $\langle f'(\mathbf{x}^*), \tilde{\mathbf{y}} - \mathbf{x}^* \rangle < 0$
- ▶ Consider a point $\mathbf{z}(t) = t\tilde{\mathbf{y}} + (1 - t)\mathbf{x}^*$, $t \in [0, 1]$

Differential criterion for constrained problem

$$\min_{\mathbf{x} \in \mathcal{X}} f(\mathbf{x}) \quad (2)$$

Theorem

A point \mathbf{x}^* is a solution of the problem (2), where f is convex function, iff $\mathbf{x}^* \in \mathcal{X}$ and $\langle f'(\mathbf{x}^*), \mathbf{y} - \mathbf{x}^* \rangle \geq 0$ for all $\mathbf{y} \in \mathcal{X}$.

Proof

- ▶ Let $\mathbf{x}^* \in \mathcal{X}$ and the inequality holds. Then according to the first order criterion of convexity function f : $f(\mathbf{y}) \geq f(\mathbf{x}^*)$
- ▶ Let \mathbf{x}^* be a solution of the problem (2), but there exists $\tilde{\mathbf{y}}$ such that $\langle f'(\mathbf{x}^*), \tilde{\mathbf{y}} - \mathbf{x}^* \rangle < 0$
- ▶ Consider a point $\mathbf{z}(t) = t\tilde{\mathbf{y}} + (1 - t)\mathbf{x}^*$, $t \in [0, 1]$
- ▶ Then $\left. \frac{d}{dt} f(\mathbf{z}(t)) \right|_{t=0} = \langle f'(\mathbf{x}^*), \tilde{\mathbf{y}} - \mathbf{x}^* \rangle < 0$

Differential criterion for constrained problem

$$\min_{\mathbf{x} \in \mathcal{X}} f(\mathbf{x}) \quad (2)$$

Theorem

A point \mathbf{x}^* is a solution of the problem (2), where f is convex function, iff $\mathbf{x}^* \in \mathcal{X}$ and $\langle f'(\mathbf{x}^*), \mathbf{y} - \mathbf{x}^* \rangle \geq 0$ for all $\mathbf{y} \in \mathcal{X}$.

Proof

- ▶ Let $\mathbf{x}^* \in \mathcal{X}$ and the inequality holds. Then according to the first order criterion of convexity function f : $f(\mathbf{y}) \geq f(\mathbf{x}^*)$
- ▶ Let \mathbf{x}^* be a solution of the problem (2), but there exists $\tilde{\mathbf{y}}$ such that $\langle f'(\mathbf{x}^*), \tilde{\mathbf{y}} - \mathbf{x}^* \rangle < 0$
- ▶ Consider a point $\mathbf{z}(t) = t\tilde{\mathbf{y}} + (1 - t)\mathbf{x}^*$, $t \in [0, 1]$
- ▶ Then $\left. \frac{d}{dt} f(\mathbf{z}(t)) \right|_{t=0} = \langle f'(\mathbf{x}^*), \tilde{\mathbf{y}} - \mathbf{x}^* \rangle < 0$
- ▶ It means that for sufficiently small t the following inequality holds $f(\mathbf{z}(t)) < f(\mathbf{x}^*)$, and we have a contradiction.

Equality constraints

Problem statement

$$\begin{aligned} f(\mathbf{x}) &\rightarrow \min_{\mathbf{x} \in \mathbb{R}^n} \\ \text{s.t. } g_i(\mathbf{x}) &= 0, \quad i = 1, \dots, m \end{aligned}$$

Equality constraints

Problem statement

$$\begin{aligned} f(\mathbf{x}) &\rightarrow \min_{\mathbf{x} \in \mathbb{R}^n} \\ \text{s.t. } g_i(\mathbf{x}) &= 0, \quad i = 1, \dots, m \end{aligned}$$

Lagrangian

$$L(\mathbf{x}, \boldsymbol{\lambda}) = f(\mathbf{x}) + \sum_{i=1}^m \lambda_i g_i(\mathbf{x})$$

Geometric interpretation

From equality constraints to inequalities

Minimization problem

$$\begin{aligned} \min_{\mathbf{x} \in \mathbb{R}^n} f(\mathbf{x}) \\ \text{s.t. } g_i(\mathbf{x}) = 0, \quad i = 1, \dots, m \\ h_j(\mathbf{x}) \leq 0, \quad j = 1, \dots, p \end{aligned}$$

Lagrangian

$$L(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\mu}) = f(\mathbf{x}) + \sum_{i=1}^m \lambda_i g_i(\mathbf{x}) + \sum_{j=1}^p \mu_j h_j(\mathbf{x})$$

KKT necessary conditions

- ▶ $g_i(\mathbf{x}^*) = 0, i = 1, \dots, m$

KKT necessary conditions

- ▶ $g_i(\mathbf{x}^*) = 0, i = 1, \dots, m$
- ▶ $h_j(\mathbf{x}^*) \leq 0, j = 1, \dots, p$

KKT necessary conditions

- ▶ $g_i(\mathbf{x}^*) = 0, i = 1, \dots, m$
- ▶ $h_j(\mathbf{x}^*) \leq 0, j = 1, \dots, p$
- ▶ $\mu_j^* \geq 0, j = 1, \dots, p$

KKT necessary conditions

- ▶ $g_i(\mathbf{x}^*) = 0, i = 1, \dots, m$
- ▶ $h_j(\mathbf{x}^*) \leq 0, j = 1, \dots, p$
- ▶ $\mu_j^* \geq 0, j = 1, \dots, p$
- ▶ $\mu_j^* h_j(\mathbf{x}^*) = 0, j = 1, \dots, p$

KKT necessary conditions

- ▶ $g_i(\mathbf{x}^*) = 0, i = 1, \dots, m$
- ▶ $h_j(\mathbf{x}^*) \leq 0, j = 1, \dots, p$
- ▶ $\mu_j^* \geq 0, j = 1, \dots, p$
- ▶ $\mu_j^* h_j(\mathbf{x}^*) = 0, j = 1, \dots, p$
- ▶ $L'(\mathbf{x}^*, \boldsymbol{\lambda}^*, \boldsymbol{\mu}^*) = 0$

Slater regularity condition

Slater regularity

There exists a point $\bar{\mathbf{x}}$ inside the interior of convex feasible set such that $f_i(\bar{\mathbf{x}}) < 0$ and $\mathbf{A}\bar{\mathbf{x}} = \mathbf{b}$

Slater regularity condition

Slater regularity

There exists a point $\bar{\mathbf{x}}$ inside the interior of convex feasible set such that $f_i(\bar{\mathbf{x}}) < 0$ and $\mathbf{A}\bar{\mathbf{x}} = \mathbf{b}$

Theorem

If a problem is convex and there exists \mathbf{x} inside the interior of the feasible set, i.e. inequality constraints hold with strict inequalities, then the KKT conditions are also sufficient.

Summary on optimality conditions

- ▶ Optimality conditions: what and why?

Summary on optimality conditions

- ▶ Optimality conditions: what and why?
- ▶ Unconstrained case

Summary on optimality conditions

- ▶ Optimality conditions: what and why?
- ▶ Unconstrained case
- ▶ Equality constrained problems

Summary on optimality conditions

- ▶ Optimality conditions: what and why?
- ▶ Unconstrained case
- ▶ Equality constrained problems
- ▶ Inequality constrained problems

Summary on optimality conditions

- ▶ Optimality conditions: what and why?
- ▶ Unconstrained case
- ▶ Equality constrained problems
- ▶ Inequality constrained problems
- ▶ KKT conditions and Slater condition