

Optimization methods  
Lecture 1: Introduction.  
Convex sets. Convex functions

Alexandr Katrutsa

Modern State of Artificial Intelligence Masters Program  
Moscow Institute of Physics and Technology

# What is this course about?

## Basic theory

- ▶ Convex sets and convex functions
- ▶ Optimality conditions
- ▶ Introduction to duality

## Numerical methods

- ▶ First order methods and their accelerated versions
- ▶ Quasi-Newton methods
- ▶ Introduction to stochastic gradient methods
- ▶ Introduction to combinatorial optimization and convex relaxations

# The place of this course in the program

- ▶ When you train some neural network, you solve some optimization problem
- ▶ Possible issues in this process will be discussed in the course
- ▶ How to solve these issues we will also discuss

## Schedule and deadlines

- ▶ Lectures and webinars are once a week

## Schedule and deadlines

- ▶ Lectures and webinars are once a week
- ▶ Home assignments are after every lecture

## Schedule and deadlines

- ▶ Lectures and webinars are once a week
- ▶ Home assignments are after every lecture
- ▶ Grading policy will be announced during webinar

## Schedule and deadlines

- ▶ Lectures and webinars are once a week
- ▶ Home assignments are after every lecture
- ▶ Grading policy will be announced during webinar
- ▶ Lecture slides are here:  
[https://github.com/amkatrutsa/opt\\_modern\\_ai\\_ms](https://github.com/amkatrutsa/opt_modern_ai_ms)

## References

- ▶ S. Boyd and L. Vandenberghe *Convex Optimization*  
<https://web.stanford.edu/~boyd/cvxbook/>
- ▶ J. Nocedal, S. J. Wright *Numerical Optimization*
- ▶ I. Goodfellow et al *Deep learning book*



## Why this course is important?

- ▶ Formalization of the problem of selecting an element from a set

## Why this course is important?

- ▶ Formalization of the problem of selecting an element from a set
- ▶ Justification of the decision correctness

## Why this course is important?

- ▶ Formalization of the problem of selecting an element from a set
- ▶ Justification of the decision correctness
- ▶ Wide range of applications:

## Why this course is important?

- ▶ Formalization of the problem of selecting an element from a set
- ▶ Justification of the decision correctness
- ▶ Wide range of applications:
  - ▶ machine learning: classification, clustering, regression

## Why this course is important?

- ▶ Formalization of the problem of selecting an element from a set
- ▶ Justification of the decision correctness
- ▶ Wide range of applications:
  - ▶ machine learning: classification, clustering, regression
  - ▶ molecular modeling

## Why this course is important?

- ▶ Formalization of the problem of selecting an element from a set
- ▶ Justification of the decision correctness
- ▶ Wide range of applications:
  - ▶ machine learning: classification, clustering, regression
  - ▶ molecular modeling
  - ▶ risk analysis

## Why this course is important?

- ▶ Formalization of the problem of selecting an element from a set
- ▶ Justification of the decision correctness
- ▶ Wide range of applications:
  - ▶ machine learning: classification, clustering, regression
  - ▶ molecular modeling
  - ▶ risk analysis
  - ▶ portfolio optimization

## Why this course is important?

- ▶ Formalization of the problem of selecting an element from a set
- ▶ Justification of the decision correctness
- ▶ Wide range of applications:
  - ▶ machine learning: classification, clustering, regression
  - ▶ molecular modeling
  - ▶ risk analysis
  - ▶ portfolio optimization
  - ▶ optimal control



## Why this course is important?

- ▶ Formalization of the problem of selecting an element from a set
- ▶ Justification of the decision correctness
- ▶ Wide range of applications:
  - ▶ machine learning: classification, clustering, regression
  - ▶ molecular modeling
  - ▶ risk analysis
  - ▶ portfolio optimization
  - ▶ optimal control
  - ▶ signal processing

## Why this course is important?

- ▶ Formalization of the problem of selecting an element from a set
- ▶ Justification of the decision correctness
- ▶ Wide range of applications:
  - ▶ machine learning: classification, clustering, regression
  - ▶ molecular modeling
  - ▶ risk analysis
  - ▶ portfolio optimization
  - ▶ optimal control
  - ▶ signal processing
  - ▶ parameters estimation in statistic

## Why this course is important?

- ▶ Formalization of the problem of selecting an element from a set
- ▶ Justification of the decision correctness
- ▶ Wide range of applications:
  - ▶ machine learning: classification, clustering, regression
  - ▶ molecular modeling
  - ▶ risk analysis
  - ▶ portfolio optimization
  - ▶ optimal control
  - ▶ signal processing
  - ▶ parameters estimation in statistic
  - ▶ many-many others

# General methodology

Main steps for exploiting optimization methods in solving real-world problems:

1. Define objective function

## General methodology

Main steps for exploiting optimization methods in solving real-world problems:

1. Define objective function
2. Define feasible set

## General methodology

Main steps for exploiting optimization methods in solving real-world problems:

1. Define objective function
2. Define feasible set
3. Optimization problem statement and its analysis

## General methodology

Main steps for exploiting optimization methods in solving real-world problems:

1. Define objective function
2. Define feasible set
3. Optimization problem statement and its analysis
4. Selection of the best algorithm for the stated problem

## General methodology

Main steps for exploiting optimization methods in solving real-world problems:

1. Define objective function
2. Define feasible set
3. Optimization problem statement and its analysis
4. Selection of the best algorithm for the stated problem
5. Algorithm implementation and verification its correctness



## General problem statement

$$\begin{aligned} & \min_{\mathbf{x} \in \mathcal{X}} f_0(\mathbf{x}) \\ \text{s.t. } & f_i(\mathbf{x}) = 0, \quad i = 1, \dots, p \\ & f_j(\mathbf{x}) \leq 0, \quad j = p + 1, \dots, m, \end{aligned}$$

## General problem statement

$$\begin{aligned} & \min_{\mathbf{x} \in \mathcal{X}} f_0(\mathbf{x}) \\ \text{s.t. } & f_i(\mathbf{x}) = 0, \quad i = 1, \dots, p \\ & f_j(\mathbf{x}) \leq 0, \quad j = p + 1, \dots, m, \end{aligned}$$

- $\mathbf{x} \in \mathbb{R}^n$  — target vector

## General problem statement

$$\begin{aligned} & \min_{\mathbf{x} \in \mathcal{X}} f_0(\mathbf{x}) \\ \text{s.t. } & f_i(\mathbf{x}) = 0, \quad i = 1, \dots, p \\ & f_j(\mathbf{x}) \leq 0, \quad j = p + 1, \dots, m, \end{aligned}$$

- ▶  $\mathbf{x} \in \mathbb{R}^n$  — target vector
- ▶  $f_0(\mathbf{x}) : \mathbb{R}^n \rightarrow \mathbb{R}$  — objective function


## General problem statement

$$\begin{aligned} & \min_{\mathbf{x} \in \mathcal{X}} f_0(\mathbf{x}) \\ \text{s.t. } & f_i(\mathbf{x}) = 0, \quad i = 1, \dots, p \\ & f_j(\mathbf{x}) \leq 0, \quad j = p + 1, \dots, m, \end{aligned}$$

- ▶  $\mathbf{x} \in \mathbb{R}^n$  — target vector
- ▶  $f_0(\mathbf{x}) : \mathbb{R}^n \rightarrow \mathbb{R}$  — objective function
- ▶  $f_k(\mathbf{x}) : \mathbb{R}^n \rightarrow \mathbb{R}$  — constraint functions


## Linear classification problem

- Given dataset:  $(\mathbf{x}_i, y_i)$ ,  $\mathbf{x}_i \in \mathbb{R}^n$ ,  $y_i = \{+1, -1\}$ ,  $i = 1, \dots, m$



## Linear classification problem


- ▶ Given dataset:  $(\mathbf{x}_i, y_i)$ ,  $\mathbf{x}_i \in \mathbb{R}^n$ ,  $y_i = \{+1, -1\}$ ,  $i = 1, \dots, m$



- ▶ Linear classifier  $\hat{y} = \text{sign}(\mathbf{w}^\top \mathbf{x} + b)$

## Linear classification problem


- ▶ Given dataset:  $(\mathbf{x}_i, y_i)$ ,  $\mathbf{x}_i \in \mathbb{R}^n$ ,  $y_i = \{+1, -1\}$ ,  $i = 1, \dots, m$



- ▶ Linear classifier  $\hat{y} = \text{sign}(\mathbf{w}^\top \mathbf{x} + b)$
- ▶ 
$$\begin{cases} \mathbf{w}^\top \mathbf{x}_i + b > 1, & y_i = +1 \\ \mathbf{w}^\top \mathbf{x}_i + b < -1, & y_i = -1 \end{cases}$$

## Linear classification problem

- ▶ Given dataset:  $(\mathbf{x}_i, y_i)$ ,  $\mathbf{x}_i \in \mathbb{R}^n$ ,  $y_i = \{+1, -1\}$ ,  $i = 1, \dots, m$




- ▶ Linear classifier  $\hat{y} = \text{sign}(\mathbf{w}^\top \mathbf{x} + b)$
- ▶ 
$$\begin{cases} \mathbf{w}^\top \mathbf{x}_i + b > 1, & y_i = +1 \\ \mathbf{w}^\top \mathbf{x}_i + b < -1, & y_i = -1 \end{cases}$$
- ▶  $y_i(\mathbf{w}^\top \mathbf{x}_i + b) > 1$




## Possible separating hypelplanes


## Possible separating hypelplanes




## Possible separating hyperplanes



## Possible separating hyperplanes



## Possible separating hyperplanes



**Q:** How to define the separating hyperplane uniquely?

# Margin maximization

## Margin maximization

- For the support samples of every class the following holds

$$\begin{cases} \mathbf{w}^\top \mathbf{x}_k + b = 1, & y_k = +1 \\ \mathbf{w}^\top \mathbf{x}_j + b = -1, & y_j = -1 \end{cases}$$

## Margin maximization

- For the support samples of every class the following holds

$$\begin{cases} \mathbf{w}^\top \mathbf{x}_k + b = 1, & y_k = +1 \\ \mathbf{w}^\top \mathbf{x}_j + b = -1, & y_j = -1 \end{cases}$$

- Distance between parallel hyperplanes  $\mathbf{w}^\top \mathbf{x} + b = c_1$  and  $\mathbf{w}^\top \mathbf{x} + b = c_2$ :

$$d = \frac{|c_1 - c_2|}{\|\mathbf{w}\|_2} = \frac{2}{\|\mathbf{w}\|_2}$$



## Margin maximization

- ▶ For the support samples of every class the following holds

$$\begin{cases} \mathbf{w}^\top \mathbf{x}_k + b = 1, & y_k = +1 \\ \mathbf{w}^\top \mathbf{x}_j + b = -1, & y_j = -1 \end{cases}$$

- ▶ Distance between parallel hyperplanes  $\mathbf{w}^\top \mathbf{x} + b = c_1$  and  $\mathbf{w}^\top \mathbf{x} + b = c_2$ :

$$d = \frac{|c_1 - c_2|}{\|\mathbf{w}\|_2} = \frac{2}{\|\mathbf{w}\|_2}$$

- ▶ We want to maximize this distance or margin between two classes

## Margin maximization

- ▶ For the support samples of every class the following holds

$$\begin{cases} \mathbf{w}^\top \mathbf{x}_k + b = 1, & y_k = +1 \\ \mathbf{w}^\top \mathbf{x}_j + b = -1, & y_j = -1 \end{cases}$$

- ▶ Distance between parallel hyperplanes  $\mathbf{w}^\top \mathbf{x} + b = c_1$  and  $\mathbf{w}^\top \mathbf{x} + b = c_2$ :


$$d = \frac{|c_1 - c_2|}{\|\mathbf{w}\|_2} = \frac{2}{\|\mathbf{w}\|_2}$$

- ▶ We want to maximize this distance or margin between two classes

## The final optimization problem

$$\begin{aligned} \min_{\mathbf{w}, b} \quad & \frac{1}{2} \|\mathbf{w}\|_2^2 \\ \text{s.t.} \quad & y_i(\mathbf{w}^\top \mathbf{x}_i + b) > 1, \quad i = 1, \dots, m \end{aligned}$$

## Optimal separating hyperplane



What solutions are possible?

## What solutions are possible?

### Definition

A point  $\mathbf{x}^*$  is called a point of **global** minimum, if  $f(\mathbf{x}) \geq f(\mathbf{x}^*)$  for all  $\mathbf{x}$  from the feasible set.

## What solutions are possible?

### Definition

A point  $\mathbf{x}^*$  is called a point of **global** minimum, if  $f(\mathbf{x}) \geq f(\mathbf{x}^*)$  for all  $\mathbf{x}$  from the feasible set.

### Definition

A point  $\mathbf{x}^*$  is called a point of **local** minimum, if  $f(\mathbf{x}) \geq f(\mathbf{x}^*)$  for all  $\mathbf{x}$  from some neighborhood of point  $\mathbf{x}^*$  and from feasible set.

## What solutions are possible?

### Definition

A point  $\mathbf{x}^*$  is called a point of **global** minimum, if  $f(\mathbf{x}) \geq f(\mathbf{x}^*)$  for all  $\mathbf{x}$  from the feasible set.

### Definition

A point  $\mathbf{x}^*$  is called a point of **local** minimum, if  $f(\mathbf{x}) \geq f(\mathbf{x}^*)$  for all  $\mathbf{x}$  from some neighborhood of point  $\mathbf{x}^*$  and from feasible set.

### Another form of problem statement

$$\begin{aligned} \mathbf{x}^* &= \arg \min_{\mathbf{x} \in \mathcal{X}} f_0(\mathbf{x}) \\ \text{s.t. } f_i(\mathbf{x}) &= 0, \quad i = 1, \dots, p \\ f_j(\mathbf{x}) &\leq 0, \quad j = p + 1, \dots, m, \end{aligned}$$

## How to solve such problems?

In general case:

- ▶ NP-complete, i.e. very hard to solve
- ▶ randomized algorithms give a trade-off between running time and robustness of approximate solution



## How to solve such problems?

In general case:

- ▶ NP-complete, i.e. very hard to solve
- ▶ randomized algorithms give a trade-off between running time and robustness of approximate solution

However, some classes of optimization problems can be solved very efficiently

## How to solve such problems?

In general case:

- ▶ NP-complete, i.e. very hard to solve
- ▶ randomized algorithms give a trade-off between running time and robustness of approximate solution

However, some classes of optimization problems can be solved very efficiently

- ▶ Linear programming
- ▶ Linear least-squares problems
- ▶ Low-rank approximation problem
- ▶ Convex optimization

## Main stages in optimization theory development

- ▶ 1940s — linear programming
- ▶ 1950s — quadratic programming
- ▶ 1960s — geometric programming
- ▶ 1990s — polynomial interior point methods for convex conic optimization problems

## Modern topics

- ▶ Huge-scale optimization problems ( $\sim 10^8 - 10^{12}$ )

## Modern topics

- ▶ Huge-scale optimization problems ( $\sim 10^8 - 10^{12}$ )
- ▶ Distributed optimization methods

## Modern topics

- ▶ Huge-scale optimization problems ( $\sim 10^8 - 10^{12}$ )
- ▶ Distributed optimization methods
- ▶ Fast tensor methods

## Modern topics

- ▶ Huge-scale optimization problems ( $\sim 10^8 - 10^{12}$ )
- ▶ Distributed optimization methods
- ▶ Fast tensor methods
- ▶ Stochastic methods

## Modern topics

- ▶ Huge-scale optimization problems ( $\sim 10^8 - 10^{12}$ )
- ▶ Distributed optimization methods
- ▶ Fast tensor methods
- ▶ Stochastic methods
- ▶ Non-convex structured optimization problems



## Modern topics

- ▶ Huge-scale optimization problems ( $\sim 10^8 - 10^{12}$ )
- ▶ Distributed optimization methods
- ▶ Fast tensor methods
- ▶ Stochastic methods
- ▶ Non-convex structured optimization problems
- ▶ Applications of convex optimization

## Convex optimization problem

$$\begin{array}{ll}\min_{\mathbf{x} \in \mathbb{R}^n} & f_0(\mathbf{x}) \\ \text{s.t.} & f_i(\mathbf{x}) \leq 0, \ i = 1, \dots, m \\ & \mathbf{Ax} = \mathbf{b}\end{array}$$

## Convex optimization problem

$$\begin{aligned} \min_{\mathbf{x} \in \mathbb{R}^n} & f_0(\mathbf{x}) \\ \text{s.t.} & f_i(\mathbf{x}) \leq 0, \quad i = 1, \dots, m \\ & \mathbf{Ax} = \mathbf{b} \end{aligned}$$

- ▶  $f_0, f_i$  — convex functions:

$$f(\alpha \mathbf{x}_1 + \beta \mathbf{x}_2) \leq \alpha f(\mathbf{x}_1) + \beta f(\mathbf{x}_2),$$

where  $\alpha, \beta \geq 0$  and  $\alpha + \beta = 1$ .

## Convex optimization problem

$$\begin{aligned} \min_{\mathbf{x} \in \mathbb{R}^n} & f_0(\mathbf{x}) \\ \text{s.t. } & f_i(\mathbf{x}) \leq 0, \quad i = 1, \dots, m \\ & \mathbf{Ax} = \mathbf{b} \end{aligned}$$

- ▶  $f_0, f_i$  — convex functions:

$$f(\alpha \mathbf{x}_1 + \beta \mathbf{x}_2) \leq \alpha f(\mathbf{x}_1) + \beta f(\mathbf{x}_2),$$

where  $\alpha, \beta \geq 0$  and  $\alpha + \beta = 1$ .

- ▶ no analytical solution

## Convex optimization problem

$$\begin{aligned} \min_{\mathbf{x} \in \mathbb{R}^n} f_0(\mathbf{x}) \\ \text{s.t. } f_i(\mathbf{x}) \leq 0, \quad i = 1, \dots, m \\ \mathbf{Ax} = \mathbf{b} \end{aligned}$$

- ▶  $f_0, f_i$  — convex functions:

$$f(\alpha \mathbf{x}_1 + \beta \mathbf{x}_2) \leq \alpha f(\mathbf{x}_1) + \beta f(\mathbf{x}_2),$$

where  $\alpha, \beta \geq 0$  and  $\alpha + \beta = 1$ .

- ▶ no analytical solution
- ▶ efficient algorithms

## Convex optimization problem

$$\begin{aligned} \min_{\mathbf{x} \in \mathbb{R}^n} & f_0(\mathbf{x}) \\ \text{s.t. } & f_i(\mathbf{x}) \leq 0, \quad i = 1, \dots, m \\ & \mathbf{Ax} = \mathbf{b} \end{aligned}$$

- ▶  $f_0, f_i$  — convex functions:

$$f(\alpha \mathbf{x}_1 + \beta \mathbf{x}_2) \leq \alpha f(\mathbf{x}_1) + \beta f(\mathbf{x}_2),$$

where  $\alpha, \beta \geq 0$  and  $\alpha + \beta = 1$ .

- ▶ no analytical solution
- ▶ efficient algorithms
- ▶ special modeling helps to convert such problems to some standard form

# Why convexity is so important?

Ralph Tyrrell Rockafellar (born 1935)

The great watershed in optimization is not between linearity and non-linearity, but convexity and non-convexity.

# Why convexity is so important?

Ralph Tyrrell Rockafellar (born 1935)

The great watershed in optimization is not between linearity and non-linearity, but convexity and non-convexity.

- ▶ Local minimum is also global minimum



# Why convexity is so important?

Ralph Tyrrell Rockafellar (born 1935)

The great watershed in optimization is not between linearity and non-linearity, but convexity and non-convexity.

- ▶ Local minimum is also global minimum
- ▶ Necessary optimality condition is also sufficient

# Why convexity is so important?

Ralph Tyrrell Rockafellar (born 1935)

The great watershed in optimization is not between linearity and non-linearity, but convexity and non-convexity.

- ▶ Local minimum is also global minimum
- ▶ Necessary optimality condition is also sufficient

Questions:

# Why convexity is so important?

Ralph Tyrrell Rockafellar (born 1935)

The great watershed in optimization is not between linearity and non-linearity, but convexity and non-convexity.

- ▶ Local minimum is also global minimum
- ▶ Necessary optimality condition is also sufficient

Questions:

- ▶ Can any convex optimization problem be efficiently solved?

# Why convexity is so important?

Ralph Tyrrell Rockafellar (born 1935)

The great watershed in optimization is not between linearity and non-linearity, but convexity and non-convexity.

- ▶ Local minimum is also global minimum
- ▶ Necessary optimality condition is also sufficient

Questions:

- ▶ Can any convex optimization problem be efficiently solved?
- ▶ Is it possible to solve non-convex optimization problems efficiently?

## Convex sets

### Definition

*A set  $\mathcal{X} \subseteq \mathbb{R}^n$  is convex if for all  $\alpha \in [0, 1]$  and for any  $\mathbf{x}, \mathbf{y} \in \mathcal{X}$  the following holds*

$$\alpha \mathbf{x} + (1 - \alpha) \mathbf{y} \in \mathcal{X}.$$

# Convex sets

## Definition

*A set  $\mathcal{X} \subseteq \mathbb{R}^n$  is convex if for all  $\alpha \in [0, 1]$  and for any  $\mathbf{x}, \mathbf{y} \in \mathcal{X}$  the following holds*

$$\alpha \mathbf{x} + (1 - \alpha) \mathbf{y} \in \mathcal{X}.$$

## Examples

- Polyhedron

# Convex sets

## Definition

*A set  $\mathcal{X} \subseteq \mathbb{R}^n$  is convex if for all  $\alpha \in [0, 1]$  and for any  $\mathbf{x}, \mathbf{y} \in \mathcal{X}$  the following holds*

$$\alpha \mathbf{x} + (1 - \alpha) \mathbf{y} \in \mathcal{X}.$$

## Examples

- ▶ Polyhedron
- ▶ Hyperplanes

# Convex sets

## Definition

A set  $\mathcal{X} \subseteq \mathbb{R}^n$  is convex if for all  $\alpha \in [0, 1]$  and for any  $\mathbf{x}, \mathbf{y} \in \mathcal{X}$  the following holds

$$\alpha \mathbf{x} + (1 - \alpha) \mathbf{y} \in \mathcal{X}.$$

## Examples

- ▶ Polyhedron
- ▶ Hyperplanes
- ▶ Balls in *any proper* norm and ellipsoids



# Convex sets

## Definition

A set  $\mathcal{X} \subseteq \mathbb{R}^n$  is convex if for all  $\alpha \in [0, 1]$  and for any  $\mathbf{x}, \mathbf{y} \in \mathcal{X}$  the following holds

$$\alpha \mathbf{x} + (1 - \alpha) \mathbf{y} \in \mathcal{X}.$$

## Examples

- ▶ Polyhedron
- ▶ Hyperplanes
- ▶ Balls in *any proper* norm and ellipsoids
- ▶ Set of symmetric and non-negative definite matrices

## Intersection of convex sets

### Theorem

*Intersection of finite or infinite number of convex sets  $\mathcal{X}_i$  is a convex set:*

$$\mathcal{X} = \bigcap_{i \in \mathcal{I}} \mathcal{X}_i.$$

# Intersection of convex sets

## Theorem

*Intersection of finite or infinite number of convex sets  $\mathcal{X}_i$  is a convex set:*

$$\mathcal{X} = \bigcap_{i \in \mathcal{I}} \mathcal{X}_i.$$

## Proof

- Consider  $\mathbf{x}, \mathbf{y} \in \mathcal{X} \rightarrow \mathbf{x}, \mathbf{y} \in \mathcal{X}_i, \forall i \in \mathcal{I}$

## Intersection of convex sets

### Theorem

*Intersection of finite or infinite number of convex sets  $\mathcal{X}_i$  is a convex set:*

$$\mathcal{X} = \bigcap_{i \in \mathcal{I}} \mathcal{X}_i.$$

### Proof

- ▶ Consider  $\mathbf{x}, \mathbf{y} \in \mathcal{X} \rightarrow \mathbf{x}, \mathbf{y} \in \mathcal{X}_i, \forall i \in \mathcal{I}$
- ▶ Consider point  $\mathbf{z} = \alpha \mathbf{x} + (1 - \alpha) \mathbf{y}$ ,  $\alpha \in [0, 1]$

# Intersection of convex sets

## Theorem

*Intersection of finite or infinite number of convex sets  $\mathcal{X}_i$  is a convex set:*

$$\mathcal{X} = \bigcap_{i \in \mathcal{I}} \mathcal{X}_i.$$

## Proof

- ▶ Consider  $\mathbf{x}, \mathbf{y} \in \mathcal{X} \rightarrow \mathbf{x}, \mathbf{y} \in \mathcal{X}_i, \forall i \in \mathcal{I}$
- ▶ Consider point  $\mathbf{z} = \alpha \mathbf{x} + (1 - \alpha) \mathbf{y}$ ,  $\alpha \in [0, 1]$
- ▶ Since  $\mathcal{X}_i$  is convex for all  $i \in \mathcal{I}$ ,  $\mathbf{z} \in \mathcal{X}_i, \forall i \in \mathcal{I}$

# Intersection of convex sets

## Theorem

*Intersection of finite or infinite number of convex sets  $\mathcal{X}_i$  is a convex set:*

$$\mathcal{X} = \bigcap_{i \in \mathcal{I}} \mathcal{X}_i.$$

## Proof

- ▶ Consider  $\mathbf{x}, \mathbf{y} \in \mathcal{X} \rightarrow \mathbf{x}, \mathbf{y} \in \mathcal{X}_i, \forall i \in \mathcal{I}$
- ▶ Consider point  $\mathbf{z} = \alpha \mathbf{x} + (1 - \alpha) \mathbf{y}, \alpha \in [0, 1]$
- ▶ Since  $\mathcal{X}_i$  is convex for all  $i \in \mathcal{I}, \mathbf{z} \in \mathcal{X}_i, \forall i \in \mathcal{I}$
- ▶ Therefore,  $\mathbf{z} \in \mathcal{X}$  and  $\mathcal{X}$  is convex set

## Linear map of convex set

### Theorem

*If the domain of any linear map is convex, then the image of this map is also convex.*

### Proof

- Let  $\mathcal{X}$  be a convex set and  $\mathbf{x}, \mathbf{y} \in \mathcal{X}$

## Linear map of convex set

### Theorem

*If the domain of any linear map is convex, then the image of this map is also convex.*

### Proof

- ▶ Let  $\mathcal{X}$  be a convex set and  $\mathbf{x}, \mathbf{y} \in \mathcal{X}$
- ▶ Let  $f$  be a linear map:  $f(\mathbf{x}) = \mathbf{A}\mathbf{x} + \mathbf{b}$



## Linear map of convex set

### Theorem

*If the domain of any linear map is convex, then the image of this map is also convex.*

### Proof

- ▶ Let  $\mathcal{X}$  be a convex set and  $\mathbf{x}, \mathbf{y} \in \mathcal{X}$
- ▶ Let  $f$  be a linear map:  $f(\mathbf{x}) = \mathbf{A}\mathbf{x} + \mathbf{b}$
- ▶ Show that  $\alpha f(\mathbf{x}) + (1 - \alpha)f(\mathbf{y}) \in f(\mathcal{X})$ , where  $\alpha \in [0, 1]$

## Linear map of convex set

### Theorem

*If the domain of any linear map is convex, then the image of this map is also convex.*

### Proof

- ▶ Let  $\mathcal{X}$  be a convex set and  $\mathbf{x}, \mathbf{y} \in \mathcal{X}$
- ▶ Let  $f$  be a linear map:  $f(\mathbf{x}) = \mathbf{A}\mathbf{x} + \mathbf{b}$
- ▶ Show that  $\alpha f(\mathbf{x}) + (1 - \alpha)f(\mathbf{y}) \in f(\mathcal{X})$ , where  $\alpha \in [0, 1]$
- ▶ Indeed,

$$\begin{aligned}\alpha f(\mathbf{x}) + (1 - \alpha)f(\mathbf{y}) &= \alpha(\mathbf{A}\mathbf{x} + \mathbf{b}) + (1 - \alpha)(\mathbf{A}\mathbf{y} + \mathbf{b}) = \\ &\mathbf{A}(\alpha\mathbf{x} + (1 - \alpha)\mathbf{y}) + \mathbf{b} = \mathbf{A}\mathbf{z} + \mathbf{b} = f(\mathbf{z}),\end{aligned}$$

where  $\mathbf{z} = \alpha\mathbf{x} + (1 - \alpha)\mathbf{y} \in \mathcal{X}$

## Arithmetic operations under convex sets

### Theorem

*Minkowski sum of two convex sets is convex set.*

## Arithmetic operations under convex sets

### Theorem

*Minkowski sum of two convex sets is convex set.*

### Proof

- ▶ Let  $\mathcal{X}_1, \mathcal{X}_2$  be convex sets. Consider  
$$\mathcal{X} = \mathcal{X}_1 + \mathcal{X}_2 = \{\mathbf{x}_1 + \mathbf{x}_2 \mid \mathbf{x}_1 \in \mathcal{X}_1, \mathbf{x}_2 \in \mathcal{X}_2\}$$

## Arithmetic operations under convex sets

### Theorem

*Minkowski sum of two convex sets is convex set.*

### Proof

- ▶ Let  $\mathcal{X}_1, \mathcal{X}_2$  be convex sets. Consider
$$\mathcal{X} = \mathcal{X}_1 + \mathcal{X}_2 = \{\mathbf{x}_1 + \mathbf{x}_2 \mid \mathbf{x}_1 \in \mathcal{X}_1, \mathbf{x}_2 \in \mathcal{X}_2\}$$
- ▶ Let  $\hat{\mathbf{x}} = \hat{\mathbf{x}}_1 + \hat{\mathbf{x}}_2$  and  $\tilde{\mathbf{x}} = \tilde{\mathbf{x}}_1 + \tilde{\mathbf{x}}_2$  belong to  $\mathcal{X}$ . Show that
$$\alpha \hat{\mathbf{x}} + (1 - \alpha) \tilde{\mathbf{x}} \in \mathcal{X}$$

## Arithmetic operations under convex sets

### Theorem

*Minkowski sum of two convex sets is convex set.*

### Proof

- ▶ Let  $\mathcal{X}_1, \mathcal{X}_2$  be convex sets. Consider  
 $\mathcal{X} = \mathcal{X}_1 + \mathcal{X}_2 = \{\mathbf{x}_1 + \mathbf{x}_2 \mid \mathbf{x}_1 \in \mathcal{X}_1, \mathbf{x}_2 \in \mathcal{X}_2\}$
- ▶ Let  $\hat{\mathbf{x}} = \hat{\mathbf{x}}_1 + \hat{\mathbf{x}}_2$  and  $\tilde{\mathbf{x}} = \tilde{\mathbf{x}}_1 + \tilde{\mathbf{x}}_2$  belong to  $\mathcal{X}$ . Show that  
 $\alpha\hat{\mathbf{x}} + (1-\alpha)\tilde{\mathbf{x}} \in \mathcal{X}$
- ▶ Indeed,  
$$\alpha\hat{\mathbf{x}} + (1-\alpha)\tilde{\mathbf{x}} = [\alpha\hat{\mathbf{x}}_1 + (1-\alpha)\tilde{\mathbf{x}}_1] + [\alpha\hat{\mathbf{x}}_2 + (1-\alpha)\tilde{\mathbf{x}}_2] = \mathbf{y}_1 + \mathbf{y}_2,$$
where  $\mathbf{y}_1 \in C_1$  and  $\mathbf{y}_2 \in C_2$  since sets  $C_1, C_2$  are convex.

## Arithmetic operations under convex sets

### Theorem

*Minkowski sum of two convex sets is convex set.*

### Proof

- ▶ Let  $\mathcal{X}_1, \mathcal{X}_2$  be convex sets. Consider  
 $\mathcal{X} = \mathcal{X}_1 + \mathcal{X}_2 = \{\mathbf{x}_1 + \mathbf{x}_2 \mid \mathbf{x}_1 \in \mathcal{X}_1, \mathbf{x}_2 \in \mathcal{X}_2\}$
- ▶ Let  $\hat{\mathbf{x}} = \hat{\mathbf{x}}_1 + \hat{\mathbf{x}}_2$  and  $\tilde{\mathbf{x}} = \tilde{\mathbf{x}}_1 + \tilde{\mathbf{x}}_2$  belong to  $\mathcal{X}$ . Show that  
 $\alpha\hat{\mathbf{x}} + (1 - \alpha)\tilde{\mathbf{x}} \in \mathcal{X}$
- ▶ Indeed,  
$$\alpha\hat{\mathbf{x}} + (1 - \alpha)\tilde{\mathbf{x}} = [\alpha\hat{\mathbf{x}}_1 + (1 - \alpha)\tilde{\mathbf{x}}_1] + [\alpha\hat{\mathbf{x}}_2 + (1 - \alpha)\tilde{\mathbf{x}}_2] = \mathbf{y}_1 + \mathbf{y}_2,$$
  
where  $\mathbf{y}_1 \in C_1$  and  $\mathbf{y}_2 \in C_2$  since sets  $C_1, C_2$  are convex.

### Corollary

Linear combination of convex sets is convex set

## Arithmetic operations under convex sets

### Theorem

*Minkowski sum of two convex sets is convex set.*

### Proof

- ▶ Let  $\mathcal{X}_1, \mathcal{X}_2$  be convex sets. Consider  
 $\mathcal{X} = \mathcal{X}_1 + \mathcal{X}_2 = \{\mathbf{x}_1 + \mathbf{x}_2 \mid \mathbf{x}_1 \in \mathcal{X}_1, \mathbf{x}_2 \in \mathcal{X}_2\}$
- ▶ Let  $\hat{\mathbf{x}} = \hat{\mathbf{x}}_1 + \hat{\mathbf{x}}_2$  and  $\tilde{\mathbf{x}} = \tilde{\mathbf{x}}_1 + \tilde{\mathbf{x}}_2$  belong to  $\mathcal{X}$ . Show that  
 $\alpha\hat{\mathbf{x}} + (1 - \alpha)\tilde{\mathbf{x}} \in \mathcal{X}$
- ▶ Indeed,  
$$\alpha\hat{\mathbf{x}} + (1 - \alpha)\tilde{\mathbf{x}} = [\alpha\hat{\mathbf{x}}_1 + (1 - \alpha)\tilde{\mathbf{x}}_1] + [\alpha\hat{\mathbf{x}}_2 + (1 - \alpha)\tilde{\mathbf{x}}_2] = \mathbf{y}_1 + \mathbf{y}_2,$$
  
where  $\mathbf{y}_1 \in C_1$  and  $\mathbf{y}_2 \in C_2$  since sets  $C_1, C_2$  are convex.

### Corollary

Linear combination of convex sets is convex set

### Exercise

Proof that Cartesian product of convex sets is convex



### Definition

A set  $\mathcal{K}$  is a cone if for any  $\mathbf{x} \in \mathcal{K}$  and arbitrary number  $\theta \geq 0$  we have  $\theta\mathbf{x} \in \mathcal{K}$ .

### Definition

A set  $\mathcal{K}$  is called **convex** cone if for any points  $\mathbf{x}_1, \mathbf{x}_2 \in \mathcal{K}$  and any numbers  $\theta_1 \geq 0, \theta_2 \geq 0$  we have  $\theta_1\mathbf{x}_1 + \theta_2\mathbf{x}_2 \in \mathcal{K}$ .

## Definition

A set  $\mathcal{K}$  is a cone if for any  $\mathbf{x} \in \mathcal{K}$  and arbitrary number  $\theta \geq 0$  we have  $\theta\mathbf{x} \in \mathcal{K}$ .

## Definition

A set  $\mathcal{K}$  is called **convex** cone if for any points  $\mathbf{x}_1, \mathbf{x}_2 \in \mathcal{K}$  and any numbers  $\theta_1 \geq 0, \theta_2 \geq 0$  we have  $\theta_1\mathbf{x}_1 + \theta_2\mathbf{x}_2 \in \mathcal{K}$ .

## Important cones

## Definition

A set  $\mathcal{K}$  is a cone if for any  $\mathbf{x} \in \mathcal{K}$  and arbitrary number  $\theta \geq 0$  we have  $\theta\mathbf{x} \in \mathcal{K}$ .

## Definition

A set  $\mathcal{K}$  is called **convex** cone if for any points  $\mathbf{x}_1, \mathbf{x}_2 \in \mathcal{K}$  and any numbers  $\theta_1 \geq 0, \theta_2 \geq 0$  we have  $\theta_1\mathbf{x}_1 + \theta_2\mathbf{x}_2 \in \mathcal{K}$ .

## Important cones

- ▶ Nonnegative orthant  $\mathbb{R}_+^n = \{\mathbf{x} \in \mathbb{R}^n \mid x_i \geq 0, i = 1, \dots, n\} \rightarrow$  Linear programming (LP)

## Definition

A set  $\mathcal{K}$  is a cone if for any  $\mathbf{x} \in \mathcal{K}$  and arbitrary number  $\theta \geq 0$  we have  $\theta\mathbf{x} \in \mathcal{K}$ .

## Definition

A set  $\mathcal{K}$  is called **convex** cone if for any points  $\mathbf{x}_1, \mathbf{x}_2 \in \mathcal{K}$  and any numbers  $\theta_1 \geq 0, \theta_2 \geq 0$  we have  $\theta_1\mathbf{x}_1 + \theta_2\mathbf{x}_2 \in \mathcal{K}$ .

## Important cones

- ▶ Nonnegative orthant  $\mathbb{R}_+^n = \{\mathbf{x} \in \mathbb{R}^n \mid x_i \geq 0, i = 1, \dots, n\} \rightarrow$  Linear programming (LP)
- ▶ Second-order cone  $\{(\mathbf{x}, t) \in \mathbb{R}^{n+1} \mid \|\mathbf{x}\|_2 \leq t\} \rightarrow$  Second-order cone programming (SOCP)

## Definition

A set  $\mathcal{K}$  is a cone if for any  $\mathbf{x} \in \mathcal{K}$  and arbitrary number  $\theta \geq 0$  we have  $\theta\mathbf{x} \in \mathcal{K}$ .

## Definition

A set  $\mathcal{K}$  is called **convex** cone if for any points  $\mathbf{x}_1, \mathbf{x}_2 \in \mathcal{K}$  and any numbers  $\theta_1 \geq 0, \theta_2 \geq 0$  we have  $\theta_1\mathbf{x}_1 + \theta_2\mathbf{x}_2 \in \mathcal{K}$ .

## Important cones

- ▶ Nonnegative orthant  $\mathbb{R}_+^n = \{\mathbf{x} \in \mathbb{R}^n \mid x_i \geq 0, i = 1, \dots, n\} \rightarrow$  Linear programming (LP)
- ▶ Second-order cone  $\{(\mathbf{x}, t) \in \mathbb{R}^{n+1} \mid \|\mathbf{x}\|_2 \leq t\} \rightarrow$  Second-order cone programming (SOCP)
- ▶ Symmetric positive semi-definite matrices  $\mathbf{S}_+^n \rightarrow$  Semidefinite programming (SDP)

## Definition

*Convex hull of the set  $\mathcal{G}$  is called such set  $\text{conv}(\mathcal{G})$  that*

- ▶ it is an intersection of all convex sets containing  $\mathcal{G}$*
- ▶ it is a set of all convex combinations of points from  $\mathcal{G}$*

$$\text{conv}(\mathcal{G}) = \left\{ \sum_{i=1}^k \theta_i \mathbf{x}_i \mid \mathbf{x}_i \in \mathcal{G}, \sum_{i=1}^k \theta_i = 1, \theta_i \geq 0 \right\}$$

- ▶ it is a minimal convex set containing  $\mathcal{G}$*

## How to use convex hulls

- ▶ Assume that you face with optimization problem with **non-convex** feasible set

## How to use convex hulls

- ▶ Assume that you face with optimization problem with **non-convex** feasible set
- ▶ You can convexify feasible set with its convex hull



## How to use convex hulls

- ▶ Assume that you face with optimization problem with **non-convex** feasible set
- ▶ You can convexify feasible set with its convex hull
- ▶ Solve the problem in the new feasible set

## How to use convex hulls

- ▶ Assume that you face with optimization problem with **non-convex** feasible set
- ▶ You can convexify feasible set with its convex hull
- ▶ Solve the problem in the new feasible set
- ▶ Recover approximate solution of the original problem from the solution of the problem with convex feasible set

## Separation of convex sets

### Definition

Sets  $\mathcal{A}, \mathcal{B}$  are called separated if there exists vector  $\mathbf{a} \neq 0$  and a number  $b$  such that

- ▶  $\mathbf{a}^\top \mathbf{x} + b \geq 0$  for all  $\mathbf{x} \in \mathcal{A}$
- ▶  $\mathbf{a}^\top \mathbf{y} + b \leq 0$  for all  $\mathbf{y} \in \mathcal{B}$ .

## Separation of convex sets

### Definition

Sets  $\mathcal{A}, \mathcal{B}$  are called separated if there exists vector  $\mathbf{a} \neq 0$  and a number  $b$  such that

- ▶  $\mathbf{a}^\top \mathbf{x} + b \geq 0$  for all  $\mathbf{x} \in \mathcal{A}$
- ▶  $\mathbf{a}^\top \mathbf{y} + b \leq 0$  for all  $\mathbf{y} \in \mathcal{B}$ .

### Theorem

Let  $A$  and  $B$  convex and non-intersected sets. Then there exists separating hyperplane.

## Separation of convex sets

### Definition

Sets  $\mathcal{A}, \mathcal{B}$  are called separated if there exists vector  $\mathbf{a} \neq 0$  and a number  $b$  such that

- ▶  $\mathbf{a}^\top \mathbf{x} + b \geq 0$  for all  $\mathbf{x} \in \mathcal{A}$
- ▶  $\mathbf{a}^\top \mathbf{y} + b \leq 0$  for all  $\mathbf{y} \in \mathcal{B}$ .

### Theorem

Let  $A$  and  $B$  convex and non-intersected sets. Then there exists separating hyperplane.

### Proof

- ▶ Assume that the distance between  $\mathcal{A}$  and  $\mathcal{B}$  is positive:

$$\inf_{\mathbf{x} \in \mathcal{A}, \mathbf{y} \in \mathcal{B}} \|\mathbf{x} - \mathbf{y}\|_2 > 0$$

## Separation of convex sets

### Definition

Sets  $\mathcal{A}, \mathcal{B}$  are called separated if there exists vector  $\mathbf{a} \neq 0$  and a number  $b$  such that

- ▶  $\mathbf{a}^\top \mathbf{x} + b \geq 0$  for all  $\mathbf{x} \in \mathcal{A}$
- ▶  $\mathbf{a}^\top \mathbf{y} + b \leq 0$  for all  $\mathbf{y} \in \mathcal{B}$ .

### Theorem

Let  $A$  and  $B$  convex and non-intersected sets. Then there exists separating hyperplane.

### Proof

- ▶ Assume that the distance between  $\mathcal{A}$  and  $\mathcal{B}$  is positive:

$$\inf_{\mathbf{x} \in A, \mathbf{y} \in B} \|\mathbf{x} - \mathbf{y}\|_2 > 0$$

- ▶ Let  $\mathbf{c} \in A$  and  $\mathbf{d} \in B$  are points where infimum is attained

## Separation of convex sets

### Definition

Sets  $\mathcal{A}, \mathcal{B}$  are called separated if there exists vector  $\mathbf{a} \neq 0$  and a number  $b$  such that

- ▶  $\mathbf{a}^\top \mathbf{x} + b \geq 0$  for all  $\mathbf{x} \in \mathcal{A}$
- ▶  $\mathbf{a}^\top \mathbf{y} + b \leq 0$  for all  $\mathbf{y} \in \mathcal{B}$ .

### Theorem

Let  $A$  and  $B$  convex and non-intersected sets. Then there exists separating hyperplane.

### Proof

- ▶ Assume that the distance between  $\mathcal{A}$  and  $\mathcal{B}$  is positive:

$$\inf_{\mathbf{x} \in A, \mathbf{y} \in B} \|\mathbf{x} - \mathbf{y}\|_2 > 0$$

- ▶ Let  $\mathbf{c} \in A$  and  $\mathbf{d} \in B$  are points where infimum is attained
- ▶ Consider  $f(\mathbf{x}) = \mathbf{a}^\top \mathbf{x} + b$ , where  $\mathbf{a} = \mathbf{d} - \mathbf{c}$  and  $b = \frac{\|\mathbf{d}\|_2^2 - \|\mathbf{c}\|_2^2}{2}$

- ▶ Show that  $f(\mathbf{y}) \geq 0$  for all  $\mathbf{y} \in B$
- ▶ Assume we have  $\mathbf{u} \in B$  such that  $f(\mathbf{u}) < 0$
- ▶  $f(\mathbf{u}) = (\mathbf{d} - \mathbf{c})^\top (\mathbf{u} - \frac{1}{2}(\mathbf{d} + \mathbf{c})) = (\mathbf{d} - \mathbf{c})^\top (\mathbf{u} - \mathbf{d}) + \frac{1}{2}\|\mathbf{d} - \mathbf{c}\|_2^2$
- ▶  $(\mathbf{d} - \mathbf{c})^\top (\mathbf{u} - \mathbf{d}) < 0$
- ▶ Note that

$$\left. \frac{d}{dt} \|\mathbf{d} - \mathbf{c} + t(\mathbf{u} - \mathbf{d})\|_2^2 \right|_{t=0} = 2(\mathbf{d} - \mathbf{c})^\top (\mathbf{u} - \mathbf{d}) < 0$$

therefore, for  $t \in (0, 1]$

$$\|\mathbf{d} - \mathbf{c} + t(\mathbf{u} - \mathbf{d})\|_2 \leq \|\mathbf{d} - \mathbf{c}\|_2.$$

- ▶ A point  $\mathbf{d} + t(\mathbf{u} - \mathbf{d}) \in B$  is closer to  $\mathbf{c}$ , than  $\mathbf{d}$ , we have a contradiction.

**Q:** does the existence of separating hyperplane imply the non-intersection of convex sets?



## Summary on the convex sets

- ▶ Definition and geometric interpretation of convex set
- ▶ Three main cones
- ▶ Operations that preserve convexity
- ▶ Separating hyperplane theorem

## Convex function

### Definition

Function  $f : \mathcal{X} \subset \mathbb{R}^n \rightarrow \mathbb{R}$  is called convex (*strictly convex*), if  $\mathcal{X}$  is *convex set* and  $\forall \mathbf{x}_1, \mathbf{x}_2 \in X$  and  $\alpha \in [0, 1]$  ( $\alpha \in (0, 1)$ ) we have:

$$f(\alpha \mathbf{x}_1 + (1 - \alpha) \mathbf{x}_2) \leq (<) \alpha f(\mathbf{x}_1) + (1 - \alpha) f(\mathbf{x}_2)$$

## Convex function

### Definition

Function  $f : \mathcal{X} \subset \mathbb{R}^n \rightarrow \mathbb{R}$  is called convex (*strictly convex*), if  $\mathcal{X}$  is *convex set* and  $\forall \mathbf{x}_1, \mathbf{x}_2 \in X$  and  $\alpha \in [0, 1]$  ( $\alpha \in (0, 1)$ ) we have:

$$f(\alpha \mathbf{x}_1 + (1 - \alpha) \mathbf{x}_2) \leq (<) \alpha f(\mathbf{x}_1) + (1 - \alpha) f(\mathbf{x}_2)$$

### Definition

Function  $f$  is concave, if function  $-f$  is convex.

# Convex function

## Definition

Function  $f : \mathcal{X} \subset \mathbb{R}^n \rightarrow \mathbb{R}$  is called convex (*strictly convex*), if  $\mathcal{X}$  is *convex set* and  $\forall \mathbf{x}_1, \mathbf{x}_2 \in X$  and  $\alpha \in [0, 1]$  ( $\alpha \in (0, 1)$ ) we have:

$$f(\alpha \mathbf{x}_1 + (1 - \alpha) \mathbf{x}_2) \leq (<) \alpha f(\mathbf{x}_1) + (1 - \alpha) f(\mathbf{x}_2)$$

## Definition

Function  $f$  is concave, if function  $-f$  is convex.

## Examples of convex functions

- ▶  $x^p$  for  $x \geq 0$  and  $p \geq 1$
- ▶  $x \log x$ , where  $x > 0$
- ▶  $\max\{x_1, \dots, x_n\}$
- ▶  $\|\mathbf{x}\|$
- ▶  $\log \left( \sum_{i=1}^n e^{x_i} \right)$
- ▶  $-\log \det \mathbf{X}$  for  $\mathbf{X} \in \mathbf{S}_{++}^n$

## Epigraph and convexity

### Definition

A set  $\text{epi } f = \{(\mathbf{x}, t) \in \mathbb{R}^{n+1} \mid t \geq f(\mathbf{x})\}$  is called *epigraph* of  $f$ .

## Epigraph and convexity

### Definition

A set  $\text{epi } f = \{(\mathbf{x}, t) \in \mathbb{R}^{n+1} \mid t \geq f(\mathbf{x})\}$  is called *epigraph* of  $f$ .

### Theorem

Function  $f$  is convex  $\Leftrightarrow \text{epi } f$  is convex set.

## Epigraph and convexity

### Definition

A set  $\text{epi } f = \{(\mathbf{x}, t) \in \mathbb{R}^{n+1} \mid t \geq f(\mathbf{x})\}$  is called *epigraph* of  $f$ .

### Theorem

Function  $f$  is convex  $\Leftrightarrow \text{epi } f$  is convex set.

### Proof

1. Let  $f$  be a convex function

## Epigraph and convexity

### Definition

A set  $\text{epi } f = \{(\mathbf{x}, t) \in \mathbb{R}^{n+1} \mid t \geq f(\mathbf{x})\}$  is called *epigraph* of  $f$ .

### Theorem

Function  $f$  is convex  $\Leftrightarrow \text{epi } f$  is convex set.

### Proof

1. Let  $f$  be a convex function

- Consider any two points from epigraph  $(\mathbf{x}_1, t_1)$  and  $(\mathbf{x}_2, t_2)$ , where  $t_1 \geq f(\mathbf{x}_1)$  and  $t_2 \geq f(\mathbf{x}_2)$



## Epigraph and convexity

### Definition

A set  $\text{epi } f = \{(\mathbf{x}, t) \in \mathbb{R}^{n+1} \mid t \geq f(\mathbf{x})\}$  is called *epigraph* of  $f$ .

### Theorem

Function  $f$  is convex  $\Leftrightarrow$   $\text{epi } f$  is convex set.

### Proof

1. Let  $f$  be a convex function

- ▶ Consider any two points from epigraph  $(\mathbf{x}_1, t_1)$  and  $(\mathbf{x}_2, t_2)$ , where  $t_1 \geq f(\mathbf{x}_1)$  and  $t_2 \geq f(\mathbf{x}_2)$
- ▶ Check that the point  $(\alpha\mathbf{x}_1 + (1 - \alpha)\mathbf{x}_2, \alpha t_1 + (1 - \alpha)t_2)$  also belongs to epigraph

## Epigraph and convexity

### Definition

A set  $\text{epi } f = \{(\mathbf{x}, t) \in \mathbb{R}^{n+1} \mid t \geq f(\mathbf{x})\}$  is called *epigraph* of  $f$ .

### Theorem

Function  $f$  is convex  $\Leftrightarrow$   $\text{epi } f$  is convex set.

### Proof

1. Let  $f$  be a convex function

- ▶ Consider any two points from epigraph  $(\mathbf{x}_1, t_1)$  and  $(\mathbf{x}_2, t_2)$ , where  $t_1 \geq f(\mathbf{x}_1)$  and  $t_2 \geq f(\mathbf{x}_2)$
- ▶ Check that the point  $(\alpha\mathbf{x}_1 + (1 - \alpha)\mathbf{x}_2, \alpha t_1 + (1 - \alpha)t_2)$  also belongs to epigraph
- ▶ From the convexity of  $f$  follows
$$\alpha t_1 + (1 - \alpha)t_2 \geq \alpha f(\mathbf{x}_1) + (1 - \alpha)f(\mathbf{x}_2) \geq f(\alpha\mathbf{x}_1 + (1 - \alpha)\mathbf{x}_2).$$

## Epigraph and convexity

### Definition

A set  $\text{epi } f = \{(\mathbf{x}, t) \in \mathbb{R}^{n+1} \mid t \geq f(\mathbf{x})\}$  is called *epigraph* of  $f$ .

### Theorem

Function  $f$  is convex  $\Leftrightarrow$   $\text{epi } f$  is convex set.

### Proof

1. Let  $f$  be a convex function

- ▶ Consider any two points from epigraph  $(\mathbf{x}_1, t_1)$  and  $(\mathbf{x}_2, t_2)$ , where  $t_1 \geq f(\mathbf{x}_1)$  and  $t_2 \geq f(\mathbf{x}_2)$
- ▶ Check that the point  $(\alpha \mathbf{x}_1 + (1 - \alpha) \mathbf{x}_2, \alpha t_1 + (1 - \alpha) t_2)$  also belongs to epigraph
- ▶ From the convexity of  $f$  follows
$$\alpha t_1 + (1 - \alpha) t_2 \geq \alpha f(\mathbf{x}_1) + (1 - \alpha) f(\mathbf{x}_2) \geq f(\alpha \mathbf{x}_1 + (1 - \alpha) \mathbf{x}_2).$$

2. Let epigraph  $\text{epi } f$  is convex set

# Epigraph and convexity

## Definition

A set  $\text{epi } f = \{(\mathbf{x}, t) \in \mathbb{R}^{n+1} \mid t \geq f(\mathbf{x})\}$  is called *epigraph* of  $f$ .

## Theorem

Function  $f$  is convex  $\Leftrightarrow$   $\text{epi } f$  is convex set.

## Proof

1. Let  $f$  be a convex function

- ▶ Consider any two points from epigraph  $(\mathbf{x}_1, t_1)$  and  $(\mathbf{x}_2, t_2)$ , where  $t_1 \geq f(\mathbf{x}_1)$  and  $t_2 \geq f(\mathbf{x}_2)$
- ▶ Check that the point  $(\alpha\mathbf{x}_1 + (1 - \alpha)\mathbf{x}_2, \alpha t_1 + (1 - \alpha)t_2)$  also belongs to epigraph
- ▶ From the convexity of  $f$  follows
$$\alpha t_1 + (1 - \alpha)t_2 \geq \alpha f(\mathbf{x}_1) + (1 - \alpha)f(\mathbf{x}_2) \geq f(\alpha\mathbf{x}_1 + (1 - \alpha)\mathbf{x}_2).$$

2. Let epigraph  $\text{epi } f$  is convex set

- ▶  $(\mathbf{x}_1, f(\mathbf{x}_1))$  and  $(\mathbf{x}_2, f(\mathbf{x}_2)) \in \text{epi } f$ , then
$$(\alpha\mathbf{x}_1 + (1 - \alpha)\mathbf{x}_2, \alpha f(\mathbf{x}_1) + (1 - \alpha)f(\mathbf{x}_2)) \in \text{epi } f$$

# Epigraph and convexity

## Definition

A set  $\text{epi } f = \{(\mathbf{x}, t) \in \mathbb{R}^{n+1} \mid t \geq f(\mathbf{x})\}$  is called *epigraph* of  $f$ .

## Theorem

Function  $f$  is convex  $\Leftrightarrow$   $\text{epi } f$  is convex set.

## Proof

1. Let  $f$  be a convex function

- ▶ Consider any two points from epigraph  $(\mathbf{x}_1, t_1)$  and  $(\mathbf{x}_2, t_2)$ , where  $t_1 \geq f(\mathbf{x}_1)$  and  $t_2 \geq f(\mathbf{x}_2)$
- ▶ Check that the point  $(\alpha\mathbf{x}_1 + (1 - \alpha)\mathbf{x}_2, \alpha t_1 + (1 - \alpha)t_2)$  also belongs to epigraph
- ▶ From the convexity of  $f$  follows
$$\alpha t_1 + (1 - \alpha)t_2 \geq \alpha f(\mathbf{x}_1) + (1 - \alpha)f(\mathbf{x}_2) \geq f(\alpha\mathbf{x}_1 + (1 - \alpha)\mathbf{x}_2).$$

2. Let epigraph  $\text{epi } f$  is convex set

- ▶  $(\mathbf{x}_1, f(\mathbf{x}_1))$  and  $(\mathbf{x}_2, f(\mathbf{x}_2)) \in \text{epi } f$ , then
$$(\alpha\mathbf{x}_1 + (1 - \alpha)\mathbf{x}_2, \alpha f(\mathbf{x}_1) + (1 - \alpha)f(\mathbf{x}_2)) \in \text{epi } f$$
- ▶ From the definition of epigraph follows convexity of  $f$

## Strongly convex function

### Definition

Function  $f : \mathcal{X} \subset \mathbb{R}^n \rightarrow \mathbb{R}$  is called **strongly** convex with constant  $m > 0$ , if  $\mathcal{X}$  is convex set and  $\forall \mathbf{x}_1, \mathbf{x}_2 \in \mathcal{X}$  and  $\alpha \in [0, 1]$  we have:

$$f(\alpha \mathbf{x}_1 + (1-\alpha) \mathbf{x}_2) \leq \alpha f(\mathbf{x}_1) + (1-\alpha) f(\mathbf{x}_2) - \frac{m}{2} \alpha (1-\alpha) \|\mathbf{x}_1 - \mathbf{x}_2\|_2^2$$

## Strongly convex function

### Definition

Function  $f : \mathcal{X} \subset \mathbb{R}^n \rightarrow \mathbb{R}$  is called **strongly** convex with constant  $m > 0$ , if  $\mathcal{X}$  is convex set and  $\forall \mathbf{x}_1, \mathbf{x}_2 \in \mathcal{X}$  and  $\alpha \in [0, 1]$  we have:

$$f(\alpha \mathbf{x}_1 + (1-\alpha)\mathbf{x}_2) \leq \alpha f(\mathbf{x}_1) + (1-\alpha)f(\mathbf{x}_2) - \frac{m}{2}\alpha(1-\alpha)\|\mathbf{x}_1 - \mathbf{x}_2\|_2^2$$

- Convexity  $\supset$  strict convexity  $\supset$  strong convexity

## Strongly convex function

### Definition

Function  $f : \mathcal{X} \subset \mathbb{R}^n \rightarrow \mathbb{R}$  is called **strongly** convex with constant  $m > 0$ , if  $\mathcal{X}$  is convex set and  $\forall \mathbf{x}_1, \mathbf{x}_2 \in \mathcal{X}$  and  $\alpha \in [0, 1]$  we have:

$$f(\alpha \mathbf{x}_1 + (1-\alpha) \mathbf{x}_2) \leq \alpha f(\mathbf{x}_1) + (1-\alpha) f(\mathbf{x}_2) - \frac{m}{2} \alpha (1-\alpha) \|\mathbf{x}_1 - \mathbf{x}_2\|_2^2$$

- ▶ Convexity  $\supset$  strict convexity  $\supset$  strong convexity
- ▶ Theoretical analysis of methods in the case of strongly convex functions significantly differs from the one for convex functions



## Gradient and hessian: preliminaries

Consider  $f : \mathbb{R}^n \rightarrow \mathbb{R}$

- ▶ Directional derivative

$$f'_{\mathbf{d}}(\mathbf{x}) = \lim_{\alpha \rightarrow 0} \frac{f(\mathbf{x} + \alpha \mathbf{d}) - f(\mathbf{x})}{\alpha}$$

- ▶ Gradient  $f'(\mathbf{x})$  is a vector such that  $[f'(\mathbf{x})]_i = \frac{\partial f}{\partial x_i}$
- ▶ Hessian is a square matrix  $f''(\mathbf{x})$  such that  $[f''(\mathbf{x})]_{ij} = \frac{\partial^2 f}{\partial x_i \partial x_j}$

## Differential criteria of convexity

We consider convex function as strongly convex function with  $m = 0$ .

## Differential criteria of convexity

We consider convex function as strongly convex function with  $m = 0$ .

### Theorem (First order criterion)

*Let function  $f(\mathbf{x})$  is differentiable and its domain is a convex set  $\mathcal{X} \subseteq \mathbb{R}^n$ . Then  $f(\mathbf{x})$  is strongly convex with  $m \geq 0$  iff*

$$f(\mathbf{x}) - f(\mathbf{x}^*) \geq \langle f'(\mathbf{x}^*), \mathbf{x} - \mathbf{x}^* \rangle + \frac{m}{2} \|\mathbf{x} - \mathbf{x}^*\|_2^2, \quad \forall \mathbf{x}, \mathbf{x}^* \in X.$$

## Differential criteria of convexity

We consider convex function as strongly convex function with  $m = 0$ .

### Theorem (First order criterion)

*Let function  $f(\mathbf{x})$  is differentiable and its domain is a convex set  $\mathcal{X} \subseteq \mathbb{R}^n$ . Then  $f(\mathbf{x})$  is strongly convex with  $m \geq 0$  iff*

$$f(\mathbf{x}) - f(\mathbf{x}^*) \geq \langle f'(\mathbf{x}^*), \mathbf{x} - \mathbf{x}^* \rangle + \frac{m}{2} \|\mathbf{x} - \mathbf{x}^*\|_2^2, \quad \forall \mathbf{x}, \mathbf{x}^* \in X.$$

Proof: if  $f$  is convex

- By definition:  $f(\alpha \mathbf{x}_1 + (1 - \alpha) \mathbf{x}_2) \leq \alpha f(\mathbf{x}_1) + (1 - \alpha) f(\mathbf{x}_2)$

## Differential criteria of convexity

We consider convex function as strongly convex function with  $m = 0$ .

### Theorem (First order criterion)

*Let function  $f(\mathbf{x})$  is differentiable and its domain is a convex set  $\mathcal{X} \subseteq \mathbb{R}^n$ . Then  $f(\mathbf{x})$  is strongly convex with  $m \geq 0$  iff*

$$f(\mathbf{x}) - f(\mathbf{x}^*) \geq \langle f'(\mathbf{x}^*), \mathbf{x} - \mathbf{x}^* \rangle + \frac{m}{2} \|\mathbf{x} - \mathbf{x}^*\|_2^2, \quad \forall \mathbf{x}, \mathbf{x}^* \in X.$$

**Proof:** if  $f$  is convex

- By definition:  $f(\alpha \mathbf{x}_1 + (1 - \alpha) \mathbf{x}_2) \leq \alpha f(\mathbf{x}_1) + (1 - \alpha) f(\mathbf{x}_2)$
- Re-write in the form

$$f(\mathbf{x}_2 + \alpha(\mathbf{x}_1 - \mathbf{x}_2)) \leq f(\mathbf{x}_2) + \alpha(f(\mathbf{x}_1) - f(\mathbf{x}_2)) \text{ or}$$
$$\frac{f(\mathbf{x}_2 + \alpha(\mathbf{x}_1 - \mathbf{x}_2)) - f(\mathbf{x}_2)}{\alpha} \leq f(\mathbf{x}_1) - f(\mathbf{x}_2)$$

## Differential criteria of convexity

We consider convex function as strongly convex function with  $m = 0$ .

### Theorem (First order criterion)

*Let function  $f(\mathbf{x})$  is differentiable and its domain is a convex set  $\mathcal{X} \subseteq \mathbb{R}^n$ . Then  $f(\mathbf{x})$  is strongly convex with  $m \geq 0$  iff*

$$f(\mathbf{x}) - f(\mathbf{x}^*) \geq \langle f'(\mathbf{x}^*), \mathbf{x} - \mathbf{x}^* \rangle + \frac{m}{2} \|\mathbf{x} - \mathbf{x}^*\|_2^2, \quad \forall \mathbf{x}, \mathbf{x}^* \in X.$$

**Proof:** if  $f$  is convex

► By definition:  $f(\alpha \mathbf{x}_1 + (1 - \alpha) \mathbf{x}_2) \leq \alpha f(\mathbf{x}_1) + (1 - \alpha) f(\mathbf{x}_2)$

► Re-write in the form

$$f(\mathbf{x}_2 + \alpha(\mathbf{x}_1 - \mathbf{x}_2)) \leq f(\mathbf{x}_2) + \alpha(f(\mathbf{x}_1) - f(\mathbf{x}_2)) \text{ or}$$

$$\frac{f(\mathbf{x}_2 + \alpha(\mathbf{x}_1 - \mathbf{x}_2)) - f(\mathbf{x}_2)}{\alpha} \leq f(\mathbf{x}_1) - f(\mathbf{x}_2)$$

► If  $\alpha \rightarrow 0$ , then

$$\langle f'(\mathbf{x}_2), \mathbf{x}_1 - \mathbf{x}_2 \rangle \leq f(\mathbf{x}_1) - f(\mathbf{x}_2)$$

Proof: if the inequality holds

- ▶ Consider  $\mathbf{z} = \alpha \mathbf{x}_1 + (1 - \alpha) \mathbf{x}_2$

## Proof: if the inequality holds

- ▶ Consider  $\mathbf{z} = \alpha \mathbf{x}_1 + (1 - \alpha) \mathbf{x}_2$
- ▶ Write two inequalities for  $\mathbf{z}, \mathbf{x}_1$  and  $\mathbf{z}, \mathbf{x}_2$

$$f(\mathbf{x}_1) \geq f(\mathbf{z}) + \langle f'(\mathbf{z}), \mathbf{z} - \mathbf{x}_1 \rangle \mid \cdot \alpha$$

$$f(\mathbf{x}_2) \geq f(\mathbf{z}) + \langle f'(\mathbf{z}), \mathbf{z} - \mathbf{x}_2 \rangle \mid \cdot (1 - \alpha)$$



## Proof: if the inequality holds

- ▶ Consider  $\mathbf{z} = \alpha\mathbf{x}_1 + (1 - \alpha)\mathbf{x}_2$
- ▶ Write two inequalities for  $\mathbf{z}, \mathbf{x}_1$  and  $\mathbf{z}, \mathbf{x}_2$

$$f(\mathbf{x}_1) \geq f(\mathbf{z}) + \langle f'(\mathbf{z}), \mathbf{z} - \mathbf{x}_1 \rangle \mid \cdot \alpha$$

$$f(\mathbf{x}_2) \geq f(\mathbf{z}) + \langle f'(\mathbf{z}), \mathbf{z} - \mathbf{x}_2 \rangle \mid \cdot (1 - \alpha)$$

- ▶ Sum these inequalities

$$f(\mathbf{z}) = f(\alpha\mathbf{x}_1 + (1 - \alpha)\mathbf{x}_2) \leq \alpha f(\mathbf{x}_1) + (1 - \alpha)f(\mathbf{x}_2)$$

## Proof: if the inequality holds

- ▶ Consider  $\mathbf{z} = \alpha\mathbf{x}_1 + (1 - \alpha)\mathbf{x}_2$
- ▶ Write two inequalities for  $\mathbf{z}, \mathbf{x}_1$  and  $\mathbf{z}, \mathbf{x}_2$

$$f(\mathbf{x}_1) \geq f(\mathbf{z}) + \langle f'(\mathbf{z}), \mathbf{z} - \mathbf{x}_1 \rangle \mid \cdot \alpha$$

$$f(\mathbf{x}_2) \geq f(\mathbf{z}) + \langle f'(\mathbf{z}), \mathbf{z} - \mathbf{x}_2 \rangle \mid \cdot (1 - \alpha)$$

- ▶ Sum these inequalities

$$f(\mathbf{z}) = f(\alpha\mathbf{x}_1 + (1 - \alpha)\mathbf{x}_2) \leq \alpha f(\mathbf{x}_1) + (1 - \alpha)f(\mathbf{x}_2)$$

## Strongly convex case

To prove the claim for the strongly convex case, the same arguments can be provided for function  $f(\mathbf{x}) - \frac{m}{2}\|\mathbf{x}\|_2^2$ .

## Proof: if the inequality holds

- ▶ Consider  $\mathbf{z} = \alpha\mathbf{x}_1 + (1 - \alpha)\mathbf{x}_2$
- ▶ Write two inequalities for  $\mathbf{z}, \mathbf{x}_1$  and  $\mathbf{z}, \mathbf{x}_2$

$$f(\mathbf{x}_1) \geq f(\mathbf{z}) + \langle f'(\mathbf{z}), \mathbf{z} - \mathbf{x}_1 \rangle \mid \cdot \alpha$$

$$f(\mathbf{x}_2) \geq f(\mathbf{z}) + \langle f'(\mathbf{z}), \mathbf{z} - \mathbf{x}_2 \rangle \mid \cdot (1 - \alpha)$$

- ▶ Sum these inequalities

$$f(\mathbf{z}) = f(\alpha\mathbf{x}_1 + (1 - \alpha)\mathbf{x}_2) \leq \alpha f(\mathbf{x}_1) + (1 - \alpha)f(\mathbf{x}_2)$$


## Strongly convex case

To prove the claim for the strongly convex case, the same arguments can be provided for function  $f(\mathbf{x}) - \frac{m}{2}\|\mathbf{x}\|_2^2$ .

## Exercise

Prove that  $f$  is strongly convex  $\Leftrightarrow f(\mathbf{x}) - \frac{m}{2}\|\mathbf{x}\|_2^2$  is convex.

## Illustration for the first order criterion



## Theorem (Second order criterion)

*Twice continuously differentiable function  $f$  is convex  $\Leftrightarrow$*   
 $f''(\mathbf{x}) \succeq m\mathbf{I}$

## Theorem (Second order criterion)

*Twice continuously differentiable function  $f$  is convex  $\Leftrightarrow$*   
 $f''(\mathbf{x}) \succeq m\mathbf{I}$

### Proof

- Consider expansion in Taylor series up to the second order

$$f(\mathbf{y}) = f(\mathbf{x}) + \langle f'(\mathbf{x}), \mathbf{y} - \mathbf{x} \rangle + \frac{1}{2} \langle \mathbf{y} - \mathbf{x}, f''(\mathbf{x}_\alpha) \mathbf{y} - \mathbf{x} \rangle$$

## Theorem (Second order criterion)

*Twice continuously differentiable function  $f$  is convex  $\Leftrightarrow$*   
 $f''(\mathbf{x}) \succeq m\mathbf{I}$

### Proof

- Consider expansion in Taylor series up to the second order

$$f(\mathbf{y}) = f(\mathbf{x}) + \langle f'(\mathbf{x}), \mathbf{y} - \mathbf{x} \rangle + \frac{1}{2} \langle \mathbf{y} - \mathbf{x}, f''(\mathbf{x}_\alpha) \mathbf{y} - \mathbf{x} \rangle$$

- If  $f''(\mathbf{x}) \succeq m\mathbf{I}$ , then  $\frac{1}{2} \langle \mathbf{y} - \mathbf{x}, f''(\mathbf{x}_\alpha) \mathbf{y} - \mathbf{x} \rangle \geq \frac{m}{2} \|\mathbf{y} - \mathbf{x}\|_2^2$ ,  
and according to the first order criterion  $f$  is convex

## Theorem (Second order criterion)

Twice continuously differentiable function  $f$  is convex  $\Leftrightarrow$   
 $f''(\mathbf{x}) \succeq m\mathbf{I}$

### Proof

- Consider expansion in Taylor series up to the second order

$$f(\mathbf{y}) = f(\mathbf{x}) + \langle f'(\mathbf{x}), \mathbf{y} - \mathbf{x} \rangle + \frac{1}{2} \langle \mathbf{y} - \mathbf{x}, f''(\mathbf{x}_\alpha) \mathbf{y} - \mathbf{x} \rangle$$

- If  $f''(\mathbf{x}) \succeq m\mathbf{I}$ , then  $\frac{1}{2} \langle \mathbf{y} - \mathbf{x}, f''(\mathbf{x}_\alpha) \mathbf{y} - \mathbf{x} \rangle \geq \frac{m}{2} \|\mathbf{y} - \mathbf{x}\|_2^2$ ,  
and according to the first order criterion  $f$  is convex
- If there exists a point  $\mathbf{z}$  such that  $f''(\mathbf{z}) \not\succeq m\mathbf{I}$ , then we can  
find direction  $\mathbf{d}$  such that  $\mathbf{d}^\top f''(\mathbf{z}) \mathbf{d} \leq m$



## Theorem (Second order criterion)

Twice continuously differentiable function  $f$  is convex  $\Leftrightarrow$   
 $f''(\mathbf{x}) \succeq m\mathbf{I}$

### Proof

- Consider expansion in Taylor series up to the second order

$$f(\mathbf{y}) = f(\mathbf{x}) + \langle f'(\mathbf{x}), \mathbf{y} - \mathbf{x} \rangle + \frac{1}{2} \langle \mathbf{y} - \mathbf{x}, f''(\mathbf{x}_\alpha) \mathbf{y} - \mathbf{x} \rangle$$

- If  $f''(\mathbf{x}) \succeq m\mathbf{I}$ , then  $\frac{1}{2} \langle \mathbf{y} - \mathbf{x}, f''(\mathbf{x}_\alpha) \mathbf{y} - \mathbf{x} \rangle \geq \frac{m}{2} \|\mathbf{y} - \mathbf{x}\|_2^2$ ,  
and according to the first order criterion  $f$  is convex
- If there exists a point  $\mathbf{z}$  such that  $f''(\mathbf{z}) \not\succeq m\mathbf{I}$ , then we can  
find direction  $\mathbf{d}$  such that  $\mathbf{d}^\top f''(\mathbf{z}) \mathbf{d} \leq m$
- In this case according to the first order criterion  $f$  is  
non-convex. Thus, we get a contradiction.

## Theorem (Second order criterion)

Twice continuously differentiable function  $f$  is convex  $\Leftrightarrow$   
 $f''(\mathbf{x}) \succeq m\mathbf{I}$

### Proof

- ▶ Consider expansion in Taylor series up to the second order

$$f(\mathbf{y}) = f(\mathbf{x}) + \langle f'(\mathbf{x}), \mathbf{y} - \mathbf{x} \rangle + \frac{1}{2} \langle \mathbf{y} - \mathbf{x}, f''(\mathbf{x}_\alpha) \mathbf{y} - \mathbf{x} \rangle$$

- ▶ If  $f''(\mathbf{x}) \succeq m\mathbf{I}$ , then  $\frac{1}{2} \langle \mathbf{y} - \mathbf{x}, f''(\mathbf{x}_\alpha) \mathbf{y} - \mathbf{x} \rangle \geq \frac{m}{2} \|\mathbf{y} - \mathbf{x}\|_2^2$ ,  
and according to the first order criterion  $f$  is convex
- ▶ If there exists a point  $\mathbf{z}$  such that  $f''(\mathbf{z}) \not\succeq m\mathbf{I}$ , then we can  
find direction  $\mathbf{d}$  such that  $\mathbf{d}^\top f''(\mathbf{z}) \mathbf{d} \leq m$
- ▶ In this case according to the first order criterion  $f$  is  
non-convex. Thus, we get a contradiction.

### Reminder

If you want to check the definiteness of a square symmetric matrix,  
you should use definition or Sylvester criterion

## What function compositions preserve convexity?

- ▶ If  $f(\mathbf{x})$  is convex, then  $g(\mathbf{x}) = f(\mathbf{A}\mathbf{x} + \mathbf{b})$  is convex

## What function compositions preserve convexity?

- ▶ If  $f(\mathbf{x})$  is convex, then  $g(\mathbf{x}) = f(\mathbf{Ax} + \mathbf{b})$  is convex
- ▶ If  $f(\mathbf{x})$  is convex, then  $g(t) = f(\mathbf{x} + t\mathbf{y})$  is convex

## What function compositions preserve convexity?

- ▶ If  $f(\mathbf{x})$  is convex, then  $g(\mathbf{x}) = f(\mathbf{A}\mathbf{x} + \mathbf{b})$  is convex
- ▶ If  $f(\mathbf{x})$  is convex, then  $g(t) = f(\mathbf{x} + t\mathbf{y})$  is convex
- ▶ If  $f_i(\mathbf{x})$  are convex, then  $f(\mathbf{x}) = \max_{i=1,\dots,m} f_i(\mathbf{x})$  is convex

## What function compositions preserve convexity?

- ▶ If  $f(\mathbf{x})$  is convex, then  $g(\mathbf{x}) = f(\mathbf{A}\mathbf{x} + \mathbf{b})$  is convex
- ▶ If  $f(\mathbf{x})$  is convex, then  $g(t) = f(\mathbf{x} + t\mathbf{y})$  is convex
- ▶ If  $f_i(\mathbf{x})$  are convex, then  $f(\mathbf{x}) = \max_{i=1,\dots,m} f_i(\mathbf{x})$  is convex
- ▶ The sum of convex functions with non-negative coefficients is convex function

## What function compositions preserve convexity?

- ▶ If  $f(\mathbf{x})$  is convex, then  $g(\mathbf{x}) = f(\mathbf{A}\mathbf{x} + \mathbf{b})$  is convex
- ▶ If  $f(\mathbf{x})$  is convex, then  $g(t) = f(\mathbf{x} + t\mathbf{y})$  is convex
- ▶ If  $f_i(\mathbf{x})$  are convex, then  $f(\mathbf{x}) = \max_{i=1,\dots,m} f_i(\mathbf{x})$  is convex
- ▶ The sum of convex functions with non-negative coefficients is convex function
- ▶ Scalar composition  $h(f(\mathbf{x}))$

Local minimum of convex function is also a  
global minimum

### Theorem

*If  $f$  is a convex function and  $\mathbf{x}^*$  is a point of **local** minimum, the  $\mathbf{x}^*$  is a point of **global** minimum.*



# Local minimum of convex function is also a global minimum

## Theorem

*If  $f$  is a convex function and  $\mathbf{x}^*$  is a point of **local** minimum, the  $\mathbf{x}^*$  is a point of **global** minimum.*

## Proof

- ▶ Assume that there exists a point  $\mathbf{y}^*$  such that  $\mathbf{y}^* \neq \mathbf{x}^*$  and  $\mathbf{y}^*$  is a point of global minimum:  $f(\mathbf{y}^*) < f(\mathbf{x}^*)$

# Local minimum of convex function is also a global minimum

## Theorem

*If  $f$  is a convex function and  $\mathbf{x}^*$  is a point of **local** minimum, the  $\mathbf{x}^*$  is a point of **global** minimum.*

## Proof

- ▶ Assume that there exists a point  $\mathbf{y}^*$  such that  $\mathbf{y}^* \neq \mathbf{x}^*$  and  $\mathbf{y}^*$  is a point of global minimum:  $f(\mathbf{y}^*) < f(\mathbf{x}^*)$
- ▶ By definition of a point of local minimum:  $f(\mathbf{x}^*) \leq f(\mathbf{x})$ , where  $\|\mathbf{x}^* - \mathbf{x}\|_2 \leq \delta$

# Local minimum of convex function is also a global minimum

## Theorem

If  $f$  is a convex function and  $\mathbf{x}^*$  is a point of *local* minimum, the  $\mathbf{x}^*$  is a point of *global* minimum.

## Proof

- ▶ Assume that there exists a point  $\mathbf{y}^*$  such that  $\mathbf{y}^* \neq \mathbf{x}^*$  and  $\mathbf{y}^*$  is a point of global minimum:  $f(\mathbf{y}^*) < f(\mathbf{x}^*)$
- ▶ By definition of a point of local minimum:  $f(\mathbf{x}^*) \leq f(\mathbf{x})$ , where  $\|\mathbf{x}^* - \mathbf{x}\|_2 \leq \delta$
- ▶ Choose sufficiently small  $\alpha \in (0, 1)$  and consider a point  $\mathbf{z} = (1 - \alpha)\mathbf{x}^* + \alpha\mathbf{y}^*$  such that  $\|\mathbf{x}^* - \mathbf{z}\|_2 \leq \delta$

## Local minimum of convex function is also a global minimum

### Theorem

If  $f$  is a convex function and  $\mathbf{x}^*$  is a point of *local* minimum, the  $\mathbf{x}^*$  is a point of *global* minimum.

### Proof

- ▶ Assume that there exists a point  $\mathbf{y}^*$  such that  $\mathbf{y}^* \neq \mathbf{x}^*$  and  $\mathbf{y}^*$  is a point of global minimum:  $f(\mathbf{y}^*) < f(\mathbf{x}^*)$
- ▶ By definition of a point of local minimum:  $f(\mathbf{x}^*) \leq f(\mathbf{x})$ , where  $\|\mathbf{x}^* - \mathbf{x}\|_2 \leq \delta$
- ▶ Choose sufficiently small  $\alpha \in (0, 1)$  and consider a point  $\mathbf{z} = (1 - \alpha)\mathbf{x}^* + \alpha\mathbf{y}^*$  such that  $\|\mathbf{x}^* - \mathbf{z}\|_2 \leq \delta$
- ▶  $f(\mathbf{x}^*) \leq f(\mathbf{z}) \leq \alpha f(\mathbf{y}^*) + (1 - \alpha)f(\mathbf{x}^*) < f(\mathbf{x}^*)$

## Local minimum of convex function is also a global minimum

### Theorem

If  $f$  is a convex function and  $\mathbf{x}^*$  is a point of *local* minimum, the  $\mathbf{x}^*$  is a point of *global* minimum.

### Proof

- ▶ Assume that there exists a point  $\mathbf{y}^*$  such that  $\mathbf{y}^* \neq \mathbf{x}^*$  and  $\mathbf{y}^*$  is a point of global minimum:  $f(\mathbf{y}^*) < f(\mathbf{x}^*)$
- ▶ By definition of a point of local minimum:  $f(\mathbf{x}^*) \leq f(\mathbf{x})$ , where  $\|\mathbf{x}^* - \mathbf{x}\|_2 \leq \delta$
- ▶ Choose sufficiently small  $\alpha \in (0, 1)$  and consider a point  $\mathbf{z} = (1 - \alpha)\mathbf{x}^* + \alpha\mathbf{y}^*$  such that  $\|\mathbf{x}^* - \mathbf{z}\|_2 \leq \delta$
- ▶  $f(\mathbf{x}^*) \leq f(\mathbf{z}) \leq \alpha f(\mathbf{y}^*) + (1 - \alpha)f(\mathbf{x}^*) < f(\mathbf{x}^*)$
- ▶ We get a contradiction, therefore assumption is incorrect and  $\mathbf{x}^*$  is a point of global minimum

## Example of the difficult convex optimization problem

### Definition

A set  $\mathcal{C}^n = \{\mathbf{A} \in \mathbf{S}^n \mid \mathbf{x}^\top \mathbf{A} \mathbf{x} \geq 0, \mathbf{x} \geq 0\}$  is called copositive cone.

## Example of the difficult convex optimization problem

### Definition

A set  $\mathcal{C}^n = \{\mathbf{A} \in \mathbf{S}^n \mid \mathbf{x}^\top \mathbf{A} \mathbf{x} \geq 0, \mathbf{x} \geq 0\}$  is called *copositive cone*.

- $\mathcal{C}^n$  is convex

## Example of the difficult convex optimization problem

### Definition

A set  $\mathcal{C}^n = \{\mathbf{A} \in \mathbf{S}^n \mid \mathbf{x}^\top \mathbf{A} \mathbf{x} \geq 0, \mathbf{x} \geq 0\}$  is called *copositive cone*.

- ▶  $\mathcal{C}^n$  is convex
- ▶  $\mathbf{S}_+^n \subset \mathcal{C}^n$



## Example of the difficult convex optimization problem

### Definition

A set  $\mathcal{C}^n = \{\mathbf{A} \in \mathbf{S}^n \mid \mathbf{x}^\top \mathbf{A} \mathbf{x} \geq 0, \mathbf{x} \geq 0\}$  is called *copositive cone*.

- ▶  $\mathcal{C}^n$  is convex
- ▶  $\mathbf{S}_+^n \subset \mathcal{C}^n$
- ▶ The problem of check  $\mathbf{X} \notin \mathcal{C}^n$  is co-NP complete!

## Example of the difficult convex optimization problem

### Definition

A set  $\mathcal{C}^n = \{\mathbf{A} \in \mathbf{S}^n \mid \mathbf{x}^\top \mathbf{A} \mathbf{x} \geq 0, \mathbf{x} \geq 0\}$  is called copositive cone.

- ▶  $\mathcal{C}^n$  is convex
- ▶  $\mathbf{S}_+^n \subset \mathcal{C}^n$
- ▶ The problem of check  $\mathbf{X} \notin \mathcal{C}^n$  is co-NP complete!
- ▶ The conic optimization problem with the feasible set  $\mathcal{C}^n$  is NP-hard

## Example of the difficult convex optimization problem

### Definition

A set  $\mathcal{C}^n = \{\mathbf{A} \in \mathbf{S}^n \mid \mathbf{x}^\top \mathbf{A} \mathbf{x} \geq 0, \mathbf{x} \geq 0\}$  is called *copositive cone*.

- ▶  $\mathcal{C}^n$  is convex
- ▶  $\mathbf{S}_+^n \subset \mathcal{C}^n$
- ▶ The problem of check  $\mathbf{X} \notin \mathcal{C}^n$  is co-NP complete!
- ▶ The conic optimization problem with the feasible set  $\mathcal{C}^n$  is NP-hard

## Example of the difficult convex optimization problem

### Definition

A set  $\mathcal{C}^n = \{\mathbf{A} \in \mathbf{S}^n \mid \mathbf{x}^\top \mathbf{A} \mathbf{x} \geq 0, \mathbf{x} \geq 0\}$  is called *copositive cone*.

- ▶  $\mathcal{C}^n$  is convex
- ▶  $\mathbf{S}_+^n \subset \mathcal{C}^n$
- ▶ The problem of check  $\mathbf{X} \notin \mathcal{C}^n$  is co-NP complete!
- ▶ The conic optimization problem with the feasible set  $\mathcal{C}^n$  is NP-hard

### Example

The problem of search the maximum independence set of vertices in graph reduces to the convex optimization problem with feasible set  $\mathcal{C}^n$ . More details see [here](#)

## Example of the easy to solve non-convex optimization problem

Given the matrix  $\mathbf{Q} \in \mathbf{S}_{++}^n$ .

$$\begin{aligned} \min_{\mathbf{x} \in \mathbb{R}^n} \quad & \mathbf{x}^\top \mathbf{Q} \mathbf{x} \\ \text{s.t.} \quad & \|\mathbf{x}\|_2 = 1 \end{aligned}$$

## Example of the easy to solve non-convex optimization problem

Given the matrix  $\mathbf{Q} \in \mathbf{S}_{++}^n$ .

$$\begin{aligned} \min_{\mathbf{x} \in \mathbb{R}^n} \mathbf{x}^\top \mathbf{Q} \mathbf{x} \\ \text{s.t. } \|\mathbf{x}\|_2 = 1 \end{aligned}$$

- Feasible set is non-convex

## Example of the easy to solve non-convex optimization problem

Given the matrix  $\mathbf{Q} \in \mathbf{S}_{++}^n$ .

$$\begin{aligned} \min_{\mathbf{x} \in \mathbb{R}^n} \quad & \mathbf{x}^\top \mathbf{Q} \mathbf{x} \\ \text{s.t.} \quad & \|\mathbf{x}\|_2 = 1 \end{aligned}$$

- ▶ Feasible set is non-convex
- ▶ Objective function is convex

## Example of the easy to solve non-convex optimization problem

Given the matrix  $\mathbf{Q} \in \mathbf{S}_{++}^n$ .

$$\begin{aligned} \min_{\mathbf{x} \in \mathbb{R}^n} \quad & \mathbf{x}^\top \mathbf{Q} \mathbf{x} \\ \text{s.t.} \quad & \|\mathbf{x}\|_2 = 1 \end{aligned}$$

- ▶ Feasible set is non-convex
- ▶ Objective function is convex



## Example of the easy to solve non-convex optimization problem

Given the matrix  $\mathbf{Q} \in \mathbf{S}_{++}^n$ .

$$\begin{aligned} \min_{\mathbf{x} \in \mathbb{R}^n} \mathbf{x}^\top \mathbf{Q} \mathbf{x} \\ \text{s.t. } \|\mathbf{x}\|_2 = 1 \end{aligned}$$

- ▶ Feasible set is non-convex
- ▶ Objective function is convex

**Q:** what meaning of  $\mathbf{x}^*$  and  $f(\mathbf{x}^*)$ ?

## Jensen's inequality

### Theorem

If function  $f$  is convex, then  $f\left(\sum_{i=1}^k \alpha \mathbf{x}_i\right) \leq \sum_{i=1}^k \alpha_i f(\mathbf{x}_i)$ , where

$$\sum_{i=1}^k \alpha_i = 1, \alpha_i \geq 0.$$

## Jensen's inequality

### Theorem

If function  $f$  is convex, then  $f\left(\sum_{i=1}^k \alpha \mathbf{x}_i\right) \leq \sum_{i=1}^k \alpha_i f(\mathbf{x}_i)$ , where

$$\sum_{i=1}^k \alpha_i = 1, \alpha_i \geq 0.$$

### Proof by induction

- Base  $k = 2$  holds according to the definition

## Jensen's inequality

### Theorem

If function  $f$  is convex, then  $f\left(\sum_{i=1}^k \alpha \mathbf{x}_i\right) \leq \sum_{i=1}^k \alpha_i f(\mathbf{x}_i)$ , where

$$\sum_{i=1}^k \alpha_i = 1, \alpha_i \geq 0.$$

### Proof by induction

- ▶ Base  $k = 2$  holds according to the definition
- ▶ Assume the inequality holds for  $k = m - 1$ :

$$f\left(\sum_{i=1}^{m-1} \alpha \mathbf{x}_i\right) \leq \sum_{i=1}^{m-1} \alpha_i f(\mathbf{x}_i) \text{ and } \sum_{i=1}^{m-1} \alpha_i = 1, \alpha_i \geq 0$$

# Jensen's inequality

## Theorem

If function  $f$  is convex, then  $f\left(\sum_{i=1}^k \alpha \mathbf{x}_i\right) \leq \sum_{i=1}^k \alpha_i f(\mathbf{x}_i)$ , where

$$\sum_{i=1}^k \alpha_i = 1, \alpha_i \geq 0.$$

## Proof by induction

- Base  $k = 2$  holds according to the definition

- Assume the inequality holds for  $k = m - 1$ :

$$f\left(\sum_{i=1}^{m-1} \alpha \mathbf{x}_i\right) \leq \sum_{i=1}^{m-1} \alpha_i f(\mathbf{x}_i) \text{ and } \sum_{i=1}^{m-1} \alpha_i = 1, \alpha_i \geq 0$$

- Consider  $k = m$ :  $f\left(\sum_{i=1}^m \hat{\alpha}_i \mathbf{x}_i\right) = f\left(\sum_{i=1}^{m-1} \hat{\alpha} \mathbf{x}_i + \hat{\alpha}_m \mathbf{x}_m\right) =$

$$f\left((1 - \hat{\alpha}_m) \sum_{i=1}^{m-1} \frac{\hat{\alpha}_i}{1 - \hat{\alpha}_m} \mathbf{x}_i + \hat{\alpha}_m \mathbf{x}_m\right) \leq$$

$$(1 - \hat{\alpha}_m) f\left(\sum_{i=1}^{m-1} \frac{\hat{\alpha}_i}{1 - \hat{\alpha}_m} \mathbf{x}_i\right) + \hat{\alpha}_m f(\mathbf{x}_m) \leq \sum_{i=1}^k \alpha_i f(\mathbf{x}_i)$$

## Corollaries and generalizations

- If we write Jensen's inequality for the function  $-\log x$ , we get inequality for geometric and arithmetic means

$$\frac{1}{m} \sum_{i=1}^m x_i \geq \sqrt[m]{x_1 \cdot \dots \cdot x_m}$$

## Corollaries and generalizations

- If we write Jensen's inequality for the function  $-\log x$ , we get inequality for geometric and arithmetic means

$$\frac{1}{m} \sum_{i=1}^m x_i \geq \sqrt[m]{x_1 \cdot \dots \cdot x_m}$$

- Hölder's inequality

$$\sum_{i=1}^n x_i y_i \leq \left( \sum_{i=1}^n |x_i|^p \right)^{1/p} \left( \sum_{i=1}^n |y_i|^q \right)^{1/q}, \quad \frac{1}{p} + \frac{1}{q} = 1$$

## Corollaries and generalizations

- ▶ If we write Jensen's inequality for the function  $-\log x$ , we get inequality for geometric and arithmetic means

$$\frac{1}{m} \sum_{i=1}^m x_i \geq \sqrt[m]{x_1 \cdot \dots \cdot x_m}$$

- ▶ Hölder's inequality

$$\sum_{i=1}^n x_i y_i \leq \left( \sum_{i=1}^n |x_i|^p \right)^{1/p} \left( \sum_{i=1}^n |y_i|^q \right)^{1/q}, \quad \frac{1}{p} + \frac{1}{q} = 1$$

- ▶ The generalization of Jensen's inequality gives the inequality for the convex function of the expected value

$$f(\mathbb{E}(\mathbf{x})) \leq \mathbb{E}(f(\mathbf{x}))$$



## Summary on convex functions

- ▶ Convex, strictly convex and strongly convex functions

## Summary on convex functions

- ▶ Convex, strictly convex and strongly convex functions
- ▶ Examples and how to verify convexity of function

## Summary on convex functions

- ▶ Convex, strictly convex and strongly convex functions
- ▶ Examples and how to verify convexity of function
- ▶ Operations that preserve convexity

## Summary on convex functions

- ▶ Convex, strictly convex and strongly convex functions
- ▶ Examples and how to verify convexity of function
- ▶ Operations that preserve convexity
- ▶ Jensen inequality