

Optimization methods

Lecture 4: Introduction to duality theory

Alexandr Katrutsa

Modern State of Artificial Intelligence Masters Program
Moscow Institute of Physics and Technology

Brief reminder of the previous lecture

- ▶ Optimality conditions for unconstrained optimization problems

Brief reminder of the previous lecture

- ▶ Optimality conditions for unconstrained optimization problems
- ▶ Saddle points

Brief reminder of the previous lecture

- ▶ Optimality conditions for unconstrained optimization problems
- ▶ Saddle points
- ▶ Optimality conditions for constrained optimization problem in the general form

Brief reminder of the previous lecture

- ▶ Optimality conditions for unconstrained optimization problems
- ▶ Saddle points
- ▶ Optimality conditions for constrained optimization problem in the general form
- ▶ Karush-Kuhn-Tucker conditions

Brief reminder of the previous lecture

- ▶ Optimality conditions for unconstrained optimization problems
- ▶ Saddle points
- ▶ Optimality conditions for constrained optimization problem in the general form
- ▶ Karush-Kuhn-Tucker conditions
- ▶ Slater regularity

Optimization problem with functional constraints

$$\begin{aligned} & \min_{\mathbf{x} \in \mathcal{D}} f_0(\mathbf{x}) \\ & \text{s.t. } g_i(\mathbf{x}) = 0, \quad i = 1, \dots, m \\ & \quad h_j(\mathbf{x}) \leq 0, \quad j = 1, \dots, p \end{aligned}$$

$$\text{dom } f_0 = \mathcal{D} \subseteq \mathbb{R}^n, \quad f_0(\mathbf{x}^*) = p^*$$

Optimization problem with functional constraints

$$\begin{aligned} \min_{\mathbf{x} \in \mathcal{D}} f_0(\mathbf{x}) \\ \text{s.t. } g_i(\mathbf{x}) = 0, \quad i = 1, \dots, m \\ h_j(\mathbf{x}) \leq 0, \quad j = 1, \dots, p \end{aligned}$$

$$\text{dom } f_0 = \mathcal{D} \subseteq \mathbb{R}^n, \quad f_0(\mathbf{x}^*) = p^*$$

Lagrangian $L : \mathcal{D} \times \mathbb{R}^m \times \mathbb{R}^p \rightarrow \mathbb{R}$

$$L(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\mu}) = f_0(\mathbf{x}) + \sum_{i=1}^m \lambda_i g_i(\mathbf{x}) + \sum_{j=1}^p \mu_j h_j(\mathbf{x})$$

- ▶ λ_i – Lagrange multipliers for constraints $g_i(\mathbf{x}) = 0, \quad i = 1, \dots, m$
- ▶ μ_j – Lagrange multipliers for constraints $h_j(\mathbf{x}) \leq 0, \quad j = 1, \dots, p$

Dual function

Definition

A function $g : \mathbb{R}^m \times \mathbb{R}^p \rightarrow \mathbb{R}$ such that

$$g(\boldsymbol{\lambda}, \boldsymbol{\mu}) = \inf_{\mathbf{x} \in \mathcal{D}} \left(f_0(\mathbf{x}) + \sum_{i=1}^m \lambda_i g_i(\mathbf{x}) + \sum_{j=1}^p \mu_j h_j(\mathbf{x}) \right)$$

is called *dual function*

Dual function

Definition

A function $g : \mathbb{R}^m \times \mathbb{R}^p \rightarrow \mathbb{R}$ such that

$$g(\boldsymbol{\lambda}, \boldsymbol{\mu}) = \inf_{\mathbf{x} \in \mathcal{D}} \left(f_0(\mathbf{x}) + \sum_{i=1}^m \lambda_i g_i(\mathbf{x}) + \sum_{j=1}^p \mu_j h_j(\mathbf{x}) \right)$$

is called *dual function*

Properties

- ▶ Concave
- ▶ Can be equal to $-\infty$ for some pairs $(\boldsymbol{\lambda}, \boldsymbol{\mu})$

Lower bound for the optimal objective function value

Claim

If $\boldsymbol{\mu} \geq 0$, then $p^* \geq g(\boldsymbol{\lambda}, \boldsymbol{\mu})$

Lower bound for the optimal objective function value

Claim

If $\boldsymbol{\mu} \geq 0$, then $p^* \geq g(\boldsymbol{\lambda}, \boldsymbol{\mu})$

Proof

Lower bound for the optimal objective function value

Claim

If $\mu \geq 0$, then $p^* \geq g(\lambda, \mu)$

Proof

- If $\hat{\mathbf{x}} \in \mathcal{D}$ and belongs to the feasible set and $\mu \geq 0$, then

$$f_0(\hat{\mathbf{x}}) \geq L(\hat{\mathbf{x}}, \lambda, \mu) \geq \inf_{\mathbf{x} \in \mathcal{D}} L(\mathbf{x}, \lambda, \mu) = g(\lambda, \mu)$$

Lower bound for the optimal objective function value

Claim

If $\mu \geq 0$, then $p^* \geq g(\lambda, \mu)$

Proof

- ▶ If $\hat{\mathbf{x}} \in \mathcal{D}$ and belongs to the feasible set and $\mu \geq 0$, then

$$f_0(\hat{\mathbf{x}}) \geq L(\hat{\mathbf{x}}, \lambda, \mu) \geq \inf_{\mathbf{x} \in \mathcal{D}} L(\mathbf{x}, \lambda, \mu) = g(\lambda, \mu)$$

- ▶ Minimizing both sides over all feasible $\hat{\mathbf{x}}$ gives

$$p^* \geq g(\lambda, \mu)$$

Lower bound for the optimal objective function value

Claim

If $\mu \geq 0$, then $p^* \geq g(\lambda, \mu)$

Proof

- ▶ If $\hat{\mathbf{x}} \in \mathcal{D}$ and belongs to the feasible set and $\mu \geq 0$, then

$$f_0(\hat{\mathbf{x}}) \geq L(\hat{\mathbf{x}}, \lambda, \mu) \geq \inf_{\mathbf{x} \in \mathcal{D}} L(\mathbf{x}, \lambda, \mu) = g(\lambda, \mu)$$

- ▶ Minimizing both sides over all feasible $\hat{\mathbf{x}}$ gives

$$p^* \geq g(\lambda, \mu)$$

Q: what can we do now to get the best approximation of the optimal p^* ?

Dual problem

Definition

Dual problem is called the following optimization problem

$$\begin{aligned} & \max g(\boldsymbol{\lambda}, \boldsymbol{\mu}) \\ \text{s.t. } & \boldsymbol{\mu} \geq 0 \end{aligned}$$

Dual problem

Definition

Dual problem is called the following optimization problem

$$\begin{aligned} & \max g(\boldsymbol{\lambda}, \boldsymbol{\mu}) \\ & \text{s.t. } \boldsymbol{\mu} \geq 0 \end{aligned}$$

- Dual problem is always convex

Dual problem

Definition

Dual problem is called the following optimization problem

$$\begin{aligned} & \max g(\boldsymbol{\lambda}, \boldsymbol{\mu}) \\ \text{s.t. } & \boldsymbol{\mu} \geq 0 \end{aligned}$$

- ▶ Dual problem is always convex
- ▶ Denote by $d^* = g(\boldsymbol{\lambda}^*, \boldsymbol{\mu}^*)$ the optimal value of the dual objective function

Dual problem

Definition

Dual problem is called the following optimization problem

$$\begin{aligned} & \max g(\boldsymbol{\lambda}, \boldsymbol{\mu}) \\ & \text{s.t. } \boldsymbol{\mu} \geq 0 \end{aligned}$$

- ▶ Dual problem is always convex
- ▶ Denote by $d^* = g(\boldsymbol{\lambda}^*, \boldsymbol{\mu}^*)$ the optimal value of the dual objective function
- ▶ It is the best lower bound of p^* that can be given by dual function

Dual problem

Definition

Dual problem is called the following optimization problem

$$\begin{aligned} & \max g(\boldsymbol{\lambda}, \boldsymbol{\mu}) \\ & \text{s.t. } \boldsymbol{\mu} \geq 0 \end{aligned}$$

- ▶ Dual problem is always convex
- ▶ Denote by $d^* = g(\boldsymbol{\lambda}^*, \boldsymbol{\mu}^*)$ the optimal value of the dual objective function
- ▶ It is the best lower bound of p^* that can be given by dual function
- ▶ Vectors $(\boldsymbol{\lambda}, \boldsymbol{\mu})$ are called dual feasible if $\boldsymbol{\mu} \geq 0$ and $(\boldsymbol{\lambda}, \boldsymbol{\mu}) \in \text{dom } g$

Weak and strong duality

Weak duality: $d^* \leq p^*$

Weak and strong duality

Weak duality: $d^* \leq p^*$

- It holds by construction

Weak and strong duality

Weak duality: $d^* \leq p^*$

- ▶ It holds by construction
- ▶ Non-trivial lower bounds for NP-hard problems

Weak and strong duality

Weak duality: $d^* \leq p^*$

- ▶ It holds by construction
- ▶ Non-trivial lower bounds for NP-hard problems

Strong duality: $d^* = p^*$

Weak and strong duality

Weak duality: $d^* \leq p^*$

- ▶ It holds by construction
- ▶ Non-trivial lower bounds for NP-hard problems

Strong duality: $d^* = p^*$

- ▶ It **does not** hold in general case

Weak and strong duality

Weak duality: $d^* \leq p^*$

- ▶ It holds by construction
- ▶ Non-trivial lower bounds for NP-hard problems

Strong duality: $d^* = p^*$

- ▶ It **does not** hold in general case
- ▶ *Typically* it holds for convex problems

Weak and strong duality

Weak duality: $d^* \leq p^*$

- ▶ It holds by construction
- ▶ Non-trivial lower bounds for NP-hard problems

Strong duality: $d^* = p^*$

- ▶ It **does not** hold in general case
- ▶ *Typically* it holds for convex problems
- ▶ But regularity conditions are important

Weak and strong duality

Weak duality: $d^* \leq p^*$

- ▶ It holds by construction
- ▶ Non-trivial lower bounds for NP-hard problems

Strong duality: $d^* = p^*$

- ▶ It **does not** hold in general case
- ▶ *Typically* it holds for convex problems
- ▶ But regularity conditions are important
- ▶ It may hold for **non-convex** problems

Weak and strong duality

Weak duality: $d^* \leq p^*$

- ▶ It holds by construction
- ▶ Non-trivial lower bounds for NP-hard problems

Strong duality: $d^* = p^*$

- ▶ It **does not** hold in general case
- ▶ *Typically* it holds for convex problems
- ▶ But regularity conditions are important
- ▶ It may hold for **non-convex** problems
- ▶ A solution of the dual problem can help to derive solution of the primal problem

Weak and strong duality

Weak duality: $d^* \leq p^*$

- ▶ It holds by construction
- ▶ Non-trivial lower bounds for NP-hard problems

Strong duality: $d^* = p^*$

- ▶ It **does not** hold in general case
- ▶ *Typically* it holds for convex problems
- ▶ But regularity conditions are important
- ▶ It may hold for **non-convex** problems
- ▶ A solution of the dual problem can help to derive solution of the primal problem

Duality gap: $f_0(\mathbf{x}) - g(\boldsymbol{\lambda}, \boldsymbol{\mu})$

Weak and strong duality

Weak duality: $d^* \leq p^*$

- ▶ It holds by construction
- ▶ Non-trivial lower bounds for NP-hard problems

Strong duality: $d^* = p^*$

- ▶ It **does not** hold in general case
- ▶ *Typically* it holds for convex problems
- ▶ But regularity conditions are important
- ▶ It may hold for **non-convex** problems
- ▶ A solution of the dual problem can help to derive solution of the primal problem

Duality gap: $f_0(\mathbf{x}) - g(\boldsymbol{\lambda}, \boldsymbol{\mu})$

- ▶ How current approximations $(\mathbf{x}_k, \boldsymbol{\lambda}_k, \boldsymbol{\mu}_k)$ are accurate

Weak and strong duality

Weak duality: $d^* \leq p^*$

- ▶ It holds by construction
- ▶ Non-trivial lower bounds for NP-hard problems

Strong duality: $d^* = p^*$


- ▶ It **does not** hold in general case
- ▶ *Typically* it holds for convex problems
- ▶ But regularity conditions are important
- ▶ It may hold for **non-convex** problems
- ▶ A solution of the dual problem can help to derive solution of the primal problem

Duality gap: $f_0(\mathbf{x}) - g(\boldsymbol{\lambda}, \boldsymbol{\mu})$

- ▶ How current approximations $(\mathbf{x}_k, \boldsymbol{\lambda}_k, \boldsymbol{\mu}_k)$ are accurate
- ▶ Proof correctness and convergence of numerical methods

Geometric interpretation

$$\begin{array}{ll} \min_{\mathbf{x} \in \mathcal{D}} f_0(\mathbf{x}) & g(\lambda) = \inf_{(u,t) \in \mathcal{G}} (t + \lambda u) \\ \text{s.t. } f_1(\mathbf{x}) \leq 0 & \mathcal{G} = \{(f_1(\mathbf{x}), f_0(\mathbf{x})) \mid \mathbf{x} \in \mathcal{D}\} \end{array}$$



Slater condition and strong duality

Slater condition: reminder

The Slater condition holds if $\exists \bar{\mathbf{x}} \in \text{int } \mathcal{D} : f_i(\bar{\mathbf{x}}) < 0, \mathbf{A}\bar{\mathbf{x}} = \mathbf{b}$

Theorem

Strong duality holds for the convex optimization problem

$$\begin{aligned} & \min f_0(\mathbf{x}) \\ & \text{s.t. } f_i(\mathbf{x}) \leq 0, \quad i = 1, \dots, m \\ & \quad \mathbf{A}\mathbf{x} = \mathbf{b}, \quad \mathbf{A} \in \mathbb{R}^{p \times n} \end{aligned}$$

if the Slater condition holds.

Complementary slackness

Assume that strong duality holds, \mathbf{x}^* is a primal solution, $(\boldsymbol{\lambda}^*, \boldsymbol{\mu}^*)$ is a dual solution, then

Complementary slackness

Assume that strong duality holds, \mathbf{x}^* is a primal solution, $(\boldsymbol{\lambda}^*, \boldsymbol{\mu}^*)$ is a dual solution, then

$$\begin{aligned} f_0(\mathbf{x}^*) = g(\boldsymbol{\lambda}^*, \boldsymbol{\mu}^*) &= \inf_{\mathbf{x} \in \mathcal{D}} \left(f_0(\mathbf{x}) + \sum_{i=1}^m \lambda_i^* g_i(\mathbf{x}) + \sum_{j=1}^p \mu_j^* h_j(\mathbf{x}) \right) \\ &\leq f_0(\mathbf{x}^*) + \sum_{i=1}^m \lambda_i^* g_i(\mathbf{x}^*) + \sum_{j=1}^p \mu_j^* h_j(\mathbf{x}^*) \\ &\leq f_0(\mathbf{x}^*) \end{aligned}$$

Complementary slackness

Assume that strong duality holds, \mathbf{x}^* is a primal solution, $(\boldsymbol{\lambda}^*, \boldsymbol{\mu}^*)$ is a dual solution, then

$$\begin{aligned} f_0(\mathbf{x}^*) = g(\boldsymbol{\lambda}^*, \boldsymbol{\mu}^*) &= \inf_{\mathbf{x} \in \mathcal{D}} \left(f_0(\mathbf{x}) + \sum_{i=1}^m \lambda_i^* g_i(\mathbf{x}) + \sum_{j=1}^p \mu_j^* h_j(\mathbf{x}) \right) \\ &\leq f_0(\mathbf{x}^*) + \sum_{i=1}^m \lambda_i^* g_i(\mathbf{x}^*) + \sum_{j=1}^p \mu_j^* h_j(\mathbf{x}^*) \\ &\leq f_0(\mathbf{x}^*) \end{aligned}$$

- \mathbf{x}^* is a minimizer of $L(\mathbf{x}, \boldsymbol{\lambda}^*, \boldsymbol{\mu}^*)$

Complementary slackness

Assume that strong duality holds, \mathbf{x}^* is a primal solution, $(\boldsymbol{\lambda}^*, \boldsymbol{\mu}^*)$ is a dual solution, then

$$\begin{aligned} f_0(\mathbf{x}^*) = g(\boldsymbol{\lambda}^*, \boldsymbol{\mu}^*) &= \inf_{\mathbf{x} \in \mathcal{D}} \left(f_0(\mathbf{x}) + \sum_{i=1}^m \lambda_i^* g_i(\mathbf{x}) + \sum_{j=1}^p \mu_j^* h_j(\mathbf{x}) \right) \\ &\leq f_0(\mathbf{x}^*) + \sum_{i=1}^m \lambda_i^* g_i(\mathbf{x}^*) + \sum_{j=1}^p \mu_j^* h_j(\mathbf{x}^*) \\ &\leq f_0(\mathbf{x}^*) \end{aligned}$$

- ▶ \mathbf{x}^* is a minimizer of $L(\mathbf{x}, \boldsymbol{\lambda}^*, \boldsymbol{\mu}^*)$
- ▶ Complementary slackness conditions

$$\mu_j^* h_j(\mathbf{x}^*) = 0, \quad j = 1, \dots, p$$

$$\mu_j^* > 0 \Rightarrow h_j(\mathbf{x}^*) = 0, \quad h_j(\mathbf{x}^*) < 0 \Rightarrow \mu_j^* = 0$$

Karush-Kuhn-Tucker conditions: convex and non-convex problems

If $(\mathbf{x}^*, \boldsymbol{\lambda}^*, \boldsymbol{\mu}^*)$ are solutions of primal and dual problems and the strong duality holds, then

Karush-Kuhn-Tucker conditions: convex and non-convex problems

If $(\mathbf{x}^*, \boldsymbol{\lambda}^*, \boldsymbol{\mu}^*)$ are solutions of primal and dual problems and the strong duality holds, then

1. $h_j(\mathbf{x}^*) \leq 0$

Karush-Kuhn-Tucker conditions: convex and non-convex problems

If $(\mathbf{x}^*, \boldsymbol{\lambda}^*, \boldsymbol{\mu}^*)$ are solutions of primal and dual problems and the strong duality holds, then

1. $h_j(\mathbf{x}^*) \leq 0$
2. $g_i(\mathbf{x}^*) = 0$

Karush-Kuhn-Tucker conditions: convex and non-convex problems

If $(\mathbf{x}^*, \boldsymbol{\lambda}^*, \boldsymbol{\mu}^*)$ are solutions of primal and dual problems and the strong duality holds, then

1. $h_j(\mathbf{x}^*) \leq 0$
2. $g_i(\mathbf{x}^*) = 0$
3. $\boldsymbol{\mu}^* \geq 0$

Karush-Kuhn-Tucker conditions: convex and non-convex problems

If $(\mathbf{x}^*, \boldsymbol{\lambda}^*, \boldsymbol{\mu}^*)$ are solutions of primal and dual problems and the strong duality holds, then

1. $h_j(\mathbf{x}^*) \leq 0$
2. $g_i(\mathbf{x}^*) = 0$
3. $\boldsymbol{\mu}^* \geq 0$
4. $\mu_j^* h_j(\mathbf{x}^*) = 0$

Karush-Kuhn-Tucker conditions: convex and non-convex problems

If $(\mathbf{x}^*, \boldsymbol{\lambda}^*, \boldsymbol{\mu}^*)$ are solutions of primal and dual problems and the strong duality holds, then

1. $h_j(\mathbf{x}^*) \leq 0$

2. $g_i(\mathbf{x}^*) = 0$

3. $\boldsymbol{\mu}^* \geq 0$

4. $\mu_j^* h_j(\mathbf{x}^*) = 0$

5. $f'_0(\mathbf{x}^*) + \sum_{i=1}^m \lambda_i^* g'_i(\mathbf{x}^*) + \sum_{j=1}^p \mu_j^* h'_j(\mathbf{x}^*) = 0$

Karush-Kuhn-Tucker conditions: convex and non-convex problems

If $(\mathbf{x}^*, \boldsymbol{\lambda}^*, \boldsymbol{\mu}^*)$ are solutions of primal and dual problems and the strong duality holds, then

1. $h_j(\mathbf{x}^*) \leq 0$
2. $g_i(\mathbf{x}^*) = 0$
3. $\boldsymbol{\mu}^* \geq 0$
4. $\mu_j^* h_j(\mathbf{x}^*) = 0$
5. $f'_0(\mathbf{x}^*) + \sum_{i=1}^m \lambda_i^* g'_i(\mathbf{x}^*) + \sum_{j=1}^p \mu_j^* h'_j(\mathbf{x}^*) = 0$

The last equality holds since

$$\min_{\mathbf{x}} L(\mathbf{x}, \boldsymbol{\lambda}^*, \boldsymbol{\mu}^*) = L(\mathbf{x}^*, \boldsymbol{\lambda}^*, \boldsymbol{\mu}^*)$$

and first-order necessary condition.

KKT conditions for convex problems

Claim 1

Assume that the primal problem is convex (f_0, h_j are convex, g_i are affine) and for $(\hat{\mathbf{x}}, \hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}})$ the KKT conditions hold, then

- ▶ the strong duality holds
- ▶ $(\hat{\mathbf{x}}, \hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}})$ are primal and dual solutions

KKT conditions for convex problems

Claim 1

Assume that the primal problem is convex (f_0, h_j are convex, g_i are affine) and for $(\hat{\mathbf{x}}, \hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}})$ the KKT conditions hold, then

- ▶ the strong duality holds
- ▶ $(\hat{\mathbf{x}}, \hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}})$ are primal and dual solutions

Proof

KKT conditions for convex problems

Claim 1

Assume that the primal problem is convex (f_0, h_j are convex, g_i are affine) and for $(\hat{\mathbf{x}}, \hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}})$ the KKT conditions hold, then

- ▶ the strong duality holds
- ▶ $(\hat{\mathbf{x}}, \hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}})$ are primal and dual solutions

Proof

- ▶ 1 and 2 KKT conditions $\rightarrow \hat{\mathbf{x}}$ belongs to a feasible set, i.e.
 $g_i(\hat{\mathbf{x}}) = 0$ and $h_j(\hat{\mathbf{x}}) \leq 0$

KKT conditions for convex problems

Claim 1

Assume that the primal problem is convex (f_0, h_j are convex, g_i are affine) and for $(\hat{\mathbf{x}}, \hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}})$ the KKT conditions hold, then

- ▶ the strong duality holds
- ▶ $(\hat{\mathbf{x}}, \hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}})$ are primal and dual solutions

Proof

- ▶ 1 and 2 KKT conditions $\rightarrow \hat{\mathbf{x}}$ belongs to a feasible set, i.e.
 $g_i(\hat{\mathbf{x}}) = 0$ and $h_j(\hat{\mathbf{x}}) \leq 0$
- ▶ $\hat{\boldsymbol{\mu}} \geq 0 \rightarrow L(\mathbf{x}, \hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}})$ is convex of \mathbf{x}

KKT conditions for convex problems

Claim 1

Assume that the primal problem is convex (f_0, h_j are convex, g_i are affine) and for $(\hat{\mathbf{x}}, \hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}})$ the KKT conditions hold, then

- ▶ the strong duality holds
- ▶ $(\hat{\mathbf{x}}, \hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}})$ are primal and dual solutions

Proof

- ▶ 1 and 2 KKT conditions $\rightarrow \hat{\mathbf{x}}$ belongs to a feasible set, i.e.
 $g_i(\hat{\mathbf{x}}) = 0$ and $h_j(\hat{\mathbf{x}}) \leq 0$
- ▶ $\hat{\boldsymbol{\mu}} \geq 0 \rightarrow L(\mathbf{x}, \hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}})$ is convex of \mathbf{x}
- ▶ The last condition $\rightarrow \hat{\mathbf{x}}$ is a minimizer of L :
$$g(\hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}}) = L(\hat{\mathbf{x}}, \hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}}) = f_0(\hat{\mathbf{x}}) + \sum_{i=1}^m \hat{\lambda}_i g_i(\hat{\mathbf{x}}) + \sum_{j=1}^p \hat{\mu}_j h_j(\hat{\mathbf{x}})$$

KKT conditions for convex problems

Claim 1

Assume that the primal problem is convex (f_0, h_j are convex, g_i are affine) and for $(\hat{\mathbf{x}}, \hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}})$ the KKT conditions hold, then

- ▶ the strong duality holds
- ▶ $(\hat{\mathbf{x}}, \hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}})$ are primal and dual solutions

Proof

- ▶ 1 and 2 KKT conditions $\rightarrow \hat{\mathbf{x}}$ belongs to a feasible set, i.e.
 $g_i(\hat{\mathbf{x}}) = 0$ and $h_j(\hat{\mathbf{x}}) \leq 0$
- ▶ $\hat{\boldsymbol{\mu}} \geq 0 \rightarrow L(\mathbf{x}, \hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}})$ is convex of \mathbf{x}
- ▶ The last condition $\rightarrow \hat{\mathbf{x}}$ is a minimizer of L :
 $g(\hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}}) = L(\hat{\mathbf{x}}, \hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}}) = f_0(\hat{\mathbf{x}}) + \sum_{i=1}^m \hat{\lambda}_i g_i(\hat{\mathbf{x}}) + \sum_{j=1}^p \hat{\mu}_j h_j(\hat{\mathbf{x}})$
- ▶ From the complementary slackness conditions $\hat{\mu}_j h_j(\hat{\mathbf{x}}) = 0$ follow that $g(\hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}}) = f_0(\hat{\mathbf{x}})$

KKT conditions for convex problems

Claim 1

Assume that the primal problem is convex (f_0, h_j are convex, g_i are affine) and for $(\hat{\mathbf{x}}, \hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}})$ the KKT conditions hold, then

- ▶ the strong duality holds
- ▶ $(\hat{\mathbf{x}}, \hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}})$ are primal and dual solutions

Proof

- ▶ 1 and 2 KKT conditions $\rightarrow \hat{\mathbf{x}}$ belongs to a feasible set, i.e.
 $g_i(\hat{\mathbf{x}}) = 0$ and $h_j(\hat{\mathbf{x}}) \leq 0$
- ▶ $\hat{\boldsymbol{\mu}} \geq 0 \rightarrow L(\mathbf{x}, \hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}})$ is convex of \mathbf{x}
- ▶ The last condition $\rightarrow \hat{\mathbf{x}}$ is a minimizer of L :
 $g(\hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}}) = L(\hat{\mathbf{x}}, \hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}}) = f_0(\hat{\mathbf{x}}) + \sum_{i=1}^m \hat{\lambda}_i g_i(\hat{\mathbf{x}}) + \sum_{j=1}^p \hat{\mu}_j h_j(\hat{\mathbf{x}})$
- ▶ From the complementary slackness conditions $\hat{\mu}_j h_j(\hat{\mathbf{x}}) = 0$
follow that $g(\hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}}) = f_0(\hat{\mathbf{x}})$
- ▶ Thus, the strong duality holds

KKT conditions for convex problems

Claim 1

Assume that the primal problem is convex (f_0, h_j are convex, g_i are affine) and for $(\hat{\mathbf{x}}, \hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}})$ the KKT conditions hold, then

- ▶ the strong duality holds
- ▶ $(\hat{\mathbf{x}}, \hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}})$ are primal and dual solutions

Proof

- ▶ 1 and 2 KKT conditions $\rightarrow \hat{\mathbf{x}}$ belongs to a feasible set, i.e.
 $g_i(\hat{\mathbf{x}}) = 0$ and $h_j(\hat{\mathbf{x}}) \leq 0$
- ▶ $\hat{\boldsymbol{\mu}} \geq 0 \rightarrow L(\mathbf{x}, \hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}})$ is convex of \mathbf{x}
- ▶ The last condition $\rightarrow \hat{\mathbf{x}}$ is a minimizer of L :
 $g(\hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}}) = L(\hat{\mathbf{x}}, \hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}}) = f_0(\hat{\mathbf{x}}) + \sum_{i=1}^m \hat{\lambda}_i g_i(\hat{\mathbf{x}}) + \sum_{j=1}^p \hat{\mu}_j h_j(\hat{\mathbf{x}})$
- ▶ From the complementary slackness conditions $\hat{\mu}_j h_j(\hat{\mathbf{x}}) = 0$ follow that $g(\hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}}) = f_0(\hat{\mathbf{x}})$
- ▶ Thus, the strong duality holds
- ▶ $(\hat{\mathbf{x}}, \hat{\boldsymbol{\lambda}}, \hat{\boldsymbol{\mu}})$ are primal and dual solutions

KKT conditions for convex problems

Claim 2

Assume that the Slater condition holds for convex optimization problem. Then \mathbf{x} is a primal solution iff there exist $(\boldsymbol{\lambda}, \boldsymbol{\mu})$ such that the KKT conditions hold for this triplet.

KKT conditions for convex problems

Claim 2

Assume that the Slater condition holds for convex optimization problem. Then \mathbf{x} is a primal solution iff there exist $(\boldsymbol{\lambda}, \boldsymbol{\mu})$ such that the KKT conditions hold for this triplet.

Proof

KKT conditions for convex problems

Claim 2

Assume that the Slater condition holds for convex optimization problem. Then \mathbf{x} is a primal solution iff there exist $(\boldsymbol{\lambda}, \boldsymbol{\mu})$ such that the KKT conditions hold for this triplet.

Proof

- From the convexity and Slater condition follow the strong duality

KKT conditions for convex problems

Claim 2

Assume that the Slater condition holds for convex optimization problem. Then \mathbf{x} is a primal solution iff there exist $(\boldsymbol{\lambda}, \boldsymbol{\mu})$ such that the KKT conditions hold for this triplet.

Proof

- ▶ From the convexity and Slater condition follow the strong duality
- ▶ Necessity of the KKT follows from the general fact.

KKT conditions for convex problems

Claim 2

Assume that the Slater condition holds for convex optimization problem. Then \mathbf{x} is a primal solution iff there exist $(\boldsymbol{\lambda}, \boldsymbol{\mu})$ such that the KKT conditions hold for this triplet.

Proof

- ▶ From the convexity and Slater condition follow the strong duality
- ▶ Necessity of the KKT follows from the general fact.
- ▶ Sufficiency follows from the claim 1

Summary on duality theory

- ▶ Dual function

Summary on duality theory

- ▶ Dual function
- ▶ Dual problem

Summary on duality theory

- ▶ Dual function
- ▶ Dual problem
- ▶ Lower bounds

Summary on duality theory

- ▶ Dual function
- ▶ Dual problem
- ▶ Lower bounds
- ▶ KKT and strong duality

Summary on duality theory

- ▶ Dual function
- ▶ Dual problem
- ▶ Lower bounds
- ▶ KKT and strong duality
- ▶ KKT for convex optimization problem with Slater regularity